Daniel Kastler

Centre de Physique Théorique CNRS - Luminy, Case 907 F-13288 Marseille Cedex 09, France and Université de la Méditerrannée

Dedicated to Sergio Doplicher and John Roberts

Abstract. These notes hopefully provide an aid to the comprehension of the Connes-Moscovici and Connes-Kreimer works, by isolating common mathematical features of the Connes-Moscovici, rooted trees, and Feynman-graph Hopf algebras (as a new special branch of the theory of Hopf algebras expected to become important). We discuss in particular the dual Milnor-Moore situation.

1 Characters and infinitesimal characters of Hopf algebras

In what follows **k** is a commutative field of zero characteristic (e.g. $\mathbf{k} = \mathbb{C}, \mathbb{R}$).

Reminder 1.1 (the algebraic dual of Hopf algebras). Let \mathbf{H} $(m, \mathbf{1} = e_1, \Delta, \varepsilon, S)$ be a Hopf algebra over \mathbf{k} . Consider the algebraic dual $\mathbf{H}^* = (m_*, \mathbf{1}_* = e_*1, \Delta_* = m^t, \varepsilon_* = e^t, S_* = S^t)$ of \mathbf{H} endowed with the topology of simple convergence on \mathbf{H} . The value of $\xi \in \mathbf{H}^*$ for $h \in \mathbf{H}$ is denoted by $\langle \xi, h \rangle$. Then:

 (i) As the dual of the coalgebra H, H* is a genuine algebra with the following product * and unit 1*:

$$\begin{cases} <\xi *\eta, h>=<\xi\otimes\eta, \Delta h>\\ <\mathbf{1}_*, h>=\varepsilon(h) \end{cases}, \quad \xi,\eta\in\mathbf{H}^*, \ h\in\mathbf{H}.$$
(1.1)

(ii) \mathbf{H}^* is however not a Hopf algebra: we do have $S_* = S^t : \mathbf{H}^* \to \mathbf{H}^*$ and $\varepsilon_* = e^t : \mathbf{H}^* \to \mathbb{C}$:

$$\begin{cases} <\xi, Sh > = < S_*\xi, h > \\ <\xi, \mathbf{1} > = \varepsilon_*(\xi) \end{cases}, \quad \xi \in \mathbf{H}^*, \ h \in \mathbf{H},$$
(1.2)

the $\sigma(\mathbf{H}^*, \mathbf{H})$ -topology.

We note this product with a * because it is the convolution product of $\operatorname{End}_{\mathbb{C}}(\mathbf{H},\mathbb{C})$, cf. definition (2,1) in [**3**], namely, denoting the covalue $\langle \xi, h \rangle$ by $\xi(h)$: $(\xi * \eta)(h) = [m_k(\xi \otimes \eta)\Delta](h)$, where we can omit m_k owing to $\mathbf{k} \otimes \mathbf{k} \cong \mathbf{k}$.

©2001 American Mathematical Society

but in general $\Delta_* = m^t : \mathbf{H}^* \to (\mathbf{H} \otimes \mathbf{H})^* \underset{\neq}{\supset} \mathbf{H}^* \otimes \mathbf{H}^*.$

We now investigate two subsets of \mathbf{H}^* which Δ_* maps into $\mathbf{H}^* \otimes \mathbf{H}^*$: The characters and infinitesimal characters of \mathbf{H} :

Definition 1.2 Let $\mathbf{H}(m, e, \Delta, \varepsilon)$ be a Hopf k-algebra.

(i) a **character of H** is a non vanishing **k**-linear map $\chi : \mathbf{H} \to \mathbf{k}$ which is multiplicative:

$$\langle \chi, hk \rangle = \langle \chi, h \rangle \langle \chi, k \rangle, \qquad h, k \in \mathbf{H}$$
 (1.3)

(note that, as a consequence, one has

$$\langle \chi, \mathbf{1} \rangle = 1, \tag{1.4}$$

indeed (1.3) entails that $\langle \chi, \mathbf{1} \rangle [\langle \chi, \mathbf{1} \rangle -1]$ vanishes, whence (1.4) if χ does not vanish). We denote by Char**H** the set of characters of **H**. Note that the counit is a character of **H**.

(ii) An infinitesimal character of **H** is a **k**-linear map $Z : \mathbf{H} \to \mathbf{k}$ fulfilling:

$$\langle Z, hk \rangle = \langle Z, h \rangle \varepsilon(k) + \varepsilon(h) \langle Z, k \rangle, \qquad h, k \in \mathbf{H}$$
 (1.5)

(note that, as a consequence, one has

$$< Z, 1 >= 0,$$
 (1.6)

indeed one has

$$<\!Z, \mathbf{1}> = <\!Z, \mathbf{1} \cdot \mathbf{1}>) = <\!Z, \mathbf{1} > \varepsilon(\mathbf{1}) + \varepsilon(\mathbf{1}) < Z, \mathbf{1}> = 2 < Z, \mathbf{1}>;$$

and that Z vanishes on the square \mathbf{H}^{+2} of the augmentation ideal $\mathbf{H}^{+} = \text{Ker}\varepsilon$). Note also that (1.5) is the infinitesimal form of (1.3) written for $\chi = \varepsilon + Z$ with Z considered as small).

We denote by $\partial \text{Char}\mathbf{H}$ the set of infinitesimal characters of \mathbf{H} .

Proposition 1.3 With the definitions in 1.2 we have that:

(i) Equipped with the convolution product (1.1), the unit $\mathbf{1}_*$:

$$\begin{cases} <\psi *\psi', h>=<\psi \otimes \psi', \Delta h> \\ <\mathbf{1}_*, h>=\varepsilon(h) \end{cases}, \quad \psi, \psi' \in Char\mathbf{H}, \ h \in \mathbf{H}.$$
(1.7)

the inverse:

$$\chi^{-1} = \chi \circ S, \qquad x \in Char\mathbf{H},\tag{1.8}$$

and the $\sigma(\mathbf{H}^*, \mathbf{H})$ -topology, $\mathbf{G} = Char\mathbf{H}$ is a topological group, the group of characters of \mathbf{H} .

(ia) **G** is dually definable as the group of "group-like elements" of \mathbf{H}^* :

$$\mathbf{G} = \{ \psi \in \mathbf{H}^*, \psi \neq 0, \Delta_* \psi = \psi \otimes \psi \}$$
(1.9)

^(1.5) defines Z as a derivation of the **H**-bimodule $_{\varepsilon}\mathbf{H}_{\varepsilon}$ consisting of **H** with multiplication by $a \in \mathbf{H}$ from the left and from the right multiplication by $\varepsilon(a)$: $\begin{cases} h \cdot \alpha = \varepsilon(h)\alpha \\ \alpha \cdot h = \varepsilon(h)\alpha \end{cases}, \alpha \in \mathbb{C}, h \in \mathbf{H}. \end{cases}$

in fact a subgroup of the group of invertibles of the algebra \mathbf{H}^* with product *.

[&]quot;group-like elements" between quotation marks because \mathbf{H}^* is not a Hopf algebra cf. (ii).

(ii) Equipped with the bracket:

$$[Z, Z'] = Z * Z' - Z' * Z, \qquad Z, Z' \in \partial Char \mathbf{H},$$
(1.10)

where * is the convolution product (1.1):

$$Z * Z' = \langle Z \otimes Z, \Delta h \rangle, \qquad Z, Z' \in \partial Char \mathbf{H}, \ h \in \mathbf{H},$$
(1.11)

 $\mathbf{L} = \partial Char \mathbf{H}$ is a Lie algebra, the Lie algebra of infinitesimal characters of \mathbf{H} .

(iia) **L** is dually definable as the Lie algebra of "primitive elements" of \mathbf{H}^* :

$$\mathbf{L} = \{ Z \in \mathbf{H}^*, \Delta_* Z = Z \otimes \mathbf{1}_* + \mathbf{1}_* \otimes Z \}.$$
(1.12)

Proof The product first line (1.7) is known to be associative. To assert that $\chi * \psi \in \text{Char}\mathbf{H}$ we need to check multiplicativity: now, for $h, k \in \mathbf{H}$:

$$\begin{array}{lll} <\chi\ast\psi,hk> &=& <\chi\otimes\psi,\Delta(hk)>=<\chi\otimes\psi,(\Delta h)(\Delta k)>\\ &=& <\chi\otimes\psi,\Delta h)><\chi\otimes\psi,\Delta k>=<\chi\ast\psi,h><\chi\ast\psi,k> \end{array}$$

where we used the multiplicativity of $\chi \otimes \psi$ due to the commutativity of **k**: indeed, one has, for x, y, x', y':

$$\begin{aligned} <\chi\otimes\psi,(x\otimes y)(x'\otimes y')> &= <\chi\otimes\psi,xx'\otimes yy'> =<\chi,xx'><\psi,yy'> \\ &= <\chi,x><\chi,x'><\psi,y><\psi,y'> \\ &= <\chi,x><\psi,y><\chi,x'><\psi,y'> \\ &= <\chi\otimes\psi,x\otimes\psi,x\otimes y><\chi\otimes\psi,(x'\otimes y')> \end{aligned}$$

Check that $\chi * \mathbf{1}_* = \mathbf{1}_* * \chi = \chi$: we have by the first line (1.7):

$$\begin{aligned} <\chi*\mathbf{1}_{*},h> &= <\chi\otimes\mathbf{1}_{*},\Delta h>=<\chi\otimes\mathbf{1}_{*},h_{(1)}\otimes_{(2)}> \\ &= <\chi,h_{(1)}><\varepsilon(h_{(2)})>=<\chi,\varepsilon(h_{(2)})h_{(1)}> \\ &= <\chi,h>. \\ <\mathbf{1}_{*}*\chi,h> &= <\mathbf{1}_{*}\otimes\chi,\Delta h>=<\mathbf{1}_{*}\otimes\chi,h_{(1)}\otimes_{(2)}> \\ &= <\mathbf{1}_{*},h_{(1)}><\chi,h_{(2)}> \\ &= <\varepsilon(h_{(1)})><\chi,h_{(2)}>=<\chi,\varepsilon(h_{(1)})h_{(2)}>=<\chi,h>. \end{aligned}$$

Check of (1.8) we have:

$$\begin{aligned} [(\chi \circ S) * \chi](h) &= [(\chi \circ S) \otimes \chi](\Delta h) = \chi(Sh_{(1)})\chi(h_{(2)}) = \chi(Sh_{(1)}a_{(2)}) \\ &= \chi(\varepsilon(h)\mathbf{1}) = \varepsilon(h)\chi(\mathbf{1}) = \varepsilon(h) \end{aligned}$$

$$\begin{aligned} [\chi * (\chi \circ S)](h) &= [\chi \otimes (\chi \circ S)](\Delta h) = \chi(h_{(1)})\chi(Sh_{(2)}) = \chi(h_{(1)})Sh_{(2)}) \\ &= \chi(\varepsilon(h)\mathbf{1}) = \varepsilon(h)\chi(\mathbf{1}) = \varepsilon(h). \end{aligned}$$

The facts that product and inverse are continuous in the $\sigma(\mathbf{H}^*, \mathbf{H})$ -topology is clear.

(ia) For the proof we write $\mathbf{G} = \operatorname{Char} \mathbf{H}$ and $\mathbf{G}' = \{\psi \in \mathbf{H}^*, \Delta_* \psi = \psi \otimes \psi\}$. Check of $\mathbf{G}' \subset \mathbf{G}$: for $\psi \in \mathbf{G}', h, h' \in \mathbf{H}$, we have:

$$\langle \psi, h \rangle \langle \psi, h' \rangle = \langle \psi \otimes \psi, h \otimes h' \rangle = \langle \Delta_* \psi, h \otimes h' \rangle$$
$$= \langle \psi, m(h \otimes h') \rangle = \langle \psi, hh' \rangle$$

in fact a Lie-subalgebra of the Lie algebra ${\rm Lie} {\bf H}^*$ associated with the algebra ${\bf H}^*.$

[&]quot;primitive elements" between quotation marks because \mathbf{H}^* is not a Hopf algebra cf. (ii).

Check of $\mathbf{G} \subset \mathbf{G}'$: $\psi \in \mathbf{G}$ implies:

$$\langle \psi, m(h \otimes h') \rangle = \langle \psi, hh' \rangle = \langle \psi, h \rangle \langle \psi, h' \rangle = \langle \psi \otimes \psi, h \otimes h' \rangle$$

hence we have $\langle \psi, m(h \otimes h') \rangle = \langle \Delta_* \psi, h \otimes h' \rangle$ with $\Delta_* \psi = \psi \otimes \psi$.

The Lie algebra property of \mathbf{L} can be checked directly from (1.5), but will immediately result from (iia), which we now check.

(iia) For the proof we write $\mathbf{L} = \partial \operatorname{Char} \mathbf{H}$ and $\mathbf{L}' = \{Z \in \mathbf{H}^*, \Delta_* Z = Z \otimes \mathbf{1}_* + \mathbf{1}_* \otimes Z\}.$

Check of $\mathbf{L}' \subset \mathbf{L}$: $Z \in \mathbf{L}'$ implies for $h, h' \in \mathbf{H}$:

$$\begin{aligned} < Z, h > < \mathbf{1}_*, h' > + < \mathbf{1}_*, h > < Z, h' > \\ = < Z \otimes \mathbf{1}_* + \mathbf{1}_* \otimes Z, h \otimes h' > \\ = < \Delta_* Z, h \otimes h' > = < Z, hh' > = < Z, m(h \otimes h') > = < Z, hh' > . \end{aligned}$$

Check of $\mathbf{L} \subset \mathbf{L}'$: for $Z \in \mathbf{L}$, $h, h' \in \mathbf{H}$ one has,

$$< Z, m(h \otimes h') > = < Z, hh' > = < Z, h > < \mathbf{1}_*, h' > + < \mathbf{1}_*, h > < Z, h' > = < Z \otimes \mathbf{1}_* + \mathbf{1}_* \otimes Z, h \otimes h' > .$$

hence we have $\langle Z, m(h \otimes h') \rangle = \langle \Delta_* Z, h \otimes h' \rangle$ with $\Delta_* \psi = Z \otimes \mathbf{1}_* + \mathbf{1}_* \otimes Z$. Check of (ii): it is clear that \mathbf{L}' is a Lie-subalgebra of the Lie algebra Lie \mathbf{H}^* . \Box

Definition-Lemma 1.4 Let H_{\ast} be the subalgebra of H^{\ast} generated by 1_{\ast} and L.

- (o) One has $\varepsilon_*(\xi * \eta) = \varepsilon_*(\xi)\varepsilon_*(\eta)$, $\xi, \eta \in \mathbf{H}^* \supset \mathbf{H}_*$, $\varepsilon_*(\mathbf{1}_*) = 1$ and ε vanishes on **L**.
- (i) One has $S_*\mathbf{H}_* \subset \mathbf{H}_*$, $S_*\mathbf{L} \subset \mathbf{L}$, and

$$\begin{cases} S_*(\xi * \eta) = (S_*\eta) * (S_*\xi), \xi, \ \eta \in \mathbf{H}^* \supset \mathbf{H}_* \\ S_*\mathbf{1}_* = \mathbf{1}_*. \end{cases}$$
(1.13)

(ii) One has $\Delta_* \mathbf{H}_* \subset \mathbf{H}_* \otimes \mathbf{H}_*$ and Δ_* is multiplicative:

$$\Delta_* m_* = (m_* \otimes m_*) P_{12}(\Delta_* \otimes \Delta_*), \tag{1.14}$$

and cocommutative in the sense:

$$\Delta_* = P_{12}\Delta_*,\tag{1.15}$$

Proof (o) One has $\varepsilon_*(\mathbf{1}_*) = \langle \mathbf{1}_*, \mathbf{1} \rangle = 1$, for $Z \in \mathbf{L}$, $\varepsilon_*(Z) = \langle Z, \mathbf{1} \rangle = 0$. For $\xi \eta \in \mathbf{H}^*$ one has $\varepsilon_*(\xi * \eta) = \langle \xi * \eta, \mathbf{1} \rangle = \langle \xi \otimes \eta, \Delta \mathbf{1} \rangle = \langle \xi, \mathbf{1} \rangle \langle \eta, \mathbf{1} \rangle = \varepsilon_*(\xi)\varepsilon_*(\eta)$.

(i) We first notice the property

$$\Delta_* S_* = (S_* \otimes S_*) P_{12} \Delta_*, \tag{1.16}$$

transposed of the known property $Sm = mP_{12}(S \otimes S)$. Check of $S_* \mathbf{L} \subset \mathbf{L}$: for $Z \in \mathbf{L}$ one has $S_* Z \in \mathbf{L}$, indeed one has:

$$(\Delta_*S_*)Z = (S_* \otimes S_*)P_{12}\Delta_*Z$$

= $(S_* \otimes S_*)P_{12}(Z \otimes \mathbf{1} + \mathbf{1} \otimes Z)$
= $S_*Z \otimes \mathbf{1} + \mathbf{1} \otimes S_*Z.$

It follows that $S_*\mathbf{H}_* \subset \mathbf{H}_*$. Now S_* turns the unit $\mathbf{1}_*$ into $\mathbf{1}_*$. And, for $\xi, \eta \in \mathbf{H}^*$ and $h \in \mathbf{H}$, one has using the known property $(H\Delta)$: $\Delta S = (S \otimes S)P_{12}\Delta$:

$$\langle S_*(\xi * \eta), h \rangle = \langle \xi * \eta, Sh \rangle = \langle \xi \otimes \eta, \Delta Sh \rangle$$

$$= \langle \xi \otimes \eta, (S \otimes S)P_{12}\Delta h \rangle$$

$$= \langle S_*\eta \otimes S_*\xi, \Delta h \rangle = \langle (S_*\eta) * (S_*\xi), h \rangle .$$

(ii) (1.14) reads $m^t \Delta^t = (\Delta^t \otimes \Delta^t) P_{12}(m^t \otimes m^t)$, transposed of the axiom $\Delta m = (m \otimes m) P_{12}(\Delta \otimes \Delta)$.

Check of the cocommutativity property (1.15): holds obviously on the generators **1** and $Z \in \mathbf{L}$. Now (1.15) propagates multiplicatively: if $\xi, \eta \in \mathbf{L}$ fulfill $\Delta \xi = P_{12}\Delta\xi$, $\Delta \eta = P_{12}\Delta\eta$, $\Delta(\xi\eta) = (\Delta\xi)(\Delta\eta) = (P_{12}\Delta\xi)(P_{12}\Delta\eta) = P_{12}(\Delta\xi)(\Delta\eta) = P_{12}\Delta(\xi\eta)$, owing to multiplicatively of $P_{12}: P_{12}(a \otimes a')P_{12}(b \otimes b') = (a' \otimes a)(b' \otimes b) = a'a \otimes b'b = P_{12}; (a \otimes a')(b \otimes b')].$

Proposition 1.5 $\mathbf{H}_*(m_*, \mathbf{1}_*, \Delta_*, \varepsilon_*, S_*)$ is a cocommutative Hopf algebra. The Hopf algebras \mathbf{H} and \mathbf{H}_* are set in duality through the bilinear form $\langle \cdot, \cdot \rangle$.

Proof H_{*} is a unital algebra with unit $\mathbf{1}_*$ by definition. Using the terminology of Appendix A, we have that the coalgebra axioms $(C\Delta)$, $(C\varepsilon)$ for \mathbf{H}_* result by transposition from the algebra axioms (Am), resp. (Ae) for \mathbf{H} . Bialgebra axioms for \mathbf{H}_* : we already met (Bm) in (1.14), (Be) is the known fact that $\mathbf{1}_* \in \mathbf{G}$, we have met $(B\varepsilon)$ and $(Be\varepsilon)$ in Definition-Lemma 1.4(o) above. Finally the axiom (H) for \mathbf{H} and \mathbf{H}_* are transposed of each other.

2 Graded Hopf algebras

Notation 2.1 With H a C-vector space direct sum of C-vector subspaces:

$$\mathbf{H} = \otimes_{n \in \mathbb{N}} \mathbf{H}^n \tag{2.1}$$

we let $\theta_z, z \in \mathbb{C}$, resp. Y, be the \mathbb{C} -linear maps: $\mathbf{H} \to \mathbf{H}$ respectively specified by:

$$\theta_{z}|_{\mathbf{H}^{n}} = e^{nz} \qquad \left(\Rightarrow \left\{ \begin{array}{l} \theta_{z+z'} = \theta_{z^{o}}\theta_{z} \\ \theta_{z} = e^{zY} \end{array} \right., \ z, z' \in \mathbf{k} \right), \tag{2.2}$$

$$Y_{\mathbf{H}^n} = n \qquad \left(\Leftrightarrow \frac{d}{dt} \theta_z |_{t=0} \right), \tag{2.3}$$

and set, for $a \in \mathbf{H}$ homogeneous:

$$\partial_Y(a) = n \text{ if } a \in \mathbf{H}^n, \qquad n \in \mathbb{N}.$$
 (2.4)

Definition-Lemma 2.2 (N-graded algebras, coalgebras and bialgebras).

(i) The \mathbb{C} -algebra $\mathbf{H} = \mathbf{H}(m, \mathbf{1} = e(1)) = \bigoplus_{n \in \mathbb{N}} \mathbf{H}^n$ is \mathbb{N} -graded (with algebra \mathbb{N} -grading $\bigoplus_{n \in \mathbb{N}} \mathbf{H}^n$) whenever one of the following equivalent requirements (2.5), (2.6), (2.7) or (2.8) prevails:

$$\mathbf{H}^{p} \cdot \mathbf{H}^{q} \subset \mathbf{H}^{p+q} \qquad , p, q \in \mathbb{N}$$

$$(2.5)$$

In the graded connected case isomorphic to the enveloping algebra $\mathbf{U}(\mathbf{L})$ of the Lie algebra \mathbf{L} of its primitive elements (special case of the Milnor-Moore Theorem).

duality generally not separating ${\bf H}$ – certainly not for ${\bf H}$ non-commutative., separating however in the case of CKM-Hopf algebras, cf. 3.5.

 $a \in H$ is **homogeneous** whenever $a \in \bigcup_{n \in \mathbb{N}} \mathbf{H}^n$ ($\Leftrightarrow a$ is an eigenvector of Y – then with eigenvalue $\partial_Y(a)$).

Daniel Kastler

$$\begin{cases} m \circ (\theta_z \otimes \theta_z) = \theta_z \circ m\\ \text{i.e., } \theta_z(ab) = (\theta_z a)(\theta_z b), \ a, b \in \mathbf{H} \end{cases}, \qquad z \in \mathbb{C}, \tag{2.6}$$

(saying that θ_z is a one-parameter automorphism group of the algebra $\mathbf{H}(m, \mathbf{1} = e(1))$, and implying $\theta_z(\mathbf{1}) = \mathbf{1}, z \in \mathbb{C}$)

$$\begin{cases} Y \circ m = m \circ (Y \otimes \operatorname{id}_{\mathbf{H}} + \operatorname{id}_{\mathbf{H}} \otimes Y) \\ \text{i.e., } Y(ab) = (Ya)b + a(Yb), \ a, b \in \mathbf{H} \end{cases}$$
(2.7)

(saying that Y is a derivation of the algebra $\mathbf{H}(m, \mathbf{1} = e(1))$, and implying $Y(\mathbf{1}) = 0$),

$$\partial_Y(ab) = \partial_Y(a) + \partial_Y(b) \tag{2.8}$$

for $a, b \in \mathbf{H}$ homogeneous, which implies $ab \in \mathbf{H}$ homogeneous.

(ii) The C-coalgebra $\mathbf{H} = \mathbf{H}(\Delta, \varepsilon) = \bigoplus_{n \in \mathbb{N}} \mathbf{H}^n$ is *N*-graded (with coalgebra *N*-grading $\bigoplus_{n \in \mathbb{N}} \mathbf{H}^n$) whenever one of the following equivalent requirements (2.9), (2.10), (2.11), or (2.12) prevails:

$$\Delta(\mathbf{H}^n) \subset \sum_{i+j=n} \mathbf{H}^i \otimes \mathbf{H}^j, \qquad n \in \mathbb{N},$$
(2.9)

$$(\theta_z \otimes \theta_z) \circ \Delta = \Delta \circ \theta_z, \qquad z \in \mathbb{C}, \tag{2.10}$$

(saying that θ_z is a one-parameter automorphism group of the coalgebra $\mathbf{H}(\Delta, \varepsilon)$, and implying $\varepsilon(\theta_z(a)) = \varepsilon(a), \ a \in H(\Delta, \varepsilon)$)

$$Y \otimes \mathrm{id}_{\mathbf{H}} + \mathrm{id}_{\mathbf{H}} \otimes Y) \circ \Delta = \Delta \circ Y.$$
(2.11)

(saying that Y is a coalgebra coderivation, and implying $\epsilon(Ya) = 0, a \in H(\Delta, \varepsilon)$)

$$\partial_Y(a) = \partial_Y(a_{(1)}) + \partial_Y(a_{(2)}) \quad \text{if } \Delta a = a_{(1)} \otimes a_{(2)}, \ a \in \mathbf{H}, \text{ a homogeneous.}$$
(2.12)

(iii) The \mathbb{C} -bialgebra $\mathbf{H}(m, \Delta, e, \varepsilon) = \bigoplus_{n \in \mathbb{N}} \mathbf{H}^n$ is \mathbb{N} -graded (with bialgebra \mathbb{N} -grading $\bigoplus_{n \in \mathbb{N}} \mathbf{H}^n$) whenever it is \mathbb{N} -graded both as an algebra and as a coalgebra. We say that \mathbf{H} is \mathbb{N} -graded connected whenever one has in addition:

$$\mathbf{H}^0 = \mathbb{C}\mathbf{1}.\tag{2.13}$$

(iv) A Hopf algebra $\mathbf{H}(m, \Delta, e, \varepsilon, S)$ is called N-graded whenever it is N-graded as a bialgebra (cf. Definition-Lemma 2.2(i)). S then automatically commutes with $\theta_z, z \in \mathbb{C}$, and with Y:

$$\begin{cases} \theta_z \circ S \circ \theta_z^{-1} = S\\ Y \circ S = S \circ Y \end{cases}$$
(2.14)

For the N-graded algebra (resp. coalgebra, bialgebra) **H** we refer to θ_z as the **automorphism group**, to Y as the **derivation** (resp. **coderivation**, **biderivation**), and to ∂_Y as the **degree**, of the N-grading.

Proof (i) (2.5) \Leftrightarrow (2.6): \Rightarrow : it suffices to take $a \in \mathbf{H}^p$ and $b \in \mathbf{H}^q$, then $ab \in \mathbf{H}^{p+q}$, then:

$$\theta_z(ab) = e^{z(p+q)}ab = (e^{zp}a)(e^{zq}b)b = (\theta_z a)\theta_z b).$$

 \Leftarrow : for $a \in \mathbf{H}^p$ and $b \in \mathbf{H}^q$,

$$\theta_z(ab) = (\theta_z a)(\theta_z b) = (e^{zp}a)(e^{zq}b)b = e^{z(p+q)}ab$$

thus $ab \in \mathbf{H}^{p+q}$.

Note that (2.1) obviously implies $\mathbf{H} \otimes \mathbf{H} = \sum_{i+j} \mathbf{H}^i \otimes \mathbf{H}^j$.

 $(2.5) \Leftrightarrow (2.7)$: \Rightarrow : it suffices to take $a \in \mathbf{H}^p$ and $b \in \mathbf{H}^q$, then

$$Y(ab) = (p+q)ab = (pa)b + a(qb) = (Ya)b + a(Yb).$$

 \Leftarrow : for $a \in \mathbf{H}^p$ and $b \in \mathbf{H}^q$,

$$Y(ab) = (Ya)b + a(Yb) = (pa)b + a(qb) = (p+q)ab$$

thus $ab \in \mathbf{H}^{p+q}$.

 $(2.7) \Leftrightarrow (2.8)$: $Y(ab) = \partial_Y(ab)ab$, whilst

$$(Ya)b + a(Yb) = \partial_Y(a)ab + a\partial_Y(b)b = [\partial_Y(a) + \partial_Y(b)]ab$$

(ii) (2.9) \Leftrightarrow (2.10): \Rightarrow : it suffices to take $a \in \mathbf{H}^p$, then

$$\begin{aligned} [(\theta_z \otimes \theta_z) \circ \Delta] a &= (\theta_z \otimes \theta_z)(a_{(1)} \otimes a_{(2)}) \\ &= (\theta_z a_{(1)}) \otimes (\theta_z a_{(2)}) = e^{p_1 z} a_{(1)} \otimes e^{p_2 z} a_{(2)} \\ &= e^{p z} \Delta a = \Delta (e^{p z} a) = [\Delta \circ \theta_z] a. \end{aligned}$$

 \Leftarrow : for $a \in \mathbf{H}^p$,

$$\begin{aligned} [(\theta_z \otimes \theta_z) \circ \Delta] a &= (\theta_z \otimes \theta_z)(a_{(1)} \otimes a_{(2)}) \\ &= e^{p_1 z} a_{(1)} \otimes e^{p_2 z} a_{(2)} \\ &= e^{(p_1 + p_2) z} \Delta a = [\Delta \circ \theta_z] a = e^{p z} \Delta a, \end{aligned}$$

implying $p_1 + p_2 = p$.

(2.9) \Leftrightarrow (2.11): \Rightarrow : it suffices to take $a \in \mathbf{H}^p$, then:

$$\begin{split} [(Y \otimes \mathrm{id}_{\mathbf{H}} + \mathrm{id}_{\mathbf{H}} \otimes Y) \circ \Delta] a &= (Y \otimes \mathrm{id}_{\mathbf{H}} + \mathrm{id}_{\mathbf{H}} \otimes Y)(a_{(1)} \otimes a_{(2)}) \\ &= Ya_{(1)} \otimes a_{(2)} + a_{(1)} \otimes (Ya_{(2)}) \\ &= (p_1 + p_2)(a_{(1)} \otimes a_{(2)}) \\ &= p\Delta(a) = \Delta(pa) = [\Delta \circ Y]a. \end{split}$$

 $\Leftarrow: \text{ for } a \in \mathbf{H}^p,$

$$[(Y \otimes \mathrm{id}_{\mathbf{H}} + \mathrm{id}_{\mathbf{H}} \otimes Y) \circ \Delta]a = (Y \otimes \mathrm{id}_{\mathbf{H}} + \mathrm{id}_{\mathbf{H}} \otimes Y)(a_{(1)} \otimes a_{(2)})$$
$$= p_1 a_{(1)} \otimes a_{(2)} + a_{(1)} \otimes p_2 a_{(2)}$$
$$= (p_1 + p_2)(a_{(1)} \otimes a_{(2)})$$
$$= (p_1 + p_2)\Delta a = (\Delta \circ Y)a = p\Delta a$$

implying $p_1 + p_2 = p$ as above.

 $(2.11) \Leftrightarrow (2.12): \Delta(Ya) = \Delta[\partial_Y(a)a] = \partial_Y(a)\Delta a$, whilst

$$(Y \otimes \mathrm{id}_{\mathbf{H}} + \mathrm{id}_{\mathbf{H}} \otimes Y) \circ \Delta a = (Y \otimes \mathrm{id}_{\mathbf{H}} + \mathrm{id}_{\mathbf{H}} \otimes Y)(a_{(1)} \otimes a_{(2)})$$

$$= Ya_{(1)} \otimes a_{(2)} + a_{(1)} \otimes Ya_{(2)}$$

$$= \partial_Y(a_{(1)})a_{(1)} \otimes a_{(2)} + a_{(1)} \otimes \partial_Y(a_{(2)})a_{(2)}$$

$$= [\partial_Y(a_{(1)}) + \partial_Y(a_{(2)})]\Delta a.$$

We chose the Sweedler basis $a_{(1)} \otimes a_{(2)}$ such that $a_{(1)}$, resp. the $a_{(2)}$, are linearly independent.

(iv) Check of the first line (2.14): we check that $\theta_z \circ S \circ \theta_z^{-1}$ is a convolution inverse of $e\varepsilon$:

$$\begin{aligned} (\theta_z \circ S \circ \theta_z^{-1}) * \mathrm{id}_{\mathbf{H}} &= m(\theta_z \circ S \circ \theta_z^{-1} \otimes \mathrm{id}_{\mathbf{H}}) \Delta \\ &= m(\theta_z \circ S \circ \theta_z^{-1} \otimes \mathrm{id}_{\mathbf{H}})(\theta_z \otimes \theta_z) \Delta \theta_z^{-1} \\ &= m(\theta_z \otimes \mathrm{id}_{\mathbf{H}}(S \otimes \mathrm{id}_{\mathbf{H}})(\theta_z^{-1} \otimes \mathrm{id}_{\mathbf{H}})(\theta_z \otimes \theta_z) \Delta \theta_z^{-1} \\ &= m(\theta_z \otimes \mathrm{id}_{\mathbf{H}})(S \otimes \mathrm{id}_{\mathbf{H}})(\mathrm{id}_{\mathbf{H}} \otimes \theta_z) \Delta \theta_z^{-1} \\ &= m(\theta_z \otimes \mathrm{id}_{\mathbf{H}})(\mathrm{id}_{\mathbf{H}} \otimes \theta_z)(S \otimes \mathrm{id}_{\mathbf{H}}) \Delta \theta_z^{-1} \\ &= m(\theta_z \otimes \theta_z)(S \otimes \mathrm{id}_{\mathbf{H}}) \Delta \theta_z^{-1} = \theta_z^{-1} m(S \otimes \mathrm{id}_{\mathbf{H}}) \\ &= \theta_z \varepsilon \varepsilon \theta_z^{-1} = \varepsilon \varepsilon, \end{aligned}$$

where we used (2.10), (2.6), $\theta_z(\mathbf{1}) = \mathbf{1}$, and $\varepsilon \circ \theta_z = \varepsilon$.

Check of the second line (2.14): equate the derivations at 0 of $\theta_z \circ S = S \circ \theta_z$. \Box

In practice recognizing a biderivation often occurs by first establishing the derivation property, and afterwards the coderivation property, using

Lemma 2.3 Assume that one knows that a bialgebra \mathbf{H} is \mathbb{N} -graded as an algebra. For investigating the possible coderivation property of Y one needs only look at the restriction of Y on a set of generators.

Proof From

$$\begin{cases} \partial_Y(a) = \partial_Y(a_{(1)}) + \partial_Y(a_{(2)}) & \text{if} \quad \Delta a = a_{(1)} \otimes a_{(2)} \\ \partial_Y(b) = \partial_Y(b_{(1)}) + \partial_Y(b_{(2)}) & \text{if} \quad \Delta a = a_{(1)} \otimes a_{(2)} \end{cases}, \ a, b \in \mathbf{H},$$

a, b homogeneous, we conclude for the product ab with coproduct $\Delta(ab) = a_{(1)}b_{(3)} \otimes a_{(2)}b_{(4)}$ that $\partial_Y(ab) = \partial_Y(a) + \partial_Y(b)$ equals

$$\partial_Y(a_{(1)}b_{(3)}) + \partial_Y(a_{(2)}b_{(4)}) = \partial_Y(a_{(1)}) + \partial_Y(b_{(3)}) + \partial_Y(a_{(2)}) + \partial_Y(b_{(4)}) = \partial_Y(a_{(1)}b_{(3)}) + \partial_Y(a_{(2)}b_{(4)}) + \partial_Y(a_{(2)}b_{(4)}) = \partial_Y(a_{(1)}b_{(3)}) + \partial_Y(a_{(2)}b_{(4)}) + \partial_Y($$

It is instructive to check instead the property (2.11), showing that

 $(Y \otimes \mathrm{id}_{\mathbf{H}} + \mathrm{id}_{\mathbf{H}} \otimes Y) \Delta(ab)$

$$= [(Y \otimes \mathrm{id}_{\mathbf{H}} + \mathrm{id}_{\mathbf{H}} \otimes Y)\Delta(a)]\Delta(b) + \Delta(a)[Y \otimes \mathrm{id}_{\mathbf{H}} + \mathrm{id}_{\mathbf{H}} \otimes Y)\Delta(b)]$$

$$= \Delta(Ya)\Delta(b) + \Delta(a)\Delta(Yb) = \Delta[(Ya)b + a(Yb)] = \Delta[Y(ab)].$$

We next show that the algebraïc dual of a N-graded coalgebra is a N-graded algebra.

Proposition 2.4 (the dual N-grading). With $\mathbf{H}(m, \Delta, e, \varepsilon)$, $\mathbf{H} = \bigoplus_{n \in \mathbb{N}} \mathbf{H}^n$, a \mathbb{N} - graded bialgebra, let \mathbf{H}^* be the algebraic dual of \mathbf{H} endowed with the algebra-structure dual of the coalgebra-structure of \mathbf{H} , thus with product:

$$\langle \xi * \eta, h \rangle = \langle \xi \otimes \eta, \Delta h \rangle, \quad \xi, \eta \in \mathbf{H}^*.$$
 (2.15)

The definitions:

$$\begin{cases} < Y_*\xi, h > = <\xi, Yh > \\ < \theta_{*z}\xi, h > = <\xi, \theta_z h > \end{cases}, \ \xi \in \mathbf{H}^*, \ h \in \mathbf{H}, \end{cases}$$
(2.16)

Here the convolution product is that of $\operatorname{End}_{\mathbb{C}}(\mathbf{H}, \mathbf{H}^*)$, and $\langle \cdot, \cdot \rangle$ is the duality bilinear form, cf. (1.1)

then yield a \mathbb{N} -grading $\mathbf{H}^* = \bigoplus_{n \in \mathbb{N}} \mathbf{H}^{*n}$ of the dual algebra \mathbf{H}^* where $\mathbf{H}^{*n} = \bigcap_{p \neq n} \mathbf{H}^{p\perp}$. One has the inclusions $Y_* "P(\mathbf{H}^*)" \subset "P(\mathbf{H}^*)"$ and $\theta_{*z} "G(\mathbf{H}^*)" \subset "G(\mathbf{H}^*)"$, $z \in \mathbb{C}$. \mathbf{H}_* thus becomes a graded bialgebra.

Proof We check that one has $Y_*\xi = n\xi$ for $\xi \in \mathbf{H}^{*n}$. With $\mathrm{id}_{\mathbf{H}} = \sum_{n \in \mathbb{N}} P_n$ the decomposition of the unit associated to the direct sum $\mathbf{H} = \bigoplus_{n \in \mathbb{N}} \mathbf{H}^n$, we have $Y = \sum_{n \in \mathbb{N}} n P_n$, thus for $\xi \in \mathbf{H}^{*n}$ and $h \in \mathbf{H}$,

$$<\xi, Yh> = <\xi, P_nYh> = <\xi, YP_nh> = n <\xi, P_nh> = n <\xi, h> = ,$$

hence $Y_*\xi = n\xi$. The fact that (2.7) and (2.11) are transposed of each other then implies that Y_* is a derivation of the algebra \mathbf{H}^* . The proof of $\theta_{*z}\xi = e^{nz}\xi$ for $\xi \in \mathbf{H}^{*n}$ is analogous. Check of the two last claims:

$$\begin{aligned} -\operatorname{for} \,\xi &\in \, {}^{\circ}P(\mathbf{H}^{*})^{"}:\\ \Delta_{*}Y_{*}\xi &= \, (\operatorname{id} \otimes Y_{*} + Y_{*} \otimes \operatorname{id})\Delta_{*}\xi = (\operatorname{id} \otimes Y_{*} + Y_{*} \otimes \operatorname{id})(\mathbf{1} \otimes \xi + \xi \otimes \mathbf{1})\\ &= \, (\mathbf{1} \otimes Y_{*}\xi + Y_{*}\xi \otimes \mathbf{1}).\\ -\operatorname{for} \,\xi &\in \, {}^{\circ}G(\mathbf{H}^{*})^{"} \text{ and } z \in \mathbb{C}: \end{aligned}$$

$$\Delta_*\theta_{*z}\xi = (\theta_{*z}\otimes\theta_{*z})\Delta_*\xi = (\theta_{*z}\otimes\theta_{*z})(\xi\otimes\xi) = (\theta_{*z}\xi\otimes\theta_{*z}\xi).$$

Proposition 2.5 (N-graded connected bialgebras.) Let

$$\mathbf{H}(m,\Delta,e,\varepsilon) = \oplus_{n \in \mathbb{N}} \mathbf{H}^n$$

be a \mathbb{N} -graded connected bialgebra (i.e. $\mathbf{H}^0 = \mathbb{C}\mathbf{1}$) and let $\mathbf{H}^+ = Ker\varepsilon$ (called the **augmentation ideal** – it is known that $\mathbf{H} = \mathbb{C}\mathbf{1}\oplus\mathbf{H}^+$ cf. Proposition A.3). Then: (i) one has:

$$\mathbf{H}^{+} = Ker\varepsilon = \bigoplus_{n \ge 1} \mathbf{H}^{n} \tag{2.17}$$

(ii) one has the implication:

$$h \in \mathbf{H}^+, \ h \ homogeneous \ \Rightarrow \Delta h = h \otimes \mathbf{1} + \mathbf{1} \otimes h + h' \otimes h'',$$
 (2.18)

with $h' \otimes h''$ a Sweedler-type sum of terms fulfilling either (and thus both) of the equivalent requirements:

- (a) all $\delta_Y(h')$, $\delta_Y(h'')$ are strictly smaller than $\delta_Y(h)$.
- (b) all $\delta_Y(h')$, $\delta_Y(h'')$ are strictly positive.

(Equivalence of (a) and (b): since the bialgebra N-grading entails that one has

$$\delta_Y(h') + \delta_Y(h'') = \delta_Y(h), \quad \delta_Y(h') = \delta_Y(h) - \delta_Y(h'')$$

and

$$\delta_Y(h'') = \delta_Y(h) - \delta_Y(h')$$

are simultaneously strictly positive).

The consequence (2.18) of N-graded connectedness plays a prominent technical role both in the Connes-Moscovici index theory and in the Connes-Kreimer theory

We define " $P(\mathbf{H}^*)$ " = { $\xi \in \mathbf{H}^*, \Delta_* \xi = \xi \otimes \mathbf{1} + \mathbf{1} \otimes \xi$ }, " $G(\mathbf{H}^*)$ " = { $\xi \in \mathbf{H}^*, \Delta_* \xi = \xi \otimes \xi$ } (written with quotation marks because \mathbf{H}^* is not a bona-fide coalgebra – this notation becomes bona-fide in \mathbf{H}_* , cf. Section 1).

Note that the property $\delta_Y(h') + \delta_Y(h'') = \delta_Y(h)$ of the Hopf N-grading merely entails that $\delta_Y(h'), \delta_Y(h'') \leq \delta_Y(h)$, whilst progressivity requires strict inequalities. Note also that applying $\varepsilon \otimes \operatorname{id}_H$ and $\operatorname{id}_H \otimes \varepsilon$ to both sides of (2.18) one proves $\varepsilon(h') = \varepsilon(h'') = 0$, hence that all h' and h'' belong to \mathbf{H}^+ .

of renormalization. We therefore cast the special name of **progressiveness** for this property, 2.5(ii) above then says that a \mathbb{N} -graded connected bialgebra is progressive.

Proof (i) Let $h \in \mathbf{H}^n$, $n \ge 1$: by (2.12) we have $\Delta h = h_n \otimes 1 + \sum_{i=1,\dots,n-1} \sum_k h_{n-i,k} \otimes h'_{i,k} + 1 \otimes h'_n$

(first subscript indicating the ∂_Y -grade).

Hence:

$$h = (\varepsilon \otimes \operatorname{id})\Delta h = \varepsilon(h_n)\mathbf{1} + \Sigma_{i=1,\dots,n-1}\Sigma_k\varepsilon(h_{n-i,k})h'_{i,k} + h'_n$$

whence $\varepsilon(h_n) = 0$ and $h = h'_n$
$$= (\operatorname{id} \otimes \varepsilon)\Delta h = h_n + \Sigma_{i=1,\dots,n-1}\Sigma_k h_{n-i}\varepsilon(h'_{i,k}) + \varepsilon(h'_n)\mathbf{1}$$

whence $h = h_n$ and $\varepsilon(h'_n) = 0$,

implying $\varepsilon(h) = \varepsilon(h_n) = 0$. We proved that $\bigoplus_{n \ge 1} \mathbf{H}^n \subset \mathbf{H}^+ = \text{Ker}\varepsilon$. (In fact we found that:

$$\Delta h = h \otimes \mathbf{1} + \mathbf{1} \otimes h'_n + \Sigma_{i=1,\dots,n-1} \Sigma_k h_{n-i,k} \otimes h'_{i,k}, \quad h \in \mathbf{H}^n,$$
(2.19)

first subscript indicating the ∂_Y -grade).

Conversely let $h \in \mathbf{H}^+ = \operatorname{Ker} \varepsilon$, with $h = \lambda \mathbf{1} + \sum_{n \ge 1} h_n$, $h_n \in \mathbf{H}^n$. By what precedes we have $0 = \varepsilon(h) = \lambda$, hence $h = \sum_{n \ge 1} h_n$, proving that $\mathbf{H}^+ \subset \bigoplus_{n \ge 1} \mathbf{H}^n$.

(ii) For $h \in \mathbf{H}^n$, (2.19) implies (2.18). However (2.18) is linear w.r.t. h, hence holds for $h \in \mathbf{H}^+$ by (i).

Remark 2.6 (i) Progressiveness at large follows from progressiveness for generators of \mathbf{H}^+ . Indeed:

(ii) Let $h, k \in \mathbf{H}^+$: if (2.18) holds for h and k, it holds for the product hk.

Proof (i) follows from (ii) and the multiplicativity of Δ .

(ii) assume that $\Delta h = h \otimes \mathbf{1} + \mathbf{1} \otimes h + h' \otimes h''$ with all $\delta_Y(h')$, $\delta_Y(h'')$ strictly positive, and $\Delta k = k \otimes \mathbf{1} + \mathbf{1} \otimes k + k' \otimes k''$ with all $\delta_Y(k')$, $\delta_Y(k'')$ strictly positive. It follows that

$$\Delta(hk) = (hk) \otimes \mathbf{1} + \mathbf{1} \otimes (hk) + h \otimes k + k \otimes h + hk' \otimes k'' + k' \otimes hk'' + h'k \otimes h'' + h' \otimes h''k + k'k \otimes k'' + k' \otimes k''k + h'k' \otimes h''k'',$$

where all the tensor products but the two first have both factors of strictly positive degree. $\hfill \Box$

We now present two important consequences of progressiveness. The first is that it makes automatic the existence of a (recursively defined) antipode:

Proposition 2.7 Assume that the \mathbb{N} -graded bialgebra $\mathbf{H}(m, \Delta, e, \varepsilon) = \bigoplus_{n \in \mathbb{N}} \mathbf{H}^n$ is \mathbb{N} -graded connected (thus progressive). Then \mathbf{H} is a Hopf algebra.

Proof The requirements, for the \mathbb{C} -linear S_r , resp. S_l : $\mathbf{H} \rightarrow \mathbf{H}$:

$$\begin{cases} S_r \mathbf{1} = \mathbf{1} \\ S_r h = -h - h' S_r h'', \quad h \in \mathbf{H}^+ = \ker \varepsilon \\ (\text{saying that } \operatorname{id}_{\mathbf{H}} * S_r = m(\operatorname{id}_{\mathbf{H}} \otimes S_r) \Delta = e\varepsilon), \end{cases}$$
(2.20)

In fact it already played the role in the early days of the theory of Hopf algebras, through the corresponding inductive definition of the antipode, cf. Proposition 2.7.

resp.

$$\begin{cases} S_l \mathbf{1} = \mathbf{1} \\ S_l h = -h - (S_l h') h'', \quad h \in \mathbf{H}^+ = \ker \varepsilon \\ (\text{saying that id}_{\mathbf{H}} * S_l = m(S_l \otimes \operatorname{id}_{\mathbf{H}}) \Delta = e\varepsilon), \end{cases}$$
(2.21)

determine S_r and S_l by induction w.r.t. the degree δ_Y as respective right- and left convolution inverses of $\mathrm{id}_{\mathbf{H}}$ for the convolution product of $\mathrm{End}_{\mathbb{C}}(\mathbf{H}, \mathbf{H})$: but these then coincide because $S_l * \mathrm{id}_{\mathbf{H}} * S_r = e\varepsilon * S_r = S_r = S_l * e\varepsilon = S_l$: thus $S = S_r = S_r$ is a bilateral convolution inverse of $\mathrm{id}_{\mathbf{H}}$, thus the (unique) antipode.

The second consequence of progressiveness of **H** is that, given $h \in \mathbf{H}$, there is a drastic limitation (depending on h) of non-vanishing values $\langle \xi, h \rangle, \xi \in \mathbf{H}_*$. We recall that (cf. Section 1):

– the algebraic dual \mathbf{H}^* of \mathbf{H} becomes a topological algebra if equipped with the $\sigma(\mathbf{H}^*, \mathbf{H})$ -topology and the algebra-structure $(m_* =^*, \mathbf{1}_* = e_*(1))$ stemming from the coalgebra-structure of \mathbf{H} ,

– the subalgebra \mathbf{H}_* generated in \mathbf{H}^* by the unit $\mathbf{1}_*$ and the Lie-algebra \mathbf{L} of infinitesimal characters becomes by transposition a Hopf algebra $\mathbf{H}_*(m_* =^*)$, $\mathbf{1}_* = e_*(1), \Delta_*, \varepsilon_*, S_*)$.

Lemma 2.8 Let the bialgebra $\mathbf{H}(m, \Delta, e, \varepsilon) = \bigoplus_{n \in \mathbb{N}} \mathbf{H}^n$ be \mathbb{N} -graded connected (thus progressive) and let P^+ be the projection on \mathbf{H}^+ parallel to $\mathbb{C}\mathbf{1}$ in \mathbf{H} . Then:

(i) One has for $n \in \mathbb{N}$ the implication:

$$\partial_Y(h) \le n \Rightarrow P^{+\otimes(n+1)} \Delta^{(n)} h = 0. \tag{2.22}$$

(ii) One has consequently for $n \in \mathbb{N}$ the implication:

$$\partial_{Y}(h) \leq n \ (i.e. \ h \in \bigoplus_{p \leq n} \mathbf{H}^{p}) \quad \Rightarrow \quad \langle Z_{1} \ast \cdots \ast Z_{n+1}, h \rangle = 0$$

for all $Z_{1}, \ldots, Z_{n+1} \in \mathbf{L}.$ (2.23)

alternatively formulated as:

$$Z_1, \dots, Z_{n+1} \in \mathbf{L} \Rightarrow Z_1 * \dots * Z_{n+1} \in (\bigoplus_{p \le n} \mathbf{H}^p)^{\perp}, \quad n \in \mathbb{N},$$
(2.24)

where \perp indicates the annihilator in the dual).

Proof (i) Since by (2.17) a general element of **H** is of the form $\lambda \mathbf{1} + h$ with $h \in \mathbf{H}^+$, and we have obviously $P^{+\otimes(n+1)}\Delta^{(n)}\mathbf{1} = 0$, it is no restriction for the proof of the implication (2.22) to assume that h belongs to \mathbf{H}^+ . The projection $P^{+\otimes(n+1)}$ applied to $\Delta^{(n)}h = h_{(1)} \otimes h_{(2)} \otimes \cdots \otimes h_{(n+1)}$ then suppresses all the products $h_{(1)} \otimes h_{(2)} \otimes \cdots \otimes h_{(n+1)}$ containing at least one factor in $\mathbf{H}^0 = \mathbb{C}\mathbf{1}$: the remaining products $h_{(1)} \otimes h_{(2)} \otimes \cdots \otimes h_{(n+1)}$ stem from the successive Sweedler-type summations $h' \otimes h''$ in (2.18), thus have by progressiveness all their factors $h^{(p)}$ such that $\delta_Y(h^{(p)} \ge 1$ (observe, as noticed in footnote 6, that all h', h'' belong to \mathbf{H}^+). The equality

$$\delta_Y(h^{(1)}) + \delta_Y(h^{(2)}) + \dots + \delta_Y(h^{(n+1)}) = \delta_Y(h)$$

requires $n + 1 \leq \delta_Y(h) \Leftrightarrow n < \delta_Y(h)$, otherwise $P^{+\otimes(n+1)}\Delta^{(n)}h$ vanishes, whence (2.20) and (2.22) for $h \in \mathbf{H}^+$, the extension to $h \in \mathbf{H}$ being trivial.

hence a Hopf algebra, cf. Proposition 2.7

One has $H = \mathbb{C}\mathbf{1} \oplus \mathbf{H}^+$, cf. (2.17), and $P^+ = \mathrm{id}_{\mathbf{H}} - e\varepsilon$, cf. Proposition A.3. Correlatively $P^{+\otimes p}$ projects $\mathbf{H}^{\otimes p}$ onto $\mathbf{H}^{+\otimes p}$.

Daniel Kastler

(ii) The expression of the n-fold convolution product of \mathbf{H}^* :

$$\langle \xi_1 * \cdots \xi_n, h \rangle = \langle \xi_1 \otimes \cdots \otimes \xi_n, \Delta^{(n)}h \rangle$$

= $\langle \xi_1 \otimes \cdots \otimes \xi_n, h_{(1)} \otimes h_{(2)} \otimes \cdots \otimes h_{(n+1)} \rangle$ (2.25)

yields for products of elements of **L** vanishing on **1**:

$$\langle Z_1 * \cdots * Z_{n+1}, h \rangle = \langle Z_1 \otimes \cdots \otimes Z_n, P^{+ \otimes (n+1)} \Delta^{(n)} h \rangle$$

$$Z_1, \ldots, Z_{n+1} \in \mathbf{L}, \ h \in \mathbf{H},$$
(2.26)

(2.23) thus follows from (2.22). Note that (2.23) holds for n = 0 (obvious directly - embodying this into our proof would require the convention $\Delta^{(0)} = id_{\mathbf{H}}$).

Lemma 2.9 Assume that the \mathbb{N} -graded connected bialgebra $\mathbf{H}(m, \Delta, e, \varepsilon) = \bigoplus_{n \in \mathbb{N}} \mathbf{H}^n$ has a countable vector space-basis $\{x_i\}_{i \in \mathbb{N}}$ consisting of homogeneous elements of non-decreasing degree, and let \mathbf{H}_* be the closure of \mathbf{H}_* in \mathbf{H} . Then:

(i) \mathbf{H}^* is metrizable with the distance

$$\begin{cases} d(\xi,\eta) = f(\xi - \eta) \\ where \ f(\xi) = \sum_{i \in \mathbb{N}} 2^{-i} \inf(|<\xi, x_i > |, 1) \end{cases}, \ \xi, \eta \in \mathbf{H}^*, \tag{2.27}$$

(ii) \mathbf{H}_* is a closed topological subalgebra of \mathbf{H} .

(iii) $\overline{\mathbf{H}}_*$ is stable under S_* acting in \mathbf{H}^* .

 \rightarrow Note that at this point the question of whether $\Delta * \mathbf{H}_* \subset \mathbf{H}_* \otimes \mathbf{H}_*$, and whether, if so,

$$\mathbf{H}_{*}(m_{*} =^{*}, \mathbf{1}_{*} = e_{*}(1), \Delta_{*}, \varepsilon_{*}, S_{*})$$

is a Hopf algebra, remains open.

Proof (i) is the known device for proving metrizability of a topological vectorspace whose topology is specified by a countable set of semi-norms. One checks all the properties stated in footnote 9.

(ii) follows from the known fact that the closure of a topological algebra is an algebra.

(iii) Check of $S_* \overline{\mathbf{H}}_* \subset \mathbf{H}_*$: let $\xi \in \overline{\mathbf{H}}_*$, $\xi = \lim_{n \to \infty} \zeta_n$ with $\zeta_n \in \mathbf{H}_*$, we have, for $h \in \mathbf{H}$:

$$\langle S_*\zeta_n, h \rangle_{n=\infty}, \langle \xi, Sh \rangle = \langle S_*\xi, h \rangle,$$

hence $S_*\zeta_n \xrightarrow[n=\infty]{} S_*\xi$, hence $S_*\xi \in \overline{\mathbf{H}}_*$.

Proposition 2.10 Assuming that the \mathbb{N} -graded connected bialgebra $\mathbf{H}(m, \Delta, e, \varepsilon) = \bigoplus_{n \in \mathbb{N}} \mathbf{H}^n$ has a countable vector space-basis $\{x_i\}_{i \in \mathbb{N}}$ such that $\partial(x_i) \to \infty$ for $n \to \infty$. Let $\overline{\mathbf{H}}_*$ be the closure of \mathbf{H}_* in \mathbf{H} , we have that:

(i) with $Z \in \mathbf{L}$, and $a_i \in \mathbb{C}$, $i \in \mathbb{N}$, the sequence $\Omega_n = \sum_{i=1,\dots,n} a_i Z^{*i}$ is a Cauchysequence of \mathbf{H}^* which converges for $n \to \infty$ towards an element $\Omega \in \overline{\mathbf{H}}_*$.

(ia) the same holds for multi-variable power-series of elements of L.

d is a translation-invariant distance: one has $d(\xi, \eta) \ge 0$, $d(\xi, \eta) = d(\eta, \xi)$, $d(\xi, \eta) \le d(\xi, \zeta) + d(\zeta, \eta)$, and $d(\xi + \zeta, \eta + \zeta) = d(\xi, \eta)$ for all $\eta, \zeta \in \bar{\mathbf{H}}_*$, and $d(\xi, \eta) = 0$, $\xi, \eta \in \bar{\mathbf{H}}_*$ implies $\xi = \eta$. These properties follow from the properties $f(\xi + \eta) \le f(\xi) + f(\eta)$, $f(-\xi) = f(\xi)$ for all $\xi, \eta \in \bar{\mathbf{H}}_*$ plus the implication $f(\xi) = 0 \Rightarrow \langle \xi, x_i \rangle = 0$ for all $i \in \mathbb{N}$. The equality of topologies arises as follows: the $V_i = \{\xi \in \bar{\mathbf{H}}_*; f(\xi) < 2^{-i}\}$ are a basis of neighbourhoods of zero in the *d*-topology, with $U_i\{\xi \in \bar{\mathbf{H}}_*; | < \xi, x_i > | < 1\} \supseteq V_i$.

(ii) in particular the Mac-Laurin power-series of e^Z , $Z \in \mathbf{L}$, yields an element, denoted by e^Z , of $\bar{\mathbf{H}}_*$ which is a character of \mathbf{H} .

 \rightarrow Note that at this point the question of whether the e^Z , $Z \in \mathbf{L}$, generate the group **G** of characters of **H** remains open.

Proof (i) we have from (2.27), with
$$i(n) = \inf\{i \in \mathbb{N}; \partial_Y(x_i) \ge n\}$$
:

$$d(\Omega_n, \Omega_{n+p}) = \sum_{i \in \mathbb{N}} 2^{-i} \inf(|\Omega_{n+p} - \Omega_n, x_i > |, 1)$$

$$= \sum_{i \in \mathbb{N}} 2^{-i} \inf(| < a_n Z^{*n} + \dots + a_{n+p} Z^{*n+p}, x_i > |1)$$

$$= \sum_{i \ge i(n)} 2^{-i} \inf(| < a_n Z^{*n} + \dots + a_{n+p} Z^{*n+p}, x_i > |, 1)$$

$$\leq \sum_{i \ge i(n)} 2^{-i} = 2^{-i(n)+1}$$

whence our claim since $i(n) \rightarrow \infty$ for $n \rightarrow \infty$.

(ia) the proof is the same up to notation.

(ii) Particular case of what precedes. Now the exponential of an infinitesimal character, if it makes sense, is a character. $\hfill \Box$

3 Connes-Moscovici-Kreimer Hopf algebras

We cast a special name for a type of Hopf algebras playing a prevalent role in index theory and in renormalized quantum field theory. In most of what follows \mathbb{C} could be replaced by any commutative field **k** of characteristics 0.

Definition 3.1 H is a **Connes-Moscovici-Kreimer Hopf algebra** (for shortness a **CMK Hopf algebra**) whenever

(i) $\mathbf{H}(m = \vee, \mathbf{1} = e\mathbf{1})$ is the \mathbb{C} -algebra of (commutative) polynomials of countably many variables x_i : in other words \mathbf{H} is the symmetric algebra over the (countably-dimensional) \mathbb{C} -vector space \mathbf{V} with a specified basis $\mathbf{X} = \{x_i\}_{i \in \mathbb{N}}$:

$$\mathbf{H} = \forall \mathbf{V} = \bigoplus_{p \in \mathbb{N}} \mathbf{V}^{\vee p} \qquad (\mathbf{V}^{\vee 0} = \mathbb{C})$$
(3.1)

(we shall write indifferently $h \lor h'$ (symmetrized tensor product) or hh' (product of polynomials) for the product of elements $h, h' \in \mathbf{H}$).

The degree of monomials yields a N-grading of the algebra **H** called its **polynomial grading**. We denote the corresponding degree by deg: $\deg|_{\mathbf{V}}^{\vee p} = p$ (thus $\deg x_i = 1$).

(ii) This algebra-structure is part of a Hopf structure $\mathbf{H}(m,\mathbf{1}=e(1),\Delta,\varepsilon,S)$ such that

$$\begin{cases} \Delta x_i - x_i \otimes \mathbf{1} - \mathbf{1} \otimes x_i \in \mathbf{H} \otimes \mathbf{V} \\ \varepsilon(x_i) = 0 \end{cases}, \ i \in \mathbb{N}$$
(3.2)

(the second line implying that $\operatorname{Ker} \varepsilon = \mathbf{H}^+ \supset \bigoplus_{n \ge 1} V^{\vee n}$).

(iii) In addition \mathbf{H} is endowed with a Hopf \mathbb{N} -grading

$$\mathbf{H} = \oplus_{n \in \mathbb{N}} \mathbf{H}^n \tag{3.3}$$

The algebra **H** is also definable as the universal envelope $\mathbf{U}(\mathbf{V})$ of the Lie-algebra \mathbf{V} with trivial brackets $[x_i, x_j], i, j \in \mathbb{N}$.

An essential feature for the sequel of the polynomial grading is that **H** is generated in degrees 0 and 1. The upper index p of the pth-grade summand $\mathbf{V}^{\vee p}$ r.h. of (3.1) indicates a p-fold power of the product of **H**.

Daniel Kastler

(distinct from the (not Hopf) polynomial grading) for which all x_i (thus all monomials) are homogeneous of strictly positive degree. Denoting respectively by Y, δ_Y , and θ_z , $z \in \mathbb{C}$, the corresponding biderivation:

$$Y|_{\mathbf{H}^n} = n, \qquad n \in \mathbb{N},\tag{3.4}$$

degree:

$$\delta_Y(h) = n, \qquad h \in \mathbf{H}^n, \ n \in \mathbb{N}, \tag{3.5}$$

and automorphism group:

$$\theta_z|_{\mathbf{H}^n} = e^{nz}, \qquad n \in \mathbb{N},\tag{3.6}$$

we thus have that

$$Yx_i = \delta_Y(x_i)x_i \quad \text{with} \quad \delta_Y(x_i) \ge 1, \qquad n \in \mathbb{N}, \tag{3.7}$$

hence Y and $\theta_z = e^{Yz}$, $z \in \mathbb{C}$, leave stable each of the subspaces $V^{\vee p}$, $p \in \mathbb{N}$ (the Hopf grading is **subordinate** to the polynomial grading).

As a consequence (cf. (2.17) in Section 2), and using a previous remark, we have:

$$\oplus_{n>1} \mathbf{H}^n = \mathbf{H}^+ = \operatorname{Ker} \varepsilon = \oplus_{n>1} V^{\vee n}.$$
(3.8)

(iv) the $\mathbb N\text{-}\mathrm{graded}$ Hopf algebra $\mathbf H$ is $\mathbb N\text{-}\mathrm{graded}\text{-}\mathrm{connected}\text{:}$

$$\mathbf{H}^0 = \mathbb{C}\mathbf{1}.\tag{3.9}$$

thus progressive, i.e., such that, using a Sweedler-like notation, we have the implication:

$$h \in \mathbf{H}^+, \ h \text{ homogeneous } \Rightarrow \Delta h = h \otimes \mathbf{1} + \mathbf{1} \otimes h + h' \otimes h''$$
 (3.10)

whereby all $\delta_Y(h')$, $\delta_Y(h'')$ are strictly positive and strictly smaller than $\delta_Y(h)$.

Remark 3.2 We could replace in (ii) "Hopf algebra" by "bialgebra" since (3.10) makes the existence of the antipode automatic, cf. Proposition 2.7.

Proposition 3.3 Let **H** be a CMK Hopf algebra. The Lie algebra $\mathbf{L} = \partial Char\mathbf{H}$ is in linear bijection with the algebraic dual of **V**: a $Z \in \mathbf{L}$ is of the type:

$$Z = \begin{cases} 0 & on \quad \mathbb{C}\mathbf{1} \\ \phi & on \quad \mathbf{V} \\ 0 & on \quad \bigoplus_{n \ge 2} \mathbf{V}^{\vee n} = \mathbf{H}^{+2} \end{cases}, \ \phi \ a \ linear \ form \ on \ \mathbf{V}, \tag{3.11}$$

all elements of **L** being obtained in this way (we then write $Z \leftrightarrow \phi$).

Proof Equality $\bigoplus_{n\geq 1} \mathbf{V}^{\vee n} = \mathbf{H}^+$: Since $\bigoplus_{n\geq 1} \mathbf{V}^{\vee n} = \mathbf{H}^+$, cf. (3.8), the inclusion \supset is obvious, the inclusion \subset following from the fact that the algebra \mathbf{H} is generated in polynomial grade 0 and 1. The definition property

$$Z \in \mathbf{L} = \partial \operatorname{Char}(\mathbf{H}) \Leftrightarrow \langle Z, hh' \rangle = \langle Z, h \rangle \varepsilon(h') + \varepsilon(h) \langle Z, h \rangle,$$

 $h, h' \in \mathbf{H}$ entails for h = h' = 1 that $\langle Z, \mathbf{1} \rangle = 0$, and that $\langle Z, hh' \rangle = 0$ for $h, h' \in \mathbf{H}^+ = \operatorname{Ker} \varepsilon$. Conversely for Z as in (3.11) one has $Z \in \mathbf{L} = \partial \operatorname{Char}(\mathbf{H})$ indeed: $-\langle Z, \mathbf{11} \rangle = \langle Z, \mathbf{1} \rangle = 0$ whilst $\varepsilon(\mathbf{1}) \langle Z, \mathbf{1} \rangle + \langle Z, \mathbf{1} \rangle \varepsilon(\mathbf{1}) = 0$,

- for $h \in \mathbf{H}^1$: $\langle Z, \mathbf{1}h \rangle = \langle Z, h \rangle = \phi(h)$ whilst $\varepsilon(\mathbf{1}) \langle Z, h \rangle + \langle Z, \mathbf{1} \rangle$ $\varepsilon(h) = 1\phi(h) = \phi(h),$

The Hopf N-grading is such that $\delta_Y(h') + \delta_Y(h'') = \delta_Y(h)$, entailing merely $\delta_Y(h')$, $\delta_Y(h'') \leq \delta_Y(h'')$. Note that we have that all $h', h'' \in \mathbf{H}^+$ (cf. footnote 6 of Section 2).

- for $h, h' \in \bigoplus_{n \ge 1} \mathbf{H}^n = \mathbf{H}^+$: $hh' \in \mathbf{H}^{+2}$ hence $\langle Z, hh' \rangle = 0$ whilst $\varepsilon(h) \langle Z, h' \rangle + \langle Z, h \rangle \varepsilon(h') = 0$ since $\varepsilon(h')$ and $\varepsilon(h)$ both vanish.

The next result, a partial n-fold generalization of Proposition 3.3, is a tool for proving Theorem 3.5

Corollary 3.4 (i) Let **H** be a CMK Hopf algebra with $Z_i \leftrightarrow \phi_i \in \mathbf{L}$, i = 1, ..., n, and $x_{i_p} \in \mathbf{V}$, p = 1, ..., n, $n \ge 1$, we have that:

$$\langle Z_1 * \cdots * Z_n, x_{i_1} \dots x_{i_n} \rangle = \Sigma_\sigma \phi_{\sigma_1}(x_{i_l}) \cdots \phi_{\sigma_n}(x_{i_n})$$
(3.12)

(the sum Σ_{σ} is over the set Π_n of permutations of the *n* first integers), consequently $\mathbf{V}^{\vee n}$ is separated by the *n*-fold product $\mathbf{L} * \cdots * \mathbf{L}$: one has the implication:

$$h \in \mathbf{V}^{\vee n}, \quad \langle Z_1 * \cdots * Z_n, h \rangle = 0 \text{ for all } Z_1, \dots, Z_n \in \mathbf{L} \Rightarrow h = 0.$$
 (3.13)

(ii) We have, for $Z_1, \ldots, Z_n, n \in \mathbb{N}, n \geq 1$:

$$< Z_1 * \dots * Z_n, x_1 \dots x_m >= 0, \quad x_{i_p} \in \mathbf{V}, \ p = 1, \dots, m, \ m > n \ge 1,$$
 (3.14)

(in other terms:

$$Z_1 * \dots * Z_n |_{\mathbf{H}^{+m}} = 0, \qquad m > n \ge 1).$$
 (3.15)

(Warning: (3.14) with $1 \le m < n$ generally does not hold: for instance, for $h \in \mathbf{H}^+$ one has $\langle Z_1 * Z_2, h \rangle = \langle Z_1, h' \rangle \phi_2(h'')$.)

Proof (i) Before proving (3.12) and (3.14) for general n, we look for orientation at the cases n = 1, 2, 3, and 4.

Case n = 1: (3.12) and (3.14) amount to the second, resp. third line (3.11).

Case n = 2: We first prove

$$\begin{cases} < Z_1 * Z_2, x_{i_1} x_{i_2} > \\ = < Z_1 \otimes Z_2, \Delta(x_{i_1} x_{i_2}) > \\ = \phi_1(x_{i_1})\phi_2(x_{i_2}) + \phi_2(x_{i_1})\phi_1(x_{i_2}), \ Z_1 \leftrightarrow \phi_1, \ Z_2 \leftrightarrow \phi_2 \in \mathbf{L}. \end{cases}$$
(3.16)

In order to simultaneously prepare the next steps we compute $\Delta(\lambda_1\lambda_2) = \Delta(\lambda_1)\Delta(\lambda_2)$ for general $\lambda_1, \lambda_2 \in \mathbf{H}^+$: computation indicated by the table (with the terms of $\Delta(\lambda_1)$) in the first column, the terms of $\Delta(\lambda_2)$ in the first line, and the products at the intersection of the other lines and columns.)

Table T_2 : Computation of $\Delta(\lambda_1 \lambda_2)$

) 01

$$\lambda_2 \otimes \mathbf{1} \qquad \mathbf{1} \otimes \lambda_2 \qquad \lambda_2 \otimes \lambda_2'$$
$$\lambda_1 \otimes \mathbf{1} \qquad \$ \lambda_1 \lambda_2 \otimes \mathbf{1} \qquad \lambda_1 \otimes \lambda_2 \qquad \$ \lambda_1 \lambda_2' \otimes \lambda_2''$$
$$\mathbf{1} \otimes \lambda_2 \qquad \lambda_2 \otimes \lambda_1 \otimes \lambda_2' \qquad (1 \otimes \lambda_2) \qquad (1 \otimes \lambda_2) \otimes \lambda_1 \otimes \lambda_2''$$

$$\begin{array}{cccc} \mathbf{1} \otimes \lambda_1 & \lambda_2 \otimes \lambda_1 & \# \mathbf{1} \otimes \lambda_1 \lambda_2 & \# \lambda_2' \otimes \lambda_1 \lambda_2'' \\ \lambda_1' \otimes \lambda_1'' & \$ \lambda_1' \lambda_2 \otimes \lambda_1'' & \# \lambda_1' \otimes \lambda_1'' \lambda_2 & \# \lambda_1' \lambda_2' \otimes \lambda_1'' \lambda_2'' \end{array}$$

yields 9 tensor products:

– the 4 = 4(2 – 1) indicated by # belong to $\mathbf{H} \otimes \mathbf{H}^{+2}$, thus vanish under $\langle Z_1 \otimes Z_2, \cdot \rangle$,

- the 3 = 2 + 1 indicated by \$ belong to $\mathbf{H}^{+2} \otimes \mathbf{H}$, thus also vanish under $\langle Z_1 \otimes Z_2, \cdot \rangle$,

– the remaining 2: $\lambda_1 \otimes \lambda_2$ and $\lambda_2 \otimes \lambda_1$, yield (3.16) upon making $\lambda_1 = x_{i_1}$, $\lambda_2 = x_{i_2}$.

We next prove

$$\begin{cases} < Z_1 * Z_2, x_{i_1} \dots x_{i_m} > \\ = < Z_1 \otimes Z_2, \Delta(x_{i_1} \dots x_{i_m}) > \end{cases} = 0 \text{ if } m > 2.$$

$$(3.17)$$

Since $x_{i_1} \ldots x_{i_m}$, m > 2, is of the form $\lambda_1 \lambda_2 \lambda_3$ with $\lambda_1, \lambda_2, \lambda_3 \in \mathbf{H}^+$, we compute the relevant part of $\Delta(\lambda_1, \lambda_2, \lambda_3) = \Delta(\lambda_1, \lambda_2) \Delta \lambda_3$ whereby the #-terms of $\Delta(\lambda_1 \lambda_2)$ can be discarded since multiplications by $\Delta \lambda_3$ leave $\mathbf{H} \otimes \mathbf{H}^{+2}$ invariant: we thus have the table:

Table T_3 : Computation of $\Delta(\lambda_1 \lambda_2 \lambda_3)$

 $\lambda_3 \otimes \mathbf{1} \qquad \mathbf{1} \otimes \lambda_3 \qquad \lambda_3' \otimes \lambda_3''$

$\lambda_1\lambda_2\otimes {f 1}$	$\lambda_1\lambda_2\lambda_3\otimes 1$	$\lambda_1\lambda_2\otimes\lambda_3$	$\lambda_1\lambda_2\lambda_3'\otimes\lambda_3''$
$\lambda_1\otimes\lambda_2$	$\lambda_1\lambda_3\otimes\lambda_2$	$\#\lambda_1\otimes\lambda_2\lambda_3$	$\#\lambda_1\lambda_3'\otimes\lambda_2\lambda_3''$
$\lambda_1\lambda_2'\otimes\lambda_2''$	$\lambda_1\lambda_2'\lambda_3\otimes\lambda_2''$	$\#\lambda_1\lambda_2'\otimes\lambda_2''\lambda_3$	$\#\lambda_1\lambda_2'\lambda_3'\otimes\lambda_2''\lambda_3''$
$\lambda_2\otimes\lambda_1$	$\lambda_2\lambda_3\otimes\lambda_1$	$\#\lambda_2\otimes\lambda_1\lambda_3$	$\#\lambda_2\lambda_3'\otimes\lambda_1\lambda_3''$
$\lambda_1'\lambda_2\otimes\lambda_1''$	$\lambda_1^\prime\lambda_2\lambda_3\otimes\lambda_1^{\prime\prime}$	$\#\lambda_1'\lambda_2\otimes\lambda_1''\lambda_3$	$\#\lambda'_1\lambda_2\lambda'_3\otimes\lambda''_1\lambda''_3$

yielding 15 tensor products, all in $\mathbf{H}^{+2} \otimes \mathbf{H}$ or $\mathbf{H} \otimes \mathbf{H}^{+2}$, thus vanishing under $Z_1 \otimes Z_2$.

Case n = 3: We first prove

$$\begin{cases} < Z_1 * Z_2 * Z_3, x_{i_1} x_{i_2} x_{i_3} > \\ = < Z_1 \otimes Z_2 \otimes Z_3, (\Delta \otimes \operatorname{id}) \Delta(x_{i_1} x_{i_2} x_{i_3}) > \\ = \Sigma_{\sigma} \phi_{\sigma_1}(x_{i_1}) \phi_{\sigma_2}(x_{i_2}) \phi_{\sigma_3}(x_{i_3}), \ Z_i \leftrightarrow \phi_i \in \mathbf{L}, \ i = 1, 2, 3. \end{cases}$$
(3.18)

We need to classify the tensor products in Table T_3 : They consist of:

– the 8 = 4(3 – 1) indicated by # belong to $\mathbf{H} \otimes \mathbf{H}^{+2}$, thus vanish under $Z_1 \otimes Z_2 \otimes Z_3^{23}$,

- the 4 = 3 + 1 indicated by \$ which belong to $\mathbf{H}^{+3} \otimes \mathbf{H}$, turned by $\Delta \otimes \mathrm{id}$ into $(\Delta \mathbf{H}^{+3}) \otimes \mathbf{H}$ whose first factor vanishes under $Z_1 \otimes Z_2$,

- the remaining 3: $\lambda_{\sigma_1}, \lambda_{\sigma_2} \otimes \lambda_{\sigma_3}, \sigma$ in the set Π_3^0 of cyclic permutations of $\{1, 2, 3\}$, whose sum is turned by $\Delta \otimes \text{id}$ into $\sum_{\sigma \in \Pi_3^0} \Delta(\lambda_{\sigma_1} \lambda_{\sigma_2}) \otimes \lambda_{\sigma_3}$, by Table T_2 equal modulo $\mathbf{H} \otimes \mathbf{H}^{+2} \otimes \mathbf{H} + \mathbf{H}^{+2} \otimes \mathbf{H} \otimes \mathbf{H}$ to $\sum_{\sigma \in \Pi_3^0} (\lambda_{\sigma_1} \otimes \lambda_{\sigma_2} + \lambda_{\sigma_2} \otimes \lambda_{\sigma_1}) \otimes \lambda_{\sigma_3}$ (in other terms $\sum_{\sigma \in \Pi_\sigma} \lambda_{\sigma_1} \otimes \lambda_{\sigma_2} \otimes \lambda_{\sigma_3}$: we proved (3.18).

in fact, as we will see, these can be discarded for the whole sequel.

We next prove

$$\begin{cases} < Z_1 * Z_2 * Z_3, x_{i_1} \dots x_{i_m} > \\ = < Z_1 \otimes Z_2 \otimes Z_3, (\Delta \otimes \operatorname{id})(x_{i_1} \dots x_{i_m}) > \end{cases} = 0 \text{ if } m > 3.$$
(3.19)

Since $x_{i_1} \ldots x_{i_m}$, m > 3, is of the form $\lambda_1 \lambda_2 \lambda_3 \lambda_4$ with $\lambda_1, \lambda_2, \lambda_3, \lambda_4 \in \mathbf{H}^+$, we compute the relevant part of $\Delta(\lambda_1 \lambda_2 \lambda_3, \lambda_4) = \Delta(\lambda_1 \lambda_2 \lambda_3) \Delta \lambda_4$ whereby the #terms of $\Delta(\lambda_1 \lambda_2 \lambda_3)$ can be discarded since multiplications by $\Delta \lambda_4$ leave $\mathbf{H} \otimes \mathbf{H}^{+2}$ invariant: we thus have the table:

Table T_4 : Computation of $\Delta(\lambda_1 \lambda_2 \lambda_3 \lambda_4)$

$\lambda_1\lambda_2\lambda_3\otimes {f 1}$	$\lambda_1\lambda_2\lambda_3\lambda_4\otimes {f 1}$	$\lambda_1\lambda_2\lambda_3\otimes\lambda_4$	$\lambda_1\lambda_2\lambda_3\lambda_4'\otimes\lambda_4''$
$\lambda_1\lambda_2\otimes\lambda_3$	$\lambda_2\lambda_2\lambda_4\otimes\lambda_3$	$\#\lambda_1\lambda_2\otimes\lambda_3\lambda_4$	$\#\lambda_1\lambda_2\lambda_4'\otimes\lambda_3\lambda_4''$
$\lambda_1\lambda_2\lambda_3'\otimes\lambda_3''$	$\lambda_1\lambda_2\lambda_3^\prime\lambda_4\otimes\lambda_3^{\prime\prime}$	$\#\lambda_1\lambda_2\lambda_3'\otimes\lambda_4\lambda_3''$	$\#\lambda_1\lambda_2\lambda'_3\lambda'_4\otimes\lambda''_3\lambda''_4$
$\lambda_1\lambda_3\otimes\lambda_2$	$\lambda_1\lambda_3\lambda_4\otimes\lambda_2$	$\#\lambda_1\lambda_3\otimes\lambda_2\lambda_4$	$\#\lambda_1\lambda_3\lambda_4'\otimes\lambda_2\lambda_4''$
$\lambda_1\lambda_2'\lambda_3\otimes\lambda_2''$	$\lambda_1\lambda_2^\prime\lambda_3\lambda_4\otimes\lambda_2^{\prime\prime}$	$\#\lambda_1\lambda_2'\lambda_3\otimes\lambda_2''\lambda_4$	$\#\lambda_1\lambda_2'\lambda_3\lambda_4'\otimes\lambda_2''\lambda_4''$
$\lambda_2\lambda_3\otimes\lambda_1$	$\lambda_2\lambda_3\lambda_4\otimes\lambda_1$	$\#\lambda_2\lambda_3\otimes\lambda_1\lambda_4$	$\#\lambda_2\lambda_3\lambda_4'\otimes\lambda_1\lambda_4''$
$\lambda_1'\lambda_2\lambda_3\otimes\lambda_1''$	$\lambda_1^\prime\lambda_2\lambda_3\lambda_4\otimes\lambda_1^{\prime\prime}$	$\#\lambda_1'\lambda_2\lambda_3\otimes\lambda_1''\lambda_4$	$\#\lambda_1'\lambda_2\lambda_3\lambda_4'\otimes\lambda_1''\lambda_4''$

yielding 21 tensor products, all in $\mathbf{H}^{+3} \otimes \mathbf{H}$ or $\mathbf{H}^{+2} \otimes \mathbf{H}^{+2}$, thus vanishing under $Z_1 \otimes Z_2 \otimes Z_3$, because $\Delta \mathbf{H}^{+3}$ vanishes under $Z_1 \otimes Z_2$ by (3.17).

A more detailed analysis of this table registrates, amongst its tensor products: – the 12 = 4(4-1) indicated by # which belong to $\mathbf{H} \otimes \mathbf{H^{+}}^2$ turned by $(\Delta \otimes \mathrm{id})(\Delta \otimes \mathrm{id})$ into $\mathbf{H} \otimes \mathbf{H} \otimes \mathbf{H} \otimes \mathbf{H^{+2}}$ vanishing under $Z_1 \otimes Z_2 \otimes Z_3 \otimes Z_4^{23}$,

- the 5 = 4 + 1 indicated by \$ which belong to $\mathbf{H}^{+4} \otimes \mathbf{H}$ turned by $(\Delta \otimes \mathrm{id} \otimes \mathrm{id})(\Delta \otimes \mathrm{id})$ into $(\Delta \otimes \mathrm{id})(\Delta \mathbf{H}^{+4}) \otimes \mathbf{H}$ vanishing under $Z_1 \otimes Z_2 \otimes Z_3 \otimes Z_4$ because $(\Delta \otimes \mathrm{id})\Delta \mathbf{H}^{+4}$ vanishes under $Z_1 \otimes Z_2 \otimes Z_3$: indeed $\Delta \otimes \mathrm{id}$ acting on the tensor products in T_4 contained in $\mathbf{H}^{+3} \otimes \mathbf{H}$, and \mathbf{H}^{+3} vanishes under $Z_1 \otimes Z_2$ by (3.17).

- the remaining 4: $\lambda_{\sigma_1}\lambda_{\sigma_2}\lambda_{\sigma_3} \otimes \lambda_{\sigma_4}$, σ in the set Π_4^0 of cyclic permutations of $\{1, 2, 3, 4\}$.

We now reexamine (3.18). We have, \cong denoting equality up to negligible terms:

$$\Delta(\lambda_1\lambda_2\lambda_3)\cong \Sigma_{\tau\in\Pi^0_3}\lambda_{\tau_1}\lambda_{\tau_2}\otimes\lambda_{\tau_3},$$

hence $(\Delta \otimes \mathrm{id})\Delta(\lambda_1\lambda_2\lambda_3) \cong \sum_{\tau \in \Pi_2^0} \Delta(\lambda_{\tau_1}\lambda_{\tau_2}) \otimes \lambda_{\tau_3}$, hence

$$< Z_{1} * Z_{2} * Z_{3} *, x_{i_{1}} x_{i_{2}} x_{i_{3}} >$$

$$= < Z_{1} \otimes Z_{2} \otimes Z_{3}, (\Delta \otimes \mathrm{id}) \Delta(x_{i_{1}} x_{i_{2}} x_{i_{3}}) >$$

$$= \Sigma_{\tau \in \Pi_{3}^{0}} < Z_{1} \otimes Z_{2}, \Delta(x_{i_{\tau_{1}}} x_{i_{\tau_{2}}}) < Z_{3}, x_{i_{3}} >$$

$$= \Sigma_{\tau \in \Pi_{3}^{0}} \Sigma_{\sigma \in \Pi_{2}} < \phi_{1}, x_{i_{\tau_{\sigma_{1}}}} > < \phi_{2}, x_{i_{\tau_{\sigma_{2}}}} > < \phi_{3}, x_{\tau_{3}} >$$

$$= \Sigma_{\sigma \in \Pi_{3}} < \phi_{1}, x_{i_{\tau_{\sigma_{1}}}} > < \phi_{2}, x_{i_{\tau_{\sigma_{2}}}} > < \phi_{3}, x_{i_{\tau_{\sigma_{3}}}} >$$

$$= \Sigma_{\sigma \in \Pi_{3}} < \phi_{\sigma_{1}}, x_{i_{1}} > < \phi_{\sigma_{2}}, x_{i_{2}} > < \phi_{\sigma_{3}}, x_{i_{3}} > .$$

It should now be clear how things propagate recursively to yield a proof for general n. We assume that Table T_n is as follows: its 3[2(n-1)+1] = 3(2n-1) tensor products consist of:

- the 4(n-1) indicated by # vanishing under $(Z_1 \otimes \cdots \otimes Z_n) \circ \Delta^{(n-1)}$

- the n+1 indicated by \$ vanishing under $(Z_1 \otimes \cdots \otimes Z_n) \circ \Delta^{(n-1)}$

- the remaining *n* with sum $S_n = \sum_{\tau \in \Pi_n^0} \lambda_{\tau_1} \lambda_{\tau_2} \dots \lambda_{\tau_{(n-1)}} \otimes \lambda_{\tau_n}$, τ in the set Π_n^0 of cyclic permutations of $\{1, \dots, n\}$, with (3.18) holding for $\lambda_n = x_{i_n} \in \mathbf{V}$.

From what precedes it is clear that these features propagate from n to n + 1. In particular we have:

$$S_{n+1} = S_n(\lambda_{n+1} \otimes \mathbf{1}) + (\lambda_1 \dots \lambda_n \otimes \mathbf{1})(\mathbf{1} \otimes \lambda_{n+1})$$

= $\Sigma_{\tau \in \Pi_n^0} \lambda_{\tau_1} \lambda_{\tau_2} \lambda_{\tau_{(n-1)}} \lambda_{n+1} \otimes \lambda_{\tau_n} + (\lambda_1 \dots \lambda_n \otimes \lambda_{n+1})$
= $\Sigma_{\tau \in \Pi_{n+1}^0} \lambda_{\tau_1} \lambda_{\tau_2} \dots \lambda_{\tau_n} \otimes \lambda_{\tau_{(n+1)}}$

and

$$\langle Z_1 * \cdots * Z_{n+1}, x_{i_1} \cdots x_{i_{n+1}} \rangle$$

$$= \langle Z_1 \otimes \cdots \otimes Z_{n+1} (\Delta \otimes \operatorname{id}) \Delta^{(n-2)} (x_{i_1} \cdots x_{i_{n+1}}) \rangle$$

$$= \Sigma_{\tau \in \Pi_n^0} \langle Z_1 \otimes \cdots \otimes Z_n, \Delta^{(n-2)} (x_{i_{\tau_1}} \cdots x_{i_{\tau_n}}) \rangle \langle Z_{n+1}, x_{i_{n+1}}) \rangle$$

$$= \Sigma_{\tau \in \Pi_n^0} \Sigma_{\sigma \in \Pi_n} \langle \phi_1, x_{i_{\tau_{\sigma_1}}} \rangle \cdots \langle \phi_n, x_{i_{\tau_{\sigma_n}}} \rangle \langle \phi_{n+1}, x_{i_{(n+1)}} \rangle$$

$$= \Sigma_{\sigma \in \Pi_{n+1}} \langle \phi_1, x_{i_{\sigma_1}} \rangle \cdots \langle \phi_{\sigma_1}, x_{i_n} \rangle$$

$$= \Sigma_{\sigma \in \Pi_{n+1}} \langle \phi_{\sigma_1}, x_{i_1} \rangle \cdots \langle \phi_{\sigma_1}, x_{i_n} \rangle .$$

Our final result elucidates the structure of \mathbf{H} by displaying it as in strict Hopfalgebra-duality with the enveloping (Hopf) algebra $\mathbf{U}(\mathbf{L})$ of the Lie algebra \mathbf{L} , the latter isomorphic to the Hopf algebra \mathbf{H}_* by the Milnor-Moore theorem.

Theorem 3.5 Let **H** be a CMK Hopf algebra. **H** is separated by \mathbf{H}_* : for $h \in \mathbf{H}$, $\langle \xi, h \rangle = 0$ for all $\xi \in \mathbf{H}_*$ entails h = 0.

Proof Each $h \in \mathbf{H}$ is of the form $h = \lambda \mathbf{1} + h^{(1)} + h^{(2)} + \dots + h^{(n)}$ with $h^{(k)} \in \mathbf{V}^{\vee k}$, $k = 1, \dots, n, n \in \mathbb{N}$. We assume that h vanishes under all elements $\xi \in \mathbf{H}_*$ and show that h = 0: indeed, let $Z_i \leftrightarrow \phi_i \in \mathbf{L}, i = 1, \dots, n$: then:

- since by (3.8) all the terms of h but the first lie in $\mathbf{H}^+ = \ker \varepsilon$ one has $\langle \mathbf{1}_*, h \rangle = \varepsilon(h) = \lambda$: the requirement $\langle \mathbf{1}_*, h \rangle = 0$ thus entails the vanishing of the first term $\lambda \mathbf{1}$.

- next by Corollary 3.4(ii) one has $\langle Z_i, h \rangle = \phi_i(h^{(1)})$: asking this to vanish for all $Z_i \in \mathbf{H}_*$ thus entails the vanishing of the second term $h^{(1)}$.

– assume that we have shown that $h^{(k)}$ vanishes for $k \le p < n$ we have, by Corollary 3.4(ii) and (iii):

$$< Z_1 * \cdots Z_{p+1}, h > = < Z_1 * \cdots Z_{p+1}, h^{(p+1)} + \cdots + h^{(n)} >$$

= $< Z_1 * \cdots Z_{p+1}, h^{(p+1)} >$
= $(\phi_1 \vee \cdots \vee \phi_{p+1})(h^{(p+1)})$

whose vanishing under all $\phi_1 \lor \cdots \lor \phi_{p+1}$ entails $h^{(p+1)} = 0$: we thus proved Theorem 3.5 inductively.

4 Loops of CMK Hopf algebras

In what follows $\mathbf{H}(m, \mathbf{1} = e_1, \Delta, \varepsilon, S)$ is a CMK Hopf algebra, cf. Sections 1 and 3 of which we adopt the notation.

Definition 4.1 In what follows the ground field is \mathbb{C} and we will handle algebras with and without unit. Since our general practice was to use the word: algebra to mean: unital algebra, we shall write "algebra" (quote-unquote) to mean: not necessarily unital algebra over \mathbb{C} .

18

(i) We consider the following "algebras" of analytic functions on the Riemann sphere \mathbf{PC}^1 with north pole $N = \infty$ and south pole S = 0.

 $\begin{cases} \mathbf{A} = \{ f \in \mathbf{Holom}(\mathbb{C} - \{0\}) \text{ with } 0 \text{ a pole of finite order} \\ \mathbf{A}_{-} = \{ \text{polynomials in } \frac{1}{z} \text{ without constant term} \} \\ \mathbf{A}_{+} = \{ \text{Restrictions to } (\mathbb{C} - \{0\}) \text{ of functions in } \mathbf{Holom}(\mathbb{C}) \} \\ \mathbf{A}_{-}^{\infty} = \{ \text{Series in } \frac{1}{z} \text{ without constant term} \} \end{cases}$ (4.1)

(ii) A **loop** is a map γ : $\mathbf{PC}^1 - \{0, \infty\} \rightarrow \mathbf{G} = Char \mathbf{H}$. The set of loops is denoted by $L(\mathbf{H})$.

(iii) With **B** one of the algebras **A**, **A**₋, **A**₊, or **A**^{∞}₋, a **B**-loop is a map γ : $\mathbf{PC}^1 - \{0, \infty\} \rightarrow \mathbf{G}$ such that the functions $\mathbf{PC}^1 - \{0, \infty\} \ni z \rightarrow \langle \gamma(z), h \rangle$, $h \in \mathbf{H}^+ = \text{Ker}\varepsilon$, belongs to **B**. The set of **B**-loops is denoted by $L_{\mathbf{B}}(\mathbf{H})$.

We recall that a A-loop γ has a unique **Birkhoff decomposition**:

$$\gamma = \gamma_{-}^{-1} \gamma_{+}, \tag{4.2}$$

where

$$\begin{cases} \mathbb{C} - \{0\} \ni z \to <\gamma_{-}(z), h > \text{ belongs to } \mathbf{A}_{-} \\ \mathbb{C} - \{0\} \ni z \to <\gamma_{+}(z), h > \text{ belongs to } \mathbf{A}_{+} \end{cases}, h \in \mathbf{H}^{+}.$$

$$(4.3)$$

Lemma 4.2 (i) The B-loops build a group under the operations:

$$\begin{cases} product \ \gamma'\gamma'': (\gamma'\gamma'')(z) = \gamma'(z) * \gamma''(z), \ z \in \mathbf{PC}^1 - \{\infty\}\\ inverse \ : \gamma^{-1}(z) = \gamma(z)^{-1} = S_*\gamma(z) \end{cases}$$
(4.4)

with the unit the constant loop: $z \rightarrow < \mathbf{1}_*, \cdot > = \varepsilon$.

(ii) The **B**-loops γ are one-to-one with elements $\underline{\gamma} \in Hom \, "alg"(\mathbf{H}, \mathbf{B}')$ through the bijection:

$$\gamma \leftrightarrow \underline{\gamma} :< \gamma(z), h \ge \underline{\gamma}(h)(z), \quad h \in \mathbf{H}, \ z \in \mathbb{C} - \{0\},$$

$$(4.5)$$

a group isomorphism with the following correspondence of group products:

$$\gamma'\gamma'' \leftrightarrow \underline{\gamma} = \underline{\gamma}' * \underline{\gamma}'', \tag{4.6}$$

where the product on the left is that of the group **G** (first line (1.7)), whilst * on the right is the convolution product of Hom "alg" (**H**, **B**'):

$$\begin{cases} (\underline{\gamma}' * \underline{\gamma}'')(h) = \underline{\gamma}'(h_{(1)})\underline{\gamma}''(h_{(2)})\\ (\underline{\gamma})^{-1}(h) = \underline{\gamma}^{-1}(h) = \underline{\gamma}(Sh) \end{cases}$$
(4.7)

Notation using the chart of \mathbf{PC}^1 obtained by stereographic projection from the north pole on the equator coordinatized by z: in phrasing the Definitions (4.3) below, this chart is used to identify $\mathbf{PC}^1 - \{0, \infty\}$ with $\mathbb{C} - \{0\}$.

Definition implying that $\langle \gamma(z), \mathbf{1} \rangle = 1$ for all z.

Hom "alg" $(\mathbf{H}, \mathbf{B}')$ is the set of homomorphisms between "algebras", i.e. \mathbb{C} -linear multiplicative maps: $\mathbf{H} \rightarrow \mathbf{B}$.

Proof One has for $h, h' \in \mathbf{H}$:

$$\langle \gamma(z), hh' \rangle = \langle \gamma(z), h \rangle \cdot \langle \gamma(z), h' \rangle = \underline{\gamma}(h)(z)\underline{\gamma}(h')(z)$$

$$= [\underline{\gamma}(h)\underline{\gamma}(h')](z) = \underline{\gamma}(hh')(z),$$

$$\langle \gamma(z), \mathbf{1} \rangle = \mathbf{1} = \underline{\gamma}(\mathbf{1})(z) = \mathbf{1}_{\mathbf{B}} = \mathbf{1},$$

$$(\underline{\gamma}' * \underline{\gamma}'')(h)(z) = \langle (\gamma'\gamma'')(z), h \rangle$$

$$= \langle \gamma'(z) * \gamma''(z), h \rangle = \langle \gamma'(z) \otimes \gamma''(z), \Delta h \rangle$$

$$= \langle \gamma'(z) \otimes \gamma''(z), h_{(1)} \otimes h_{(2)} \rangle = \langle \gamma'(z), h_{(1)} \rangle \langle \gamma''(z), h_{(2)} \rangle$$

$$= \underline{\gamma}'(h_{(1)})(z)\underline{\gamma}''(h_{(2)})(z) = [\underline{\gamma}'(h_{(1)})\underline{\gamma}''(h_{(2)})](z).$$

In the sequel we are concerned with the "special loops" defined as follows:

Definition 4.3 The B-loop γ , $\mathbf{B} = \mathbf{A}_{-}$ or \mathbf{A}_{-}^{∞} , is special whenever its inverse ϕ :

$$\phi_{\varepsilon} = \gamma(\varepsilon)^{*-1}$$
 (inverse in the group **G**) (4.8)

is such that $\phi_{\varepsilon}^{*-1} * \theta_{t\varepsilon}(\phi_{\varepsilon})$, (product in the group $\mathbf{G}, t, \varepsilon \in \mathbb{R}$) has a limit for $\varepsilon \to 0$ for all t:

$$\phi_{\varepsilon}^{*-1} * \theta_{t\varepsilon}(\phi_{\varepsilon}) \xrightarrow[]{\to}]{} F_t, \tag{4.9}$$

and moreover F_t is differentiable in t for t = 0.

Remark 4.4 In the case of \mathbf{A}_{-} -loops, as soon as condition (4.9) is fulfilled, F_t is differentiable as shown by the following computation. Continuity of the limit needs only be examined for the values on the kernel of ε since $\langle \phi_{\varepsilon}^{*-1} * \theta_{t\varepsilon}(\phi_{\varepsilon}), \mathbf{1} \rangle =$ 1 is constant. Now for $h \in \mathbf{H}^+$ with $\delta_Y(h) = p$:

This makes it clear that the limit F_t is a polynomial in t if it exists.

Lemma 4.5 Let γ be a special loop as in Definition 4.3. Then F_t in (4.9) is a flow, one has:

$$F_{t+s} = F_t * F_s, (4.10)$$

thus

$$F_t = e^{*\beta t},\tag{4.11}$$

To alleviate writing we write the complex variable of the loop ϕ as an index. Note that the limit in (4.9) is taken in the topology of **G**: simple convergence on **H**.

with the β -function the following element of **L**:

$$\beta = \frac{\partial e^{*\beta t}}{\partial t} \bigg|_{t=0}.$$
(4.12)

Proof (i) Let $t, s \in \mathbb{R}$. First

$$\theta_{*t\varepsilon}[\phi_{\varepsilon}^{*}{}^{-1}\theta_{*s\varepsilon}(\phi_{\varepsilon})] \xrightarrow[\varepsilon=0]{} F_s$$

indeed, for $h \in \mathbf{H}$ of degree n: first

$$< \theta_{*t\varepsilon}[\phi_{\varepsilon}^{*-1} * \theta_{*s\varepsilon}(\phi_{\varepsilon})], h > = < [\phi_{\varepsilon}^{*-1} * \theta_{*s\varepsilon}(\phi_{\varepsilon})], \theta_{*t\varepsilon}h >$$
$$= e^{int\varepsilon} < \phi_{\varepsilon}^{*-1} * \theta_{*s\varepsilon}(\phi_{\varepsilon}), h > \to F_{s}$$

since $e^{int\varepsilon} \to 1$. Then:

$$F_{t+s} = \lim_{\varepsilon = 0} [\phi_{\varepsilon}^{*-1} * \theta_{*(t+s)\varepsilon}(\phi_{\varepsilon})]$$

$$= \lim_{\varepsilon = 0} [\phi_{\varepsilon}^{*-1} * \theta_{*t\varepsilon} [\theta_{*s\varepsilon}(\phi_{\varepsilon})]]$$

$$= \lim_{\varepsilon = 0} [\phi_{\varepsilon}^{*-1} * \theta_{*t\varepsilon} \{\phi_{\varepsilon} * \phi_{\varepsilon}^{*-1} * \theta_{*s\varepsilon}(\phi_{\varepsilon})\}]$$

$$= \lim_{\varepsilon = 0} [\phi_{\varepsilon}^{*-1} * \theta_{*t\varepsilon} \phi_{\varepsilon} * \theta_{*t\varepsilon} [\phi_{\varepsilon}^{*-1} * \theta_{*s\varepsilon}(\phi_{\varepsilon})]]$$

$$= \{\lim_{\varepsilon = 0} \phi_{\varepsilon}^{*-1} * \theta_{*t\varepsilon} \phi_{\varepsilon}\} * \{\lim_{\varepsilon = 0} \theta_{*t\varepsilon} [\phi_{\varepsilon}^{*-1} \theta_{*s\varepsilon}(\phi_{\varepsilon})]]$$

$$= F_t * F_s.$$

Lemma 4.6 Let γ be a special \mathbf{A}^{∞}_{-} -loop with:

$$\phi_{\varepsilon} = \mathbf{1}_{*} + \Sigma_{n \ge 1} \frac{d_{n}}{\varepsilon^{n}} \quad \text{with } d_{n} \in \mathbf{H}^{*+}, \text{ i.e., } < d_{n}, \mathbf{1} >= 0, \ n \ge 1$$
(4.13)

(we set $d_1 = Res\phi$, called the residue of ϕ). One has:

$$\begin{cases} Y_*d_1 = Y_*(Res\phi) = \beta \\ Y_*d_{n+1} = d_n * \beta, \ n > 1 \end{cases},$$
(4.14)

rewritten

$$\begin{cases} d_1 = \underline{Y}_*^{-1}\beta \\ d_{n+1} = \underline{Y}_*^{-1}d_n * \beta, \ n > 1 \end{cases}, \\ \underline{Y}_* \text{ the injective restriction of } Y_* \text{ to } \mathbf{H}^{*+} = Ker\varepsilon_*, \quad (4.15)$$

explicitly:

$$d_n = \int_{s_1 \ge s_2 \ge \dots \ge s_n \ge 0} \theta_{*-s_1}(\beta) * \theta_{*-s_2}(\beta) * \dots * \theta_{*-s_n}(\beta) \ ds_1 \dots ds_n \qquad (4.16)$$

(using

$$\underline{Y}_{*}^{-1} = \int_{0}^{\infty} \theta_{*-s} \, ds \text{ in restriction to } \mathbf{H}^{*+}). \tag{4.17}$$

The morale of this is that the loop ϕ is determined by its residue, thus by β , with dependence expressed explicitly by (4.16).

terminology borrowed from renormalization theory.

Daniel Kastler

Proof Performing a licit exchange of limits in:

we first show that we have

$$<\beta, h>=\lim_{\varepsilon=0}\varepsilon<\phi_{\varepsilon}^{*-1}*Y_{*}(\phi_{\varepsilon}), h>, \quad h\in\mathbf{H},$$

$$(4.19)$$

Indeed, the horizontal limit comes from (4.9) with $F_t = e^{*\beta t}$; the right vertical limit is (4.12). As for the left vertical limit, we have that $\frac{\partial}{\partial t}|_{t=0}$ applied to:

yields

$$\langle \phi_{\varepsilon}, Sh_{(1)} \rangle \langle \phi_{\varepsilon}, Yh_{(2)} \rangle = \varepsilon \langle S_{*}\phi_{\varepsilon}, h_{(1)} \rangle \langle Y\phi_{\varepsilon}, Yh_{(2)} \rangle$$
$$= e \langle \phi_{\varepsilon}^{*-1} \otimes (Y_{*}\phi_{\varepsilon}), \Delta h \rangle$$
$$= \varepsilon \langle \phi_{\varepsilon}^{*-1} * (Y_{*}\phi_{\varepsilon}), h \rangle.$$

Next the function $\{\varepsilon \to \varepsilon < \phi_{\varepsilon}^{*-1} * Y_*(\phi_{\varepsilon}), h >\}$ is holomorphic on the whole Riemann sphere, thus must be constant: we have $\varepsilon < \phi_{\varepsilon}^{*-1} * Y_*(\phi_{\varepsilon}), h > = < \beta, h >, \phi_{\varepsilon}^{*-1} * Y_*(\phi_{\varepsilon}) = \frac{1}{\varepsilon}\beta$, i.e.:

$$Y_*\phi_\varepsilon = \phi_\varepsilon * \frac{1}{\varepsilon}\beta. \tag{4.20}$$

Feeding (4.13) into this yields:

$$\sum_{n\geq 1} \frac{1}{\varepsilon^n} Y_* d_n = \frac{1}{\varepsilon} \mathbf{1} * \beta + \sum_{n\geq 1} \frac{d_n}{\varepsilon^{n+1}} * \beta$$
$$= \frac{1}{\varepsilon} \beta + \sum_{n\geq 2} \frac{d_{n-1}}{\varepsilon^n} * \beta,$$

equating coefficients of $\frac{1}{\varepsilon^n}$ then yields (4.14). Passage from (4.14) to (4.15): Y_* a is injective in restriction to the augmentation ideal $\mathbf{H}^{*+} = \bigoplus_{n>1} \mathbf{H}^{*n}$. Thus β (thus also $\operatorname{Res}\phi$) determines F, the key to the explicit dependence being formula (4.17), itself obtained as follows: integrating from 0 to ∞ ,

$$\frac{\partial \theta_{*-s}}{\partial t} = \frac{\partial e^{*-tY_*}}{\partial t} = -Y_*e^{*-tY_*} = -Y_*\theta_{*-s}$$

yields $-id_{\mathbf{H}^*} = Y_* \int_0^\infty \theta_{*-s} \, ds$, it suffices now to inverse Y_* in restriction to \mathbf{H}^{*+} . Applying (4.17) to β yields (4.16) for n = 1. Inductive iteration then yields (4.16):

assume it holds for n: applying \underline{Y}^{-1} on both sides yields:

$$d_{n+1} = \underline{Y}^{-1}d_n * \beta$$

$$= \int_0^\infty \theta_{*-s} \left[\int_{s_1 \ge s_2 \ge \cdots \ge s_n \ge 0} \theta_{*-s_1}(\beta) * \theta_{*-s_2}(\beta) \\ * \cdots * \theta_{*-s_n}(\beta) \ ds_1 \dots ds_n \right] ds$$

$$= \int_0^\infty \left[\int_{s_1 \ge s_2 \ge \cdots \ge s_n \ge 0} \theta_{*-(s+s_1)}(\beta) * \theta_{*-(s+s_2)}(\beta) \\ * \cdots * \theta_{*-(s+s_n)}(\beta) \ ds_1 \dots ds_n \right] ds$$

$$= \int_{s \ge s_1 \ge s_2 \ge \cdots \ge s_n \ge 0} \theta_{*-s}(\beta) * \theta_{*-s_1}(\beta) * \theta_{*-s_2}(\beta) \\ * \cdots * \theta_{*-s_n}(\beta) \ dsds_1 \dots ds_n$$

up to numbering of variables identical to (4.16).

Extension of the Lie algebra \mathbf{L} by an additional element Z_0 enables us now to give a compact expression of the last result, and, as a byproduct, a characterization of "specialty" for \mathbf{A}_{-}^{∞} -loops.

Definition-Lemma 4.7 (i) $\tilde{\mathbf{L}}$ is $\mathbf{L} \oplus \mathbb{C}\{Z_0\}$ with the Lie-brackets of \mathbf{L} plus:

$$[Z_0, Z] = -[Z, Z_0] = Y_*(Z), \qquad Z \in \mathbf{L}.$$
(4.21)

(ii) The so defined $\tilde{\mathbf{L}}$ is a Lie algebra.

Proof We need only check the additional cases of the Jacobi requirement. Now, for $Z, Z' \in \mathbf{L}$, we have

$$[Z_0, [Z_0, Z]] + [Z_0, [Z, Z_0]] + [Z, [Z_0, Z_0]] = 0,$$

and

 ϕ

$$[Z_0, [Z, Z']] + [Z, [Z', Z_0]] + [Z'[Z_0, Z]] = Y_*([Z, Z']) - [Z, Y_*(Z')] + [Z', Y_*(Z)] = 0$$

by the fact that the algebra-derivation Y_* of \mathbf{H}^* is a Lie algebra-derivation of the Lie algebra Lie \mathbf{H}^* : for $\xi, \eta \in \mathbf{H}^*$ we have $Y_*(\xi\eta) = (Y_*\xi)\eta + \xi(\eta Y_*)$ and $Y_*(\eta\xi) = (Y_*\eta)\xi + \eta(\xi Y_*)$ thus $Y_*[\xi,\eta] = [Y_*\xi,\eta] + [\xi,\eta Y_*]$.

Theorem 4.8 With $\tilde{\mathbf{L}}$ as in Definition-Lemma 4.7, one has for each \mathbf{A}_{-}^{∞} -loop

$$\phi_{\varepsilon} = \lim_{t \to \infty} e^{*-tZ_0} * e^{*-t(\beta/\varepsilon + Z_0)} \qquad \text{where } \beta = Y(\text{Res}\phi). \tag{4.22}$$

Proof We apply the expansional formula

$$e^{*A+B} = \sum_{n \in \mathbb{N}} \int_{\Sigma u_i = 1, u_i > 0} e^{*u_0 A} * B * e^{*u_1 A} * \dots * B * e^{*u_n A} du_1 \dots du_n \quad (4.23)$$

to $A = tZ_0, B = t\beta, t > 0$, yielding: $e^{*t(\beta+Z_0)}$ $= \sum_{n \in \mathbb{N}} \int_{\Sigma u_i = 1, u_i > 0} e^{*u_0 tZ_0} * \beta * e^{*u_1 tZ_0} * \cdots$ $*e^{*u_{n-1} tZ_0} * \beta * e^{*u_n tZ_0} t^n du_1 \cdots du_n$ $(v_i = tu_i, i = 1, \dots, n)$ $= \sum_{n \in \mathbb{N}} \int_{\Sigma v_i = t, v_i > 0} e^{*v_0 Z_0} * \beta * e^{*v_1 Z_0} * \cdots * \beta * e^{*v_n Z_0} dv_0 dv_1 \cdots dv_n$ $(v_0 = t - s_1, v_1 = s_1 - s_2, \dots, v_{n-1} = s_{n-1} - s_n, v_n = s_n, \beta \to \frac{1}{\varepsilon}\beta)$ $e^{*-t(\beta/\varepsilon + Z_0)}$ $= \frac{1}{\varepsilon^n} \sum_{n \in \mathbb{N}} \int_{t \ge s_1 \ge s_2 \ge \cdots \ge s_n \ge 0} e^{*(t-s_1)Z_0} * \beta * e^{*(s_1 - s_2)Z_0} * \cdots$

$$= e^{*tZ_0} \sum_{n \in \mathbb{N}} \frac{1}{\varepsilon^n} \int_{t \ge s_1 \ge s_2 \ge \dots \ge s_n \ge 0} \theta_{*-s_1}(\beta) * \theta_{*-s_2}(\beta) * \dots \\ * \theta_{*-s_n}(\beta) dv_0 dv_1 \dots dv_n$$

We thus have

$$e^{*-tZ_0} * e^{*-t(\beta/\varepsilon+Z_0)}$$

$$= \sum_{n\in\mathbb{N}} \frac{1}{\varepsilon^n} \int_{t\geq s_1\geq s_2\geq\cdots\geq s_n\geq 0} \theta_{*-s_1}(\beta) * \theta_{*-s_1}(\beta) *\cdots$$

$$*\theta_{*-s_n}(\beta) dv_0 dv_1 \cdots, dv_n,$$

whence (4.22)

$$\lim_{t=\infty} e^{*-tZ_0} * e^{*-t(\beta/\varepsilon+Z_0)}$$

$$= \sum_{n\in\mathbb{N}} \frac{1}{\varepsilon^n} \int_{s_1 \ge s_2 \ge \dots \ge s_n \ge 0} \theta_{*-s_1}(\beta) * \theta_{*-s_1}(\beta) * \dots * \theta_{*-s_n}(\beta) dv_0 dv_1 \dots dv_n$$

$$= \sum_{n\in\mathbb{N}} \frac{1}{\varepsilon^n} d_n = \phi_{\varepsilon}.$$

5 Algebraïc Birkhoff decomposition

Definition 5.1 A **Birkhoff sum** (over **k**) is a commutative (unital) **k**-algebra \mathbb{A} which is the direct sum $\mathbb{A} = \mathbb{A}_{-} \oplus \mathbb{A}_{+}$ of two **k**-linear multiplicative subspaces \mathbb{A}_{-} and \mathbb{A}_{+} . The projection $T : \mathbb{A} \to \mathbb{A}_{-}$ parallel to \mathbb{A}_{+} thus fulfills:

$$T(ab) + (Ta)(Tb) = T[(Ta)b + a(Tb)], \quad a, b \in \mathbb{A}.$$
 (5.1)

We do not say: subalgebras because according to our general terminological practice a subalgebra contains the unit. Note that Definition (5.1) follows from the fact that both \mathbb{A}_{-} and A_{+} are closed under multiplications.

Proposition 5.2 (Algebraic Birkhoff decomposition). With $\mathbf{H} \ a \ \mathbb{N}$ -graded, connected, progressive, commutative Hopf algebra over \mathbf{k} , and $\mathbb{A} = \mathbb{A}_{-} \oplus \mathbb{A}_{+}$ a Birkhoff sum, denote by $Hom_{\mathbf{k}-alg}(\mathbf{H}, \mathbb{A})$ the set of unital \mathbf{k} -algebra homomorphisms: $\mathbf{H} \rightarrow \mathbb{A}$, considered as a \mathbf{k} -algebra under the convolution product:

$$(\phi\phi')(h) = m_{\mathbb{A}}(\phi \otimes \phi')(\Delta h), \quad \phi, \phi' \in Hom_{\mathbf{k}-alg}(\mathbf{H}, \mathbb{A}), \ h \in H.$$
(5.2)

Let $\phi \in Hom_{\mathbf{k}-alg}(\mathbf{H}, \mathbb{A})$ be given. Requiring, for a \mathbf{k} -linear $\phi_{-} : \mathbf{H} \to \mathbb{A}$:

$$\phi_{-}(\mathbf{1}) = \mathbf{1}_{\mathbb{A}} \tag{5.3}$$

$$\phi_{-}(h) = -T[\phi(h) + \phi_{-}(h')\phi(h'')], \qquad (5.4)$$

 $h \in \mathbf{H}^+ = Ker\varepsilon$, specifies inductively a $\phi_- \in Hom_{\mathbf{k}-alg}(\mathbf{H}, \mathbb{A})$ having its range in \mathbb{A}_- :

$$\phi_{-}(h) = T\phi_{-}(h), \tag{5.5}$$

such that the product $\phi_+ = \phi_- \phi \in \operatorname{Hom}_{\mathbf{k}-alg}(\mathbf{H}, \mathbb{A})$ has its range in \mathbb{A}_+ :

$$\phi_{+}(h) = (id_{\mathbb{A}} - T)[\phi(h) + \phi_{-}(h')\phi(h'')], \qquad h \in \mathbf{H}^{p},$$
(5.6)

this yielding the "algebraic Birkhoff decomposition":

$$\phi = \phi_{-}^{-1} \phi_{+}. \tag{5.7}$$

Proof Progressiveness of **H** implies that (5.3) and (5.4) specifies inductively a **k**-linear map ϕ_{-} : $\mathbf{H} \rightarrow \mathbb{A}$.

Check of (5.5): obvious from (5.4) and $T^2 = T$.

Check of (5.6): we have, by (5.2) and (5.3), (5.4), for $h \in \mathbf{H}$, omitting parentheses:

$$\begin{aligned} (\phi_{-}\phi')(h) &= m_{\mathbb{A}}(\phi_{-}\otimes\phi)(\Delta h) \\ &= m_{\mathbb{A}}(\phi_{-}\otimes\phi)(h\otimes\mathbf{1}+\mathbf{1}\otimes h+h'\otimes h'') \\ &= \phi_{-}h+\phi h+\phi_{-}h'\cdot\phi h'', \end{aligned}$$

$$T[(\phi_{-}\phi')(h)] = \phi_{-}h + T[\phi_{-}h + \phi_{h} + \phi_{-}h' \cdot \phi_{h}h'] = 0,$$

whence (5.6) owing to $T^2 = T$.

What is not obvious from (5.4) is that ϕ_{-} is multiplicative. We prove this by induction: for $x \in \mathbf{H}^p$, $y \in \mathbf{H}^p$ we have, still omitting parentheses:

$$\begin{array}{lll} \Delta xy &=& xy \otimes \mathbf{1} + \mathbf{1} \otimes xy + x \otimes y + y \otimes x + x'y \otimes x'' + x' \otimes x''y + xy' \otimes y'' \\ && +y' \otimes xy'' + x'y' \otimes x''y'' \end{array}$$

i.e., **H** has a \mathbb{N} -grading commuting with the Hopf structure, such that $\mathbf{H}_0 = \mathbb{C}\mathbf{1}$, and moreover such that $\Delta h = h \otimes \mathbf{1} + \mathbf{1} \otimes h + h' \otimes h''$ with max(degh',degh'') $\langle p, h \in H^p, p \rangle 0$.

The multiplicativity of ϕ and ϕ' implies that of $\phi\phi'$ owing to the commutativity of \mathbb{A} . The (commutative) product $\phi_{-}(h')\phi(h'')$ r.h.s. is in \mathbb{A} .

The product and the inverse r.h.s. are taken w.r.t. the convolution product (5.2).

hence, by the induction hypothesis:

$$\begin{split} \phi_{-}(xy) &= -T[\phi xy + \phi_{-}x \cdot \phi y + \phi_{-}y \cdot \phi x + \phi_{-}x'y \cdot \phi x'' + \phi_{-}x' \cdot \phi x''y \\ &+ \phi_{-}xy' \cdot \phi y'' + \phi_{-}y' \cdot \phi xy'' + \phi_{-}x'y' \cdot \phi x''y''] \\ 1 & 2 & 3 & 4 & 5 \\ &= -T[\phi x \cdot \phi y + \phi_{-}x \cdot \phi y + \phi_{-}y \cdot \phi x + \phi_{-}y \cdot \phi_{-}x' \cdot \phi x'' + \phi y \cdot \phi_{-}x' \cdot \phi x'' \\ & 6 & 7 & 8 \\ &+ \phi_{-}x \cdot \phi_{-}y' \phi y'' + \phi x \cdot \phi_{-}y' \cdot \phi y'' + \phi_{-}x' \cdot \phi_{-}y' \cdot \phi x'' \cdot y'']. \end{split}$$

On the other hand we have, using (5.1) and the commutativity of A:

$$\begin{split} \phi_{-}x \cdot \phi_{-}y &= TA \cdot TB = T[ATB + BTA - AB] = -T[A\phi_{-}y + B\phi_{-}x + AB] \\ &= -T[(\phi x + \phi_{-}x' \cdot \phi x'')\phi_{-}y + (\phi y + \phi_{-}y' \cdot \phi y'')\phi_{-}x \\ &+ (\phi x + \phi_{-}x' \cdot \phi x'')(\phi y + \phi_{-}y' \cdot \phi y'')] \\ &3 & 4 & 2 & 6 \\ &= -T[\phi_{-}y \cdot \phi x + \phi_{-}y \cdot \phi_{-}x' \cdot \phi x'' + \phi_{-}x \cdot \phi_{-}y' \cdot \phi y'' \\ &1 & 7 & 5 \\ &+ \phi x \cdot \phi y + \phi x \cdot \phi_{-}y' \cdot \phi y'' + \phi y \cdot \phi_{-}x' \cdot \phi x'' \\ &8 \\ &+ \phi_{-}x' \cdot \phi x'' \cdot \phi x'' \cdot \phi y'']. \end{split}$$

the two expressions containing the same terms.

Appendix A Algebras, coalgebras, bialgebras, Hopf algebras

In what follows ${\bf k}$ is a commutative field of characteristics zero (e.g. $\mathbb{C},\,\mathbb{R},\, {\rm or}\,\,\mathbb{Q}).$

Definitions and notation A.1 (Algebras, coalgebras, bialgebras, Hopf algebras). Let **H** be a **k**-vector space: with the following notation for the maps $m, \Delta, e, \varepsilon, S$:

Notation:

$$\begin{array}{ll} m: \mathbf{H} \otimes \mathbf{H} \to \mathbf{H} & \Delta: \mathbf{H} \to \mathbf{H} \otimes \mathbf{H} & e: \mathbf{k} \to \mathbf{H} & \varepsilon: \mathbf{H} \to \mathbf{k} & S: \mathbf{H} \to \mathbf{H} \\ \text{multiplication} & \text{comultiplication} & \text{unit} & \text{counit} & \text{antipode} \\ & \Delta a = a_{(1)} \otimes a_{(2)} & e(1) = \mathbf{1} \end{array}$$

the following axioms successively define ${\bf H}$ as an algebra, coalgebra, bialgebra, and Hopf algebra:

Axioms:

Algebra

$$(Am) \quad m(\mathrm{id}_{\mathbf{H}} \otimes m) = m(m \otimes \mathrm{id}_{\mathbf{H}}) \qquad (\mathbf{H} \otimes \mathbf{H} \to \mathbf{H}) \qquad a(bc) = (ab)c$$

$$(Ae) \quad \left\{ \begin{array}{l} m(\mathrm{id}_{\mathbf{H}} \otimes e) = \mathrm{id}_{\mathbf{H}} \\ m(e \otimes \mathrm{id}_{\mathbf{H}}) = \end{array} \qquad \left(\left\{ \begin{array}{l} \mathbf{H} \otimes \mathbf{k} \cong \mathbf{H} \to \mathbf{H} \\ \mathbf{k} \otimes \mathbf{H} \cong \mathbf{H} \to \mathbf{H} \end{array} \right) \quad \left\{ \begin{array}{l} a(\mathbf{1}k) = ka \\ (\mathbf{1}k)a = \end{array} \right. \right.$$

Coalgebra

$$\begin{array}{ll} (C\Delta) & \Delta^{(2)} = (\mathrm{id}_{\mathbf{H}} \otimes \Delta)\Delta & (\mathbf{H} \to \mathbf{H} \otimes \mathbf{H}) \\ & = (\Delta \otimes \mathrm{id}_{\mathbf{H}})\Delta & = a_{(1)}(3) \otimes a_{(2)}(3) \otimes a_{(2)}(4) \\ & = a_{(1)}(3) \otimes a_{(1)}(4) \otimes a_{(2)} \end{array}$$

$$(C\varepsilon) \quad \left\{ \begin{array}{ll} (\mathrm{id}_{\mathbf{H}} \otimes \varepsilon) \Delta = \mathrm{id}_{\mathbf{H}} \\ (\varepsilon \otimes \mathrm{id}_{\mathbf{H}}) \Delta = \end{array} \quad \left(\left\{ \begin{array}{ll} \mathbf{H} \to \mathbf{H} \otimes \mathbf{k} \cong \mathbf{H} \\ \mathbf{H} \to \mathbf{k} \otimes \mathbf{H} \cong \mathbf{H} \end{array} \right) \quad \left\{ \begin{array}{ll} a_{(1)} \varepsilon(a_{(2)}) = a \\ \varepsilon(a_{(1)})a_{(2)} = a \end{array} \right. \right.$$

Bialgebra: add to the previous axioms:

$$(Bm) \quad \Delta m = m_{\otimes}(\Delta \otimes \Delta) \qquad (\mathbf{H} \otimes \mathbf{H} \to \mathbf{H} \otimes \mathbf{H}) \qquad \Delta(ab) = (\Delta a)(\Delta b)$$
$$(Be) \qquad \Delta e = e \otimes e \qquad (\mathbf{k} \cong \mathbf{k} \otimes \mathbf{k} \to \mathbf{H} \otimes \mathbf{H}) \qquad \Delta(k\mathbf{1} = k\mathbf{1} \otimes \mathbf{1})$$

$$(B\varepsilon) \qquad \varepsilon m = \varepsilon \otimes \varepsilon \qquad (\mathbf{H} \otimes \mathbf{H} \to \mathbf{k} \cong \mathbf{k} \otimes \mathbf{k}) \qquad \varepsilon(ab) = (\varepsilon a)(\varepsilon b)$$

$$(B\varepsilon e) \qquad \varepsilon e = \mathrm{id}_{\mathbf{k}} \qquad (\mathbf{k} \to \mathbf{k}) \qquad \varepsilon [e(k)] = k$$

Hopf algebra: add to the previous axioms:

$$\begin{array}{ll} (H) & p = m(\mathrm{id}_{\mathbf{H}} \otimes S)\Delta = m(S \otimes \mathrm{id}_{\mathbf{H}})\Delta & (\mathbf{H} \to \mathbf{H}) & a_{(1)}(Sa_{(2)}) = (Sa_{(1)})a_{(2)} \\ & = e\varepsilon & = (\varepsilon a)\mathbf{1} \\ S * \mathrm{id} = \mathrm{id} * S = e\varepsilon \end{array}$$

Proposition A.2 Let $\mathbf{H}(\cdot, \Delta, e, \varepsilon, S)$ be a Hopf algebra over \mathbf{k} . One has the following properties:

$$\begin{array}{ll} (i) & (Hm) & Sm = mP_{12}(S \otimes S) & i.e. \quad S(ab) = (Sb)(Sa), \ a,b, \in H, \\ (ii) & (H\Delta) & \Delta S = (S \otimes S)P_{12}\Delta & i.e. \quad \Delta Sa = Sa_{(2)} \otimes Sa_{(1)}, \ a \in H, \\ (iii) & \left\{ \begin{array}{ll} (He) & S\mathbf{1} = \mathbf{1} \\ (H\varepsilon) & \varepsilon S = \varepsilon \\ (Hp) & Sp = pS = p \end{array} \right. .$$

Proof of Proposition A.2 (i) resp. (ii) see below B.3 resp. B.4. (iii) (He): by (H), (Be) and $(B\varepsilon e)$:

$$m(S \otimes \operatorname{id}_{\mathbf{H}})(\Delta \mathbf{1} = \mathbf{1} \otimes \mathbf{1}) = S\mathbf{1} = (e\varepsilon)\mathbf{1} = (\varepsilon\mathbf{1})\mathbf{1} = \mathbf{1}.$$

(Hp): for $a \in \mathbf{H}$:

$$Spa = S[(\varepsilon a)\mathbf{1}] = (\varepsilon a)S\mathbf{1} = (\varepsilon a)\mathbf{1} = pa,$$

moreover, by (H), $(H\Delta)$, (Hm):

$$pS = m(S \otimes \mathrm{id}_{\mathbf{H}})\Delta S = m(S \otimes \mathrm{id}_{\mathbf{H}})(S \otimes S)P_{12}\Delta$$
$$= mP_{12}(S \otimes S)(\mathrm{id}_{\mathbf{H}} \otimes S)\Delta$$
$$= Sm(\mathrm{id}_{\mathbf{H}} \otimes S)\Delta = Sp.$$

 $(H\varepsilon)$: by (Hp) for $a \in \mathbf{H}$:

$$pSa = \varepsilon(Sa)\mathbf{1} = pa = \varepsilon(a)\mathbf{1},$$

hence $\varepsilon(Sa) = \varepsilon(a)$.

Proposition A.3 (The short exact sequence of ε .) (i) The short exact sequence of ε is split with lift e:

$$0 \to \mathbf{H}^{+} = Ker\varepsilon \ \hookrightarrow \mathbf{H} \ \stackrel{\varepsilon}{\underset{e}{\to}} \mathbf{k} \to 0 \tag{A.1}$$

(where \mathbf{H}^+ is a notation for $Ker\varepsilon$) corresponding to the following direct sum splitting into a "vertical" and a "horizontal" part:

$$\mathbf{H} = \mathbf{H}^+ \oplus \mathbf{k}\mathbf{1}.\tag{A.2}$$

The splitting of short exact sequence (1,1) can alternatively be specified by:

- the "horizontal" projection $p = e\varepsilon$

- the "vertical" projection $p^+ = id_{\mathbf{H}} - p = id_{\mathbf{H}} - e\varepsilon$ (with $Imp_+ = \mathbf{H}^+ = Ker\varepsilon$).

Proof of Proposition A.3 (i) one has $\varepsilon e = id_k$, indeed, for $\lambda \in k$:

$$\varepsilon e \lambda = \varepsilon(\lambda \mathbf{1}) = \lambda \varepsilon(\mathbf{1}) = \lambda.$$

The other claims follow from known facts about split short exact sequences. $\hfill \square$

Appendix B Convolution

In what follows the ground field is \mathbf{k} (with possible choices $\mathbb{C}, \mathbb{R}, \mathbb{Q}$). We write \otimes , Hom, and End for $\otimes_{\mathbf{k}}$, Hom_{**k**}, and End_{**k**}.

Definition-Proposition B.1 With $\mathbf{C}(\Delta, \varepsilon)$ a coalgebra, $\mathbb{A}(m, e)$ an algebra; and $Hom(\mathbf{C}, \mathbb{A})$ the vector space of homomomorphisms: $\mathbf{C} \to \mathbb{A}$ (as **k**-vector spaces), define as follows $\Phi * \Phi' \in Hom(\mathbf{C}, \mathbb{A})$:

$$\Phi * \Phi' = m(\Phi \otimes \Phi')\Delta, \qquad \Phi, \Phi' \in \operatorname{Hom}(\mathbf{C}, \mathbb{A}). \tag{B.1}$$

Under this product called **convolution** and with $e\varepsilon$ as a unit, $Hom(\mathbf{C}, \mathbb{A})$ becomes an algebra. Specifically one has :

$$\Phi * (\Phi' * \Phi'') = (\Phi * \Phi') * \Phi'' = m^{(2)} (\Phi \otimes \Phi' \otimes \Phi'') \Delta^{(2)},$$

$$\Phi, \Phi', \Phi'' \in Hom(\mathbf{C}, \mathbb{A})$$
(B.2)

(more generally

$$\Phi_1 * \cdots * \Phi_{n+1} = m^{(n)} (\Phi_1 \otimes \cdots \otimes \Phi_{n+1}) \Delta^{(n)},$$

$$\Phi_1, \dots, \Phi_{n+1} \in Hom(\mathbf{C}, \mathbb{A}), \ n \in \mathbb{N}),$$
(B.3)

in Sweedlers' notation:

$$(\Phi * \Phi')a = (\Phi a_{(1)})(\Phi' a_{(2)}), \quad \Phi, \Phi' \in \operatorname{Hom}(\mathbf{C}, \mathbb{A}), \ a \in \mathbb{A},$$
(B.4)

(more generally

$$(\Phi_1 * \dots * \Phi_{n+1}'')a = m^{(n)}(\Phi_1 a_{(1)}) \dots (\Phi_{n+1} a_{(n+1)}),$$

$$1, \dots, \Phi_{n+1} \in Hom(\mathbf{C}, \mathbb{A}), \ a \in \mathbb{A}),$$
(B.5)

finally

Φ

$$\Phi * e\varepsilon = e\varepsilon * \Phi = \Phi, \qquad \Phi \in \operatorname{Hom}(\mathbf{C}, \mathbb{A}), \ a \in \mathbb{A}.$$
(B.6)

Proof We have $\Phi * \Phi' \in \text{Hom}(\mathbf{C}, \mathbb{A})$ because $\Delta \in \text{Hom}(\mathbf{C}, \mathbf{C} \otimes \mathbf{C})$, $\Phi \otimes \Phi' \in \text{Hom}(\mathbf{C} \otimes \mathbf{C}, \mathbf{C} \otimes \mathbf{C})$, $m \in \text{Hom}(\mathbf{C} \otimes \mathbf{C}, \mathbf{C})$. Check of (B.2), we have:

$$(\Phi * \Phi') * \Phi'' = m(m((\Phi \otimes \Phi')\Delta) \otimes \Phi'')\Delta = m(m \otimes \mathrm{id})((\Phi \otimes \Phi') \otimes \Phi'')(\Delta \otimes \mathrm{id})\Delta = m^{(2)}(\Phi \otimes \Phi' \otimes \Phi'')\Delta^{(2)},$$

whilst

$$\begin{aligned} \Phi * (\Phi' * \Phi'') &= m(\Phi \otimes (m(\Phi \otimes \Phi')\Delta)\Delta \\ &= m(\mathrm{id} \otimes m)(\Phi \otimes (\Phi' \otimes \Phi''))(\mathrm{id} \otimes \Delta)\Delta. \end{aligned}$$

Check of (B.4), we have:

$$\begin{aligned} (\Phi * \Phi')a &= m(\Phi \otimes \Phi')\Delta a = m(\Phi \otimes \Phi')(a_{(1)} \otimes a_{(2)}) \\ &= m[(\Phi a_{(1)}) \otimes (\Phi' a_{(2)})] \\ &= (\Phi a_{(1)}) \otimes (\Phi' a_{(2)}). \end{aligned}$$

Check of (B.6),

$$(\Phi * e\varepsilon)a = (\Phi'a_{(1)})(\varepsilon(a_{(2)}))\mathbf{1} = \Phi[a_{(1)}\varepsilon(a_{(2)})]\mathbf{1} = (\Phi a)\mathbf{1} = \Phi a_{(2)}$$

and

(

$$e\varepsilon * \Phi)a = (\varepsilon(a_{(1)})\mathbf{1})(\Phi a_{(2)}) = \Phi[\varepsilon(a_{(1)})a_{(2)}]\mathbf{1} = (\Phi a)\mathbf{1} = \Phi a, \quad a \in \mathbb{A}.$$

Example B.2 (i) Result B.1 applies in particular to the case where $\mathbf{C} = \mathbb{A}(m, e, \Delta, \varepsilon)$ is a bialgebra (with unit element **1**). Hom $(\mathbb{A}, \mathbb{A}) = \operatorname{End}(\mathbb{A})$ then possesses an algebra structures, with product * for which $p = e\varepsilon$ acting as $p * \Phi = \Phi$, $\Phi \in \operatorname{End}(\mathbb{A})$, is the unit.

(ii) The further specialization $\mathbf{C} = \mathbb{A} = \mathbf{H}$ with $H(m, e, \Delta, \varepsilon, S)$ a Hopf algebra interprets the Hopf axiom as the fact that (H): $S * \mathrm{id}_{\mathbf{H}} = \mathrm{id}_{\mathbf{H}} * S = e\varepsilon$ (this immediately implying uniqueness of the antipode, $S * \mathrm{id}_{\mathbf{H}} = e\varepsilon$ and $\mathrm{id}_{\mathbf{H}} * S = e\varepsilon$ implying $S * \mathrm{id}_{\mathbf{H}} * S' = S = S'$).

Example B.3 With $\mathbf{H}(m, e, \Delta, \varepsilon, S)$ a Hopf algebra, taking for \mathbf{C} the tensor product coalgebra $\mathbf{H} \otimes \mathbf{H}(\Delta_{\otimes}, \varepsilon_{\otimes})$ and for \mathbf{A} the algebra $\mathbf{H}(m, e)$, we get for Hom $(\mathbf{H} \otimes \mathbf{H}, \mathbf{H})$ an algebra-structure with product * and unit $e\varepsilon_{\otimes}$. For the latter Sm is a left, and $mP_{12}(S \otimes S)$ is a right inverse of $m : Sm * m = e\varepsilon_{\otimes} = m * mP_{12}(S \otimes S)$ (they thus coincide, this proving property (Hm)).

Example B.4 With $\mathbf{H}(m, e, \Delta, \varepsilon, S)$ a Hopf algebra, taking for \mathbf{C} the coalgebra $\mathbf{H}(\Delta, \varepsilon)$ and for \mathbf{A} the tensor product algebra $\mathbf{H} \otimes \mathbf{H}(m_{\otimes}, \varepsilon_{\otimes})$ we get for Hom $(\mathbf{H}, \mathbf{H} \otimes \mathbf{H})$ an algebra structure with product * and unit $e_{\otimes}\varepsilon$. For the latter ΔS is a left, and $(S \otimes S)P_{12}\Delta$ is a right inverse of $\Delta : \Delta S * \Delta = e_{\otimes}\varepsilon = \Delta * (S \otimes S)P_{12}\Delta$ (they thus coincide, this proving property (H Δ)).

Appendix C Primitive and group-like elements

Definition C.1 Let $\mathbf{H}(\Delta, \varepsilon)$ be a bialgebra over the field \mathbf{k} . (i) The element $a \in \mathbf{H}$ is called **primitive** whenever

$$\Delta a = a \otimes \mathbf{1} + \mathbf{1} \otimes a. \tag{C.1}$$

The set of primitive elements of **H** is denoted by $P(\mathbf{H})$.

(ii) The element $a \in \mathbf{H}$ is called **group-like** whenever

$$a \neq 0$$
 and $\Delta a = a \otimes a$. (C.2)

The set of group-like elements of **H** is denoted by $G(\mathbf{H})$.

Proposition C.2 Let $\mathbf{H}(\Delta, \varepsilon)$ be a k-bialgebra.

(i) P(**H**) is a **k**-linear subspace of **H**. Moreover if a, b∈P(**H**), then [a,b] = ab - ba∈P(**H**). Consequently P(**H**) is a Lie subalgebra of the Lie algebra Lie**H**.
 (ii) One has P(**H**)⊂ Kerε:

$$a \in P(\mathbf{H}) \Rightarrow \varepsilon(a) = 0,$$
 (C.3)

(iii) $G(\mathbf{H})$ is multiplicative: if $a, b \in G(\mathbf{H})$, then $ab \in G(\mathbf{H})$.

Daniel Kastler

(iv) One has

$$a \in G(\mathbf{H}) \Rightarrow \varepsilon(a) = 1.$$
 (C.4)

If furthermore \mathbf{H} is a Hopf algebra with antipode S, then:

$$a \in P(\mathbf{H}) \implies Sa = -a. \tag{C.5}$$

$$a \in G(\mathbf{H}) \Rightarrow Sa = a^{-1},$$
 (C.6)

showing that $G(\mathbf{H})$ is then a group.

Proof (i) Let $a, b \in P(\mathbf{H}), \lambda \in \mathbf{k}$, then $a + b, \lambda a \in P(\mathbf{H})$. Furthermore:

$$\begin{aligned} \Delta[a,b] &= \Delta(ab-ba) = (\Delta a)(\Delta b) - (\Delta b)(\Delta a) \\ &= (a \otimes \mathbf{1} + \mathbf{1} \otimes a)(b \otimes \mathbf{1} + \mathbf{1} \otimes b) - (b \otimes \mathbf{1} + \mathbf{1} \otimes b)(a \otimes \mathbf{1} + \mathbf{1} \otimes a) \\ &= ab \otimes \mathbf{1} + a \otimes b + b \otimes a + \mathbf{1} \otimes ab - (ba \otimes \mathbf{1} + b \otimes a + a \otimes b + \mathbf{1} \otimes ba) \\ &= (ab-ba) \otimes \mathbf{1} + \mathbf{1} \otimes (ab-ba) = [a,b] \otimes \mathbf{1} + \mathbf{1}[a,b]. \end{aligned}$$

(ii) $(id \otimes \varepsilon) \Delta = id$ entails

$$(\mathrm{id} \otimes \varepsilon) \Delta a = (\mathrm{id} \otimes \varepsilon) (a \otimes \mathbf{1} + \mathbf{1} \otimes a) \\ = a \otimes \mathbf{1} + \mathbf{1} \otimes \varepsilon(a) \\ = a + \varepsilon(a) \mathbf{1} = a.$$

(iii) $a, b \in G(\mathbf{H})$ entails

$$\Delta(ab) = (\Delta a)(\Delta b) = (a \otimes a)(b \otimes b) = ab \otimes ab.$$

(iv) $(id \otimes \varepsilon) \Delta = id$ entails

$$(\mathrm{id}\otimes\varepsilon)\Delta a = (\mathrm{id}\otimes\varepsilon)(a\otimes a) = a\otimes\varepsilon(a) = a\varepsilon(a) = a.$$

Assume that \mathbf{H} is a Hopf algebra with antipode S.

Check of (C.5): $m(\mathrm{id} \otimes S)\Delta = e\varepsilon$ entails

$$0 = m(\mathrm{id} \otimes S)(a \otimes \mathbf{1} + \mathbf{1} \otimes a) = m(a \otimes \mathbf{1} + \mathbf{1} \otimes Sa) = a + Sa.$$

Check of (C.6): $m(\mathrm{id}\otimes S)\Delta = e\varepsilon$ entails

r

$$m(\operatorname{id}\otimes S)(a\otimes a) = m(a\otimes Sa) = a(Sa) = 1.$$

whilst $m(S \otimes id) \Delta = e\varepsilon$ entails

$$m(S \otimes \mathrm{id})(a \otimes a) = m(Sa \otimes a) = (Sa)a = \mathbf{1}.$$

References

- A. Connes and D. Kreimer. Renormalization in quantum field theory and the Riemann-Hilbert Problem. I: the Hopf algebra structure of graphs and the main theorem. hep-th/9912092 13 Dec. 1999.
- A. Connes and D. Kreimer. Renormalization in quantum field theory and the Riemann-Hilbert Problem. II: the β -function, diffeomorphisms and the renormalization group. arXiv: hep-th/0003 188 21 Mars 2000.
- N. Bourbaki. Algèbre. Chapitre II. Par. 6.