Commun.Math.Phys. 81,239-241(1981)

Lax Representation for the Systems of S.Kowalevskaya Type

A.M. Perelomov

Institute of Theoretical and Experimental Physics, 117259 Moscow, USSR

Abstract

We describe a number of dynamical systems that are generalizations of the S. Kowalevskaya system and admit the Lax representation.

It is well known (see for example [1,2]) that the equations of motion of a threedimensional heavy rigid body rotated about a fixed point is a completely integrable dynamical system only in the cases of Euler [3], Lagrange [4] and Kowalevskaya [5]. In the present note we describe a number of systems that are of the Kowalevskaya type and admit the Lax representation¹.

1. Let \mathcal{G} be the Lie algebra of a group G, \mathcal{G}^* be the space dual to \mathcal{G} and $\{x_{\alpha}\}$ be the coordinates of a point in the space \mathcal{G}^* . In the space $\mathcal{F}(\mathcal{G}^*)$ of smooth functions on \mathcal{G}^* , let us define the Poisson bracket ²

$$\{f,g\} = C_{\alpha\beta}^{\gamma} x_{\gamma} \partial^{\alpha} f \partial^{\beta} g, \qquad \partial^{\alpha} = \partial/\partial x_{\alpha}.$$
 (1)

Here $C_{\alpha\beta}^{\gamma}$ are the structure constants of the Lie algebra. The space $\mathcal{F}(\mathcal{G}^*)$ is endowed by the above formula with the structure of the Lie algebra. A

¹ Note that the Lax representation for equations of motion of a rigid body was discovered first by Manakov [8] for some particular cases.

² In this note we use the tensor notations; in particular, everywhere repeated indices imply summation.

dynamical system in \mathcal{G}^* is determined by a Hamiltonian function $H(x) \in \mathcal{F}$, so that the equations of motion have the form

$$\dot{x}_{\alpha} = \{H, x_{\alpha}\}. \tag{2}$$

The coadjoint representation of the group G acts on the space \mathcal{G}^* . Orbits of this representation are invariant with respect to an arbitrary Hamiltonian H, and are the phase spaces of the considered systems.

2. Let G be a compact simple Lie group, K be its subgroup such that the factor space G/K is a symmetric space [6]. Then $\mathcal{G} = \mathcal{K} \oplus \mathcal{S}$, \mathcal{K} is the Lie algebra of the group K, \mathcal{S} is the orthogonal complement to \mathcal{K} in \mathcal{G} relative to the Killing-Cartan form.

It is well known that a certain irreducible representation $T^0(k)$ of the group K acts on the space \mathcal{S} . Let us consider an irreducible representation T(g) of the group G, such that under the restriction of this representation on the subgroup K, the irreducible representation $T^0(k)$ is contained in T(g) with the unit multiplicity. Let V be a vector space in which the representation T(g) acts. Then it is possible to define the group $\tilde{G} = G \cdot V$ that is a semidirect product of the group G and the abelian vector group V. Let us denote by V^0 the subspace of V on which the representation $T^0(k)$ acts, and by A the projection operator on the subspace V^0 . Let V^1 be the orthogonal complement to $V^0: V = V^0 \oplus V^1$. Note that the spaces V^0 and S are isomorphic because T(g) contains $T^0(k)$ only once. Therefore we have the decomposition

$$\tilde{\mathcal{G}} = \mathcal{K} \oplus \mathcal{S} \oplus \mathcal{V}^0 \oplus \mathcal{V}^1$$
,

and the analogous decomposition for the space $\tilde{\mathcal{G}}^*$:

$$\tilde{\mathcal{G}}^* = \mathcal{L} \oplus \mathcal{N} \oplus \mathcal{P} \oplus \mathcal{T}. \tag{3}$$

Here $\mathcal{L} = \mathcal{K}^*$, $\mathcal{N} = \mathcal{S}^*$, $\mathcal{P} = \mathcal{V}^{(0)*}$, $\mathcal{T} = \mathcal{V}^{(1)*}$. In addition, dim $\mathcal{N} = \dim \mathcal{P} = n$, and \mathcal{N} and \mathcal{P} are isomorphic relative to the action of the group K.

Let \hat{l} be the matrix of the representation T of the Lie algebra \mathcal{G} , $\hat{l} = \{l_{jk}\}$ corresponds to the matrix of the representation T^0 of the algebra \mathcal{K} acting in the space \mathcal{P} . Let us introduce the matrix, which is important in the sequel,

$$L = A \left[-\hat{l}^2 + (\gamma \otimes p + p \otimes \gamma) \right] A, \tag{4}$$

where $p \in \mathcal{P} \oplus \mathcal{T}$ and $\gamma \in \mathcal{P}$ is the constant vector in \mathcal{P} . We identify \mathcal{P} and \mathcal{P}^* by means of the K-invariant scalar product on \mathcal{P} .

The system of S. Kowalevskaya type is defined by the Hamiltonian

$$H = \frac{1}{2} \operatorname{tr} L = \alpha I_2(l) + \beta \mathcal{J}_2(n) + (\gamma p).$$
 (5)

Here $I_2(l)$ and $\mathcal{J}_2(n)$ are the quadratic functions on \mathcal{L} and \mathcal{N} respectively that are invariant relative to the coadjoint representation of the group K and $p \in \mathcal{P}$.

3. Let us consider in detail the case: G = SO(n+1), K = SO(n), $\tilde{G} = G \cdot V = E(n+1)$ is the motion group of the (n+1)-dimensional Euclidean space $V = \mathbb{R}^{n+1}$, $V^0 = \mathbb{R}^n$, $V^1 = \mathbb{R}^1$. Let $\hat{l}_{jk} = -\hat{l}_{kj}$ and \hat{p}_m (j, k = 1, ..., (n+1)) be the standard basis in the space of linear functions on \mathcal{G}^* and \mathcal{V}^* respectively with the standard Poisson brackets.

The Hamiltonian (5) takes now the form

$$2H = 2\sum_{j < k}^{n} l_{jk}^{2} + \sum_{j=1}^{n} n_{j}^{2} + 2\sum_{j=1}^{n} \gamma_{j} p_{j}.$$
 (6)

Theorem. The equations of motion (2) of the system with Hamiltonian (6) are equivalent to the Lax equation

$$\dot{L} = [L, M], \tag{7}$$

where L is given by the formula (4), and $M = c l = c A \hat{l} A$, c is a constant. The theorem is verified by a direct calculation.

From the Lax representation (7), it follows immediately that the eigenvalues of the matrix L or the quantities $I_{2k} = k^{-1} \operatorname{tr}(L^k)$, $k = 1, \ldots, n$ are integrals of motion for Eqs. (2). Notice $I_2 = 2H$.

There are also (n-1)(n-2)/2 linear integrals related to the invariance of H with respect to the group $G_0 = SO(n-1)$ that leaves the vector γ invariant, and also ([n/2]+1) independent polynomials that are invariant with respect to the coadjoint representation of the group \tilde{G} . Using the theory of reduction of hamiltonian systems with symmetries [2,7] we may show that the above system is reduced to a system with 2n-dimensional phase space. It can be verifies that integrals I_2, I_4, \ldots, I_{2n} are functionally independent and in involution. Therefore, the systems under consideration are completely integrable.

Finally, we wish to note the following facts:

- 1. The system considered by S. Kowalevskaya corresponds to n=2.
- 2. Similar results are valid for other symmetric spaces. The simplest ones are the cases of the spaces of rank one: $SU(n+1)/SU(n) \times U(1)$, $Sp(n+1)/Sp(n) \times Sp(1)$ and $F_4/SO(9)$.
- 3. If $p \in \mathcal{T}$ in formula (5), then we deal with the generalization of the Lagrange case. In this case, the constants α and β may be arbitrary.
- 4. The results of this section are valid also for similar systems, related to group $\tilde{G} = SO(n+2)$.

The detailed presentation of results will be published elsewhere.

Acknowledgement. I am grateful to V.Golo for the improvement of the language of this note.

References

- [1] Motion of a rigid body around a fixed point. A collection of papers, dedicated to the memory of S. Kowalevskaya. Acad. of Sci. USSR, Moscow (1940) (in Russian)
- [2] Arnold, V.I.: Mathematical methods of classical mechanics. New York: Springer 1978
- [3] Euler, L.: Decouverte d'un nouveau principe de Mecanique. Mem. de l'Acad. des Sciences de Berlin 1758
- [4] Lagrange, J.L.: Mecanique analytique. Paris 1788
- [5] Kowalevskaya, S.: Acta Math. **12**, H.2 177–232 (1889)
- [6] Helgason, S.: Differential geometry and symmetric spaces. New York: Acad. Press 1962
- [7] Marsden, J., Weinstein, A.: Rep. Math. Phys. 5, 121–130 (1974)
- [8] Manakov, S.: Funkt. Anal. Priloz. **10**, 93–94 (1976)