# Exact Chromatic Polynomials for Toroidal Chains of Complete Graphs

Shu-Chiuan Chang\*

C. N. Yang Institute for Theoretical Physics State University of New York Stony Brook, N. Y. 11794-3840

#### Abstract

We present exact calculations of the partition function of the zero-temperature Potts antiferromagnet (equivalently, the chromatic polynomial) for graphs of arbitrarily great length composed of repeated complete subgraphs  $K_b$  with b = 5, 6 which have periodic or twisted periodic boundary condition in the longitudinal direction. In the  $L_x \to \infty$  limit, the continuous accumulation set of the chromatic zeros  $\mathcal{B}$  is determined. We give some results for arbitrary b including the extrema of the eigenvalues with coefficients of degree b - 1 and the explicit forms of some classes of eigenvalues. We prove that the maximal point where  $\mathcal{B}$ crosses the real axis,  $q_c$ , satisfies the inequality  $q_c \leq b$  for  $2 \leq b$ , the minimum value of q at which  $\mathcal{B}$  crosses the real q axis is q = 0, and we make a conjecture concerning the structure of the chromatic polynomial for Klein bottle strips.

<sup>\*</sup>email: shu-chiuan.chang@sunysb.edu

### I. INTRODUCTION

The q-state Potts antiferromagnet (AF) [1,2] exhibits nonzero ground state entropy,  $S_0 > 0$  (without frustration) for sufficiently large q on a given lattice  $\Lambda$  or, more generally, on a graph G. This is equivalent to a ground state degeneracy per site W > 1, since  $S_0 = k_B \ln W$ . There is a close connection with graph theory here, since the zero-temperature partition function of the above-mentioned q-state Potts antiferromagnet on a graph G satisfies

$$Z(G, q, T = 0)_{PAF} = P(G, q) , \qquad (1.1)$$

where P(G, q) is the chromatic polynomial expressing the number of ways of coloring the vertices of the graph G with q colors such that no two adjacent vertices have the same color (for reviews, see [3]- [5]). The minimum number of colors necessary for such a coloring of G is called the chromatic number,  $\chi(G)$ . Thus

$$W(\{G\}, q) = \lim_{n \to \infty} P(G, q)^{1/n} , \qquad (1.2)$$

where n is the number of vertices of G and  $\{G\} = \lim_{n\to\infty} G$ . Where no confusion will result, we shall sometimes write G rather than  $\{G\}$  for the infinite-length limit of a given type of strip graph. At certain special points  $q_s$  (typically  $q_s = 0, 1, ..., \chi(G)$ ), one has the noncommutativity of limits

$$\lim_{q \to q_s} \lim_{n \to \infty} P(G, q)^{1/n} \neq \lim_{n \to \infty} \lim_{q \to q_s} P(G, q)^{1/n} , \qquad (1.3)$$

and hence it is necessary to specify the order of the limits in the definition of  $W(\{G\}, q_s)$  [6]. Denoting  $W_{qn}$  and  $W_{nq}$  as the functions defined by the different order of limits on the left and right-hand sides of (1.3), we take  $W \equiv W_{qn}$  here; this has the advantage of removing certain isolated discontinuities that are present in  $W_{nq}$ .

Using the expression for P(G,q), one can generalize q from  $\mathbb{Z}_+$  to  $\mathbb{C}$ . The zeros of P(G,q)in the complex q plane are called chromatic zeros; a subset of these may form an accumulation set in the  $n \to \infty$  limit, denoted  $\mathcal{B}$ , which is the continuous locus of points where  $W(\{G\},q)$ is nonanalytic. <sup>1</sup> The maximal region in the complex q plane to which one can analytically continue the function  $W(\{G\},q)$  from physical values where there is nonzero ground state entropy is denoted  $R_1$ . The maximal value of q where  $\mathcal{B}$  intersects the (positive) real axis is labeled  $q_c(\{G\})$ . This point is important since  $W(\{G\},q)$  is a real analytic function from large values of q down to  $q_c(\{G\})$ .

<sup>&</sup>lt;sup>1</sup>For some families of graphs  $\mathcal{B}$  may be null, and W may also be nonanalytic at certain discrete points.

Here we present exact calculations of the partition function of the chromatic polynomial P(G,q) for graphs of arbitrarily great length composed of repeated complete subgraphs  $K_b$  with b = 5, 6 which have periodic or twisted periodic boundary conditions in the longitudinal direction.<sup>2</sup> Thus, consider  $L_x$  copies of the complete graph  $K_{L_y}$  with vertex set  $V(K_{L_y}) = \{1, 2, ..., L_y\}$ . Denote the edges joining each adjacent pair of  $K_{L_y}$  and  $K'_{L_y}$  graphs as  $L = \{v_i v'_j | i, j = 1, ..., L_y\}$ , where  $v_i, v'_j$  are vertices of adjacent complete graphs  $K_{L_y}$  and  $K'_{L_y}$ , and impose periodic boundary conditions in the longitudinal direction, x. In [7], the graph with  $L_x$  copies of  $K_{L_y}$  for arbitrary  $L_x$  and  $L_y$  was called the bracelet strip. The total number of vertices is  $n = L_x L_y$  for these strips.

A generic form for chromatic polynomials for recursively defined families of graphs (families of graphs that can be constructed via repeated addition of some subgraph), of which strip graphs  $G_s$  are special cases, is [8]

$$P((G_s)_m, q) = \sum_{j=1}^{N_{G_s,\lambda}} c_{G_s,j}(q) (\lambda_{G_s,j}(q))^{L_x} , \qquad (1.4)$$

where  $c_{G_{s,j}}(q)$  and the  $N_{G_{s,\lambda}}$  terms  $\lambda_{G_{s,j}}(q)$  depend on the type of strip graph  $G_s$  but are independent of  $L_x$ .

For a given type of strip graph  $G_s$ , we denote the sum of the coefficients  $c_{G_s,j}$  as

$$C(G_s, q) = \sum_{j=1}^{N_{G_s,\lambda}} c_{G_s,j}(q) .$$
(1.5)

Some works on calculations of chromatic polynomials for recursive families of graphs include [9]- [49].

### **II. FAMILY OF TOROIDAL STRIPS**

Specifically, we consider the set of edges linking two successive complete graphs on the chain is  $L = \{11, 22, 33, ..., L_y L_y\}$ , which will be denoted as a torus. For  $L_y = 2$ , this is the ladder graph with vertex set  $T_2 = K_2$  [9]. For the graph with  $L_y = 3, 4$ , which has vertex set  $C_3 = K_3$  and  $K_4$ , the chromatic polynomial was given in [10] and [11], respectively. In [58,59], these families of graphs with linking edge sets  $\{11, 22, 33, ..., bb\}$  were denoted as

 $<sup>{}^{2}</sup>K_{n}$  denotes the complete graph, i.e. the graph with *n* vertices such that each vertex is connected by an edge to all of the other vertices.

 $B_m(b)$ , where  $b = L_y$ ,  $m = L_x$ . The corresponding Klein bottle strip is the same as the toroidal strip except one of the linking edge set is  $\{1L_y, 2(L_y - 1), ... L_y 1\}$ . The form of the chromatic polynomials for the  $B_m(b)$  strips was determined to be

$$P(B_m(b),q) = \sum_{d=0}^{b} \sum_{j=1}^{N_{B_m(b),d,\lambda}} c_{B_m(b),d,j}(q) c'_{B_m(b),d,j}(\lambda_{B_m(b),d,j}(q))^m , \qquad (2.1)$$

where d is the degree of the coefficient  $c_{B_m(b),d,j}(q)$  as a polynomial in q, and the  $c'_{B_m(b),d,j}$ 's are numbers which depend on b. The structural feature that these coefficients can be grouped into sets of fixed degree d is similar to that found in [14] for cyclic strips of the square and triangular lattice, although for these bracelet chain graphs, for  $b \geq 3$ , there is more than one coefficient of a given degree (see [10] for the case b = 3), whereas in contrast, for the cyclic strip graphs studied in [14] (and the self-dual family studied in [55]) there is only one coefficient of each degree. The degree d is denoted as the 'level' in [60,58,59]. The  $\lambda_{B_m(b),d,j}(q)$ is the eigenvalue of an appropriately defined transfer matrix [37] and is a polynomial in q of degree b - d. For sufficiently large integer q, the coefficient  $c_{B_m(b),d,j}(q)c'_{B_m(b),d,j}$  can be interpreted as the multiplicity of this eigenvalue.

There are partitions associated with each d which were used to determine the forms of  $c_{B_m(b),d,j}$  in [59]. For a partition X of d such that  $x_d + x_{d-1} + \ldots + x_1 = d$  and  $x_d \leq x_{d-1} \leq \ldots \leq x_1$  where  $x_i$  for  $2 \leq i \leq d$  are non-negative integers and  $x_1$  is a positive integer, there is an associated partition Y of d(d+1)/2 such that  $y_d + y_{d-1} + \ldots + y_1 = d(d+1)/2$  and  $y_i = x_i + d - i$  for  $1 \leq i \leq d$ . Then the  $c_{B_m(b),d,j}$  for a partition X can be written as

$$c_{X,d} = \frac{1}{d!} \prod_{i=1}^{d} (q - y_i) .$$
(2.2)

For the partitions [d] and  $[1^d]$ , the associated  $\lambda_{B_m(b),d,j}(q)$ 's were given in [58].

## A. $L_y = 5$ toroidal strip

By using theorems in [58], we find the chromatic polynomial for  $B_m(5)$ , i.e., the 5 ×  $L_x$  strips with each transverse slice forming a  $K_5$  graph, and with edge linking set  $L = \{11, 22, 33, 44, 55\}$  between two adjacent slices.

It is convenient to express the  $\lambda$ 's in term of the following functions [60]

$$f_i(b,q) = \sum_{s=0}^{b-i} (-1)^s {\binom{b-i}{s}} (q-i-s)^{(b-i-s)} , \qquad (2.3)$$

where  $q^{(k)}$  is falling factorial defined as

$$q^{(k)} = \prod_{s=0}^{k-1} (q-s) .$$
 (2.4)

Here we adopt the convention that  $f_i(b,q) = 0$  if b < i.

For the case b = 5, one has

$$f_0(5,q) = q^5 - 15q^4 + 95q^3 - 325q^2 + 609q - 501$$

$$f_0(5,q) = q^4 - 14q^3 + 77q^2 - 200q + 200$$

$$(2.5)$$

$$f_1(5,q) = q^4 - 14q^3 + 77q^2 - 200q + 209$$
(2.6)

 $f_2(5,q) = q^3 - 12q^2 + 50q - 73 \tag{2.7}$ 

$$f_3(5,q) = q^2 - 9q + 21 \tag{2.8}$$

$$f_4(5,q) = q - 5 \tag{2.9}$$

$$f_5(5,q) = 1 {,} (2.10)$$

and the chromatic polynomial for the  $B_m(5)$  strip is

$$P(B_m(5),q) = \sum_{d=0}^{5} \sum_{j=1}^{N_{B_m(5),d,\lambda}} c_{B_m(5),d,j}(q) c'_{B_m(5),d,j}(\lambda_{B_m(5),d,j}(q))^m , \qquad (2.11)$$

where  $N_{B_m(5),d,\lambda}$ 's are equal to 1, 2, 5, 9, 9, 1 for d from 0 to 5. We find that the total number of distinct  $\lambda$ 's is

$$N_{P,B_m(5),\lambda} = \sum_{d=0}^{5} N_{B_m(5),d,\lambda} = 27 .$$
(2.12)

We note that our result (2.12) differs from the pattern  $N_{P,B_m(b),\lambda} = 2^b$  that was found for  $1 \le b \le 4$ , namely  $N_{P,B_m(1),\lambda} = 2$ ,  $N_{P,B_m(2),\lambda} = 2^2$  [9],  $N_{P,B_m(3),\lambda} = 2^3$  [10], and  $N_{P,B_m(4),\lambda} = 2^4$  [11].

We calculate that the eigenvalue with coefficient of degree 0 in q is unique and is given by

$$\lambda_{B_m(5),0,1} = f_0(5,q) = q^5 - 15q^4 + 95q^3 - 325q^2 + 609q - 501 .$$
(2.13)

For the eigenvalues with coefficients of degree 1 in q we find

$$\lambda_{B_m(5),1,1} = -f_1(5,q) + 4f_2(5,q) = -q^4 + 18q^3 - 125q^2 + 400q - 501$$
(2.14)

$$\lambda_{B_m(5),1,2} = -f_1(5,q) - f_2(5,q) = -(q-4)(q^3 - 9q^2 + 29q - 34) .$$
(2.15)

Next, for the eigenvalues with coefficients of degree 2 in q we obtain

$$\lambda_{B_m(5),2,1} = f_2(5,q) - 6f_3(5,q) + 6f_4(5,q) = q^3 - 18q^2 + 110q - 229$$
(2.16)

$$\lambda_{B_m(5),2,2} = f_2(5,q) - f_3(5,q) - 4f_4(5,q) = q^3 - 13q^2 + 55q - 74$$
(2.17)

$$\lambda_{B_m(5),2,3} = f_2(5,q) + 2f_3(5,q) + 2f_4(5,q) = q^3 - 10q^2 + 34q - 41$$
(2.18)

$$\lambda_{B_m(5),2,4} = f_2(5,q) - 3f_3(5,q) = q^3 - 15q^2 + 77q - 136$$
(2.19)

$$\lambda_{B_m(5),2,5} = f_2(5,q) + 2f_3(5,q) = q^3 - 10q^2 + 32q - 31 .$$
(2.20)

For  $1 \leq j \leq 3$  these  $\lambda_{B_m(5),2,j}$ 's correspond to the partition [2] and for j = 4, 5 to the partition [11].

Proceeding to the eigenvalues with coefficients of degree 3, we find

$$\lambda_{B_m(5),3,1} = -f_3(5,q) + 6f_4(5,q) - 6f_5(5,q) = -q^2 + 15q - 57$$
(2.21)

$$\lambda_{B_m(5),3,2} = -f_3(5,q) + f_4(5,q) + 4f_5(5,q) = -q^2 + 10q - 22$$
(2.22)

$$\lambda_{B_m(5),3,3} = -f_3(5,q) - 2f_4(5,q) - 2f_5(5,q) = -q^2 + 7q - 13$$
(2.23)

$$\lambda_{B_m(5),3,4} = -f_3(5,q) + 4f_4(5,q) - 2f_5(5,q) = -q^2 + 13q - 43$$
(2.24)

$$\lambda_{B_m(5),3,5} = -f_3(5,q) + f_4(5,q) + f_5(5,q) = -(q-5)^2$$
(2.25)

$$\lambda_{B_m(5),3,6} = -f_3(5,q) - f_4(5,q) + 3f_5(5,q) = -q^2 + 8q - 13$$
(2.26)

$$\lambda_{B_m(5),3,7} = -f_3(5,q) - 3f_4(5,q) - 3f_5(5,q) = -(q-3)^2$$
(2.27)

$$\lambda_{B_m(5),3,8} = -f_3(5,q) + 2f_4(5,q) = -q^2 + 11q - 31$$
(2.28)

$$\lambda_{B_m(5),3,9} = -f_3(5,q) - 3f_4(5,q) = -q^2 + 6q - 6 .$$
(2.29)

Here the  $\lambda_{B_m(5),3,j}$  for  $1 \le j \le 3$  correspond to the partition [3], for  $4 \le j \le 7$  to the partition [21], and for j = 8, 9 to the partition [111].

For the eigenvalues with coefficients of degree 4 in q, we calculate

$$\lambda_{B_m(5),4,1} = f_4(5,q) - 4f_5(5,q) = q - 9 \tag{2.30}$$

$$\lambda_{B_m(5),4,2} = f_4(5,q) + f_5(5,q) = q - 4 \tag{2.31}$$

$$\lambda_{B_m(5),4,3} = f_4(5,q) - 3f_5(5,q) = q - 8 \tag{2.32}$$

$$\lambda_{B_m(5),4,4} = f_4(5,q) + 3f_5(5,q) = q - 2 \tag{2.33}$$

$$\lambda_{B_m(5),4,5} = f_4(5,q) + 2f_5(5,q) = q - 3 \tag{2.34}$$

$$\lambda_{B_m(5),4,6} = f_4(5,q) = q - 5 \tag{2.35}$$

$$\lambda_{B_m(5),4,7} = f_4(5,q) - 2f_5(5,q) = q - 7 \tag{2.36}$$

$$\lambda_{B_m(5),4,8} = f_4(5,q) - f_5(5,q) = q - 6 \tag{2.37}$$

$$\lambda_{B_m(5),4,9} = f_4(5,q) + 4f_5(5,q) = q - 1 .$$
(2.38)

Here the  $\lambda_{B_m(5),4,j}$  for j = 1, 2 correspond to the partition [4], for j = 3, 5, 6 to the partition [31], for j = 4, 6, 7 to the partition [211], for j = 5, 7 to the partition [22], and for j = 8, 9 to the partition [1111]. Notice that the terms  $\lambda_{B_m(5),4,j}$  for  $5 \le j \le 7$  involve degeneracies from two different associated partitions.

Finally, for the eigenvalue with coefficient of degree 5, we have

$$\lambda_{B_m(5),5,1} = -f_5(5,q) = -1 . (2.39)$$

This involves a degeneracy from all the possible partitions.

The coefficient  $c_{B_m(5),d,j}(q)c'_{B_m(5),d,j}$  of degree 0 in q is

$$c_{B_m(5),0,1}c'_{B_m(5),0,1} = 1. (2.40)$$

The coefficients of degree 1 in q are

$$c_{B_m(5),1,1}c'_{B_m(5),1,1} = q - 1 \tag{2.41}$$

$$c_{B_m(5),1,2}c'_{B_m(5),1,2} = 4(q-1) , \qquad (2.42)$$

where  $c_{B_m(5),1,j} = q - 1$  for j = 1, 2, and  $c'_{B_m(5),1,j} = 1, 4$  for j = 1, 2.

For d = 2, there are partitions [2] and [11], and the associated  $c_{X,2}$  are  $c_{[2],2} = \frac{1}{2}q(q-3)$ and  $c_{[11],2} = \frac{1}{2}(q-1)(q-2)$ . The coefficients of degree 2 in q are listed in Table I

j	partition	$c^{\prime}_{B_m(5),2,j}$	$c_{B_m(5),2,j}c'_{B_m(5),2,j}$
1	[2]	1	$\frac{1}{2}q(q-3)$
<b>2</b>	[2]	4	2q(q-3)
3	[2]	5	$rac{5}{2}q(q-3)$
4	[11]	4	2(q-1)(q-2)
5	[11]	6	3(q-1)(q-2)

TABLE I. Coefficients of degree 2 in q for  $B_m(5)$  strip.

For d = 3, there are partitions [3], [21] and [111], and the associated  $c_{X,3}$  are  $c_{[3],3} = \frac{1}{6}q(q-1)(q-5)$ ,  $c_{[21],3} = \frac{1}{6}q(q-2)(q-4)$  and  $c_{[111],3} = \frac{1}{6}(q-1)(q-2)(q-3)$ . The coefficients of degree 3 in q are listed in Table II

TABLE II. Coefficients of degree 3 in q for  $B_m(5)$  strip.

j	partition	$c_{B_m(5),3,j}'$	$c_{B_m(5),3,j}c'_{B_m(5),3,j}$
1	[3]	1	$\frac{1}{6}q(q-1)(q-5)$
2	[3]	4	$\frac{2}{3}q(q-1)(q-5)$
3	[3]	5	$\frac{5}{6}q(q-1)(q-5)$
4	[21]	8	$\frac{4}{3}q(q-2)(q-4)$
5	[21]	10	$\frac{5}{3}q(q-2)(q-4)$
6	[21]	12	2q(q-2)(q-4)
7	[21]	10	$\frac{5}{3}q(q-2)(q-4)$
8	[111]	6	(q-1)(q-2)(q-3)
9	[111]	4	$\frac{2}{3}(q-1)(q-2)(q-3)$

For d = 4, there are partitions [4], [31], [22], [211] and [1111], and the associated  $c_{X,4}$  are  $c_{[4],4} = \frac{1}{24}q(q-1)(q-2)(q-7), c_{[31],4} = \frac{1}{24}q(q-1)(q-3)(q-6), c_{[22],4} = \frac{1}{24}q(q-1)(q-4)(q-5), c_{[211],4} = \frac{1}{24}q(q-2)(q-3)(q-5), \text{ and } c_{[1111],4} = \frac{1}{24}(q-1)(q-2)(q-3)(q-4).$  The coefficients of degree 4 in q are listed in Table III

j	partition	$c'_{B_m(5),4,j}$	$c_{B_m(5),4,j}c'_{B_m(5),4,j}$
1	[4]	1	$\frac{1}{24}q(q-1)(q-2)(q-7)$
2	[4]	4	$\frac{1}{6}q(q-1)(q-2)(q-7)$
3	[31]	12	$\frac{1}{2}q(q-1)(q-3)(q-6)$
4	[211]	12	$\frac{1}{2}q(q-2)(q-3)(q-5)$
5	[31]	18	$\frac{1}{6}q(q-1)(7q^2-63q+131)$
	[22]	10	
6	[31]	15	$rac{5}{4}q(q-3)(q^2-7q+8)$
	[211]	15	
7	[211]	18	$\frac{1}{6}q(q-5)(7q^2-35q+37)$
	[22]	10	
8	[1111]	4	$\frac{1}{6}(q-1)(q-2)(q-3)(q-4)$
9	[1111]	1	$\frac{1}{24}(q-1)(q-2)(q-3)(q-4)$

TABLE III. Coefficients of degree 4 in q for  $B_m(5)$  strip.

Finally, the coefficient of degree 5 in q is given by

$$c_{B_m(5),5,1}c'_{B_m(5),5,1} = q^5 - 15q^4 + 75q^3 - 145q^2 + 89q - 1.$$
(2.43)

The sum of all the coefficients is equal to

$$C(B_m(5),q) = \sum_{d=0}^{5} \sum_{j=1}^{N_{B_m(5),d,\lambda}} c_{B_m(5),d,j}(q) c'_{B_m(5),d,j}$$
  
=  $q(q-1)(q-2)(q-3)(q-4) = P(K_5,q)$ . (2.44)

The chromatic number is  $\chi(B_m(5)) = 5$ .

The locus  $\mathcal{B}$  and chromatic zeros for the  $B_m(5)$  strip with  $L_x = m = 20$  are shown in Fig. 1. The locus  $\mathcal{B}$  crosses the real q-axis at q = 0, 2, 4 and  $q_c(B_m(5))$ , where

$$q_c(B_m(5)) \simeq 4.50634 \cdots$$
 (2.45)



FIG. 1. Locus  $\mathcal{B}$  for the  $m \to \infty$  limit of the family  $B_m(5)$  with toroidal boundary conditions and chromatic zeros for  $B_m(5)$  with m = 20 (i.e., n = 100).

The locus  $\mathcal{B}$  has support for  $Re(q) \geq 0$ , and separates the q plane into four regions. The outermost one, region  $R_1$ , extends to infinite |q| and includes the intervals  $q \geq q_c(B_m(5))$  and  $q \leq 0$  on the real q axis. Region  $R_2$  includes the real interval  $4 \leq q \leq q_c(B_m(5))$ , region  $R_3$  includes the real interval  $2 \leq q \leq 4$ , while region  $R_4$  includes the real interval  $0 \leq q \leq 2$ . In regions  $R_i$ ,  $1 \leq i \leq 4$ , the dominant terms are  $\lambda_{B_m(5),0,1}$ ,  $\lambda_{B_m(5),3,1}$ ,  $\lambda_{B_m(5),2,1}$ , and  $\lambda_{B_m(5),1,1}$ , respectively. Thus, the  $q_c(B_m(5))$  given in (2.45) is the degeneracy between  $|\lambda_{B_m(5),0,1}|$  and  $|\lambda_{B_m(5),3,1}|$ , and is the real solution of  $q^5 - 15q^4 + 95q^3 - 326q^2 + 624q - 558 = 0$ .

### **B.** $L_y = 6$ toroidal strip

We have also succeeded in obtaining the chromatic polynomial for  $B_m(6)$ , i.e., the  $6 \times L_x$  strips with each transverse slice forming a  $K_6$  graph, and with edge linking set  $L = \{11, 22, 33, 44, 55, 66\}$  between two adjacent slices.

For b = 6, one has

$$f_0(6,q) = q^6 - 21q^5 + 190q^4 - 965q^3 + 2944q^2 - 5155q + 4051$$
(2.46)

$$f_1(6,q) = q^5 - 20q^4 + 165q^3 - 710q^2 + 1609q - 1546$$
(2.47)

$$f_2(6,q) = q^4 - 18q^3 + 125q^2 - 400q + 501$$
(2.48)

$$f_3(6,q) = q^3 - 15q^2 + 77q - 136 \tag{2.49}$$

$$f_4(6,q) = q^2 - 11q + 31 \tag{2.50}$$

$$f_5(6,q) = q - 6 \tag{2.51}$$

$$f_6(6,q) = 1 {,} {(2.52)}$$

and the chromatic polynomial for the  $B_m(6)$  strip is

$$P((B_m(6),q) = \sum_{d=0}^{6} \sum_{j=1}^{N_{B_m(6),d,\lambda}} c_{B_m(6),d,j}(q) c'_{B_m(6),d,j}(\lambda_{B_m(6),d,j}(q))^m , \qquad (2.53)$$

where  $N_{B_m(6),d,\lambda}$ 's are equal to 1, 2, 5, 10, 16, 11, 1 for d from 0 to 6. We find that the total number of distinct  $\lambda$ 's is

$$N_{P,B_m(6),\lambda} = 46$$
 . (2.54)

The eigenvalue with coefficient of degree 0 in q is unique and is given by

$$\lambda_{B_m(6),0,1} = f_0(6,q) = q^6 - 21q^5 + 190q^4 - 965q^3 + 2944q^2 - 5155q + 4051 .$$
 (2.55)

For the eigenvalues with coefficients of degree 1 in q we have

$$\lambda_{B_m(6),1,1} = -f_1(6,q) + 5f_2(6,q) = -q^5 + 25q^4 - 255q^3 + 1335q^2 - 3609q + 4051 \quad (2.56)$$

$$\lambda_{B_m(6),1,2} = -f_1(6,q) - f_2(6,q) = -(q-5)(q^4 - 14q^3 + 77q^2 - 200q + 209) .$$
 (2.57)

For the eigenvalues with coefficients of degree 2 in q we find

$$\lambda_{B_m(6),2,1} = f_2(6,q) - 8f_3(6,q) + 12f_4(6,q) = q^4 - 26q^3 + 257q^2 - 1148q + 1961$$
(2.58)

$$\lambda_{B_m(6),2,2} = f_2(6,q) - 2f_3(6,q) - 6f_4(6,q) = q^4 - 20q^3 + 149q^2 - 488q + 587$$
(2.59)

$$\lambda_{B_m(6),2,3} = f_2(6,q) + 2f_3(6,q) + 2f_4(6,q) = q^4 - 16q^3 + 97q^2 - 268q + 291$$
(2.60)

$$\lambda_{B_m(6),2,4} = f_2(6,q) - 4f_3(6,q) = q^4 - 22q^3 + 185q^2 - 708q + 1045$$
(2.61)

$$\lambda_{B_m(6),2,5} = f_2(6,q) + 2f_3(6,q) = q^4 - 16q^3 + 95q^2 - 246q + 229 .$$
(2.62)

For  $1 \leq j \leq 3$ , these  $\lambda_{B_m(6),2,j}$  correspond to the partition [2] and for j = 4, 5 to the partition [11].

For the eigenvalues with coefficients of degree 3 in q we obtain

$$\lambda_{B_m(6),3,1} = -f_3(6,q) + 9f_4(6,q) - 18f_5(6,q) + 6f_6(6,q) = -q^3 + 24q^2 - 194q + 529 \quad (2.63)$$

$$\lambda_{B_m(6),3,2} = -f_3(6,q) + 3f_4(6,q) + 6f_5(6,q) - 6f_6(6,q) = -q^3 + 18q^2 - 104q + 187 \quad (2.64)$$

$$\lambda_{B_m(6),3,3} = -f_3(6,q) - f_4(6,q) + 2f_5(6,q) + 6f_6(6,q) = -q^3 + 14q^2 - 64q + 99 \quad (2.65)$$

$$\lambda_{B_m(6),3,4} = -f_3(6,q) - 3f_4(6,q) - 6f_5(6,q) - 6f_6(6,q) = -q^3 + 12q^2 - 50q + 73 \quad (2.66)$$

$$\lambda_{B_m(6),3,5} = -f_3(6,q) + 6f_4(6,q) - 6f_5(6,q) = -q^3 + 21q^2 - 149q + 358$$
(2.67)

$$\lambda_{B_m(6),3,6} = -f_3(6,q) + 2f_4(6,q) + 2f_5(6,q) = -(q-6)(q^2 - 11q + 31)$$
(2.68)

$$\lambda_{B_m(6),3,7} = -f_3(6,q) + 6f_5(6,q) = -q^3 + 15q^2 - 71q + 100$$
(2.69)

$$\lambda_{B_m(6),3,8} = -f_3(6,q) - 3f_4(6,q) - 3f_5(6,q) = -q^3 + 12q^2 - 47q + 61$$
(2.70)

$$\lambda_{B_m(6),3,9} = -f_3(6,q) + 3f_4(6,q) = -q^3 + 18q^2 - 110q + 229$$
(2.71)

$$\lambda_{B_m(6),3,10} = -f_3(6,q) - 3f_4(6,q) = -q^3 + 12q^2 - 44q + 43.$$
(2.72)

Here the  $\lambda_{B_m(6),3,j}$  for  $1 \leq j \leq 4$  correspond to the partition [3], for  $5 \leq j \leq 8$  to the partition [21], and for j = 9, 10 to the partition [111].

Proceeding to the eigenvalues with coefficients of degree 4 in q we calculate

$$\lambda_{B_m(6),4,1} = f_4(6,q) - 8f_5(6,q) + 12f_6(6,q) = q^2 - 19q + 91$$
(2.73)

$$\lambda_{B_m(6),4,2} = f_4(6,q) - 2f_5(6,q) - 6f_6(6,q) = q^2 - 13q + 37$$
(2.74)

$$\lambda_{B_m(6),4,3} = f_4(6,q) + 2f_5(6,q) + 2f_6(6,q) = q^2 - 9q + 21$$
(2.75)

$$\lambda_{B_m(6),4,4} = f_4(6,q) - 6f_6(6,q) = q^2 - 11q + 25$$
(2.76)

$$\lambda_{B_m(6),4,5} = f_4(6,q) - 2f_5(6,q) - 2f_6(6,q) = q^2 - 13q + 41$$
(2.77)

$$\lambda_{B_m(6),4,6} = f_4(6,q) = q^2 - 11q + 31 \tag{2.78}$$

$$\lambda_{B_m(6),4,7} = f_4(6,q) + 3f_5(6,q) + 3f_6(6,q) = (q-4)^2$$
(2.79)

$$\lambda_{B_m(6),4,8} = f_4(6,q) - 6f_5(6,q) + 6f_6(6,q) = q^2 - 17q + 73$$
(2.80)

$$\lambda_{B_m(6),4,9} = f_4(6,q) + f_5(6,q) - 3f_6(6,q) = q^2 - 10q + 22$$
(2.81)

$$\lambda_{B_m(6),4,10} = f_4(6,q) + 4f_5(6,q) + 6f_6(6,q) = q^2 - 7q + 13$$
(2.82)

$$\lambda_{B_m(6),4,11} = f_4(6,q) - 4f_5(6,q) + 2f_6(6,q) = q^2 - 15q + 57$$
(2.83)

$$\lambda_{B_m(6),4,12} = f_4(6,q) + 2f_5(6,q) - 4f_6(6,q) = q^2 - 9q + 15$$
(2.84)

$$\lambda_{B_m(6),4,13} = f_4(6,q) - f_5(6,q) - f_6(6,q) = (q-6)^2$$
(2.85)

$$\lambda_{B_m(6),4,14} = f_4(6,q) + 4f_5(6,q) + 4f_6(6,q) = q^2 - 7q + 11$$
(2.86)

$$\lambda_{B_m(6),4,15} = f_4(6,q) - 2f_5(6,q) = q^2 - 13q + 43$$
(2.87)

$$\lambda_{B_m(6),4,16} = f_4(6,q) + 4f_5(6,q) = q^2 - 7q + 7 .$$
(2.88)

Here the  $\lambda_{B_m(6),4,j}$  for  $1 \leq j \leq 3$  correspond to the partition [4], for  $4 \leq j \leq 8$  to the partition [31], for  $9 \leq j \leq 11$  to the partition [22], for  $11 \leq j \leq 14$  to the partition [211], and for j = 15, 16 to the partition [1111]. Notice that the term  $\lambda_{B_m(6),4,11}$  involves a degeneracy from two different associated partitions.

For the eigenvalues with coefficients of degree 5 in q we find

$$\lambda_{B_m(6),5,1} = -f_5(6,q) + 5f_6(6,q) = -q + 11 \tag{2.89}$$

$$\lambda_{B_m(6),5,2} = -f_5(6,q) - f_6(6,q) = -q + 5 \tag{2.90}$$

$$\lambda_{B_m(6),5,3} = -f_5(6,q) + 4f_6(6,q) = -q + 10 \tag{2.91}$$

$$\lambda_{B_m(6),5,4} = -f_5(6,q) = -q + 6 \tag{2.92}$$

$$\lambda_{B_m(6),5,5} = -f_5(6,q) - 2f_6(6,q) = -q + 4 \tag{2.93}$$

$$\lambda_{B_m(6),5,6} = -f_5(6,q) + 3f_6(6,q) = -q + 9 \tag{2.94}$$

$$\lambda_{B_m(6),5,7} = -f_5(6,q) + f_6(6,q) = -q + 7 \tag{2.95}$$

$$\lambda_{B_m(6),5,8} = -f_5(6,q) - 3f_6(6,q) = -q + 3 \tag{2.96}$$

$$\lambda_{B_m(6),5,9} = -f_5(6,q) + 2f_6(6,q) = -q + 8 \tag{2.97}$$

$$\lambda_{B_m(6),5,10} = -f_5(6,q) - 4f_6(6,q) = -q + 2 \tag{2.98}$$

$$\lambda_{B_m(6),5,11} = -f_5(6,q) - 5f_6(6,q) = -q + 1 .$$
(2.99)

Here  $\lambda_{B_m(6),5,j}$  for j = 1, 2 correspond to the partition [5], for  $3 \leq j \leq 5$  to the partition [41], for  $5 \leq j \leq 7$  to the partition [32], for j = 4, 6, 8 to the partition [311], for j = 2, 8, 9 to the partition [221], for j = 4, 9, 10 to the partition [2111], and for j = 7, 11 to the partition [11111]. Notice that the terms  $\lambda_{B_m(6),5,j}$  for j = 2 and  $5 \leq j \leq 9$  involve degeneracies from two different associated partitions, and for j = 4 from three associated partitions.

Finally, for the eigenvalue with coefficient of degree 6 in q we have

$$\lambda_{B_m(6),6,1} = f_6(6,q) = 1 . (2.100)$$

This involves a degeneracy from all the possible partitions.

The coefficient  $c_{B_m(6),d,j}(q)c'_{B_m(6),d,j}$  of degree 0 in q is

$$c_{B_m(6),0,1}c'_{B_m(6),0,1} = 1 . (2.101)$$

The coefficients of degree 1 in q are

$$c_{B_m(6),1,1}c'_{B_m(6),1,1} = q - 1 \tag{2.102}$$

$$c_{B_m(6),1,2}c'_{B_m(6),1,2} = 5(q-1) , \qquad (2.103)$$

where  $c_{B_m(6),1,j} = q - 1$  for j = 1, 2, and  $c'_{B_m(6),1,j} = 1, 5$  for j = 1, 2.

The coefficients of degree 2 in q are listed in Table IV

TABLE IV. Coefficients of degree 2 in q for  $B_m(6)$  strip.

j	partition	$c_{B_m(6),2,j}^\prime$	$c_{B_m(6),2,j}c'_{B_m(6),2,j}$
1	[2]	1	$rac{1}{2}q(q-3)$
2	[2]	5	$rac{5}{2}q(q-3)$
3	[2]	9	$rac{9}{2}q(q-3)$
4	[11]	5	$\frac{5}{2}(q-1)(q-2)$
5	[11]	10	5(q-1)(q-2)

The coefficients of degree 3 in q are listed in Table V

j	partition	$c_{B_m(6),3,j}'$	$c_{B_m(6),3,j}c'_{B_m(6),3,j}$
1	[3]	1	$\frac{1}{6}q(q-1)(q-5)$
2	[3]	5	$\frac{5}{6}q(q-1)(q-5)$
3	[3]	9	$\frac{3}{2}q(q-1)(q-5)$
4	[3]	5	$\tfrac{5}{6}q(q-1)(q-5)$
5	[21]	10	$\frac{5}{3}q(q-2)(q-4)$
6	[21]	18	3q(q-2)(q-4)
7	[21]	20	$\frac{10}{3}q(q-2)(q-4)$
8	[21]	32	$\frac{16}{3}q(q-2)(q-4)$
9	[111]	10	$\frac{5}{3}(q-1)(q-2)(q-3)$
10	[111]	10	$\frac{5}{3}(q-1)(q-2)(q-3)$

TABLE V. Coefficients of degree 3 in q for  $B_m(6)$  strip.

# The coefficients of degree 4 in q are listed in Table VI

j	partition	$c_{B_m(6),4,j}^\prime$	$c_{B_m(6),4,j}c_{B_m(6),4,j}'$
1	[4]	1	$\frac{1}{24}q(q-1)(q-2)(q-7)$
2	[4]	5	$\frac{5}{24}q(q-1)(q-2)(q-7)$
3	[4]	9	$\frac{3}{8}q(q-1)(q-2)(q-7)$
4	[31]	30	$\frac{5}{4}q(q-1)(q-3)(q-6)$
5	[31]	27	$\frac{9}{8}q(q-1)(q-3)(q-6)$
6	[31]	15	$\frac{5}{8}q(q-1)(q-3)(q-6)$
7	[31]	48	2q(q-1)(q-3)(q-6)
8	[31]	15	$\frac{5}{8}q(q-1)(q-3)(q-6)$
9	[22]	32	$\frac{4}{3}q(q-1)(q-4)(q-5)$
10	[22]	10	$\frac{5}{12}q(q-1)(q-4)(q-5)$
11	[22]	18	$\frac{1}{2}q(q-5)(2q-3)(2q-7)$
	[211]	30	
12	[211]	30	$\frac{5}{4}q(q-2)(q-3)(q-5)$
13	[211]	48	2q(q-2)(q-3)(q-5)
14	[211]	27	$\frac{9}{8}q(q-2)(q-3)(q-5)$
15	[1111]	10	$\frac{5}{12}(q-1)(q-2)(q-3)(q-4)$
$1\overline{6}$	[1111]	5	$\frac{5}{24}(q-1)(q-2)(q-3)(q-4)$

TABLE VI. Coefficients of degree 4 in q for  $B_m(6)$  strip.

For d = 5, there are partitions [5], [41], [32], [311], [221], [2111] and [11111], and the associated  $c_{X,5}$  are  $c_{[5],5} = \frac{1}{120}q(q-1)(q-2)(q-3)(q-9)$ ,  $c_{[41],5} = \frac{1}{120}q(q-1)(q-2)(q-4)(q-8)$ ,  $c_{[32],5} = \frac{1}{120}q(q-1)(q-2)(q-5)(q-7)$ ,  $c_{[311],5} = \frac{1}{120}q(q-1)(q-3)(q-4)(q-7)$ ,  $c_{[221],5} = \frac{1}{120}q(q-1)(q-3)(q-5)(q-6)$ ,  $c_{[2111],5} = \frac{1}{120}q(q-2)(q-3)(q-4)(q-6)$ , and  $c_{[11111],5} = \frac{1}{120}(q-1)(q-2)(q-3)(q-4)(q-5)$ . We list the coefficients of degree 5 in q in Table VII

j	partition	$c_{B_m(6),5,j}^\prime$	$c_{B_m(6),5,j}c_{B_m(6),5,j}'$
1	[5]	1	$\frac{1}{120}q(q-1)(q-2)(q-3)(q-9)$
2	[5]	5	$\frac{1}{4}q(q-1)(q-3)(q-4)(q-7)$
	[221]	25	
3	[41]	20	$\frac{1}{6}q(q-1)(q-2)(q-4)(q-8)$
4	[41]	36	$\frac{1}{5}q(q-4)(7q^3-77q^2+217q-162)$
	[311]	96	
	[2111]	36	
5	[41]	40	$q(q-1)(q-2)(q^2-12q+34)$
	[32]	80	
6	[32]	45	$\frac{1}{8}q(q-1)(q-7)(7q^2-49q+78)$
	[311]	60	
7	[32]	25	$\frac{1}{4}(q-1)(q-2)(q-5)(q^2-7q+2)$
	[11111]	5	
8	[311]	60	$\frac{1}{8}q(q-1)(q-3)(7q^2-77q+202)$
	[221]	45	
9	[221]	80	$q(q-3)(q-6)(q^2-6q+6)$
	[2111]	40	
10	[2111]	20	$\frac{1}{6}q(q-2)(q-3)(q-4)(q-6)$
11	[11111]	1	$\frac{1}{120}(q-1)(q-2)(q-3)(q-4)(q-5)$

TABLE VII. Coefficients of degree 5 in q for  $B_m(6)$  strip.

Finally, the coefficient of degree 6 is

$$c_{B_m(6),6,1}c'_{B_m(6),6,1} = q^6 - 21q^5 + 160q^4 - 545q^3 + 814q^2 - 415q + 1.$$
(2.104)

The sum of all the coefficients is equal to

$$C(B_m(6),q) = \sum_{d=0}^{6} \sum_{j=1}^{N_{B_m(6),d,\lambda}} c_{B_m(6),d,j}(q) c'_{B_m(6),d,j}$$
  
=  $q(q-1)(q-2)(q-3)(q-4)(q-5) = P(K_6,q)$ . (2.105)

The chromatic number is  $\chi(B_m(6)) = 6$ .

The locus  $\mathcal{B}$  and chromatic zeros for the  $B_m(6)$  strip with  $L_x = m = 15$  are shown in Fig. 2. The locus  $\mathcal{B}$  crosses the real q-axis at q = 0, 2, 4 and  $q_c(B_m(6))$ , where

$$q_c(B_m(6)) \simeq 5.3236 \cdots$$
 (2.106)



FIG. 2. Locus  $\mathcal{B}$  for the  $m \to \infty$  limit of the family  $B_m(6)$  with toroidal boundary conditions and chromatic zeros for  $B_m(6)$  with m = 15 (i.e., n = 90).

The locus  $\mathcal{B}$  has support for  $Re(q) \geq 0$ , and separates the q plane into four regions. The outermost one, region  $R_1$ , extends to infinite |q| and includes the intervals  $q \geq q_c(B_m(6))$  and  $q \leq 0$  on the real q axis. Region  $R_2$  includes the real interval  $4 \leq q \leq q_c(B_m(6))$ , region  $R_3$  includes the real interval  $2 \leq q \leq 4$ , while region  $R_4$  includes the real interval  $0 \leq q \leq 2$ . In regions  $R_i$ ,  $1 \leq i \leq 4$ , the dominant terms are  $\lambda_{B_m(6),0,1}$ ,  $\lambda_{B_m(6),3,1}$ ,  $\lambda_{B_m(6),2,1}$ , and  $\lambda_{B_m(6),1,1}$ , respectively. Thus, the  $q_c(B_m(6))$  given in (2.106) is the degeneracy between  $|\lambda_{B_m(6),0,1}|$  and  $|\lambda_{B_m(6),3,1}|$ , and is the real solution of  $q^5 - 19q^4 + 152q^3 - 660q^2 + 1600q - 1761 = 0$ .

#### **III. FAMILY OF KLEIN BOTTLE STRIPS**

Consider the graph with  $L_x$  set of  $K_b$  with  $L_x - 1$  edge linking sets  $\{11, 22, ..., bb\}$  and one edge linking set  $\{1b, 2(b-1), ..., b1\}$ . This is the Klein bottle strip and will be denoted as  $\overline{B}_m(b)$ . The chromatic polynomials has the same form as given in eq. (2.1), and the  $\lambda$ 's are the same as those of the corresponding toroidal strip with the same b. The only difference is the coefficients.

### A. $L_y = 5$ Klein bottle strip

For b = 5 considered here, the coefficients  $c_{\bar{B}_m(5),d,j}(q)c'_{\bar{B}_m(5),d,j}$  are summarized in Table VIII

d	j	partition	$c'_{ar{B}_m(5),d,j}$	$c_{ar{B}_m(5),d,j}c'_{ar{B}_m(5),d,j}$
0	1	[0]	1	1
1	1	[1]	1	q-1
1	2	[1]	0	0
<b>2</b>	1	[2]	1	$rac{1}{2}q(q-3)$
<b>2</b>	2	[2]	0	0
<b>2</b>	3	[2]	1	$rac{1}{2}q(q-3)$
<b>2</b>	4	[11]	0	0
<b>2</b>	5	[11]	-2	-(q-1)(q-2)

TABLE VIII. Coefficients for  $B_m(5)$  strip.

3	1	[3]	1	$\frac{1}{6}q(q-1)(q-5)$
3	2	[3]	0	0
3	3	[3]	1	$\frac{1}{6}q(q-1)(q-5)$
3	4	[21]	0	0
3	5	[21]	2	$\frac{1}{3}q(q-2)(q-4)$
3	6	[21]	-4	$-\frac{2}{3}q(q-2)(q-4)$
3	7	[21]	2	$\frac{1}{3}q(q-2)(q-4)$
3	8	[111]	-2	$-\frac{1}{3}(q-1)(q-2)(q-3)$
3	9	[111]	0	0
4	1	[4]	1	$\frac{1}{24}q(q-1)(q-2)(q-7)$
4	2	[4]	0	0
4	3	[31]	0	0
4	4	[211]	0	0
4	5	[31]	-6	$-\frac{1}{6}q(q-1)(q^2-9q+17)$
		[22]	2	
4	6	[31]	3	$\frac{1}{4}q(q-3)(q^2-7q+8)$
		[211]	3	
4	7	[211]	-6	$-\frac{1}{6}q(q-5)(q^2-5q+7)$
		[22]	2	
4	8	[1111]	0	0
4	9	[1111]	1	$\frac{1}{24}(q-1)(q-2)(q-3)(q-4)$
5	1	see text		q-1

where for d = 5, it involves all possible partitions with appropriate  $c'_{\bar{B}_m(5),5,j}$  which are not listed here. Notice that the coefficients for  $\lambda_{\bar{B}_m(5),d,j}$  with (d, j) = (1, 2), (2, 2), (2, 4), (3, 2),(3, 4), (3, 9), (4, 2), (4, 3), (4, 4), (4, 8) are zero, i.e., these ten  $\lambda$ 's do not contribute to the chromatic polynomial of the Klein bottle strip. We thus find that the total number of distinct  $\lambda$ 's for the b = 5 Klein bottle strip with indicated linking is

$$N_{P,\bar{B}_m(5),\lambda} = 17 . (3.1)$$

The sum of all of the coefficients for the Klein bottle strip is

$$C(\bar{B}_m(5),q) = \sum_{d=0}^{5} \sum_{j=1}^{N_{\bar{B}_m(5),d,\lambda}} c_{\bar{B}_m(5),d,j}(q) c'_{\bar{B}_m(5),d,j} = 0 , \qquad (3.2)$$

which is easy to understand by the coloring argument.

# **B.** $L_y = 6$ Klein bottle strip

For b = 6, the coefficients  $c_{\bar{B}_m(6),d,j}(q)c'_{\bar{B}_m(6),d,j}$  are summarized in Table IX

d	j	partition	$c'_{ar{B}_m(6),d,j}$	$c_{ar{B}_m(6),d,j}c'_{ar{B}_m(6),d,j}$
0	1	[0]	1	1
1	1	[1]	1	q-1
1	2	[1]	-1	-(q-1)
2	1	[2]	1	$\frac{1}{2}q(q-3)$
2	2	[2]	-1	$-\frac{1}{2}q(q-3)$
2	3	[2]	3	$\frac{3}{2}q(q-3)$
2	4	[11]	-1	$-\frac{1}{2}(q-1)(q-2)$
2	5	[11]	-2	-(q-1)(q-2)
3	1	[3]	1	$\frac{1}{6}q(q-1)(q-5)$
3	2	[3]	-1	$-\frac{1}{6}q(q-1)(q-5)$
3	3	[3]	3	$\frac{1}{2}q(q-1)(q-5)$
3	4	[3]	-3	$-\frac{1}{2}q(q-1)(q-5)$
3	5	[21]	-2	$-\frac{1}{3}q(q-2)(q-4)$
3	6	[21]	6	q(q-2)(q-4)
3	7	[21]	-4	$-\frac{2}{3}q(q-2)(q-4)$
3	8	[21]	0	0
3	9	[111]	-2	$-\frac{1}{3}(q-1)(q-2)(q-3)$
3	10	[111]	2	$\frac{1}{3}(q-1)(q-2)(q-3)$
4	1	[4]	1	$\frac{1}{24}q(q-1)(q-2)(q-7)$
4	2	[4]	-1	$-\frac{1}{24}q(q-1)(q-2)(q-7)$
4	3	[4]	3	$\frac{1}{8}q(q-1)(q-2)(q-7)$
4	4	[31]	-6	$-\frac{1}{4}q(q-1)(q-3)(q-6)$
4	5	[31]	9	$\frac{3}{8}q(q-1)(q-3)(q-6)$
4	6	[31]	-9	$-\frac{3}{8}q(q-1)(q-3)(q-6)$
4	7	[31]	0	0
4	8	[31]	$-\overline{3}$	$-\frac{1}{8}q(q-1)(q-3)(q-6)$
4	9	[22]	0	0
4	10	[22]	6	$\frac{1}{4}q(q-1)(q-4)(q-5)$
4	11	[22]	6	$-\frac{1}{2}q(q-5)$

TABLE IX. Coefficients for $\bar{B}_m(6)$ strip.	TABLE IX.	Coefficients for	$\bar{B}_m(6)$	strip.
--	-----------	------------------	----------------	--------

		[211]	-6	
4	12	[211]	6	$\frac{1}{4}q(q-2)(q-3)(q-5)$
4	13	[211]	0	0
4	14	[211]	-9	$-\frac{3}{8}q(q-2)(q-3)(q-5)$
4	15	[1111]	2	$\frac{1}{12}(q-1)(q-2)(q-3)(q-4)$
4	16	[1111]	1	$\frac{1}{24}(q-1)(q-2)(q-3)(q-4)$
5	1	[5]	1	$\frac{1}{120}q(q-1)(q-2)(q-3)(q-9)$
5	2	[5]	-1	$\frac{1}{60}q(q-1)(q-3)(7q^2-77q+216)$
		[221]	15	
5	3	[41]	-4	$-\frac{1}{30}q(q-1)(q-2)(q-4)(q-8)$
5	4	[41]	12	-q(q-2)(q-4)
		[2111]	-12	
5	5	[41]	-8	$-\frac{1}{15}q(q-1)(q-2)(q-4)(q-8)$
5	6	[32]	15	$\frac{1}{40}q(q-1)(q-7)(q^2-7q+2)$
		[311]	-12	
5	7	[32]	-15	$-\frac{1}{60}(q-1)(q-2)(q-5)(7q^2-49q-6)$
		[11111]	1	
5	8	[311]	12	$-\frac{1}{40}q(q-1)(q-3)(q^2-11q+38)$
		[221]	-15	
5	9	[2111]	8	$\frac{1}{15}q(q-2)(q-3)(q-4)(q-6)$
5	10	[2111]	4	$\frac{1}{30}q(q-2)(q-3)(q-4)(q-6)$
5	11	[11111]	-1	$-\frac{1}{120}(q-1)(q-2)(q-3)(q-4)(q-5)$
6	1	see text		-1

where for d = 6, it involves all possible partitions with appropriate  $c'_{\bar{B}_m(6),6,j}$  which are not listed here. Notice that the coefficients for  $\lambda_{\bar{B}_m(6),d,j}$  with (d,j) = (3,8), (4,7), (4,9), (4,13)are zero, i.e., these four  $\lambda$ 's do not contribute to the chromatic polynomial of the Klein bottle strip. We thus find that the total number of distinct  $\lambda$ 's for the b = 6 Klein bottle strip with indicated linking is

$$N_{P,\bar{B}_m(6)} = 42 . (3.3)$$

The sum of all of the coefficients for the Klein bottle strip is again zero,

$$C(\bar{B}_m(6),q) = \sum_{d=0}^{6} \sum_{j=1}^{N_{\bar{B}_m(6),d,\lambda}} c_{\bar{B}_m(6),d,j}(q) c'_{\bar{B}_m(6),d,j} = 0 .$$
(3.4)

### IV. PROPERTIES OF EIGENVALUES, COEFFICIENTS, AND $\mathcal{B}$

We first list the number of eigenvalues,  $N_{B_m(b),d,\lambda}$ , for  $1 \leq b \leq 6$ ,  $0 \leq d \leq b$ , the total number of eigenvalues,  $N_{P,B_m(b),\lambda}$ , and the points at which  $\mathcal{B}$  crosses the real q axis, as well as the corresponding values for the Klein bottle strips in Table X.

TABLE X. Properties of P and  $\mathcal{B}$  for strip graphs  $B_m(b)$  and  $\overline{B}_m(b)$ . The properties apply for a given strip of size  $L_y \times L_x$ ; some apply for arbitrary  $L_x$ , such as  $N_{P,B_m(b),\lambda}$ , while others apply for the infinite-length limit, such as the properties of the locus  $\mathcal{B}$ . For the boundary conditions in the y and x directions  $(BC_y, BC_x)$ , P, and T denote periodic, and orientation-reversed (twisted) periodic.  $N_{B_m(b),d,\lambda}$ 's are abbreviated as  $N_{d,\lambda}$ ,  $N_{P,B_m(b),\lambda}$  is abbreviated as  $N_{P,\lambda}$ , and similarly for  $\overline{B}_m(b)$  strips. The blank entries are zero. The column denoted BCR lists the points at which  $\mathcal{B}$  crosses the real q axis; the largest of these is  $q_c(B_m(b))$ for the given family  $B_m(b)$  or  $\overline{B}_m(b)$ . Column labelled "SN" refers to whether  $\mathcal{B}$  has support for negative Re(q), indicated as yes (y) or no (n).

b	$BC_y$	$BC_x$	$N_{0,\lambda}$	$N_{1,\lambda}$	$N_{2,\lambda}$	$N_{3,\lambda}$	$N_{4,\lambda}$	$N_{5,\lambda}$	$N_{6,\lambda}$	$N_{P,\lambda}$	BCR	SN
1	Р	Р	1	1						2	0, 2	n
2	Р	Р	1	2	1					4	0, 2	n
2	Р	TP	1	2	1					4	0, 2	n
3	Р	Р	1	2	4	1				8	0, 2, 3	n
3	Р	TP	1	1	2	1				5	0,2,3	n
4	Р	Р	1	2	5	7	1			16	0, 2, 3.67	n
4	Р	TP	1	2	5	7	1			16	0, 2, 3.67	n
5	Р	Р	1	2	5	9	9	1		27	0, 2, 4, 4.51	n
5	Р	TP	1	1	3	6	5	1		17	0, 2, 4, 4.51	n
6	Р	Р	1	2	5	10	16	11	1	46	$0, 2, \overline{4}, 5.32$	n
6	Р	TP	1	2	5	9	13	11	1	42	0, 2, 4, 5.32	n

We conjecture the following q value at which  $\mathcal{B}$  crosses the real axis for the  $B_m(b)$  strips,

$$\mathcal{B} \supset \{q = 0, 2, 4, ..., 2[\frac{b-1}{2}], q_c(B_m(b))\} .$$
(4.1)

where  $[\nu]$  denotes the integral part of  $\nu$ , and the dominant eigenvalue in interval between 2(d'-1) and 2d' is  $\lambda_{B_m(b),d',1}$  for  $1 \leq d' \leq [\frac{b+1}{2}]$ .

The upper and lower bounds for the eigenvalues with coefficients of degree b - 1 can be determined in the following theorem.

**Theorem 1** The eigenvalues with coefficients of degree b-1 are bounded between  $(-1)^{b-1}(q-1)$  and  $(-1)^{b-1}(q-2b+1)$ .

Proof. Denote the  $b! \times b!$  matrix for the eigenvalues with coefficients of degree b-1 as  $M_{b-1}$ . By theorem 2 of [58], the diagonal elements of  $M_{b-1}$  are all the same to be  $(-1)^{b-1}f_{b-1}(b,q) = (-1)^{b-1}(q-b)$ , and other non-zero elements of  $M_{b-1}$  are all the same to be  $(-1)^b f_b(b,q) = (-1)^b$ . Now consider  $M'_{b-1} = M_{b-1} - (-1)^{b-1}(q-b)I$ , where I is the  $b! \times b!$  identity matrix. The number of non-zero elements,  $(-1)^b$ , in every row and columns of  $M'_{b-1}$  is b-1 by the construction of the matrix. If we denote the eigenvalues of  $M'_{b-1}$  as  $\lambda'$  and the normalized eigenvector of  $M'_{b-1}$  as  $W^T = (w_1, w_2, ..., w_{b!})$ , then we have

$$M'_{b-1}W = \lambda'W$$
  

$$\Rightarrow |M'_{b-1}W|^2 = |\lambda'W|^2$$
  

$$\Rightarrow (b-1)(w_1^2 + w_2^2 + \dots w_{b!}^2) + 2\sum_{i,j,i\neq j} w_i w_j = \lambda'^2 (w_1^2 + w_2^2 + \dots w_{b!}^2)$$
  

$$\Rightarrow \lambda'^2 = b - 1 + 2\sum_{i,j,i\neq j} w_i w_j ,$$
(4.2)

where we sum all the possible multiplications of two different  $w_i$  and  $w_j$  in each row of  $M'_{b-1}$ . Therefore,

$$\lambda^{\prime 2} \le b - 1 + \sum_{i,j,i \ne j} (w_i^2 + w_j^2)$$
  
= b - 1 + (b - 1)(b - 2) = (b - 1)^2, (4.3)

that is,  $-(b-1) \leq \lambda' \leq b-1$ . The upper and lower bounds for the eigenvalues of  $M_{b-1}$  are

$$(-1)^{b-1}(q-b) \pm (b-1) = \begin{cases} (-1)^{b-1}(q-1) \\ (-1)^{b-1}(q-2b+1) \end{cases}$$
(4.4)

It is easy to see that  $(-1)^{b-1}(q-2b+1)$  is one of the principal terms and  $(-1)^{b-1}(q-1)$  is one of the alternating terms given in [58]. It follows that these values are realized as the largest and smallest values for the eigenvalues of  $M_{b-1}$   $\Box$ .

We shall show that the locus  $\mathcal{B}$  crosses the real q axis at q = 0, and doesn't cross any  $q \in \mathbb{R}_{-}$ . We need the following lemma.

**Lemma 1** On the non-positive real q axis, the dominant eigenvalue with coefficient of degree d is

$$\lambda_{B_m(b),d} = (-1)^d \sum_{j=0}^d (-1)^j \binom{d}{j} (b-d)^{(j)} f_{d+j}(b,q) .$$
(4.5)

Proof. The matrix for the eigenvalues with coefficients of degree d can be written as [58]  $M_d = (-1)^d f_d(b,q)I + (-1)^{d+1}f_{d+1}(b,q)M_{d,1} + (-1)^{d+2}f_{d+2}(b,q)M_{d,2} + ...,$  where the non-zero elements are 1 in every  $M_{d,j}$ , and the number of these in each row and column is  $\binom{d}{j}(b-d)^{(j)}$ . By the same reason given in Theorem 1, the magnitudes of the eigenvalues of  $M_{d,j}$  are bounded above by  $\binom{d}{j}(b-d)^{(j)}$ . Since the signs of  $f_{d+j}(b,q)$  alternate for even j and odd j for any non-positive q value,  $\lambda_{B_m(b),d}$  given in eq. (4.5) is the upper bound for the eigenvalues with coefficients of degree d. Since  $\lambda_{B_m(b),d}$  is the eigenvalue with  $c'_{B_m(b),d,j} = 1$  given in [58], the proof is completed.  $\Box$ .

**Theorem 2** The minimum real q at which  $\mathcal{B}$  crosses the real q axis is q = 0.

*Proof.* We will begin with the proof that  $|\lambda_{B_m(b),0}| > |\lambda_{B_m(b),1}| > ...$  on the whole negative real q axis. The first few examples of  $\lambda_{B_m(b),d}$ 's are

$$\lambda_{B_m(b),0} = f_0(b,q)$$

 $\lambda_{B_m(b),1} = -f_1(b,q) + (b-1)f_2(b,q)$ 

$$\lambda_{B_m(b),2} = f_2(b,q) - 2(b-2)f_3(b,q) + (b-2)(b-3)f_4(b,q)$$

$$\lambda_{B_m(b),3} = -f_3(b,q) + 3(b-3)f_4(b,q) - 3(b-3)(b-4)f_5(b,q) + (b-3)(b-4)(b-5)f_6(b,q)$$

$$\lambda_{B_m(b),4} = f_4(b,q) - 4(b-4)f_5(b,q) + 6(b-4)(b-5)f_6(b,q) - 4(b-4)(b-5)(b-6)f_7(b,q)$$

$$+(b-4)(b-5)(b-6)(b-7)f_8(b,q) . (4.6)$$

To show that  $f_0(b,q)$  has the largest magnitude on the negative real q axis, we only need  $|f_i(b,q)| > |f_{i+1}(b,q) - (b-i-1)f_{i+2}(b,q)|$  as follows,

$$\begin{split} |f_{i}(b,q)| &= |f_{i+1}(b,q) - (b-i-1)f_{i+2}(b,q)| \\ &= (-1)^{b-i}f_{i}(b,q) - (-1)^{b-i-1}\Big[f_{i+1}(b,q) - (b-i-1)f_{i+2}(b,q)\Big] \\ &= (-1)^{b-i}\Big[\sum_{s=0}^{b-i}(-1)^{s}\binom{b-i}{s}(q-i-s)^{(b-i-s)} + f_{i+1}(b,q) - (b-i-1)f_{i+2}(b,q)\Big] \\ &= (-1)^{b-i}\Big[(q-i)^{(b-i)} + \sum_{s=0}^{b-i-1}(-1)^{s+1}\binom{b-i}{s+1}(q-i-s-1)^{(b-i-s-1)} \\ &+ \sum_{s=0}^{b-i-1}(-1)^{s}\binom{b-i-1}{s}(q-i-1-s)^{(b-i-1-s)} - (b-i-1)f_{i+2}(b,q)\Big] \\ &= (-1)^{b-i}\Big[(q-i)^{(b-i)} + (-1)^{b-i} + (-1)^{b-i-1} + \sum_{s=0}^{b-i-2}(-1)^{s+1}(q-i-1-s)^{(b-i-1-s)} \\ &\times \Big(\frac{(b-i)!}{(s+1)!(b-i-s-1)!} - \frac{(b-i-1)!}{s!(b-i-1-s)!}\Big) - (b-i-1)f_{i+2}(b,q)\Big] \\ &= (-1)^{b-i}\Big[(q-i)^{(b-i)} + \sum_{s=0}^{b-i-2}(-1)^{s+1}(q-i-1-s)^{(b-i-1-s)}\binom{b-i-1}{s+1} \\ &- (b-i-1)\sum_{s=0}^{b-i-2}(-1)^{s}\binom{b-i-2}{s}(q-i-2-s)^{(b-i-2-s)}\Big] \\ &= (-1)^{b-i}\Big[(q-i)^{(b-i)} + \sum_{s=0}^{b-i-2}(-1)^{s+1}(q-i-2-s)^{(b-i-2-s)} \\ &\times \Big[(q-i-1-s)\frac{(b-i-1)!}{(s+1)!(b-i-2-s)!} + (b-i-1)\frac{(b-i-2)!}{s!(b-i-2-s)!}\Big]\Big] \\ &= (-1)^{b-i}\Big[(q-i)^{(b-i)} + \sum_{s=0}^{b-i-2}(-1)^{s+1}(q-i-2-s)^{(b-i-2-s)} \\ &\times \Big[(q-i-1-s)\frac{(b-i-1)!}{(s+1)!(b-i-2-s)!} + (b-i-1)\frac{(b-i-2)!}{s!(b-i-2-s)!}\Big]\Big] \\ &= (-1)^{b-i}\Big[(q-i)^{(b-i)} + \sum_{s=0}^{b-i-2}(-1)^{s+1}(q-i-2-s)^{(b-i-2-s)} \\ &\times \Big[(q-i-1-s)\frac{(b-i-1)!}{(s+1)!(b-i-2-s)!} + (b-i-1)\frac{(b-i-2)!}{s!(b-i-2-s)!}\Big]\Big] \\ &= (-1)^{b-i}\Big[(q-i)^{(b-i)} + \sum_{s=0}^{b-i-2}(-1)^{s+1}(q-i-2-s)^{(b-i-2-s)} \\ &\times \Big[(q-i-1-s)\frac{(b-i-2)!}{(s+1)!(b-i-2-s)!} + (b-i-1)\frac{(b-i-2)!}{s!(b-i-2-s)!}\Big]\Big] \\ &= (-1)^{b-i}\Big[(q-i)^{(b-i)} + \sum_{s=0}^{b-i-2}(-1)^{s+1}(q-i-2-s)^{(b-i-2-s)} \\ &\times \Big[(q-i-1-s)\frac{(b-i-2)!}{(s+1)!(b-i-2-s)!} + (b-i-1)\frac{(b-i-2)!}{s!(b-i-2-s)!}\Big]\Big] \\ &= (-1)^{b-i}\Big[(q-i)^{(b-i)} + \sum_{s=0}^{b-i-2}(-1)^{s+1}(q-i-2-s)^{(b-i-2-s)} \\ &\times \Big[(b-i-2)\frac{(b-i-2)!}{s!} + \frac{(b-i-2)!}{s!} + \frac{(b-i-2)!}{s!} + \frac{(b-i-2)!}{s!}\Big] \\ &= (-1)^{b-i}\Big[(q-i)^{(b-i)} + \sum_{s=0}^{b-i-2}(-1)^{s+1}(q-i-2-s)^{(b-i-2-s)} \\ &= (-1)^{b-i}\Big[(d-i)^{(b-i)} + \frac{(b-i-2)!}{s!} + \frac{(b-i-2)!}{s!} + \frac{(b-i-2)!}{s!}\Big] \\ &= (-1)^{b-i}\Big[(b-i-2)\frac{(b-i-2)!}{s!} + \frac{(b-i-2)!}{s!} + \frac{(b-i-2)!}{s!} + \frac{(b-i-2)!}{s!} + \frac{(b-i-2)!}{s!}$$

For q = 0, all above argument are still valid, with the exception that  $|f_i(b,0)| = |f_{i+1}(b,0) - (b-i-1)f_{i+2}(b,0)|$  when i = 0, i.e.  $|\lambda_{B_m(b),0}| = |\lambda_{B_m(b),1}|$  for q = 0.  $\Box$ .

We will prove that the maximal value of q where  $\mathcal{B}$  intersects the (positive) real axis is smaller than q = b for  $b \ge 2$ . We have the following lemma.

**Lemma 2**  $f_i(b,q) \ge 0$  if  $q \ge b$ .

*Proof.* Since we have  $(q - i - s)^{(b-i-s)} > 0$  when  $q \ge b$ , one only need to show that the magnitudes of the terms in eq. (2.3) decrease for s from 0 to b - i.

$$\binom{b-i}{s} (q-i-s)^{(b-i-s)} - \binom{b-i}{s+1} (q-i-s-1)^{(b-i-s-1)}$$

$$= (q-i-s-1)^{(b-i-s-1)} \left[ \frac{(b-i)!}{s!(b-i-s)!} (q-i-s) - \frac{(b-i)!}{(s+1)!(b-i-s-1)!} \right]$$

$$= (q-i-s-1)^{(b-i-s-1)} \frac{(b-i)!}{(s+1)!(b-i-s)!} \left[ (s+1)(q-i-s) - (b-i-s) \right]$$

$$= (q-i-s-1)^{(b-i-s-1)} \frac{(b-i)!}{(s+1)!(b-i-s)!} \left[ s(q-i-s) + q-b \right]$$

$$\ge 0 \quad \text{for } 0 \le s \le b-i-1 \text{ and } q \ge b ,$$

$$(4.8)$$

where the equal sign is realized only when s = 0 and q = b. Therefore,  $f_i(b,q) = 0$  only when i = b - 1 at q = b, and  $f_i(b,q) > 0$  for any other  $0 \le i \le b$  and  $q \ge b$ .  $\Box$ .

It is known that  $q_c(B_m(1)) = 2$ ,  $q_c(B_m(2)) = 2$  [9,6], and  $q_c(B_m(3)) = 3$  [10]. Thus, for b = 1,  $q_c = b + 1$  while for b = 2 and for b = 3,  $q_c$  has the respective values  $q_c = b$ . It is of interest to ask how  $q_c(B_m(b))$  is related to b for  $b \ge 4$ . We determine this relation with the following theorem.

**Theorem 3**  $q_c(B_m(b)) < b$  for  $b \ge 4$ .

*Proof.* By a reason similar to that given in Lemma 1 and the result of Lemma 2, the magnitude of the eigenvalues with coefficients of degree d are bounded above by

$$\lambda'_{B_m(b),d} = \sum_{j=0}^d \binom{d}{j} (b-d)^{(j)} f_{d+j}(b,q) \quad \text{for } q \ge b , \qquad (4.9)$$

where it is equal to  $\lambda_{B_m(b),b-1,1}$  for d = b - 1 at q = b and equal to  $\lambda_{B_m(b),0,1}$  for d = 0 and any q. The first few examples of  $\lambda'_{B_m(b),d}$ 's are

$$\begin{split} \lambda'_{B_m(b),0} &= f_0(b,q) \\ \lambda'_{B_m(b),1} &= f_1(b,q) + (b-1)f_2(b,q) \\ \lambda'_{B_m(b),2} &= f_2(b,q) + 2(b-2)f_3(b,q) + (b-2)(b-3)f_4(b,q) \\ \lambda'_{B_m(b),3} &= f_3(b,q) + 3(b-3)f_4(b,q) + 3(b-3)(b-4)f_5(b,q) + (b-3)(b-4)(b-5)f_6(b,q) \\ \lambda'_{B_m(b),4} &= f_4(b,q) + 4(b-4)f_5(b,q) + 6(b-4)(b-5)f_6(b,q) + 4(b-4)(b-5)(b-6)f_7(b,q) \end{split}$$

$$+(b-4)(b-5)(b-6)(b-7)f_8(b,q) . (4.10)$$

To show that  $f_0(b,q) = \lambda_{B_m(b),0,1}$  has the largest magnitude for  $q \ge b$ , we first need  $f_i(b,q) > f_{i+1}(b,q) + (b-i-1)f_{i+2}(b,q)$  for  $0 \le i \le b-2$  and  $q \ge b$ ,

$$\begin{split} f_i(b,q) &= f_{i+1}(b,q) - (b-i-1)f_{i+2}(b,q) \\ &= \sum_{s=0}^{b-i} (-1)^s \binom{b-i}{s} (q-i-s)^{(b-i-s)} - \sum_{s=0}^{b-i-1} (-1)^s \binom{b-i-1}{s} (q-i-1-s)^{(b-i-1-s)} \\ &- (b-i-1)f_{i+2}(b,q) \\ &= (-1)^{b-i} + \sum_{s=0}^{b-i-1} (-1)^s (q-i-1-s)^{(b-i-1-s)} \frac{(b-i-1)!}{s!(b-i-s)!} \Big[ (b-i)(q-i-s) - (b-i-s) \Big] \\ &- (b-i-1)f_{i+2}(b,q) \\ &= (-1)^{b-i} + \sum_{s=0}^{b-i-1} (-1)^s (q-i-1-s)^{(b-i-1-s)} \Big[ \binom{b-i-1}{s} (b-i-1) + \binom{b-i}{s} (q-b) \Big] \\ &- (b-i-1)f_{i+2}(b,q) \\ &= (-1)^{b-i} + (-1)^{b-i-1} \Big[ b-i-1 + (b-i)(q-b) \Big] + \sum_{s=0}^{b-i-2} (-1)^s (q-i-2-s)^{(b-i-2-s)} \\ &\times \Big[ (q-i-1-s) \Big[ \binom{b-i-1}{s} (b-i-1) + \binom{b-i}{s} (q-b) \Big] - (b-i-1) \binom{b-i-2}{s} \Big] \\ &= (-1)^{b-i} + (-1)^{b-i-1} \Big[ b-i-1 + (b-i)(q-b) \Big] + \sum_{s=0}^{b-i-2} (-1)^s (q-i-2-s)^{(b-i-2-s)} \\ &\times \Big[ (q-i) - 1 + (b-i)(q-b) \Big] + \sum_{s=0}^{b-i-2} (-1)^s (q-i-2-s)^{(b-i-2-s)} \\ &\times \Big[ (q-b) \Big[ \binom{b-i-1}{s} (b-i-1) + \binom{b-i}{s} (q-b) \Big] + (b-i-1-s) \binom{b-i-2}{s} \Big] \\ &+ (b-i-1)(b-i-2) \binom{b-i-2}{s} \Big] \Big] \end{split}$$

 $\geq 0 \quad \text{for } 0 \leq i \leq b-2 \text{ and } q \geq b ,$ 

where the equal sign is realized only when i = b - 2 at q = b. Therefore, we have  $\lambda'_{B_m(b),0} > \lambda'_{B_m(b),1} > \ldots > \lambda'_{B_m(b),b/2}$  for even b and  $\lambda'_{B_m(b),0} > \lambda'_{B_m(b),1} > \ldots > \lambda'_{B_m(b),(b-1)/2}$  for odd b. Recall  $f_i(b,q) = 0$  if b < i. To show  $\lambda'_{B_m(b),b/2} > \ldots > \lambda'_{B_m(b),b}$  for even b and  $\lambda'_{B_m(b),(b+1)/2} > \ldots > \lambda'_{B_m(b),b}$  for odd b, we must have the coefficient of  $f_b(b,q)$  in  $\lambda'_{B_m(b),d}$  is smaller than the sum of the coefficients of  $f_{b-2}(b,q)$  and  $f_b(b,q)$  in  $\lambda'_{B_m(b),d-1}$  as follows,

(4.11)

$$\binom{d-1}{b-d-1} (b-d+1)^{(b-d-1)} + \binom{d-1}{b-d+1} (b-d+1)^{(b-d+1)} - \binom{d}{b-d} (b-d)^{(b-d)}$$

$$= \frac{(d-1)!(b-d+1)!}{(b-d-1)!(2d-b)!2} + \frac{(d-1)!(b-d+1)!}{(b-d+1)!(2d-b-2)!} - \frac{d!(b-d)!}{(b-d)!(2d-b)!}$$

$$= \frac{(d-1)!}{(2d-b)!} \frac{(b-d+1)(b-d)}{2} + \frac{(d-1)!}{(2d-b-2)!} - \frac{d!}{(2d-b)!}$$

$$= \frac{(d-1)!}{2(2d-b)!} \left[ (b-d+1)(b-d) + 2(2d-b)(2d-b-1) - 2d \right]$$

$$= \frac{(d-1)!}{2(2d-b)!} (3b^2 - 10bd + 9d^2 + 3b - 7d) ,$$

$$(4.12)$$

which is larger than 0 for even b with  $b/2 < d \le b$  and odd b with  $(b+1)/2 < d \le b$  except for (b,d) = (2,2) and (4,3) where it is 0. For odd b and d = (b+1)/2, there is no  $f_b(b,q)$ term in  $\lambda'_{B_m(b),(b-1)/2}$ , and we need the coefficient of the  $f_b(b,q)$  in  $\lambda'_{B_m(b),(b+1)/2}$  is smaller than the coefficient of  $f_{b-2}(b,q)$  in  $\lambda'_{B_m(b),(b-1)/2}$ ,

$$\binom{d-1}{b-d-1} (b-d+1)^{(b-d-1)} - \binom{d}{b-d} (b-d)^{(b-d)}$$

$$= \frac{(d-1)!(b-d+1)!}{2(b-d-1)!(2d-b)!} - \frac{d!(b-d)!}{(b-d)!(2d-b)!}$$

$$= \frac{(d-1)!}{2(2d-b)!} \Big[ (b-d+1)(b-d) - 2d \Big]$$

$$= \frac{1}{2} (\frac{b-1}{2})! \Big[ (\frac{b+1}{2})(\frac{b-1}{2}) - (b+1) \Big]$$

$$= \frac{1}{8} (\frac{b-1}{2})!(b-5)(b+1) \ge 0 \quad \text{for } b \ge 5 .$$

$$(4.13)$$

Therefore, for  $b \ge 4$ , we find  $\lambda_{B_m(b),0,1}$  is dominant for  $q \ge b$ , i.e.  $q_c(B_m(b)) < b$ .  $\Box$ .

We remark that our theorem, in combination with the known results that  $q_c(B_m(b)) = b$ for b = 2, 3, yields the theorem that

$$q_c(B_m(b)) \le b \quad \text{for} \quad b \ge 2 \ . \tag{4.14}$$

We observe that the length between b and  $q_c(B_m(b))$  are -1, 0, 0, 0.3264, 0.4937, and 0.6764 for the respective values  $1 \le b \le 6$ . These values suggest that this length might well increase monotonically up to 1 as  $b \to \infty$ , since the lower bound for  $q_c(B_m(b))$  is b-1 [62].

We observe the following general form of four  $\lambda_{B_m(b),3,j}$  corresponding to the partition [21],

$$\lambda_{B_m(b),3,j_1} = -f_3(b,q) + 2(b-3)f_4(b,q) - (b-3)(b-4)f_5(b,q)$$
(4.15)

$$\lambda_{B_m(b),3,j_2} = -f_3(b,q) + (b-4)f_4(b,q) + (b-4)f_5(b,q)$$
(4.16)

$$\lambda_{B_m(b),3,j_3} = -f_3(b,q) + (b-6)f_4(b,q) + 3(b-4)f_5(b,q)$$
(4.17)

$$\lambda_{B_m(b),3,j_4} = -f_3(b,q) - 3f_4(b,q) - 3f_5(b,q) .$$
(4.18)

These formulae are correct for  $3 \le b \le 9$ , and we conjecture that these are the only  $\lambda_{B_m(b),3,j}$  corresponding to the partition [21] for arbitrarily b. For toroidal strips, their coefficients are  $c_{B_m(b),3,j} = \frac{1}{6}q(q-2)(q-4)$ , and  $c'_{B_m(b),3,jk} = 2(b-1), b(b-3), (b-1)(b-2)$ , and  $\frac{2}{3}b(b-2)(b-4)$  for  $1 \le k \le 4$ .

From toroidal strips to Klein bottle strips, we observe the transformations from  $c'_{B_m(b),d,j}$  to  $c'_{\bar{B}_m(b),d,j}$  have certain patterns for  $0 \le d \le 4$  and  $2 \le b \le 6$ , and summarized them in table XI. Notice that the results are different for even and odd b.

partition	d	$c^{\prime}_{B_m(b),d,j}$	$c'_{\bar{B}_m(b),d,j}$ for odd $b$	$c'_{\bar{B}_m(b),d,j}$ for even $b$
[d]	$d \ge 0$	1	1	1
	$d \ge 1$	b-1	0	-1
	$d \geq 2$	b(b-3)/2	(b-3)/2	b/2
	$d \geq 3$	b(b-1)(b-5)/6	0	-b/2
	$d \geq 4$	b(b-1)(b-2)(b-7)/24	(b-1)(b-7)/8	b(b-2)/8
$[1^d]$	2	b - 1	0	-1
	$^{2,3}$	(b-1)(b-2)/2	-(b-1)/2	-(b-2)/2
	3,4	(b-1)(b-2)(b-3)/6	0	(b-2)/2
	$^{4,5}$	(b-1)(b-2)(b-3)(b-4)/24	(b-1)(b-3)/8	(b-2)(b-4)/8
[21]	3	2(b-1)	0	-2
	3	b(b-3)	b-3	b
	3	(b-1)(b-2)	-(b-1)	-(b-2)
	3	2b(b-2)(b-4)/3	2	0
[31]	4	3(b-1)	0	-3
	4	3b(b-3)/2	3(b-3)/2	3b/2
	4	3(b-1)(b-2)/2	-3(b-1)/2	-3(b-2)/2
	4	b(b-1)(b-5)/2	0	-3b/2
	4	b(b-2)(b-4)	3	0
	4	3b(b-1)(b-3)(b-6)/8	-3(b-1)(b-3)/8	-3b(b-6)/8
[22]	4	b(b-3)	b-3	b
	4	2b(b-2)(b-4)/3	2	0
	4	b(b-1)(b-4)(b-5)/6	(b-1)(b-5)/2	b(b-4)/2
[211]	4	3(b-1)(b-2)/2	-3(b-1)/2	-3(b-2)/2
	4	(b-1)(b-2)(b-3)/2	0	3(b-2)/2
	4	b(b-2)(b-4)	3	0
	4	3b(b-2)(b-3)(b-5)/8	-3(b-3)(b-5)/8	-3b(b-2)/8

TABLE XI. Relation of  $c'_{B_m(b),d,j}$  and  $c'_{\bar{B}_m(b),d,j}$ 

For the partition [d], if we label  $c'_{B_m(b),d\geq 0}$  as the  $c'_{B_m(b),d,j} = 1$  in the first row of table XI, and similarly for other  $c'_{B_m(b),d\geq d'}$ , we conjecture the following forms of the corresponding  $c'_{\bar{B}_m(b),d\geq d'}$  for general d',

$$c'_{\bar{B}_m(b),d\geq d'} = \begin{cases} \binom{(b-1)/2}{d'/2} - \binom{(b-1)/2}{d'/2-1} & \text{for even } d' \geq 2\\ 0 & \text{for odd } d' \geq 1 \end{cases} \quad \text{for odd } b , \qquad (4.19)$$

and

$$c'_{\bar{B}_m(b),d\geq d'} = \begin{cases} \binom{b/2}{d'/2} & \text{for even } d' \geq 0\\ -\binom{b/2}{(d'-1)/2} & \text{for odd } d' \geq 1 \end{cases} \quad \text{for even } b .$$

$$(4.20)$$

For the partition  $[1^d]$ , if we label  $c'_{B_m(b),2}$  as the  $c'_{B_m(b),d,j} = b - 1$  in the first row of the  $[1^d]$  partition of table XI, and similarly for other  $c'_{B_m(b),d'}$ , we conjecture the following forms of the corresponding  $c'_{\bar{B}_m(b),d'}$  for general d',

$$c'_{\bar{B}_m(b),d'} = \begin{cases} (-1)^{(d'-1)/2} \binom{(b-1)/2}{(d'-1)/2} & \text{for odd } d' \ge 3\\ 0 & \text{for even } d' \ge 2 \end{cases} \quad \text{for odd } b , \qquad (4.21)$$

and

$$c'_{\bar{B}_m(b),d'} = \begin{cases} \sum_{j=0}^{(d'-1)/2} (-1)^j {b/2 \choose j} & \text{for odd } d' \ge 3\\ \sum_{j=0}^{(d'-2)/2} (-1)^{j-1} {b/2 \choose j} & \text{for even } d' \ge 2 \end{cases}$$
 for even  $b$ , (4.22)

If the sum of the coefficients for a specific d is denoted as  $C(\bar{B}_m(b), d, q)$ , that is,

$$C(\bar{B}_m(b), d, q) = \sum_{j=1}^{N_{\bar{B}_m(b), d, \lambda}} c_{\bar{B}_m(b), d, j}(q) c'_{\bar{B}_m(b), d, j} , \qquad (4.23)$$

we have the following conjecture,

$$C(\bar{B}_m(\text{even } b), d, q) = \begin{cases} (-1)^{d/2} {b/2 \choose d/2} & \text{for even } d \\ 0 & \text{for odd } d \end{cases}$$
(4.24)

$$C(\bar{B}_m(\text{odd } b), d, q) = \begin{cases} (-1)^{d/2} \binom{(b-1)/2}{d/2} & \text{for even } d\\ (-1)^{(d-1)/2} (q-1) \binom{(b-1)/2}{(d-1)/2} & \text{for odd } d \end{cases}$$
(4.25)

### V. CONCLUSIONS

In this paper we have presented exact zero-temperature partition functions for families of chain graphs composed of repeated complete subgraphs  $K_b$ , b = 5, 6 with periodic or twisted periodic boundary condition in the longitudinal direction. In the  $L_x \to \infty$  limit, the continuous accumulation set of chromatic zeros  $\mathcal{B}$  was determined. We showed the eigenvalues with coefficients of degree b-1 are bounded between  $(-1)^{b-1}(q-1)$  and  $(-1)^{b-1}(q-2b+1)$ , and gave the explicit forms of the eigenvalues for partitions of the form [21]. The minimum real q at which  $\mathcal{B}$  crosses the real q axis was proven to be q = 0, and  $q_c(B_m(b)) < b$  for  $4 \leq b$ .

Acknowledgment: I would like to thank Prof. R. Shrock for helpful discussions.

- [1] R. B. Potts, Proc. Camb. Phil. Soc. 48 (1952) 106.
- [2] F. Y. Wu, Rev. Mod. Phys. **54** (1982) 235.
- [3] R. C. Read, J. Combin. Theory 4 (1968) 52.
- [4] R. C. Read and W. T. Tutte, "Chromatic Polynomials", in *Selected Topics in Graph Theory*, 3, eds. L. W. Beineke and R. J. Wilson (Academic Press, New York, 1988.).
- [5] N. L. Biggs, Algebraic Graph Theory (2nd ed., Cambridge Univ. Press, Cambridge, 1993).
- [6] R. Shrock and S.-H. Tsai, Phys. Rev. E55 (1997) 5165.
- [7] N. L. Biggs, LSE report LSE-CDAM-99-08 (1999).
- [8] S. Beraha, J. Kahane, and N. Weiss, J. Combin. Theory B 27 (1979) 1; *ibid.*, 28 (1980) 52.
- [9] N. L. Biggs, R. M. Damerell, and D. A. Sands, J. Combin. Theory B 12 (1972) 123.
- [10] N. L. Biggs and R. Shrock, J. Phys. A (Letts) **32**, L489 (1999).
- [11] S.-C. Chang, Physica A296, 495 (2001).
- [12] E. H. Lieb, Phys. Rev. **162** (1967) 162.
- [13] S.-C. Chang and R. Shrock, Phys. Rev. E, **62**, 4650 (2000).
- [14] S.-C. Chang and R. Shrock, Physica A 296 (2001) 131.
- [15] N. L. Biggs and G. H. Meredith, J. Combin. Theory B20 (1976) 5.
- [16] N. L. Biggs and G. H. Meredith, J. Combin. Theory B20 (1976) 5.
- [17] N. L. Biggs, Bull. London Math. Soc. 9 (1977) 54.
- [18] R. C. Read, in Proc. 3rd Caribbean Conf. on Combin. and Computing (1981).
- [19] R. C. Read, in Proc. 5th Caribbean Conf. on Combin. and Computing (1988).
- [20] R. J. Baxter, J. Phys. A **20** (1987) 5241.
- [21] R. C. Read and G. F. Royle, in *Graph Theory, Combinatorics, and Applications* (Wiley, NY, 1991), vol. 2, p. 1009.
- [22] R. Shrock and S.-H. Tsai, Phys. Rev. **E55** (1997) 6791.

- [23] R. Shrock and S.-H. Tsai, Phys. Rev. E56, 1342, 2733, 3935, 4111 (1997).
- [24] R. Shrock and S.-H. Tsai, Phys. Rev. **E56** (1997) 3935.
- [25] J. Salas and A. Sokal, J. Stat. Phys. 86 (1997) 551.
- [26] M. Roček, R. Shrock, and S.-H. Tsai, Physica A252 (1998) 505.
- [27] M. Roček, R. Shrock, and S.-H. Tsai, Physica A259 (1998) 367.
- [28] R. Shrock and S.-H. Tsai, Physica A259 (1998) 315.
- [29] R. Shrock and S.-H. Tsai, Phys. Rev. E58 (1998) 4332.
- [30] R. Shrock and S.-H. Tsai, J. Phys. A **31** (1998) 9641; Physica **A265** (1999) 186.
- [31] R. Shrock and S.-H. Tsai, J. Phys. A **32** (1999) 5053.
- [32] R. Shrock and S.-H. Tsai, J. Phys. A Lett. **32** (1999) L195.
- [33] R. Shrock and S.-H. Tsai, Phys. Rev. E60 (1999) 3512.
- [34] R. Shrock and S.-H. Tsai, Physica A **275** (2000) 429.
- [35] R. Shrock, Phys. Lett. A261 (1999) 57.
- [36] A. Sokal, Comb. Probab. Comput. 10, 41 (2001).
- [37] N. L. Biggs, J. Combin. Theory B 82, 19 (2001); Bull. London Math. Soc., in press.
- [38] R. Shrock, in the Proceedings of the 1999 British Combinatorial Conference, BCC99, Discrete Math. 231 (2001) 421.
- [39] H. Kluepfel and R. Shrock, unpublished; H. Kluepfel, Stony Brook thesis "The q-State Potts Model: Partition Functions and Their Zeros in the Complex Temperature and q Planes" (July, 1999).
- [40] R. Shrock, Physica **A281**, 221 (2000).
- [41] S.-C. Chang and R. Shrock, Phys. Rev. **E62**, 4650 (2000).
- [42] R. Shrock, Physica A283, 388 (2000).
- [43] S.-C. Chang and R. Shrock, Physica A286, 189 (2000).
- [44] S.-C. Chang and R. Shrock, Physica A **290**, 402 (2001).
- [45] S.-C. Chang and R. Shrock, Ann. Phys. **290**, 124 (2001).
- [46] S.-C. Chang and R. Shrock, Phys. Rev. E62, 4650 (2000).
- [47] J. Salas and A. Sokal, J. Stat. Phys., **104**, 611 (2001).
- [48] S.-C. Chang and R. Shrock, Physica A 286, 189 (2000).
- [49] S.-C. Chang and R. Shrock, Physica A **292**, 307 (2001).
- [50] S.-C. Chang and R. Shrock, Physica A 296, 234 (2001).
- [51] S.-C. Chang and R. Shrock, Physica A **296**, 183 (2001).
- [52] S.-C. Chang and R. Shrock, Int. J. Mod. Phys. B 15, 443 (2001).
- [53] J. L. Jacobsen and J. Salas, J. Stat. Phys. **104**, 703 (2001).
- [54] J. Salas and R. Shrock, Phys. Rev. E 64, 011111 (2001).
- [55] S.-C. Chang and R. Shrock, Physica A, **301**, 301 (2001).
- [56] S.-C. Chang and R. Shrock, Phys. Rev. E, in press; cond-mat/0107012.
- [57] J. Salas, S.-C. Chang and R. Shrock, J. Stat. Phys., in press (cond-mat/0108144).
- [58] N. L. Biggs, "Chromatic polynomials and representations of the symmetric group", AGT workshop, Edinburgh (July 2001), LSE report LSE-CDAM-01-02 (2001).
- [59] N. L. Biggs, M. Klin, and P. Reinfeld, "Algebraic methods for chromatic polynomials", AGT workshop, Edinburgh (July, 2001), LSE report LSE-CDAM-01-06 (2001).

- [60] N. L. Biggs, LSE report LSE-CDAM-00-04 (May 2000).
- [61] N. L. Biggs and P. Reinfeld, LSE report LSE-CDAM-00-07 (June 2000), http://www.cdam.lse.ac.uk/Reports/ .
- [62] N. L. Biggs, S.-C. Chang, P. Reinfeld, and R. Shrock, in preparation.