

HYDRODYNAMICS OF A DRIVEN LATTICE GAS WITH OPEN BOUNDARIES: THE ASYMMETRIC SIMPLE EXCLUSION.

O. BENOIS, R. ESPOSITO, R. MARRA, AND M. MOURRAGUI

ABSTRACT. We consider the asymmetric simple exclusion process in $d \geq 3$ with open boundaries. The particle reservoirs of constant densities are modeled by birth and death processes at the boundary. We prove that, if the initial density and the densities of the boundary reservoirs differ for order of ε from $1/2$, the density empirical field, rescaled as ε^{-1} , converges to the solution of the initial-boundary value problem for the viscous Burgers equation in a finite domain with given density on the boundary.

1. INTRODUCTION

A driven lattice gas with open boundaries is a system of particles jumping at random on a lattice, subject to the action of an external field and exchanging matter with a reservoir at his boundary. The combined action of the force field and the density gradient induced by the boundary conditions forces the system to reach a stationary non-equilibrium state. Systems of this kind show a complex behavior exhibiting non-equilibrium phase transitions [K], [SZ]. The simplest example of driven lattice gas is the asymmetric simple exclusion (ASEP). In the one-dimensional totally asymmetric case an explicit stationary solution is known [D] showing a very reach phase diagram with different behavior of the steady current depending on the values of the fixed densities on the boundaries.

In this paper we study the time-dependent measure of the ASEP with open boundaries in $d \geq 3$ in the macroscopic limit. The system is contained in a finite cylinder $\Lambda_\varepsilon = [-\varepsilon^{-1}, \varepsilon^{-1}] \times \pi_\varepsilon^{d-1}$, with π_ε^{d-1} the $(d-1)$ -dimensional microscopic torus of size $2\varepsilon^{-1} + 1$ with the axes in the direction x_1 , namely we impose periodic boundary conditions in all the directions but x_1 . In the bulk particles jump to one of the nearest neighbors if empty with jump rate p_{e_i} (p_{-e_i}) in the direction e_i ($-e_i$). We assume that the vector of components $\delta_i = p_{e_i} - p_{-e_i}$ is such that $\delta_1 > 0$. On the boundaries $x_1 = -\varepsilon^{-1}, \varepsilon^{-1}$ we allow for production and destruction of particles in the following way. Let $b(u)$ be smooth functions on $[-1, 1] \times \pi_1^{d-1}$. A particle is added independently in each site of $x_1 = -\varepsilon^{-1}$, when the site is empty, with rate $\delta_1(1/2 + \varepsilon b(\varepsilon x))$ and removed independently in each site of $x_1 = \varepsilon^{-1}$, when the site is occupied, with rate $\delta_1(1/2 - \varepsilon b(\varepsilon x))$. If $b(u)$ has constant values b_\pm , this choice of the jump rates corresponds to coupling the system to reservoirs of constant densities $1/2 + \varepsilon b_-$ and $1/2 + \varepsilon b_+$ respectively. In the paper we write the computations for this cylinder geometry, but the proof could be extended to a more general convex domain. The initial measure is the local equilibrium corresponding to a density profile which is a perturbation of order ε of a constant profile: $\rho_0(\varepsilon x) = 1/2 + \varepsilon m_0(\varepsilon x)$.

We prove a law of large numbers stating the weak convergence of the rescaled empirical field ($\eta_t(x) = 0, 1$ is the occupation number in the site x at time t)

$$\varepsilon^{d-1} \sum_{x \in \Lambda_\varepsilon} \left(\eta_{\varepsilon^{-2}t}(x) - \frac{1}{2} \right) \delta(\varepsilon x)$$

with $\delta(\cdot)$ the Dirac measure, to the solution of the viscous Burgers equation in the domain Λ_1 with density $b(u)$ on the boundary Γ

$$\begin{cases} \partial_t m(t, u) &= \delta \cdot \nabla_u (m(t, u))^2 + \sum_{i,j=1}^d D_{i,j} \partial_{u_i, u_j}^2 m(t, u) \\ m(0, \cdot) &= m_0(\cdot) \\ m(t, \cdot) \Big|_{\Gamma} &= b(\cdot) \quad \text{for } t \geq 0, \end{cases}$$

where D is a positive definite diffusion matrix whose expression is given by the Green-Kubo formula for this model. We remind here that the transport matrix is supposed to be infinite in $d \leq 2$ [Sp].

This special choice of the initial condition as well as of the boundary conditions (the difference between the top and the bottom densities is of order ε) is forced by the fact that we want to study the behavior of the system on the diffusive time scale ε^{-2} to see the effect of a finite dissipation. On this time scale the transport term, which involves first order time derivatives, is enhanced by a factor of order ε^{-1} so that if it is of order ε at time 0 its contribution stays finite in the limit. This is also called incompressible limit [EMY1] in analogy with the Navier-Stokes case [EMY2].

There are few rigorous results on the hydrodynamic limit for interacting particle systems in a bounded domain. In [ELS] it is proved the hydrodynamic limit for one-dimensional gradient systems both in the time-dependent and stationary case. In [KLO] the analogous result has been obtained in the stationary case for one-dimensional non-gradient models. Finally, in [LMS] the latter result has been extended to the d -dimensional case. All these papers deal with bulk reversible dynamics. If the lattice gas is driven by the boundary conditions, in general the stationary measure does not coincide with the invariant measure of the dynamics in the infinite volume, so that even if the generator of the bulk dynamics is reversible versus its invariant measure, the total dynamics is not. The asymmetric simple exclusion is an example of a lattice gas driven by an external field and it is not reversible w.r.t. his invariant measure in the infinite volume, which is a product measure. The case we consider in this paper is an example of a lattice gas driven by both external field and boundary conditions, so that there are two sources of non-reversibility. The effect of the asymmetry is seen in the macroscopic current as the term $\delta_i m^2$ which gives rise to the transport term in the macroscopic equation. We have to face the difficulty of non reversibility in the bulk and the fact that the stationary measure is not explicitly known.

The first problem has been solved in [EMY1] where it is proved the law of large numbers in the incompressible limit for the system without reservoirs. The method used there is based on the use of relative entropy of the true measure with respect to some suitable local equilibrium measure and the main point is to prove that this relative entropy vanishes in the limit $\varepsilon \rightarrow 0$. In this case it is natural to assume for the local equilibrium measure

a product measure because this is the invariant measure of the generator. However, for the open system it is not clear which is the good candidate for describing the system for ε small. We introduce a measure μ_ε which is a product measure with chemical potential $\varepsilon\lambda(t, \varepsilon x)$ such that $E^{\mu_\varepsilon}[\eta_x] = 1/2 + \varepsilon m(t, x)$, with m solution of the macroscopic equation. We prove that the relative entropy $s(\mu|\mu_\varepsilon)$ of the non-equilibrium measure μ w.r.t. μ_ε satisfy

$$\lim_{\varepsilon \rightarrow 0} \varepsilon^{-2} s(\mu|\mu_\varepsilon) = 0 \quad (1.1)$$

which implies the law of large numbers. One can understand the factor ε^{-2} noting that the specific entropy of μ_ε is of order ε^2 . To get this result, we introduce an auxiliary measure $\tilde{\mu}_\varepsilon$ which differs from μ_ε near the boundary and prove the previous limit for the relative entropy w.r.t $\tilde{\mu}_\varepsilon$. Then, the result easily follows from the fact that $\lim_{\varepsilon \rightarrow 0} \varepsilon^{-2} s(\mu_\varepsilon|\tilde{\mu}_\varepsilon) = 0$.

The method used in proving the hydrodynamic limit in the papers quoted above is different from ours because it is based on the martingale approach. However, in both methods it is needed an a priori estimate of the entropy (which assures also the control of the Dirichlet form) of the state of the process with respect to the invariant state of the full dynamics. Since this is not known, one uses some trial reference measure which is not invariant and as a result this entropy does not decrease in time. The difficulty to get an entropy bound in the asymmetric case is due to a diverging contribution to the flux of entropy due to the asymmetric part of the generator. This flux has to be controlled by the boundary generator. The choice of this generator is different from the one used in the case of reversible bulk dynamics, where the boundary death and birth process is required to satisfy a detailed balance condition. Our generator is a generalization of the one used by Derrida [D]. Our method would also provide the results for a reversible boundary generator, but then we have to speed it up much more than ε^{-2} .

One technical remark: the control of the terms in the bulk is done by using the replacement Lemma and the non gradient result in [EMY1]. This requires a localization procedure for the currents, which generates, in an open system, boundary terms. These terms are dealt with, in a reversible non-gradient case [LMS], by a simple use of integration by part and Schwartz inequality. The version of the integration by parts for the asymmetric case (Lemma 6.1 in [EMY1]) involves a function which is not bounded but just summable and as a consequence we need rather to use a modified version of the replacement Lemma to control these boundary terms.

We conclude this section by noticing that the extension to the stationary problem of the convergence result in the macroscopic limit could give some insight on the structure of the non-stationary states. Recent papers [DLS], [BDGJL], focus on the problem of characterizing the stationary non-equilibrium measures in terms of large deviation functional. Unfortunately, the relative entropy method does not seem to be useful for the stationary problem, because it relies on estimating the time derivative of the relative entropy in terms of the entropy production which is expressed back again in terms of the relative entropy, so that one gets a closed differential inequality for the relative entropy. We think that the

usual martingale approach could be more suited for the stationary case. We plan to refer on that in a future paper.

The plan of the paper is the following: in Section 2 we state the results. Section 3 is devoted to the strategy of the proof of the entropy bounds and to the control of the bulk terms while in Section 4 we have collected the estimates of the boundary terms. Finally, in Section 5 we discuss extensions to different geometries and different boundary conditions.

2. NOTATIONS AND RESULTS

Let $\varepsilon > 0$ such that ε^{-1} is integer and $\Lambda_\varepsilon = \{-\varepsilon^{-1}, \dots, \varepsilon^{-1}\} \times \pi_\varepsilon^{d-1}$ be the cylinder in \mathbb{Z}^d , $d \geq 3$, of length $2\varepsilon^{-1} + 1$ with basis π_ε^{d-1} , the $(d-1)$ -dimensional microscopic torus of size $2\varepsilon^{-1} + 1$. We denote by $\Gamma_\varepsilon = \{x \in \Lambda_\varepsilon \mid x_1 = \pm\varepsilon^{-1}\}$ the boundary of Λ_ε . The elements of Λ_ε will be denoted by letters x, y, \dots

A particle configuration is described as an element $\eta \in X_\varepsilon = \{0, 1\}^{\Lambda_\varepsilon}$, where $\eta(x) = 1, 0$ means that the site x is occupied by a particle or is empty. In this paper we are interested in the asymmetric simple exclusion process (ASEP) on Λ_ε with stochastic reservoirs at the boundary of Λ_ε . This Markov process is defined through its infinitesimal generator

$$\mathcal{L}_\varepsilon = \mathcal{L}_{\varepsilon,0} + \mathcal{L}_{\varepsilon,b}. \quad (2.2)$$

The generator $\mathcal{L}_{\varepsilon,0}$ is the nearest neighbor ASEP. Its action on functions $f : X_\varepsilon \rightarrow \mathbb{R}$ is

$$\mathcal{L}_{\varepsilon,0}f(\eta) = \sum_{i=1}^d \sum_{x, x+e_i \in \Lambda_\varepsilon} r_{x, x+e_i}(\eta) [f(\eta^{x, x+e_i}) - f(\eta)],$$

where (e_1, \dots, e_d) is the canonical basis of \mathbb{R}^d and the rate functions $r_{x, x+e_i}(\eta)$ are given by

$$r_{x, x+e_i}(\eta) = p_{e_i}\eta(x)(1 - \eta(x + e_i)) + p_{-e_i}\eta(x + e_i)(1 - \eta(x))$$

and $\eta^{x,y}$ is the configuration obtained by exchanging the occupations of sites x and y :

$$\eta^{x,y}(z) = \begin{cases} \eta(y) & \text{if } z = x, \\ \eta(x) & \text{if } z = y, \\ \eta(z) & \text{if } z \neq x, y. \end{cases}$$

For $1 \leq i \leq d$, p_{e_i} (p_{-e_i}) is the jump probability of a particle in the direction e_i ($-e_i$). For convenience we normalize the p_e 's so that, for $i = 1, \dots, d$, $p_{e_i} + p_{-e_i} = 2$. We call $\delta_i := p_{e_i} - p_{-e_i}$, $i = 1, \dots, d$, and we suppose that δ_1 is non vanishing. We assume it positive without loss of generality.

The currents $W_{x,i}$ are defined as

$$\begin{aligned} W_{x,i}(\eta) &= -W_{x,i}^{(s)} + W_{x,i}^{(a)} \\ &:= -[\eta(x + e_i) - \eta(x)] + \delta_i \left[\eta(x + e_i)\eta(x) - \frac{\eta(x) + \eta(x + e_i)}{2} \right]. \end{aligned}$$

Let Λ be the set $[-1, 1] \times \pi^{d-1}$, where π^{d-1} is the $(d-1)$ -dimensional torus with length 2, whose elements are denoted by u, v, \dots . We will consider as reference laws the Bernoulli

product measures ν_ρ on X_ε which are defined, for any smooth function $0 < \rho < 1$ on Λ , by $\nu_\rho(\eta(x) = 1) = \rho(\varepsilon x)$. It is well known that, in infinite volume, the product measure of Bernoulli laws with any constant parameter is invariant for the ASEP, but this is no longer valid in finite volume.

The generator $\mathcal{L}_{\varepsilon,b}$ in (2.2) is the infinitesimal generator of a birth and death process which creates particles on the “left” of Λ_ε and destroys them on the “right”. For any function f on X_ε

$$(\mathcal{L}_{\varepsilon,b}f)(\eta) = \sum_{x \in \Gamma_\varepsilon} C^b(\varepsilon x, \eta) [f(\eta^x) - f(\eta)],$$

where η^x is the configuration obtained from η by flipping the occupation number at site x

$$\eta^x(z) = \begin{cases} \eta(z) & \text{if } z \neq x \\ 1 - \eta(x) & \text{if } z = x. \end{cases}$$

The rate functions are designed to fix the value of the particle density at the boundary of Λ_ε . Let $\Gamma_{\varepsilon,+}$ (resp. $\Gamma_{\varepsilon,-}$) be the “left” (resp. “right”) side of Γ_ε :

$$\begin{aligned} \Gamma_{\varepsilon,+} &= \{(-\varepsilon^{-1}, x_2, \dots, x_d), (x_2, \dots, x_d) \in \pi_\varepsilon^{d-1}\}, \\ \Gamma_{\varepsilon,-} &= \{(\varepsilon^{-1}, x_2, \dots, x_d), (x_2, \dots, x_d) \in \pi_\varepsilon^{d-1}\}. \end{aligned}$$

For a smooth function b on the boundary Γ of Λ that we fix from now on, we choose $C^b(\varepsilon x, \eta)$ such that a particle is added on $x \in \Gamma_{\varepsilon,+}$, when the site is empty, with rate $\delta_1(1/2 + \varepsilon b(\varepsilon x))$ and removed in $\Gamma_{\varepsilon,-}$, when the site is occupied, with rate $\delta_1(1/2 - \varepsilon b(\varepsilon x))$:

$$C^b(\varepsilon x, \eta) = \delta_1(1/2 + \varepsilon b(\varepsilon x))(1 - \eta(x)) \mathbb{1}\{x \in \Gamma_{\varepsilon,+}\} + \delta_1(1/2 - \varepsilon b(\varepsilon x))\eta(x) \mathbb{1}\{x \in \Gamma_{\varepsilon,-}\}.$$

To understand this choice, remark that if we take $b = 0$ then the density $1/2$ is stationary for the full dynamics \mathcal{L}_ε , therefore the leading coefficient $1/2$ in the rate is necessary for the incompressible limit to make sense. The term εb will fix the value at the boundary of the density perturbation with respect to the stationary value $1/2$.

We will study the dynamics defined by the generator \mathcal{L}_ε under the diffusive space-time scaling $(\varepsilon x, \varepsilon^{-2}t)$. We denote by η_t the particle configuration at time t of the associated Markov process. When the process starts from the product measure $\mu_\varepsilon = \nu_{1/2 + \varepsilon m(\cdot)}$, m a smooth function on Λ such that $m|_\Gamma = b$, its law is denoted by $\mathbb{P}_\varepsilon^{b,m}$.

As in [Y] and [EMY1], the proof is based on the study of the relative entropy. Given two measures μ_1 and μ_2 on X_ε (with μ_1 absolutely continuous w.r.t. μ_2), the entropy of μ_1 with respect to μ_2 is defined as

$$H(\mu_1|\mu_2) = \int \frac{d\mu_1}{d\mu_2} \log \frac{d\mu_1}{d\mu_2} d\mu_2.$$

We also introduce the relative entropy $s(\mu_1|\mu_2) := \varepsilon^d H(\mu_1|\mu_2)$.

For simplicity we call $\nu := \nu_{1/2}$ the reference measure. Notice that the measure μ_ε has a density Ψ with respect to ν given by

$$\Psi(\eta) = \frac{1}{Z} \exp \left\{ \sum_{x \in \Lambda_\varepsilon} \varepsilon \varphi(m(\varepsilon x)) \eta(x) \right\} \nu(\eta),$$

where

$$\varphi(u) = \varepsilon^{-1} \log \left\{ \frac{1 + 2\varepsilon m(u)}{1 - 2\varepsilon m(u)} \right\} \quad (2.3)$$

and Z is a normalization constant. Then the relative entropy of the measure $f\nu$ (where f is some probability density) w.r.t. μ_ε can be written as:

$$s(f|\Psi) := s(f\nu|\mu_\varepsilon) = \varepsilon^d \int f \log \frac{f}{\Psi} d\nu.$$

The probability $\mu_\varepsilon(t)$ is the law of η_t when the initial distribution of the process is μ_ε . Its density with respect to ν is denoted by f_t . We now consider the solution $m(t, u)$ of the following partial differential equation

$$\begin{cases} \partial_t m(t, u) &= \sum_{i=1}^d \delta_i \partial_{u_i} (m(t, u))^2 + \sum_{1 \leq i, j \leq d} D_{i,j} \partial_{u_i, u_j}^2 m(t, u) \\ m(0, \cdot) &= m \\ m(t, \cdot) \Big|_{\Gamma} &= b(\cdot) \quad \text{for } t \geq 0, \end{cases} \quad (2.4)$$

where the diffusion coefficients $D_{i,j}$ will be defined later on.

We introduce the measure $\nu_{1/2+\varepsilon m(t, \cdot)}$ whose density w.r.t. ν is given by

$$\Psi_t(\eta) = \frac{1}{Z_t} \exp \left\{ \sum_{x \in \Lambda_\varepsilon} \varepsilon \varphi(m(t, \varepsilon x)) \eta(x) \right\},$$

with Z_t a normalization constant.

Our main result is:

Theorem 2.1. *Let $s(t) = s(f_t|\Psi_t)$. Then, for any $t > 0$,*

$$\lim_{\varepsilon \rightarrow 0} \varepsilon^{-2} s(t) = 0.$$

Using standard arguments based on the entropy inequality (see e.g. [Y] and [EMY1]), we get as a corollary of the previous theorem the incompressible limit of ASEP with stochastic reservoirs:

Theorem 2.2. *Let π_t^ε be the empirical measure*

$$\pi_t^\varepsilon(\eta) = \frac{1}{|\Lambda_\varepsilon|} \sum_{x \in \Lambda_\varepsilon} \left(\eta_t(x) - \frac{1}{2} \right) \delta(\varepsilon x).$$

Then π_t^ε converges weakly in $\mathbb{P}_\varepsilon^{b,m}$ -probability to the solution $m(t, u)$ to (2.4).

3. BOUNDS ON THE ENTROPY

The strategy in proving Theorem 2.1. is based on the study of the time evolution of the relative entropy. Here the difficulty relies on the fact that the invariant measure for the full dynamics is unknown.

Actually, we will not work directly with the entropy $s(t)$ but with some approximation of $s(t)$ obtained by a modification of the density Ψ_t . Let m be the solution of the equation (2.4) and γ be a smooth function defined on a neighborhood of Λ such that $\partial_{u_1}\gamma$ has a compact support included in $] -1, 1[\times \pi^{(d-1)}$ with $\gamma(\cdot)|_{\Gamma} = b$. We introduce a new parameter $0 < \theta < 1$ which will go to 0 after ε , we denote by Ω_θ the set

$$\Omega_\theta := [-1 + \theta, 1 - \theta] \times \pi^{(d-1)} \quad (3.5)$$

and we choose a smooth function χ^θ in a neighborhood of Λ such that $\chi^\theta|_{\Omega_{2\theta}} = 1$ and $\chi^\theta = 0$ outside Ω_θ . Then we define

$$\rho(t, u) = m(t, u)\chi^\theta(u) + \gamma(u)(1 - \chi^\theta(u)).$$

Notice that $\rho(t, \cdot)$ is identically equal to $m(t, \cdot)$ in $\Omega_{2\theta}$ and equal to γ on $\Omega \setminus \Omega_\theta$.

Recalling that φ was defined in (2.3), we denote by $\lambda(t, u)$ the function

$$\lambda(t, u) = \varphi(\rho(t, u)),$$

then it is easy to check that, for $u \in \Omega_{2\theta}$,

$$\partial_t \lambda(t, u) - \sum_{i=1}^d \delta_i \partial_{u_i} (\lambda(t, u))^2 - \sum_{1 \leq i, j \leq d} D_{i,j} \partial_{u_i, u_j}^2 \lambda(t, u) = o(\varepsilon), \quad (3.6)$$

where $\varepsilon^{-1}o(\varepsilon)$ is a bounded function. Moreover, with a suitable choice of χ^θ , we may assume that the derivative $\partial_{u_i} \lambda$, $i = 1, \dots, d$ are bounded, uniformly in θ .

Following [EMY1], we modify suitably the density Ψ_t in the definition of the relative entropy $s(t)$: for an integer ℓ , we set $k = \ell \varepsilon^{-2/d}$ and $\Lambda_k = \{-k, \dots, k\}^d$. The normalized indicator function ω is defined as

$$\omega(x) = |\Lambda_k|^{-1} \mathbb{1}\{x \in \Lambda_k\}.$$

Let $\{F_n\}$ be a family of local functions (i.e. which depend on the particle configuration only through a finite number of sites) on X_ε and

$$\Phi_n(\eta) = \sum_{i=1}^d \sum_{x \in \Lambda_\varepsilon^{2k}} (\partial_{u_i} \lambda)(t, \varepsilon x) (\tau_x F_n * \omega),$$

for $x \in \mathbb{Z}^d$, where τ_\cdot is the shift operator on $\mathbb{X} = \{0, 1\}^{\mathbb{Z}^d}$, $\Lambda_\varepsilon^{2k} = \{-\varepsilon^{-1} + 2k, \varepsilon^{-1} - 2k\} \times \pi^{d-1}$ and $*$ is the convolution product. We will work with n fixed and will take the limit $n \rightarrow \infty$

after the limit $\varepsilon \rightarrow 0$. Therefore the index n will be omitted for sake of shortness. The modified density $\tilde{\Psi}_t^F$ is defined as in [EMY1] by

$$\tilde{\Psi}_t^F(\eta) = \frac{1}{\tilde{Z}_t^F} \exp \left\{ \sum_{x \in \Lambda_\varepsilon} \varepsilon (\lambda * \omega)(t, \varepsilon x) \eta(x) + \varepsilon^2 \Phi(\eta) \right\},$$

where \tilde{Z}_t^F is the normalization constant.

Moreover, we define

$$s_1(t) := s(f_t | \tilde{\Psi}_t^F), \quad h_t^F = \frac{f_t}{\tilde{\Psi}_t^F}, \quad h_t = \frac{f_t}{\tilde{\Psi}_t}$$

and $\tilde{\Psi}_t$ the density

$$\tilde{\Psi}_t(\eta) = \frac{1}{\tilde{Z}_t} \exp \left\{ \sum_{x \in \Lambda_\varepsilon} \varepsilon (\lambda * \omega)(t, \varepsilon x) \eta(x) \right\}.$$

We finally introduce the functionals D_0 , D_b which are closely related to the Dirichlet forms. For any non negative function h and any measure μ on X_ε ,

$$\begin{aligned} D_0(h, \mu) &= \sum_{i=1}^d \sum_{x, x+e_i \in \Lambda_\varepsilon} \int r_{x, x+e_i}(\eta) (h(\eta^{x, x+e_i}) - h(\eta))^2 d\mu, \\ D_b(h, \mu) &= \frac{\delta_1}{2} \sum_{x \in \Gamma_\varepsilon} \int (h(\eta^x) - h(\eta))^2 d\mu. \end{aligned} \tag{3.7}$$

Then we have

Proposition 3.1. *There exists some constant $K > 0$ such that for every $t > 0$,*

$$\lim_{\theta \rightarrow 0} \overline{\lim}_{n \rightarrow \infty} \overline{\lim}_{\ell \rightarrow \infty} \overline{\lim}_{\varepsilon \rightarrow 0} \left\{ \varepsilon^{-2} s_1(t) + K \varepsilon^{d-4} \int_0^t ds \left(D_0(\sqrt{h_s^F}, \tilde{\Psi}_s^F \nu) + D_b(\sqrt{h_s}, \tilde{\Psi}_s \nu) \right) \right\} = 0.$$

Remark. Since the Dirichlet forms D_0 and D_b are positive, Theorem 2.1 will follow from Proposition 3.1 if we can show that

$$\overline{\lim}_{\theta \rightarrow 0} \overline{\lim}_{\ell \rightarrow \infty} \overline{\lim}_{\varepsilon \rightarrow 0} \varepsilon^{-2} [s(t) - s_1(t)] \leq 0. \tag{3.8}$$

Consider the density Ψ_t^1 with respect to ν given by

$$\Psi_t^1(\eta) = \frac{1}{Z_t^1} \exp \left\{ \sum_{x \in \Lambda_\varepsilon} \varepsilon \lambda(t, \varepsilon x) \eta(x) \right\}. \tag{3.9}$$

with Z_t^1 the normalization constant. It is proved in Lemma 3.2 of [EMY1] that for any $0 < \theta < 1$

$$\lim_{\varepsilon \rightarrow 0} \varepsilon^{-2} [s(f_t | \Psi_t^1) - s_1(t)] = 0. \tag{3.10}$$

Moreover a straightforward generalization of the proof of this lemma together with the entropy inequality leads to the following estimate: for any $\alpha > 0$, there exists a constant $c(\alpha)$ such that

$$\varepsilon^{-2}[s(t) - s(f_t|\Psi_t^1)] \leq \alpha\varepsilon^{-2}s(f_t|\Psi_t^1) + c(\alpha)\theta \quad (3.11)$$

where we have used the fact that $\lambda(t, u) = \varphi(m(t, u))$ for any $t > 0$ and $u \in \Omega_{2\theta}$. Then (3.10) and (3.11) imply (3.8).

Proof of Proposition 3.1. A simple computation shows that

$$\varepsilon^{-2} \frac{d}{dt} s_1(t) = \varepsilon^{d-4} \int f_t \mathcal{L}_\varepsilon \log h_t^F d\nu - \varepsilon^{d-2} \int \frac{\partial_t \tilde{\Psi}_t^F}{\tilde{\Psi}_t^F} f_t d\nu$$

Since the function $\Phi(\eta)$ depends on the configuration η only through the variables $\{\eta(x_1, \dots, x_d), |x_1| \leq (\varepsilon^{-1} - k) + s_F\}$, with s_F the support of F , then

$$\varepsilon^{d-4} \int f_t \mathcal{L}_\varepsilon \log h_t^F d\nu = \varepsilon^{d-4} \int f_t \mathcal{L}_{\varepsilon,0} \log h_t^F d\nu + \varepsilon^{d-4} \int f_t \mathcal{L}_{\varepsilon,b} \log h_t d\nu.$$

We use the basic inequality

$$a \log \frac{a}{b} \leq -(\sqrt{a} - \sqrt{b})^2 + a - b$$

for a and b positive numbers. It is applied with $a = h_t^F(\eta^{x,x+e_i})$ and $b = h_t^F(\eta)$ to manage with $\mathcal{L}_{\varepsilon,0}$ and with $a = h_t(\eta^x)$ and $b = h_t(\eta)$ to manage with $\mathcal{L}_{\varepsilon,b}$. We get

$$\begin{aligned} \varepsilon^{-2} \frac{d}{dt} s_1(t) &\leq E(\varepsilon, t) - \varepsilon^{d-4} D_0(\sqrt{h_t^F}, \tilde{\Psi}_t^F \nu) \\ &\quad + \varepsilon^{d-4} \int (\mathcal{L}_{\varepsilon,b} h_t) \tilde{\Psi}_t d\nu - \frac{1}{2} \varepsilon^{d-4} D_b(\sqrt{h_t}, \tilde{\Psi}_t \nu) \end{aligned} \quad (3.12)$$

with $E(\varepsilon, t) := E_1 + E_2$, where

$$\begin{aligned} E_1 &:= \varepsilon^{d-4} \int \frac{\mathcal{L}_{\varepsilon,0}^* \tilde{\Psi}_t^F}{\tilde{\Psi}_t^F} f_t d\nu = \varepsilon^{d-4} \int \frac{\mathcal{L}_{\varepsilon,0}^* \tilde{\Psi}_t^F}{\tilde{\Psi}_t^F} (f_t - \tilde{\Psi}_t^F) d\nu \\ E_2 &:= -\varepsilon^{d-2} \int \frac{\partial_t \tilde{\Psi}_t^F}{\tilde{\Psi}_t^F} f_t d\nu \end{aligned}$$

and $\mathcal{L}_{\varepsilon,0}^*$ denotes the adjoint operator of $\mathcal{L}_{\varepsilon,0}$ in $L^2(\nu)$. After a change of variables the term E_1 can be rewritten as

$$\begin{aligned} E_1 &= \varepsilon^{d-4} \sum_{i=1}^d \sum_{x, x+e_i \in \Lambda_\varepsilon} \int r_{x, x+e_i}^*(\eta) \left\{ \frac{\tilde{\Psi}_t^F(\eta^{x, x+e_i})}{\tilde{\Psi}_t^F(\eta)} - 1 \right\} (f_t - \tilde{\Psi}_t^F)(\eta) d\nu \\ &\quad + \varepsilon^{d-4} \sum_{i=1}^d \sum_{x, x+e_i \in \Lambda_\varepsilon} \delta_i \int (\eta(x+e_i) - \eta(x)) (f_t - \tilde{\Psi}_t^F)(\eta) d\nu \\ &:= E_1^1 + E_1^2, \end{aligned} \quad (3.13)$$

where

$$r_{x,x+e_i}^*(\eta) = p_{e_i}\eta(x+e_i)(1-\eta(x)) + p_{-e_i}\eta(x)(1-\eta(x+e_i)).$$

Using again a change of variables at the boundary, the second term of E_1 is equal to

$$\begin{aligned} E_1^2 &= \varepsilon^{d-4} \sum_{x \in \Gamma_\varepsilon} \delta_1 \int (\eta(x) - \langle \eta(x) \rangle_{\tilde{\Psi}_t^F}) n_1(\varepsilon x) f_t(\eta) d\nu \\ &= \varepsilon^{d-4} \sum_{x \in \Gamma_\varepsilon} \delta_1 \int (\eta(x) - \langle \eta(x) \rangle_{\tilde{\Psi}_t}) n_1(\varepsilon x) f_t(\eta) d\nu, \end{aligned} \quad (3.14)$$

where $n = (n_1, 0, \dots, 0)$ is the outward unit normal vector to the boundary surface Γ , and by standard manipulations can be bounded by the Dirichlet form and the generator at the boundary:

Lemma 3.1. *For all $a > 0$,*

$$\begin{aligned} \varepsilon^{d-4} \sum_{x \in \Gamma_\varepsilon} \delta_1 \int (\eta(x) - \langle \eta(x) \rangle_{\tilde{\Psi}_t}) n_1(\varepsilon x) f_t(\eta) d\nu &\leq \\ &\leq -\varepsilon^{d-4} \int (\mathcal{L}_{\varepsilon,b} h_t) \tilde{\Psi}_t d\nu + \varepsilon^{d-4} \frac{a}{2} D_b(\sqrt{h_t}, \tilde{\Psi}_t \nu) + \varepsilon^3 k^4 \frac{C_1}{a}, \end{aligned}$$

for some positive constant C_1 .

Proof. For $\varepsilon > 0$, denote by $\tilde{\mathcal{L}}_{\varepsilon,b}^k$ the following generator at the boundary

$$\begin{aligned} (\tilde{\mathcal{L}}_{\varepsilon,b}^k f)(\eta) &= \delta_1 \sum_{x \in \Gamma_{\varepsilon,+}} \tilde{C}_+^b(\varepsilon x, \eta) [f(\eta^x) - f(\eta)] \\ &\quad + \delta_1 \sum_{x \in \Gamma_{\varepsilon,-}} \tilde{C}_-^b(\varepsilon x, \eta) [f(\eta^x) - f(\eta)], \end{aligned}$$

where

$$\begin{aligned} \tilde{C}_+^b(\varepsilon x, \eta) &= (1/2 + \varepsilon(\lambda * \omega)(t, \varepsilon x)) (1 - \eta(x)) \\ \tilde{C}_-^b(\varepsilon x, \eta) &= (1/2 - \varepsilon(\lambda * \omega)(t, \varepsilon x)) \eta(x) \end{aligned}$$

Remark that the replacement of $\mathcal{L}_{\varepsilon,b}$ by $\tilde{\mathcal{L}}_{\varepsilon,b}^k$ in the derivative of the entropy produces a term that we can bound by the Dirichlet form. Indeed, we have

$$\begin{aligned} \varepsilon^{d-4} \int \left((\mathcal{L}_{\varepsilon,b} - \tilde{\mathcal{L}}_{\varepsilon,b}^k) h_t \right) \tilde{\Psi}_t d\nu &= \\ &= \varepsilon^{d-3} \sum_{x \in \Gamma_{\varepsilon,+}} \left(b(\varepsilon x) - (\lambda * \omega)(t, \varepsilon) \right) \int (1 - \eta(x)) [h_t(\eta^x) - h_t(\eta)] \tilde{\Psi}_t d\nu \\ &\quad + \varepsilon^{d-3} \sum_{x \in \Gamma_{\varepsilon,-}} \left((\lambda * \omega)(t, \varepsilon) - b(\varepsilon x) \right) \int \eta(x) [h_t(\eta^x) - h_t(\eta)] \tilde{\Psi}_t d\nu, \end{aligned}$$

by Taylor expansion at the second order, this last quantity is equal to

$$\begin{aligned} & \varepsilon^{d-1} k^2 \sum_{x \in \Gamma_{\varepsilon,+}} G(t, \varepsilon x) \int (1 - \eta(x)) [h_t(\eta^x) - h_t(\eta)] \tilde{\Psi}_t d\nu \\ & + \varepsilon^{d-1} k^2 \sum_{x \in \Gamma_{\varepsilon,-}} G(t, \varepsilon x) \int \eta(x) [h_t(\eta^x) - h_t(\eta)] \tilde{\Psi}_t d\nu \\ & \leq \varepsilon^{d-4} \frac{a}{2} D_b(\sqrt{h_t}, \tilde{\Psi}_t \nu) + \varepsilon^3 k^4 \frac{C_1}{a} \end{aligned}$$

for all $a > 0$ and some constant C_1 , where $G(t, \varepsilon x)$ is bounded and proportional to the second derivative of $\lambda(t, \cdot)$. We have used, in the last inequality, Schwartz inequality and the fact that h_t is a probability density with respect to $\tilde{\Psi}_t \nu$.

To conclude the proof of the lemma, it is easy to check that the left hand side of the inequality in Lemma 3.1 can be rewritten as $-\varepsilon^{d-4} \int (\tilde{\mathcal{L}}_{\varepsilon,b}^k h_t) \tilde{\Psi}_t d\nu$, by using the change of variable formulas valid for any measure ν_g

$$\begin{aligned} \int (1 - \eta_x) f(\eta^{x,+}) d\nu_g(\eta) &= \frac{1 - g(\varepsilon x)}{g(\varepsilon x)} \int \eta(x) f(\eta) d\nu_g(\eta) \\ \int \eta_x f(\eta^{x,-}) d\nu_g(\eta) &= \frac{g(\varepsilon x)}{1 - g(\varepsilon x)} \int (1 - \eta(x)) f(\eta) d\nu_g(\eta) \end{aligned}$$

where $\eta^{x,+}$ and $\eta^{x,-}$ are the configurations obtained from η by adding or removing a particle at the site x . \square

The term E_1^1 is shown, by using Taylor expansion and Schwarz inequality, to be less or equal to

$$\varepsilon^{d-2} \int \left\{ \sum_{i=1}^d \sum_{x \in \Lambda_\varepsilon} (\partial^{\varepsilon,i} \lambda * \omega)(t, \varepsilon x) \widehat{W}_{x,i} - \mathcal{L}_0^* \Phi \right\} (f_t - \tilde{\Psi}_t^F)(\eta) d\nu + R_\varepsilon(t) + o_\varepsilon(1). \quad (3.15)$$

where $W_{x,i}$ is the current defined in section 2, $\partial^{\varepsilon,i} \lambda$ is the discrete gradient of λ

$$\partial^{\varepsilon,i} \lambda(t, u) = \varepsilon^{-1} [\lambda(t, u + \varepsilon e_i) - \lambda(t, u)]$$

and for any function g on X_ε , we denote by $\widehat{g} = g - \langle g \rangle_\nu$. The operator \mathcal{L}_0 is the generator of the ASEP in infinite volume space $X = \mathbb{Z} \times \pi_\varepsilon^{d-1}$ and \mathcal{L}_0^* is its adjoint in $L^2(\nu)$, given for any local function f by

$$(\mathcal{L}_0^* f)(\eta) = \sum_{i=1}^d \sum_{x \in \mathbb{Z} \times \pi_\varepsilon^{d-1}} r_{x, x+e_i}^*(\eta) [f(\eta^{x, x+e_i}) - f(\eta)].$$

In equation (3.15) as in the sequel, $R_\varepsilon(t)$ is real sequence of the form

$$R_\varepsilon(t) = \varepsilon^d \sum_{x \in \Lambda_\varepsilon} G(t, \varepsilon x) \int \tau_x \widehat{g}(\eta) (f_t - \tilde{\Psi}_t^F)(\eta) d\nu \quad (3.16)$$

with some bounded functions $G(t, \cdot)$ defined on Λ and g on X_ε . Observe that, for such sequences, one can use the entropy inequality and obtain, for any $A > 0$,

$$R_\varepsilon(t) \leq A\varepsilon^{d-2}s_1(t) + A^{-1}o_\varepsilon(1), \quad (3.17)$$

for a real sequence $o_\varepsilon(1)$ which is bounded in absolute value by a constant that converges to 0 as $\varepsilon \downarrow 0$.

At this point, we would like to replace the currents $W_{x,i}$ appearing in the first term of the right hand side of (3.15) by its convolution with ω . In order to do it we need some notation. For a local function g denote by g^k the convolution

$$(\tau_x g)^k(\eta) = \frac{1}{|\Lambda_k|} \sum_{y \in \Lambda_k} (\tau_{x+y} \widehat{g})(\eta),$$

when $g(\eta) = \eta(0) - 1/2$, we shall denote $(\tau_x g)^k$ simply by $\eta^k(x)$. Observe that, $\eta^k(0) + 1/2$ is precisely the empirical density in Λ_k . Furthermore, for $\varepsilon > 0$, let $\Lambda_\varepsilon^k = \{-\varepsilon^{-1} + k, \varepsilon^{-1} - k\} \times \pi^{d-1}$ and $\Gamma_\varepsilon^k = \Lambda_\varepsilon \setminus \Lambda_\varepsilon^k$. For $x \in X = \mathbb{Z} \times \pi^{d-1}$ denote by $\overline{W}_{x,i}$ the current defined on $\mathbb{X} = \{0, 1\}^X$ by

$$\overline{W}_{x,i}(\eta) = \begin{cases} W_{x,i}(\eta) & \text{if } x, x + e_i \in \Lambda_\varepsilon \\ 0 & \text{if } x \text{ or } x + e_i \notin \Lambda_\varepsilon \end{cases}$$

with this notation we have

$$\begin{aligned} E_1^1 &= \varepsilon^{d-2} \int \left\{ \sum_{i=1}^d \sum_{x \in \Lambda_\varepsilon^k} \partial^{\varepsilon,i} \lambda(t, \varepsilon x) (\tau_x \widehat{W}_{0,i})^k - \mathcal{L}_0^* \Phi \right\} (f_t - \widetilde{\Psi}_t^F)(\eta) d\nu \\ &+ \varepsilon^{d-2} \int \left\{ \sum_{i=1}^d \sum_{x \in \Gamma_\varepsilon^k} \partial^{\varepsilon,i} \lambda(t, \varepsilon x) (\widehat{\overline{W}}_{x,i})^k \right\} (f_t - \widetilde{\Psi}_t^F)(\eta) d\nu \\ &:= E_1^3 + E_1^4. \end{aligned} \quad (3.18)$$

Let us summarize what we have done so far: for any fixed $A > 0$ and $a > 0$, we got the following bound for the entropy derivative

$$\begin{aligned} \frac{d}{dt} \varepsilon^{d-2} s_1(t) &\leq A\varepsilon^{d-2} s_1(t) + E_2 + E_1^3 + E_1^4 \\ &- \varepsilon^{d-4} \left\{ D_0(\sqrt{h_t^F}, \widetilde{\Psi}_t^F \nu_{1/2}) + \frac{1-a}{2} D_b(\sqrt{h_t}, \widetilde{\Psi}_t \nu_{1/2}) \right\} \\ &+ o_\varepsilon(1). \end{aligned} \quad (3.19)$$

From the fact that $\partial^{\varepsilon,1}\lambda(t, \varepsilon x)$ has a compact support, Taylor expansion and (3.17), the term E_1^3 can be written as

$$\begin{aligned} E_1^3 &= \varepsilon^{d-1} \sum_{i=1}^d \sum_{x \in \Lambda_\varepsilon^k} \partial_{u_i}^2 \lambda(t, \varepsilon x) \int \eta^k(x) (f_t - \tilde{\Psi}_t^F)(\eta) d\nu \\ &+ \varepsilon^{d-2} \int \left\{ \sum_{i=1}^d \sum_{x \in \Lambda_\varepsilon^k} \partial^{\varepsilon,i} \lambda(t, \varepsilon x) (\widehat{W}_{x,i}^{(a)})^k - \mathcal{L}_0^* \Phi \right\} (f_t - \tilde{\Psi}_t^F)(\eta) d\nu \\ &+ o_\varepsilon(1). \end{aligned} \tag{3.20}$$

Following the method of [EMY1] we now replace the currents $(\widehat{W}_{0,i}^{(a)})^k$ in the bulk by a linear combination of the gradients $(\eta(e_j) - \eta(0))^k$, $j = 1, \dots, d$. This requires some notation. For $x \in \Lambda_\varepsilon^k$ and $1 \leq i \leq d$, denote by $M_{x,i}^k$ the conditional expectation of $W_{x,i}^{(a)}$ given the density of particles on $\Lambda_{x,k}$:

$$M_{x,i}^k(\eta) = \mathbb{E} \left[W_{x,i}^{(a)} \mid \eta^k(x) \right],$$

where $\Lambda_{x,k} = \{x + y : y \in \Lambda_k\}$. One can compute $M_{x,i}^k(\eta)$ easily. It is given by

$$\delta_i \left(1 + \frac{1}{(2k+1)^d - 1} \right) \eta^k(x) (\eta^k(x) - 1).$$

Furthermore, for $\varepsilon > 0$, $1 \leq i \leq d$ and $\eta \in X_\varepsilon$, let

$$\mathbb{V}_i^k(\eta) = \left((\widehat{W}_{0,i}^{(a)})^k - M_{0,i}^k \right)(\eta) - (\mathcal{L}_0^* F)^k + \sum_{j=1}^d \tilde{D}_{i,j} \times (\eta^k(e_j) - \eta^k(0)),$$

where $\tilde{D}_{i,j} = D_{i,j} - \delta_{i,j}$ and the diffusion matrix $D_{i,j}$ is defined in Section 2.

The next result is the main step towards the proof of the bounds of the entropy.

Theorem 3.1. *For all $A > 0$ and any probability density f with respect to ν ,*

$$\overline{\lim}_{n \rightarrow \infty} \overline{\lim}_{\ell \rightarrow \infty} \overline{\lim}_{\varepsilon \rightarrow 0} \left\{ \varepsilon^{d-2} \int \left(\sum_{i=1}^d \sum_{x \in \Lambda_\varepsilon^k} \partial^{\varepsilon,i} \lambda(t, \varepsilon x) \tau_x \mathbb{V}_i^k(\eta) \right) f(\eta) d\nu - A \varepsilon^{d-4} D_0(\sqrt{f}, \nu) \right\} \leq 0.$$

This Theorem was proved in [EMY1], [LY]. The next lemma takes care of the terms close to the boundary (E_1^4)

Lemma 3.2. *For all $A > 0$ and any probability density f with respect to ν ,*

$$\overline{\lim}_{\ell \rightarrow \infty} \overline{\lim}_{\varepsilon \rightarrow 0} \left\{ \varepsilon^{d-2} \int \sum_{i=1}^d \sum_{x \in \Gamma_\varepsilon^k} \partial^{\varepsilon,i} \lambda(t, \varepsilon x) (\widehat{W}_{x,i}^{(a)})^k f d\nu - A \varepsilon^{d-4} D_0(\sqrt{f}, \nu) \right\} \leq 0.$$

The proof of this Lemma will be given in next section.

Notice that, in the previous results, the Dirichlet form $D_0(\sqrt{f_t}, \nu)$ appeared while, in the derivative of the entropy, we got $D_0(\sqrt{h_t^F}, \tilde{\Psi}_t^F \nu)$. The next lemma allows us to replace the one by the other. Moreover we will also need some estimate about $D_0(\sqrt{\tilde{\Psi}_t^F}, \nu)$.

Lemma 3.3. *There exist two constants C_0 and C'_0 such that*

$$\begin{aligned} -\varepsilon^{d-4} D_0(\sqrt{h_t^F}, \tilde{\Psi}_t^F \nu) &\leq -\frac{1}{2} \varepsilon^{d-4} D_0(\sqrt{f_t}, \nu) + C_0, \\ \varepsilon^{d-4} D_0(\sqrt{\tilde{\Psi}_t^F}, \nu) &\leq C'_0 \end{aligned}$$

Proof. The second inequality follows by inspection. For two nearest neighbor sites x, y in Λ_ε , denote by $S_{x,y}$ the operator defined by

$$(S_{s,y} f)(\eta) = f(\eta^{x,y}) - f(\eta).$$

and write the Dirichlet form $D_0(\sqrt{f_t}, \nu)$ as

$$\begin{aligned} D_0(\sqrt{f_t}, \nu) &= \sum_{i=1}^d \sum_{x, x+e_i \in \Lambda_\varepsilon} \int r_{x, x+e_i} \left\{ \left(S_{x, x+e_i} (\tilde{\Psi}_t^F)^{-\frac{1}{2}} \right) (\eta) \sqrt{f_t}(\eta^{x, x+e_i}) \right. \\ &\quad \left. + \left(S_{x, x+e_i} \sqrt{h_t^F} \right) (\eta) \right\}^2 \tilde{\Psi}_t^F(\eta) d\nu. \end{aligned}$$

The elementary inequality $(a+b)^2 \leq 2(a^2+b^2)$ and a change of variables give that

$$\begin{aligned} \varepsilon^{d-4} D_0(\sqrt{f_t}, \nu) &\leq 2\varepsilon^{d-4} D_0(\sqrt{h_t^F}, \tilde{\Psi}_t^F \nu) \\ &\quad + 2\varepsilon^{d-4} \sum_{i=1}^d \sum_{x, x+e_i \in \Lambda_\varepsilon} \int r_{x, x+e_i}^*(\eta) \left(\sqrt{\frac{\tilde{\Psi}_t^F(\eta^{x, x+e_i})}{\tilde{\Psi}_t^F(\eta)}} - 1 \right)^2 f_t(\eta) d\nu. \end{aligned}$$

□

Applying Lemma (3.2) with densities f_t then $\tilde{\Psi}_t^F$ and using Lemma 3.3, we obtain the following bound for E_1^4 : for any $A > 0$

$$E_1^4 \leq A\varepsilon^{d-4} D_0(\sqrt{f_t}, \nu) + AC'_0 + r(\varepsilon, k, A) \quad (3.21)$$

where

$$\overline{\lim}_{\ell \rightarrow \infty} \overline{\lim}_{\varepsilon \rightarrow 0} r(\varepsilon, k, A) \leq 0.$$

With the notations introduced before Theorem 3.1, a summation by parts and a Taylor expansion permit to rewrite the quantity E_1^3 as

$$\begin{aligned}
E_1^3 &= \varepsilon^{d-1} \int \left\{ \sum_{1 \leq i, j \leq d} \sum_{x \in \Lambda_\varepsilon^k} \partial_{u_i, u_j}^2 \lambda(t, \varepsilon x) D_{i, j} \eta^k(x) \right\} (f_t - \tilde{\Psi}_t^F)(\eta) d\nu \\
&+ \varepsilon^{d-2} \int \left\{ \sum_{i=1}^d \sum_{x \in \Lambda_\varepsilon^k} \partial^{\varepsilon, i} \lambda(t, \varepsilon x) M_{x, i}^k(\eta) \right\} (f_t - \tilde{\Psi}_t^F)(\eta) d\nu \\
&+ \varepsilon^{d-2} \int \left\{ \sum_{i=1}^d \sum_{x \in \Lambda_\varepsilon^k} \partial^{\varepsilon, i} \lambda(t, \varepsilon x) \tau_x \mathbb{V}_i^\varepsilon(\eta) \right\} (f_t - \tilde{\Psi}_t^F)(\eta) d\nu \\
&+ \varepsilon^{d-2} \int \left\{ \sum_{i=1}^d \sum_{x \in \partial \Lambda_\varepsilon^k} \partial^{\varepsilon, i} \lambda(t, \varepsilon x) (t, \varepsilon x) \eta^k(x) n_1(\varepsilon x) \right\} (f_t - \tilde{\Psi}_t^F)(\eta) d\nu \\
&+ R_\varepsilon(t) + o_\varepsilon(1),
\end{aligned} \tag{3.22}$$

where $\partial \Lambda_\varepsilon^k$ stands for the boundary of Λ_ε^k , $n(\varepsilon x) = (n_1(\varepsilon x), 0, \dots, 0)$ for the outward unit normal vector to $\varepsilon(\partial \Lambda_\varepsilon^k)$ at εx and $R_\varepsilon(t)$ has been defined in (3.16).

We first estimate the third line in the formula above. Using Theorem 3.1 and Lemma 3.3, we get, for any $A > 0$,

$$\begin{aligned}
\varepsilon^{d-2} \int \left\{ \sum_{i=1}^d \sum_{x \in \Lambda_\varepsilon^k} \partial^{\varepsilon, i} \lambda(t, \varepsilon x) \tau_x \mathbb{V}_i^\varepsilon(\eta) \right\} (f_t - \tilde{\Psi}_t^F)(\eta) d\nu &\leq \\
&\leq A \varepsilon^{d-4} D_0(\sqrt{f_t}, \nu) + A(C'_0 + C_0) + r(\varepsilon, k, n, A)
\end{aligned} \tag{3.23}$$

where

$$\overline{\lim}_{n \rightarrow \infty} \overline{\lim}_{\ell \rightarrow \infty} \overline{\lim}_{\varepsilon \rightarrow 0} r(\varepsilon, k, n, A) \leq 0.$$

We now examine the second line in (3.22). We can write $M_{x, i}^k$ as

$$\delta_i Q_{k, \varepsilon}(x) + 2\varepsilon \delta_i \lambda(t, \varepsilon x) \eta^k(x) + \text{const} + O(k^{-d}) + O(\varepsilon^3 k^2),$$

where

$$Q_{k, \varepsilon}(\eta, x) = (\eta^k(x) - \varepsilon \lambda(t, \varepsilon x))^2$$

and const stands for a term independent of the configuration. By the entropy inequality, for any bounded function $G : \mathbb{R} \times \pi^{d-1} \rightarrow \mathbb{R}$, for all $q > 0$

$$\begin{aligned}
\varepsilon^{d-2} \int \sum_{x \in \Lambda_\varepsilon^k} G(\varepsilon x) \widehat{Q}_{k, \varepsilon}(\eta, x) (f_t - \tilde{\Psi}_t^F)(\eta) d\nu \\
\leq q^{-1} \varepsilon^{d-2} s_1(t) + q^{-1} \varepsilon^{d-2} \log E^{\tilde{\Psi}_t^F} \left[\exp q \sum_{x \in \Lambda_\varepsilon^k} \widehat{Q}_{k, \varepsilon}(\eta, x) \right]
\end{aligned}$$

with

$$\widehat{Q}_{k,\varepsilon}(\eta, x) := Q_{k,\varepsilon}(\eta, x) - E^{\widetilde{\Psi}_t^F} [Q_{k,\varepsilon}(\eta, x)].$$

From large deviations estimate (Lemma 3.1 in [EMY1]) there exists $q_0 > 0$, such that, for all $q < q_0$,

$$\lim_{\ell \rightarrow \infty} \overline{\lim}_{\varepsilon \rightarrow 0} \varepsilon^{d-2} \log E^{\widetilde{\Psi}_t^F} [\exp q \sum_{x \in \Lambda_\varepsilon^k} \widehat{Q}_{k,\varepsilon}(\eta, x)] = 0.$$

In conclusion, we got the following inequality for the second line in (3.22)

$$\begin{aligned} & \varepsilon^{d-2} \int \sum_{i=1}^d \sum_{x \in \Lambda_\varepsilon^k} \partial^{\varepsilon, i} \lambda(t, \varepsilon x) M_{x,i}^k(\eta) (f_t - \widetilde{\Psi}_t^F)(\eta) d\nu \\ & \leq \varepsilon^{d-1} \int \sum_{i=1}^d \sum_{x \in \Lambda_\varepsilon^k} \partial^{\varepsilon, i} \lambda(t, \varepsilon x) (2\lambda(t, \varepsilon x)) \eta^k(x) (f_t - \widetilde{\Psi}_t^F)(\eta) d\nu \\ & \quad + q^{-1} \varepsilon^{d-2} s_1(t) + r_{\varepsilon,k}(q), \end{aligned}$$

where, for all $q < \frac{q_0}{2}$, $r_{\varepsilon,k}(q)$ converges to 0 when $\varepsilon \downarrow 0$ and $\ell \uparrow \infty$.

To deal with the fourth line in (3.22) (boundary term), we need the following lemma

Lemma 3.4. *Fix a bounded function $G : \mathbb{R} \times \pi^{d-1} \rightarrow \mathbb{R}$ and $x_1 \in [-\varepsilon^{-1} + k, -\varepsilon^{-1} + 2k] \cup [\varepsilon^{-1} - 2k, \varepsilon^{-1} - k]$. Then there exists some constant $C > 0$ such that for all $A > 0$,*

$$\begin{aligned} & \varepsilon^{d-2} \int \sum_{y \in \pi_\varepsilon^{d-1}} G(\varepsilon(x_1, y)) \eta^k(x_1, y) (f_t - \widetilde{\Psi}_t^F)(\eta) d\nu \leq \\ & \leq \varepsilon^{d-4} \frac{A}{2} \left(D_b(\sqrt{h_t}, \widetilde{\Psi}_t \nu) + D_0(\sqrt{f_t}, \nu) \right) + \varepsilon k C \|G\|_\infty \left(\frac{\|G\|_\infty}{A} + 1 \right). \end{aligned}$$

The proof of this lemma is postponed to the next section.

We now turn to the proof of Proposition 3.1 and we first consider the term E_2 . Since the function $(\partial_t \lambda)$ is equal to 0 outside Ω_θ (cf. formula (3.5) and below), a simple computation shows that

$$\begin{aligned} E_2 &= -\varepsilon^{d-2} \int \frac{\partial_t \widetilde{\Psi}_t^F}{\widetilde{\Psi}_t^F} f_t d\nu \\ &= -\varepsilon^{d-1} \int \sum_{x \in \Lambda_\varepsilon} \partial_t (\lambda * \omega)(t, \varepsilon x) \eta(x) (f_t - \widetilde{\Psi}_t^F)(\eta) d\nu \\ &= -\varepsilon^{d-1} \int \sum_{x: \varepsilon x \in \Omega_\theta} \partial_t \lambda(t, \varepsilon x) \eta^k(x) (f_t - \widetilde{\Psi}_t^F)(\eta) d\nu. \end{aligned}$$

To conclude the proof of proposition 3.1, we integrate the inequality (3.19) from 0 to t . Combining with the above estimates, we obtain that there exist constants $K_1 > 0$, $K_2 > 0$,

$K_3 > 0$ and $c > 0$ such that, for all small enough $A > 0$,

$$\begin{aligned}
 \varepsilon^{-2}s_1(t) &\leq K_1\varepsilon^{-2} \int_0^t s_1(u)du \\
 &\quad - K_2\varepsilon^{d-4} \int_0^t ds \left\{ D_0(\sqrt{h_s^F}, \tilde{\Psi}_s^F \nu) + D_b(\sqrt{h_s}, \tilde{\Psi}_s \nu) \right\} \\
 &\quad + \varepsilon^{d-1} \int_0^t ds \left\{ \sum_{x \in \Omega_{\varepsilon k}} H(s, \varepsilon x) \int \eta^k(x)(f_s - \tilde{\Psi}_s^F) d\nu \right\} \\
 &\quad + AK_3\varepsilon^{d-4} \int_0^t D_0(\sqrt{f_s}, \nu) ds + r(\varepsilon, k, n, A) + cA,
 \end{aligned} \tag{3.24}$$

where $\overline{\lim}_{n \rightarrow \infty} \overline{\lim}_{\ell \rightarrow \infty} \overline{\lim}_{\varepsilon \rightarrow 0} r(\varepsilon, k, n, A) \leq 0$ for all $A > 0$, and the function H is given by

$$H(s, \varepsilon x) = \sum_{i=1}^d \delta_i \partial_{e_i} (\lambda(s, \varepsilon x))^2 + \sum_{1 \leq i, j \leq d} D_{i,j} \partial_{u_i}^2 \lambda(s, \varepsilon x) - \partial_s \lambda(s, \varepsilon x).$$

Recalling that $\lambda(t, u)$ satisfies (3.6) in $\Omega_{2\theta}$ and using the notation R_ε defined in (3.16), we have

$$\varepsilon^{d-1} \sum_{x \in \Omega_{\varepsilon k}} H(s, \varepsilon x) \int \eta^k(x)(f_s - \tilde{\Psi}_s^F) d\nu = \varepsilon^{d-1} \sum_{x \in \Omega_{\varepsilon k} \setminus \Omega_{2\theta}} H(s, \varepsilon x) \int \eta^k(x)(f_s - \tilde{\Psi}_s^F) d\nu + R_\varepsilon(s).$$

We will prove in the next section:

Lemma 3.5. *For any bounded function G defined on Λ and for any density f with respect to ν , we have for any $A > 0$*

$$\begin{aligned}
 &\varepsilon^{d-1} \sum_{\varepsilon x \in \Omega_{\varepsilon k} \setminus \Omega_{2\theta}} G(\varepsilon x) \int \eta^k(x) (f_s - \tilde{\Psi}_s^F) (\eta) d\nu \\
 &\leq A\varepsilon^{d-4} \left\{ D_b(\sqrt{h_s}, \tilde{\Psi}_s \nu) + D_0(\sqrt{f_s}, \nu) \right\} + C\theta^3 \|G\|_\infty \left(\frac{\|G\|_\infty}{A} + 1 \right)
 \end{aligned}$$

for some positive constant C .

Therefore, noticing that $\|H\|_\infty \leq \text{const } \theta^{-2}$, we get for any small enough $A > 0$

$$\begin{aligned}
 \varepsilon^{-2}s_1(t) &\leq K_1\varepsilon^{-2} \int_0^t s_1(u)du \\
 &\quad - K_2\varepsilon^{d-4} \int_0^t ds \left\{ D_0(\sqrt{h_s^F}, \tilde{\Psi}_s^F \nu) + D_b(\sqrt{h_s}, \tilde{\Psi}_s \nu) \right\} \\
 &\quad + AK_3\varepsilon^{d-4} \int_0^t ds D_0(\sqrt{f_s}, \nu) + r(\varepsilon, k, n, A) + c \left(A + \frac{\theta}{A} \right),
 \end{aligned} \tag{3.25}$$

where the constants previously defined may have changed their values.

To conclude, we use Lemma 3.3 to control the Dirichlet form $\varepsilon^{d-4}D_0(\sqrt{f_s}, \nu)$ with $\varepsilon^{d-4}D_0(\sqrt{h_s^F}, \tilde{\Psi}_s^F \nu)$. The error obtained from this replacement is a constant but we notice that there is the small factor A in front of this term. Therefore there exists a constant $c' > 0$ such that for any small enough $A > 0$,

$$\begin{aligned} \varepsilon^{-2}s_1(t) &\leq K_1\varepsilon^{-2} \int_0^t s_1(u)du \\ &\quad - K_2\varepsilon^{d-4} \int_0^t ds \left\{ D_0(\sqrt{h_s^F}, \tilde{\Psi}_s^F \nu) + D_b(\sqrt{h_s}, \tilde{\Psi}_s \nu) \right\} \\ &\quad + r(\varepsilon, k, n, A) + c' \left(A + \frac{\theta}{A} \right), \end{aligned}$$

then we choose $A = A(\theta)$ vanishing with θ in such a way that $\lim_{\theta \rightarrow 0} \theta A^{-1}(\theta) = 0$ (e.g. $A(\theta) = \sqrt{\theta}$). Finally we consider the successive limits $\varepsilon \rightarrow 0$, $\ell \rightarrow \infty$, $n \rightarrow \infty$, $\theta \rightarrow 0$ and we apply Gronwall lemma. \square

4. ESTIMATES ON BOUNDARY TERMS

Proof of Lemma 3.2. From the definition of $\widehat{W}_{x,i}$ we have that the term E_1^4 in (3.18) can be written as

$$\varepsilon^{d-2} \int \left\{ \sum_{i=1}^d \sum_{x \in \Gamma_\varepsilon^k} H^{\varepsilon,i}(t, \varepsilon x) \frac{1}{|\Lambda_k|} \sum_{y \in \bar{\Lambda}_k(x)} \widehat{W}_{y,i} \right\} (f_t - \tilde{\Psi}_t^F)(\eta) d\nu$$

where $\bar{\Lambda}_k(x)$ is the block of rectangular shape which is the set of $y \in \Lambda_k(x)$ (cube centered in x) such that y and $y + e_1$ belong to Λ_ε . Let $\bar{M}_{x,i}^k(\eta)$ be the conditional expectation of $W_{x,i}$, $x \in \Lambda_\varepsilon$ given the density of particles on $\bar{\Lambda}_k(x)$. Repeating the argument given in the paragraph following inequality (3.23), it suffices to prove that, for all $A > 0$, for any bounded function J and any probability density f with respect to ν ,

$$\begin{aligned} \lim_{n \rightarrow \infty} \lim_{\varepsilon \rightarrow 0} \left\{ \varepsilon^{d-2} \int \left(\sum_{i=1}^d \sum_{x \in \Gamma_\varepsilon^k} J(\varepsilon x) \left[(\widehat{W}_{0,i} * \omega)(x) - \bar{M}_{x,i}^k(\eta) \right] \right) f d\nu \right. \\ \left. - A\varepsilon^{d-4}D_0(\sqrt{f}) \right\} = 0. \end{aligned}$$

We need the following definition:

Let μ_ℓ be the canonical measure in the block Λ_ℓ with given density η^ℓ . For any $\{x, y\} \subset \Lambda_\ell$, we introduce the Dirichlet form $D_0^b(h)$ as

$$D_0^{\{x,y\}}(h) = \int r_{\{x,y\}}(\eta) (h(\eta^{\{x,y\}}) - h(\eta))^2 d\mu_\ell$$

and we define the finite volume variance

$$V_\ell(G, \eta^\ell) = (2\ell + 1)^{-d} \left\langle \sum_{|x| \leq \ell} (\tau_x G - E[G|\eta^\ell]) (-\mathcal{L}_\ell^{(s)})^{-1} \sum_{|x| \leq \ell} (\tau_x G - E[G|\eta^\ell]) \right\rangle_{\mu_\ell}.$$

where $\mathcal{L}_\ell^{(s)}$ is the symmetric part of the generator $\mathcal{L}_{\varepsilon,0}$ restricted to the box Λ_ℓ . The proof of Theorem 4.6 in [EMY1] is based on the following result:

Lemma 4.1. *For any cylinder function h there exist a constant C and a function $C(h, \ell)$ vanishing for $\ell \rightarrow \infty$ such that for any positive A and $\bar{\ell} = \ell^{d+2}$*

$$\begin{aligned} \int \omega * [\tau_x h - E[h|\eta^k(x)]] f_t d\mu_k - A \frac{1}{|\Lambda_k|} \sum_{\{y,z\} \subset \Lambda_k(x)} D_0^{\{y,z\}}(\sqrt{f}) \\ \leq \frac{C}{A} \varepsilon^2 \int V_{\bar{\ell}}(h, \eta^{\bar{\ell}}) f_t d\mu_k + \varepsilon^2 C(h, \ell) \end{aligned}$$

The proof of this Lemma is given in [EMY1] for square blocks but it can be extended easily to a rectangular shape provided that the volume of the block is of order k^d . Using Lemma 4.1 we prove Lemma 3.2 by taking the expectation with respect to ν , multiplying by $\varepsilon^{d-2} J$, summing over $x \in \Gamma_\varepsilon^k$ and noting that the number of terms in Γ_ε^k is $\varepsilon^{-d+1} k$.

Proof of Lemma 3.4. Fix $\varepsilon^{-1} - 2k \leq x_1 \leq \varepsilon^{-1} - k$ and write the sum as

$$\begin{aligned} \varepsilon^{d-2} \int \sum_{y \in \pi_\varepsilon^{d-1}} G(\varepsilon(x_1, y)) \eta^k(x_1, y) (f_t - \tilde{\Psi}_t^F) d\nu \\ = \frac{\varepsilon^{d-2}}{k} \sum_{|z_1| \leq k} \int \sum_{y \in \pi_\varepsilon^{d-1}} (G(\varepsilon(x_1, y)))^k \eta(x_1 + z_1, y) (f_t - \tilde{\Psi}_t^F) d\nu, \end{aligned}$$

where $(G(\varepsilon(x_1, y)))^k$ is the $(d-1)$ -dimensional convolution in the variable $y \in \pi_\varepsilon^{d-1}$. Therefore, to prove the lemma, it is enough to show that, if $H : \mathbb{R} \rightarrow \mathbb{R}$ is a bounded function, then for every $\varepsilon^{-1} - 3k \leq z_1 \leq \varepsilon^{-1}$, and $A > 0$

$$\begin{aligned} \varepsilon^{d-2} \int \sum_{y \in \pi_\varepsilon^{d-1}} H(\varepsilon(z_1, y)) \eta(z_1, y) (f_t - \tilde{\Psi}_t^F) d\nu \\ \leq \varepsilon^{d-4} \frac{A}{2} \left(D_b(\sqrt{h_t}, \tilde{\Psi}_t \nu) + D_0(\sqrt{f_t}, \nu) \right) + \text{const } \varepsilon k \|G\|_\infty \left(\frac{\|G\|_\infty}{A} + 1 \right), \end{aligned}$$

Let $\varepsilon^{-1} - 3k \leq z_1 \leq \varepsilon^{-1}$ and decompose the left hand side of the last inequality into two terms B_1 and B_2

$$\begin{aligned} \varepsilon^{d-2} \int \sum_{y \in \pi_\varepsilon^{d-1}} H(\varepsilon(z_1, y)) (\eta(z_1, y) - \eta(\varepsilon^{-1}, y)) (f_t - \tilde{\Psi}_t^F) d\nu \\ + \varepsilon^{d-2} \int \sum_{y \in \pi_\varepsilon^{d-1}} H(\varepsilon(z_1, y)) \eta(\varepsilon^{-1}, y) (f_t - \tilde{\Psi}_t) d\nu \\ := B_1 + B_2. \end{aligned}$$

The term B_2 is the simplest one. From an integration by parts and Schwartz inequality, it is bounded, for all $A > 0$, by

$$B_2 \leq \varepsilon^{d-2} \sum_{y \in \pi_\varepsilon^{d-1}} \left\{ \frac{A}{2} \varepsilon^{-2} D_{(\varepsilon^{-1}, y)}^b(\sqrt{h_t}, \tilde{\Psi}_t \nu) + \varepsilon^2 \frac{\text{const}}{A} \|H\|_\infty^2 \right\}, \quad (4.26)$$

where for a function f , a positive measure μ and $x \in \Gamma_\varepsilon$

$$D_x^b(f, \mu) = \langle (f(\eta^x) - f(\eta))^2 \rangle_\mu.$$

We now consider B_1 . By Schwartz inequality

$$\begin{aligned} B_1 &= \varepsilon^{d-2} \int \sum_{y \in \pi_\varepsilon^{d-1}} \sum_{z=z_1}^{\varepsilon^{-1}-1} H(\varepsilon(z_1, y)) (\eta(t+z+1, y) - \eta(t+z, y)) (f_t - \tilde{\Psi}_t^F) d\nu \\ &\leq \varepsilon^{d-2} \left\{ \frac{\varepsilon^{-2} A}{2} D_0(\sqrt{f_t}, \nu) + \frac{\varepsilon^{-d+3} k}{2A} \|H\|_\infty^2 \right\} + \varepsilon k C \|H\|_\infty \end{aligned}$$

for some constant $C > 0$, where we have used in the last inequality, for the second term corresponding to the integration with respect to $\tilde{\Psi}_t^F \nu$, integration by parts and Taylor expansion. \square

Proof of Lemma 3.5. The summation over the set $\{x, \varepsilon x \in \Omega_{\varepsilon k} \setminus \Omega_{2\theta}\}$ can be divided in two similar terms. We consider the one where the first coordinate is such that $\varepsilon^{-1}(1-2\theta) \leq x_1 \leq \varepsilon^{-1} - k$, the second term is handled in the same way. If we repeat the arguments used in the proof of the Lemma 3.4, we obtain for $\varepsilon^{-1}(1-2\theta) \leq x_1 \leq \varepsilon^{-1} - k$,

$$\begin{aligned} &\varepsilon^{d-1} \sum_{y \in \pi_\varepsilon^{d-1}} G(\varepsilon(x_1, y)) \int \eta^k(x_1, y) (f_t(\eta) - \tilde{\Psi}_t) d\nu \\ &\leq \varepsilon \left\{ \varepsilon^{d-4} \frac{A}{2\theta} \left(D_0(\sqrt{f_t}, \nu) + D_b(\sqrt{h_t}, \tilde{\Psi}_t \nu) \right) + C\theta^2 \|G\|_\infty \left(\frac{\|G\|_\infty}{A} + 1 \right) \right\}. \end{aligned}$$

To conclude the proof of the lemma, we just have to take the sum over x_1 . \square

5. COMMENTS

We conclude with a few generalizations:

1) Assumption $\delta_1 > 0$.

If $\delta_1 = 0$ we need to introduce a different boundary generator to fix the density on the boundary. We make the choice which is usually done for the symmetric case, a death and birth process $\bar{\mathcal{L}}_b$ acting on each site of the boundary such that it is reversible with respect to the one site measure with density $\rho(\varepsilon x) = 1/2 + \varepsilon b(\varepsilon x)$

$$\bar{\mathcal{L}}_b f(\eta) = \sum_{x \in \Gamma_\varepsilon} \left[\rho(\varepsilon x) (1 - \eta_x) [f(\eta^{x,+}) - f(\eta)] + (1 - \rho(\varepsilon x)) \eta_x [f(\eta^{x,-}) - f(\eta)] \right]$$

where $\eta^{x,+}$ and $\eta^{x,-}$ are the configurations obtained from η by adding or removing a particle at site x .

2) General domain.

We generalize now the model to the case of a macroscopic system in a smooth bounded convex domain of \mathbb{R}^d . We introduce the boundary generator acting on the boundary Γ_ε as in Section 2 with the boundary rates $C^b(\varepsilon x, \eta)$ chosen in such a way that a particle is added at the site x on the boundary when the site is empty, with rate $|\delta \cdot n|(1/2 + \varepsilon b(\varepsilon x))$ if $\delta \cdot n$, the scalar product of the vector δ and the outward normal in x , is positive and removed when the site is occupied, with rate $|\delta \cdot n|(1/2 - \varepsilon b(\varepsilon x))$ if the scalar product is negative. For $x \in \Gamma_\varepsilon$

$$C^b(\varepsilon x, \eta) = |\delta \cdot n| \left(\frac{1}{2} + \varepsilon b(\varepsilon x) \right) (1 - \eta(x)) \mathbb{1}\{\delta \cdot n(\varepsilon x) > 0\} \\ + |\delta \cdot n| \left(\frac{1}{2} - \varepsilon b(\varepsilon x) \right) \eta(x) \mathbb{1}\{\delta \cdot n(\varepsilon x) < 0\}.$$

These rates fix the value of the particle density on the macroscopic boundary to be $1/2 + \varepsilon b(x)$. This choice is sufficient to remove the entropy flow generated by the drift of ASEP. Since $\delta \cdot n(\varepsilon x)$ can be zero in some points for a general domain we have to add, on the basis of the previous remark, also a reversible boundary generator to fix the density in these sites.

3) Different boundary conditions.

To remove the entropy flux it is possible to choose a generator $\bar{\mathcal{L}}_b$ which is reversible instead of using the non reversible generator $\mathcal{L}_{\varepsilon,b}$, but then we have to speed up it by a factor ε^{-3} with respect to the jump process. The total generator speeded up is then

$$\varepsilon^{-2} \mathcal{L}_\varepsilon = \varepsilon^{-2} \mathcal{L}_{\varepsilon,0} + \varepsilon^{-5} \bar{\mathcal{L}}_b.$$

We note that the bad boundary terms in our proof are eliminated by means of a cancellation. If we use only the reversible generator $\varepsilon^{-2} \bar{\mathcal{L}}_b$ on the boundary they will be controlled by the Dirichlet form associated to $\bar{\mathcal{L}}_b$, by a generalization of the Lemma 3.3.

Acknowledgments. O.B. and M.M. would like to thank the hospitality of the University of Roma Tor Vergata and R.E and R. M. the hospitality of the University of Rouen. This work has been partially supported by GNFM-INdAM and MURST.

REFERENCES

- [D] B. Derrida *A exactly soluble non-equilibrium system: The asymmetric simple exclusion process*, Phys. Rep. **301**, 65–83 (1998).
- [DLS] B. Derrida J. L. Lebowitz and E.R. Speer *Exact large deviation functional for the density profile in a stationary nonequilibrium open system*, preprint (2001).
- [BDGJL] L. Bertini, A. De Sole, D. Gabrielli, G. Jona-Lasinio and C. Landim. *Fluctuations in stationary nonequilibrium states*, preprint (2001).
- [ELS] G. Eyink, J. L. Lebowitz and H. Spohn *Hydrodynamics of Stationary Nonequilibrium States for Some Lattice Gas Models*, Commun. Math. Phys., Vol. **132**, 252-283 (1990). *Lattice Gas Models in*

- contact with Stochastic Reservoirs: Local Equilibrium and Relaxation to the Steady State*, Commun. Math. Phys., Vol. **140**, 119–131 (1991).
- [EMY1] R. Esposito, R. Marra and H.T. Yau *Diffusive limit of asymmetric simple exclusion*, Review in Math. Phys. **6**, 1233-1267 (1994)
- [EMY2] R. Esposito, R. Marra and H.T. Yau *Navier-Stokes equations for stochastic particle systems on the lattice*, Commun. Math. Phys., **182**, 395–456 (1996).
- [KLO] C. Kipnis, C. Landim and S.Olla *Macroscopic properties of a stationary nonequilibrium distribution for a non-gradient interacting particle system*, Ann. Inst. H. Poincaré **31**, 191–221 (1995). 1998.
- [K] J. Krug, *Boundary-Induced Phase transitions in Driven Diffusive Systems*, Phys. Rev. Lett. **67** 1882–(1991).
- [LMS] C. Landim, M. Mourragui and S. Sellami, *Hydrodynamical limit for a nongradient interacting particle system with stochastic reservoirs* Probab. Theory and Appl. **45**, N. 4, 2000.
- [LY] C. Landim, and H.T. Yau, *Fluctuation–dissipation equation of asymmetric simple exclusion processes* Probability Theory and Related Fields **108** 321–(1997)
- [SZ] B. Schmittmann and R.K.Zia, *Statistical Mechanics of Driven Diffusive Systems*, in Phase Transitions and Critical Phenomena **17**, C.Domb and J.L.Lebowitz ed.s, (1995)
- [Sp] H. Spohn, *Large scale dynamics of interacting particles*, Springer, Berlin, (1991).
- [Y] H. T. Yau, *Relative entropy and hydrodynamics of Ginsburg-Landau models* Letters Math. Phys., **22**, 63–80, (1991).

O. Benois and M. Mourragui

LABORATOIRE DE MATHÉMATIQUES RAPHAËL SALEM, UMR 6085, UNIVERSITÉ DE ROUEN, 76821 MONT SAINT AIGNAN, FRANCE

R. Esposito

CENTRO INTERDISCIPLINARE LINCEO “BENIAMINO SEGRE”, VIA DELLA LUNGARA 10, 00165 ROMA, ITALY; on leave from DIPARTIMENTO DI MATEMATICA PURA ED APPLICATA, UNIVERSITÀ DI L’AQUILA, 67100 COPPITO, AQ, ITALY

R. Marra

DIPARTIMENTO DI FISICA E UNITÀ INFN, UNIVERSITÀ DI ROMA TOR VERGATA, VIA DELLA RICERCA SCIENTIFICA, 00133 ROMA, ITALY