

# On the uncertainty principle for proper time and mass

B. RAM

Physics Department, New Mexico State University  
Las Cruces, NM 88003, USA<sup>†</sup>

and

Umrao Institute of Fundamental Research  
A2/214 Janak Puri, New Delhi, 110 058, India

## Abstract

In [J. Math. Phys. **40**, 1237 (1999)] Kudaka and Matsumoto derive the uncertainty relation  $c^2\Delta m\Delta\tau \geq \hbar/2$  between the rest mass  $m$  and the proper time  $\tau$ , by considering the Lagrangian  $M(\dot{\tau} - c^{-1}\sqrt{-g_{\mu\nu}\dot{x}^\mu\dot{x}^\nu}) + eA_\mu(x)\dot{x}^\mu$ . In this note we give an alternative derivation based on a special case of the time-like geodesic equation obtained using the general relativistic Lagrangian  $g_{\mu\nu}\frac{dx^\mu}{d\tau}\frac{dx^\nu}{d\tau}$ .

---

<sup>†</sup> correspondence address

In an interesting paper [1] published in this journal, Kudaka and Matsumoto derive the uncertainty relation

$$c^2 \Delta m \Delta \tau \geq \frac{\hbar}{2} \quad (1)$$

between the rest mass  $m$  and the proper time  $\tau$ , by considering a Lagrangian in which the proper time is included as a dynamic variable like the positions  $x^i$ . Specifically, they consider the Lagrangian

$$L = M(\dot{\tau} - c^{-1} \sqrt{-g_{\mu\nu}(x) \dot{x}^\mu \dot{x}^\nu}) + e A_\mu(x) \dot{x}^\mu \quad (2)$$

and find, as a result of their analysis, that the energy  $E = mc^2$  is the generalized momentum conjugate to the proper time, and that  $\tau$ ,  $E$ ,  $x^i$ , and  $p_i$  are canonical variables of the system. Consequently the corresponding operators

$$\hat{\tau}, \hat{E}, \hat{x}^i, \hat{p}_i \quad (i = 1, 2, 3),$$

satisfy the commutation relations

$$[\hat{E}, \hat{\tau}] = [\hat{x}^i, \hat{p}_i] = i\hbar, \quad (3)$$

the relation  $[\hat{E}, \hat{\tau}] = i\hbar$  in (3) giving the uncertainty relation (1).

In the present note we give an alternative derivation of Eq. (1).

It is well-known [2] that in general relativity the equations governing the time-like geodesics in a space-time with the line element

$$ds^2 = g_{\mu\nu} dx^\mu dx^\nu \quad (4)$$

can be derived from the Lagrangian

$$2\mathcal{L} = g_{\mu\nu} \frac{dx^\mu}{d\tau} \frac{dx^\nu}{d\tau}, \quad (5)$$

where  $\tau$  is the proper time. For the Schwarzschild space-time, the Lagrangian is (from now on we use units in which  $c = \hbar = G = 1$ )

$$\mathcal{L} = \frac{1}{2} \left[ \left(1 - \frac{2m}{r}\right) \dot{t}^2 - \frac{\dot{r}^2}{1 - \frac{2m}{r}} - r^2 \dot{\theta}^2 - (r^2 \sin^2 \theta) \dot{\phi}^2 \right], \quad (6)$$

where the dot denotes differentiation with respect to  $\tau$ . The corresponding canonical momenta are

$$p_t = \frac{\partial \mathcal{L}}{\partial \dot{t}} = \left(1 - \frac{2m}{r}\right) \dot{t}, \quad (7a)$$

$$p_r = -\frac{\partial \mathcal{L}}{\partial \dot{r}} = \left(1 - \frac{2m}{r}\right)^{-1} \dot{r}, \quad (7b)$$

$$p_\phi = -\frac{\partial \mathcal{L}}{\partial \dot{\phi}} = (r^2 \sin^2 \theta) \dot{\phi}, \quad (7c)$$

and

$$p_\theta = -\frac{\partial \mathcal{L}}{\partial \dot{\theta}} = r^2 \dot{\theta}. \quad (7d)$$

The resulting Hamiltonian is

$$H = p_t \dot{t} - (p_r \dot{r} + p_\theta \dot{\theta} + p_\phi \dot{\phi}) - \mathcal{L} = \mathcal{L}. \quad (8)$$

From the equality of the Hamiltonian and the Lagrangian it follows that

$$H = \mathcal{L} = \text{constant}. \quad (9)$$

For time-like geodesics,  $2\mathcal{L}$  has the value +1. Integrating the equations

$$\frac{dp_t}{d\tau} = \frac{\partial \mathcal{L}}{\partial t} = 0 \quad \text{and} \quad \frac{dp_\phi}{d\tau} = -\frac{\partial \mathcal{L}}{\partial \phi} = 0 \quad (10)$$

one gets

$$p_t = \left(1 - \frac{2m}{r}\right) \dot{t} = \text{constant} = \mathcal{E} \quad (\text{say}) \quad (11)$$

and

$$p_\phi = r^2 \sin^2 \theta \dot{\phi} = \text{constant}. \quad (12)$$

Moreover, from the equation

$$\frac{dp_\theta}{d\tau} = \frac{d}{d\tau}(r^2\dot{\theta}) = -\frac{\partial\mathcal{L}}{\partial\theta} = (r^2 \sin\theta \cos\theta)\dot{\phi}^2, \quad (13)$$

it follows that if we choose  $\theta = \pi/2$  when  $\dot{\theta} = 0$ , then  $\ddot{\theta}$  will also be zero; and  $\theta$  will maintain the value  $\pi/2$ . In other words, the geodesic is described in an invariant plane given by  $\theta = \pi/2$ . Equation (12) then gives

$$p_\phi = r^2\dot{\phi} = \text{constant} = L \quad (\text{say}), \quad (14)$$

where  $L$  denotes the angular momentum about an axis normal to the invariant plane, say the  $x$ - $y$  plane. With (11), (14) and  $\mathcal{L} = 1/2$ , Eq. (6) for the time-like geodesic becomes

$$\frac{1}{2} \left[ \frac{\mathcal{E}^2}{1 - \frac{2m}{r}} - \frac{\dot{r}^2}{1 - \frac{2m}{r}} - \frac{L^2}{r^2} \right] = \frac{1}{2}. \quad (15)$$

For the special case when both the constants of integration  $\mathcal{E}$  and  $L$  are zero, Eq. (15) reduces to

$$\frac{1}{2}\dot{r}^2 - \frac{m}{r} = -\frac{1}{2}. \quad (16)$$

It is easy to see that Eq. (16) describes the region  $r \leq 2m$ ,  $\dot{r}$  being zero at  $r = 2m$ .

For the Kerr space-time the Lagrangian in the equatorial plane (for which  $\dot{\theta} = 0$  and  $\theta = \text{a constant} = \pi/2$ ) is [2]

$$2\mathcal{L} = \left(1 - \frac{2m}{r}\right)\dot{t}^2 + \frac{4am}{r}\dot{t}\dot{\phi} - \frac{r^2}{\Delta}\dot{r}^2 - \left[(r^2 + a^2) + \frac{2a^2m}{r}\right]\dot{\phi}^2, \quad (17)$$

where  $a$  is the angular momentum per unit mass of the inner region and

$$\Delta = r^2 - 2mr + a^2. \quad (18)$$

Following (see pp. 326-328 of ref. [2]) in an analogous manner as in the Schwarzschild case, one obtains, in terms of the constants of integration  $\mathcal{E}$  and  $L$ ,

$$\frac{1}{2}\dot{r}^2 - \frac{m}{r} + \frac{1}{2}(1 - \mathcal{E}^2) \left(1 + \frac{a^2}{r^2}\right) + \frac{L^2}{2r^2} - \frac{m}{r^3}(L - a\mathcal{E})^2 = 0 \quad (19)$$

as the equation for the time-like geodesic. For the special case when both the constants of integration  $\mathcal{E}$  and  $L$  are zero, Eq. (19) reduces to

$$\frac{1}{2}\dot{r}^2 + \frac{a^2}{2r^2} - \frac{m}{r} = -\frac{1}{2}. \quad (20)$$

Note that Eq. (20) reduces to Eq. (16) when  $a = 0$ , signifying that the Schwarzschild solution is a special case of the Kerr solution. Note also that Eq. (20) can be rewritten as

$$\frac{1}{2}\dot{\vec{r}}^2 - \frac{m}{r} = -\frac{1}{2}, \quad (21)$$

$\vec{r}$  being given the meaning of a Euclidean vector. Equation (21) is simply a generalization of Eq. (16). It describes the region  $r \leq 2m$ ,  $\dot{\vec{r}}$  being zero at  $r = 2m$ . With

$$x_1 = r \sin \theta_{in} \cos \phi_{in}, \quad x_2 = r \sin \theta_{in} \sin \phi_{in}, \quad x_3 = r \cos \theta_{in},$$

$$r^2 = x_1^2 + x_2^2 + x_3^2, \quad \text{and} \quad v^2 = \dot{x}_1^2 + \dot{x}_2^2 + \dot{x}_3^2 = \dot{\vec{r}}^2,$$

Eq. (21) is rewritten as

$$\frac{1}{2}v^2 - \frac{m}{r} = -\frac{1}{2}. \quad (22)$$

It is also well-known [3] that the three-dimensional Kepler problem, Eq. (22), is equivalent

to a four-dimensional harmonic oscillator. Below we briefly sketch how it is so, by use of a matrix transformation known as the Kustaanheimo-Stiefel (KS) transformation [4].

Let us define the column matrices

$$X = \begin{pmatrix} x_1 \\ x_2 \\ x_3 \\ 0 \end{pmatrix} \quad \text{and} \quad S = \begin{pmatrix} s_1 \\ s_2 \\ s_3 \\ s_4 \end{pmatrix}, \quad (23)$$

$$\dot{X} = \begin{pmatrix} \dot{x}_1 \\ \dot{x}_2 \\ \dot{x}_3 \\ 0 \end{pmatrix} \quad \text{and} \quad \dot{S} = \begin{pmatrix} \dot{s}_1 \\ \dot{s}_2 \\ \dot{s}_3 \\ \dot{s}_4 \end{pmatrix}, \quad (24)$$

where  $x_1, x_2, x_3$  and  $s_1, s_2, s_3, s_4$  are, respectively the three-dimensional and four-dimensional Cartesian coordinates, and  $\dot{x}_1, \dot{x}_2, \dot{x}_3$  and  $\dot{s}_1, \dot{s}_2, \dot{s}_3, \dot{s}_4$  are the corresponding velocity components. The KS transformation which transforms the coordinates is given by

$$X = AS, \quad (25)$$

and that which transforms the velocities (or momenta) is given by

$$\dot{X} = \frac{1}{2}s^{-2}A\dot{S}, \quad (26)$$

where  $s^2 = s_1^2 + s_2^2 + s_3^2 + s_4^2$ ,

$$A = \begin{pmatrix} s_3 & -s_4 & s_1 & -s_2 \\ s_4 & s_3 & s_2 & s_1 \\ s_1 & s_2 & -s_3 & -s_4 \\ s_2 & -s_1 & -s_4 & s_3 \end{pmatrix}, \quad (27)$$

and

$$s^{-2} \tilde{A} A = \mathbf{1}, \quad (28)$$

$\tilde{A}$  being the transposed matrix and  $\mathbf{1}$  the unit matrix. It is to be emphasized that (25) and (26) are *independent* transformations. From Eqs. (25), (26) and (28), one obtains

$$r^2 = x_1^2 + x_2^2 + x_3^2 = s^4 \quad (29)$$

and

$$v^2 = \dot{x}_1^2 + \dot{x}_2^2 + \dot{x}_3^2 = \frac{1}{4s^2} \dot{s}^2 = \frac{1}{4s^2} (\dot{s}_1^2 + \dot{s}_2^2 + \dot{s}_3^2 + \dot{s}_4^2). \quad (30)$$

Using (29) and (30) it is straightforward to show that Eq. (21) transforms into

$$\frac{1}{2} m_{ho} \dot{s}^2 + \frac{1}{2} m_{ho} \omega^2 s^2 = m, \quad (31)$$

with  $m_{ho} \equiv$  mass of the four-dimensional harmonic oscillator  $=1/4$  and  $\omega = 2$ .

Note that in Eq. (31)  $m$  corresponds to the total classical energy ( $E$ ), the proper time  $\tau$  to the Newtonian time ( $t$ ), and the quantum equation that corresponds to (31) is

$$-\frac{1}{2m_{ho}} \sum_{i=1}^4 \left( \frac{\partial^2}{\partial s_i^2} - m_{ho}^2 \omega^2 s^2 \right) \psi = (n+1) \psi \quad (32)$$

In (32)  $(n+1)$ ,  $n = 0, 1, \dots$  are the eigenvalues of the operator  $m$ . Consequently the Heisen-

berg uncertainty relation [5]  $\Delta E \Delta t \geq \hbar/2$  (inserting  $c$  and  $\hbar$ ) simply translates into

$$c^2 \Delta m \Delta \tau \geq \frac{\hbar}{2}.$$

For application of the quantum equations that correspond to (16) and (21) to the black hole physics, the interested reader may refer to refs. [6-8].

### Acknowledgements

The author thanks S.R. Choudhury, N. Mukunda and S.M. Roy for conversations, R.S. Bhalerao for a critical reading of the manuscript leading to an improved version and the Tata Institute for a pleasant stay.



## References

- [1] S. Kudaka and S. Matsumoto, *J. Math. Phys.* **40**, 1237 (1999).
- [2] S. Chandrasekhar, *The Mathematical Theory of Black Holes*, (Oxford University Press, Oxford, 1983).
- [3] A.O. Barut, C.K.E. Schneider, and R. Wilson, *J. Math. Phys.* **20**, 2244 (1979); A.C. Chen, *J. Math. Phys.* **23**, 412 (1982); A.C. Chen, *Am. J. Phys.* **55**, 250 (1987); H.A. Mavromatis, *Rep. Math. Phys.* **40**, 17 (1997).
- [4] P. Kustaanheimo and E. Stiefel, *J. Reine Angew. Math.* **218**, 204 (1965).
- [5] W. Heisenberg, *Zeit. für Physik*, **43**, 172 (1927).
- [6] B. Ram, *Phys. Lett. A* **265**, 1 (2000), LANL Archive gr-qc/9908036.
- [7] B. Ram, LANL Archive gr-qc/0101056.
- [8] B. Ram, LANL Archive hep-th/0208128.