# An Interaction of An Oscillator with An One-Dimensional Scalar Field.

## Simple Exactly Solvable Models based on Finite Rank Perturbations Methods.

## II: Resolvents formulae

#### S.A. Chorošavin

#### Abstract

This paper is an electronic application to my set of lectures, subject: 'Formal methods in solving differential equations and constructing models of physical phenomena'. Addressed, mainly: postgraduates and related readers. Content: a very detailed discussion of the simple model of interaction based on the equation array:

$$z \begin{pmatrix} q \\ u \end{pmatrix} - \begin{pmatrix} -\Omega^2 & \Omega^2 < l|_2 \\ 4\gamma_c \delta_{\alpha,x_0} < 1|_1 & B - 4\gamma_c \delta_{\alpha,x_0} < l|_2 \end{pmatrix} \begin{pmatrix} q \\ u \end{pmatrix} = \begin{pmatrix} w_1 \\ w_2 \end{pmatrix}$$

Besides, less detailed discussion of related models. Central mathematical points: Finite Rank Perturbations Methods, Resolvents formulae, Donoghuelike models, Friedrichs-like models. Central physical points: phenomenon of Resonance and notion of Second Sheet.

Hereafter I use a P.A.M. Dirac's "bra-ket" syntax and suppose that B stands for an abstract linear operator, l for a linear functional,  $u, w_2, \delta_{\alpha, x_0}$  for abstract elements;  $q, w_1 z, \Omega, \gamma_c$  stand for numbers. q, u are objects to be found, the others are arbitrarily given.

#### Introduction.

#### A Harmonic Oscillator Coupled to an One-Dimensional Scalar Field.

In a previous paper I had discussed several models of an one-dimensional harmonic oscillator coupled to an one-dimensional scalar field. Primarily I was interested in the models which one can describe by the equation array

$$\begin{aligned} \ddot{q} &= -\Omega^2(q-Q) + f_0(t) \\ \ddot{u} &= Bu - 4\gamma_c \left(\delta_{\alpha,t,x_0}\right) \cdot \left(Q-q\right) + f_1(t) \\ Q &= Q(t) = \langle l | u(t) \rangle \end{aligned}$$

as well as the models which one can describe by the equation array

$$\begin{aligned} \ddot{q}_0 &= -\Omega^2 q_0 + \gamma_1 Q_\phi + f_0(t) \\ \ddot{\phi} &= B\phi + 4\gamma_{2,c} \Big(\delta_{\alpha,t,x_0}\Big) \cdot q_0 + f_1(t) \\ Q_\phi &= Q_\phi(t) = \langle l | \phi(t) \rangle \end{aligned}$$

Hereafter I use a P.A.M. Dirac's "bra-ket" syntax and suppose that q and Q,  $q_0$  and  $Q_{\phi}$  are usual (one-dimensional) functions of t:

$$q = q(t)$$
,  $Q = Q(t)$ ,  $q_0 = q_0(t)$ ,  $Q_{\phi} = Q_{\phi}(t)$ ,

*B* is an abstract linear operator, *l* is a linear functional,  $\{u(t)\}_t$ ,  $\{\phi(t)\}_t$  and  $\{\delta_{\alpha,t,x_0}\}_t$  are families of abstract elements; of course the type of  $\delta_{\alpha,t,x_0}$  must be the same as one of u(t) or resp.  $\{\phi(t)\}_t$ .

Normally I supposed that  $\delta_{\alpha,t,x_0}$  did not depend on t, i.e. was constant in t. In that case I wrote  $\delta_{\alpha,x_0}$  instead of  $\delta_{\alpha,t,x_0}$ .

In this second paper I will suppose that  $\delta_{\alpha,t,x_0}$  is constant in t as well, and rewrite the above equations as follows:

$$\begin{pmatrix} \ddot{q} \\ \ddot{u} \end{pmatrix} = \begin{pmatrix} -\Omega^2 & \Omega^2 < l|_2 \\ 4\gamma_c \delta_{\alpha,x_0} < 1|_1 & B - 4\gamma_c \delta_{\alpha,x_0} < l|_2 \end{pmatrix} \begin{pmatrix} q \\ u \end{pmatrix} + \begin{pmatrix} f_0(t) \\ f_1(t) \end{pmatrix}$$
$$\begin{pmatrix} \ddot{q}_0 \\ \ddot{\phi} \end{pmatrix} = \begin{pmatrix} -\Omega^2 & \gamma_1 < l|_2 \\ 4\gamma_{2,c} \delta_{\alpha,x_0} < 1|_1 & B \end{pmatrix} \begin{pmatrix} q_0 \\ \phi \end{pmatrix} + \begin{pmatrix} f_0(t) \\ f_1(t) \end{pmatrix}$$

where the subscribts 1 and 2 in  $< \cdots |_1$  and  $< \cdots |_2$  mean that arguments of  $< \cdots |_1$  are elements of the first component of the vector  $\begin{pmatrix} q \\ u \end{pmatrix}$  resp.  $\begin{pmatrix} q_0 \\ \phi \end{pmatrix}$  and arguments of  $< \cdots |_2$  are elements of the second component of the suitable vector. <sup>1</sup> Of course,

 $<1|q>=<1|_1q>=q$ ,  $<1|_{q_0}>=<1|_1q_0>=q_0$ 

The subject will primarily be **resolvents formulae** i.e. the formulae that resolve the equation array:

$$z \begin{pmatrix} q \\ u \end{pmatrix} - \begin{pmatrix} -\Omega^2 & \Omega^2 < l|_2 \\ 4\gamma_c \delta_{\alpha, x_0} < 1|_1 & B - 4\gamma_c \delta_{\alpha, x_0} < l|_2 \end{pmatrix} \begin{pmatrix} q \\ u \end{pmatrix} = \begin{pmatrix} w_1 \\ w_2 \end{pmatrix}$$

<sup>&</sup>lt;sup>1</sup>often we omite these subscripts, using them primarily for emphasis.

resp.

$$z \begin{pmatrix} q_0 \\ \phi \end{pmatrix} - \begin{pmatrix} -\Omega^2 & \gamma_1 < l|_2 \\ 4\gamma_{2,c}\delta_{\alpha,x_0} < 1|_1 & B \end{pmatrix} \begin{pmatrix} q_0 \\ \phi \end{pmatrix} = \begin{pmatrix} w_1 \\ w_2 \end{pmatrix}$$

Of course, here all quantities,  $q, u, w_1, w_2, \delta_{\alpha, x_0}, l$ , etc. are supposed to be constant in t.

In the previous paper I had discussed d'Alembert-like solutions to the systems and resp. phenomena of **Radiation Reaction**, **Braking Radiation** and **Resonance**.

In this paper I will discuss Donoghue-Friedrichs-like solutions to the systems  $^2$  and resp. the phenomenon of **Resonance** and notion of the **Second Sheet**.

<sup>3</sup> 

<sup>&</sup>lt;sup>2</sup>i.e., I will discuss resolvents formulae of the system

## 1 Finite Rank Perturbations. Abstract Formulae.

#### 1.1 Simple Cases. Rank-One Perturbations and Rank-Two Perturbations

#### 1.1.1 Rank-One Perturbations.

The simplest case of the problem looks like this: Suppose, we can find  $v_0$  to an equation

$$Av_0 = w_0$$

for any given  $w_0$ , where A is an invertible linear operator, and suppose, we need to find any v to

$$Av - f_a < l_a | v > = w$$

where  $\langle l_a |$  is a given linear functional, and the elements  $f_a, w$  are given, as well. One is used to saying concisely: we need

$$(A - f_a < l_a|)^{-1}w$$

Then we do as follows:

First, write

$$Av = w + f_a < l_a | v >$$

and notice,

in order to obtain 
$$v$$
 we need to obtain ONLY  $< l_a | v >$ 

Actually,

$$v = A^{-1}(w + f_a < l_a | v >)$$
  
 $v = A^{-1}w + < l_a | v > A^{-1}f_a$ 

So,

$$v = A^{-1}w + c_a A^{-1} f_a$$
, where  $c_a := < l_a | v >$ .

Apply now this form to itself and to the recent relationship, consequently:

$$v = A^{-1}w + < l_a | v > A^{-1}f_a$$

then

$$< l_a | v > = < l_a | (A^{-1}w + < l_a | v > A^{-1}f_a) >$$

i.e.,

$$c_a = < l_a | A^{-1} w + c_a A^{-1} f_a >$$

On the other hand, to find a  $c_a$ , it is sufficient to fulfil

$$A^{-1}w + c_a A^{-1}f_a = A^{-1}w + \langle l_a | A^{-1}w + c_a A^{-1}f_a \rangle A^{-1}f_a$$

i.e.,

$$c_a A^{-1} f_a = \langle l_a | A^{-1} w + c_a A^{-1} f_a \rangle A^{-1} f_a$$

This relation is fulfilled, if

$$c_a = < l_a | A^{-1} w + c_a A^{-1} f_a >$$

It is the same equation to  $c_a$  as above. Therefore, we need to solve this only scalar equation and set  $v = A^{-1}w + c_a A^{-1}f_a$ . To do it, recall, f is linear. Hence

$$c_a = < l_a | A^{-1}w > + c_a < l_a | A^{-1}f_a >$$

Thus we have seen,

$$(1 - \langle l_a | A^{-1} f_a \rangle) c_a = \langle l_a | A^{-1} w \rangle$$

and hence

$$c_a = \frac{\langle l_a | A^{-1} w \rangle}{1 - \langle l_a | A^{-1} f_a \rangle}$$
, when  $1 - \langle l_a | A^{-1} f_a \rangle \neq 0$ .

Finally

$$(A - f_a < l_a|)^{-1}w = A^{-1}w + A^{-1}f_a \frac{\langle l_a|A^{-1}w \rangle}{1 - \langle l_a|A^{-1}f_a \rangle}, \text{ when } 1 - \langle l_a|A^{-1}f_a \rangle \neq 0.$$

#### **Rank-Two Perturbations.** 1.1.2

A problem connected with Rank-Two Perturbations looks quite similarly: We know  $A^{-1}$  and need

$$A - f_a < l_a | - f_b < l_b |)^{-1}$$

( More precisely, we need to solve the associated equation.

We can do it in two ways: we can iterate the final formulae of the previous subsubsection, and we can iterate the argumentations of one. In the former way, we first calculate

$$(A - f_a < l_a|)^{-1}$$

and then

$$(B - f_b < l_b|)^{-1}$$
 where  $B := A - f_a < l_a|$ .

Describe the latter way.

The equation to be solved is this:

$$Av - f_a < l_a | v > -f_b < l_b | v > = w$$

We rewrite it firstly as

$$Av = w + f_a < l_a | v > + f_b < l_b | v >$$

and this relation, we rewrite it as

 $v = A^{-1}(w + f_a < l_a | v > + f_b < l_b | v >) = A^{-1}w + A^{-1}f_a < l_a | v > + A^{-1}f_b < l_b | v >$ where we have used the fact that  $A^{-1}$  is linear. Thus,

$$v = A^{-1}(w + c_a f_a + c_b f_b) = A^{-1}w + c_a A^{-1} f_a + c_b A^{-1} f_b ,$$
  
where  $c_a := < l_a | v >, c_b := < l_b | v >$ 

Apply now this form to itself and to the previous relationship. Then obtain,

$$\begin{aligned} c_a A^{-1} f_a + c_b A^{-1} f_b \\ &= A^{-1} f_a < l_a | A^{-1} w + c_a A^{-1} f_a + c_b A^{-1} f_b > \\ &+ A^{-1} f_b < l_b | A^{-1} w + c_a A^{-1} f_a + c_b A^{-1} f_b > \end{aligned}$$

To fulfil these relations, it is sufficient to satisfy the relations  $^{3}$ 

$$\begin{array}{rcl} c_a &=& < l_a | A^{-1} w + c_a A^{-1} f_a + c_b A^{-1} f_b > \\ c_b &=& < l_b | A^{-1} w + c_a A^{-1} f_a + c_b A^{-1} f_b > \end{array}$$

i.e., <sup>4</sup>

$$\begin{pmatrix} 1 - \langle l_a | A^{-1} f_a \rangle & - \langle l_a | A^{-1} f_b \rangle \\ - \langle l_b | A^{-1} f_a \rangle & 1 - \langle l_b | A^{-1} f_b \rangle \end{pmatrix} \begin{pmatrix} c_a \\ c_b \end{pmatrix} = \begin{pmatrix} \langle l_a | A^{-1} w \rangle \\ \langle l_b | A^{-1} w \rangle \end{pmatrix}$$

Therefore, if  $detrminant \neq 0$ ,

$$\begin{pmatrix} c_a \\ c_b \end{pmatrix} = \begin{pmatrix} 1 - \langle l_a | A^{-1} f_a \rangle & - \langle l_a | A^{-1} f_b \rangle \\ - \langle l_b | A^{-1} f_a \rangle & 1 - \langle l_b | A^{-1} f_b \rangle \end{pmatrix}^{-1} \begin{pmatrix} \langle l_a | A^{-1} w \rangle \\ \langle l_b | A^{-1} w \rangle \end{pmatrix}$$
$$\begin{pmatrix} c_a \\ c_b \end{pmatrix} = \frac{1}{determinant} \begin{pmatrix} 1 - \langle l_b | A^{-1} f_b \rangle & \langle l_a | A^{-1} f_b \rangle \\ \langle l_b | A^{-1} f_a \rangle & 1 - \langle l_a | A^{-1} f_a \rangle \end{pmatrix} \begin{pmatrix} \langle l_a | A^{-1} w \rangle \\ \langle l_b | A^{-1} w \rangle \end{pmatrix}$$
Finally,

$$(A - f_a < l_a| - f_b < l_b|)^{-1}w = A^{-1}w + c_a A^{-1}f_a + c_b A^{-1}f_b$$
$$w = A^{-1}w + (A^{-1}f_a \oplus A^{-1}f_b) \begin{pmatrix} 1 - \langle l_a|A^{-1}f_a \rangle & -\langle l_a|A^{-1}f_b \rangle \\ -\langle l_b|A^{-1}f_a \rangle & 1 - \langle l_b|A^{-1}f_b \rangle \end{pmatrix}^{-1} \begin{pmatrix} \langle l_a|A^{-1}w \rangle \\ \langle l_b|A^{-1}w \rangle \end{pmatrix}$$

 $<sup>^3</sup>$  if  $A^{-1}f_a,A^{-1}f_b$  are linearly independent, then it is also necessary, to do ones  $^4$  recall,  $< l_a|$  and  $< l_b|$  are linear

#### 1.2 Finite Rank Perturbations of Matrix-operator.

The variant of the previous problems, we will deal, looks like this: Suppose, we can solve an equation array

$$\begin{pmatrix} A_{11} & 0 \\ 0 & A_{22} \end{pmatrix} \begin{pmatrix} v_{01} \\ v_{02} \end{pmatrix} = \begin{pmatrix} w_{01} \\ w_{02} \end{pmatrix}$$

and suppose, we need to solve an equation array

$$\begin{pmatrix} A_{11} - f_{11} < l_1|_1 & -f_{12} < l_2|_2 \\ -f_{21} < l_1|_1 & A_{22} - f_{22} < l_2|_2 \end{pmatrix} \begin{pmatrix} v_1 \\ v_2 \end{pmatrix} = \begin{pmatrix} w_1 \\ w_2 \end{pmatrix}$$

i.e.

$$\begin{pmatrix} A_{11}v_1 - f_{11} < l_1 | v_1 > -f_{12} < l_2 | v_2 > \\ -f_{21} < l_1 | v_1 > +A_{22}v_2 - f_{22} < l_2 | v_2 > \end{pmatrix} = \begin{pmatrix} w_1 \\ w_2 \end{pmatrix},$$

where, of course,  $A_{11}, A_{22}$  are two given invertible linear operators, and  $\langle l_1 |, \langle l_2 |$  are two given linear functionals; the elements  $f_{11}, f_{12}, f_{21}, f_{22}, w_1, w_2$  are given, as well.

One can show, it is exactly a rank-two case, an we can use the *final formalae* of the previous subsection, but we prefer direct reasoning and we imitate the *arguing* of the previous subsection and do as follows:

Firstly, write

$$\begin{pmatrix} A_{11}v_1 \\ A_{22}v_2 \end{pmatrix} = \begin{pmatrix} w_1 + f_{11} < l_1 | v_1 > + f_{12} < l_2 | v_2 > \\ w_2 + f_{21} < l_1 | v_1 > + f_{22} < l_2 | v_2 > \end{pmatrix}$$

and define

$$\left(\begin{array}{c} c_1\\ c_2 \end{array}\right) := \left(\begin{array}{c} < l_1 | v_1 > \\ < l_2 | v_2 > \end{array}\right)$$

Then write

$$\begin{pmatrix} A_{11}v_1 \\ A_{22}v_2 \end{pmatrix} = \begin{pmatrix} w_1 + f_{11}c_1 + f_{12}c_2 \\ w_2 + f_{21}c_1 + f_{22}c_2 \end{pmatrix}$$

Then

$$\begin{pmatrix} v_1 \\ v_2 \end{pmatrix} = \begin{pmatrix} A_{11}^{-1}w_1 + A_{11}^{-1}f_{11}c_1 + A_{11}^{-1}f_{12}c_2 \\ A_{22}^{-1}w_2 + A_{22}^{-1}f_{21}c_1 + A_{22}^{-1}f_{22}c_2 \end{pmatrix}$$

Then

$$\begin{pmatrix} c_1 \\ c_2 \end{pmatrix} = \begin{pmatrix}  \\  \end{pmatrix} = \begin{pmatrix}  \\  \end{pmatrix}$$

Recall,  $l_1, l_2$  are linear. Therefore

$$\begin{pmatrix} c_1 \\ c_2 \end{pmatrix} = \begin{pmatrix}  +  c_1 +  c_2 \\  +  c_1 +  c_2 \end{pmatrix}$$

$$\begin{pmatrix} c_1 -  c_1 -  c_2 \\ -  c_1 + c_2 -  c_2 \end{pmatrix} = \begin{pmatrix}  \\  \end{pmatrix}$$

$$\begin{pmatrix} 1 -  & -  \\ -  & 1 -  \\ -  & 1 -  \end{pmatrix} \begin{pmatrix} c_1 \\ c_2 \end{pmatrix} = \begin{pmatrix}  \\  \end{pmatrix}$$

Thus we conclude

$$\begin{pmatrix} v_1 \\ v_2 \end{pmatrix} = \begin{pmatrix} A_{11}^{-1}w_1 \\ A_{22}^{-1}w_2 \end{pmatrix} + \begin{pmatrix} A_{11}^{-1}f_{11} & A_{11}^{-1}f_{12} \\ A_{22}^{-1}f_{21} & A_{22}^{-1}f_{22} \end{pmatrix} \begin{pmatrix} c_1 \\ c_2 \end{pmatrix}$$

$$\begin{pmatrix} c_1 \\ c_2 \end{pmatrix} = \frac{1}{det} \begin{pmatrix} 1 - \langle l_2 | A_{22}^{-1} f_{22} \rangle & \langle l_1 | A_{11}^{-1} f_{12} \rangle \\ \langle l_2 | A_{22}^{-1} f_{21} \rangle & 1 - \langle l_1 | A_{11}^{-1} f_{11} \rangle \end{pmatrix} \begin{pmatrix} \langle l_1 | A_{11}^{-1} w_1 \rangle \\ \langle l_2 | A_{22}^{-1} w_2 \rangle \end{pmatrix}$$

where

$$det := (1 - \langle l_1 | A_{11}^{-1} f_{11} \rangle)(1 - \langle l_2 | A_{22}^{-1} f_{22} \rangle) - \langle l_1 | A_{11}^{-1} f_{12} \rangle \langle l_2 | A_{22}^{-1} f_{21} \rangle$$

Of course, we need here  $det \neq 0$ . Note, we have obtained, that the values of  $c_1, c_2$  must be such, as they are written. But do they exist at all? Are the found  $c_1, c_2$  a solution of the problem? The answer is: yes. The way to verify it is plain: imitate the corresponding part of the rank-one case,– we omit details.

Finally we remark. If  $A_{11}$  or  $A_{22}$  are not invertible, but if  $A_{11} - f_{11} < l_1|_1$  or  $A_{22} - f_{22} < l_2|_2$ , if at least one of these operators is invertible, in this case we can use one of the following Frobenius formulae for the inverse of a block matrix:

$$\begin{pmatrix} A & B \\ C & D \end{pmatrix}^{-1}$$

$$= \begin{pmatrix} A^{-1} + A^{-1}B(D - CA^{-1}B)^{-1}CA^{-1} & -A^{-1}B(D - CA^{-1}B)^{-1} \\ -(D - CA^{-1}B)^{-1}CA^{-1} & (D - CA^{-1}B)^{-1} \end{pmatrix}$$

or

$$\begin{pmatrix} A & B \\ C & D \end{pmatrix}^{-1} \\ = \begin{pmatrix} (A - BD^{-1}C)^{-1} & -(A - BD^{-1}C)^{-1}BD^{-1} \\ -D^{-1}C(A - BD^{-1}C)^{-1} & D^{-1} + D^{-1}C(A - BD^{-1}C)^{-1}BD^{-1} \end{pmatrix},$$

respectively.

We will not now concentrate ourselves upon details.

#### $\mathbf{2}$ Resolvents Formulae for the Models of Interaction

Return to

$$z \begin{pmatrix} q \\ u \end{pmatrix} - \begin{pmatrix} -\Omega^2 & \Omega^2 < l|_2 \\ 4\gamma_c \delta_{\alpha, x_0} < 1|_1 & B - 4\gamma_c \delta_{\alpha, x_0} < l|_2 \end{pmatrix} \begin{pmatrix} q \\ u \end{pmatrix} = \begin{pmatrix} w_1 \\ w_2 \end{pmatrix}$$

which we write also as

$$\begin{pmatrix} z + \Omega^2 & -\Omega^2 < l|_2 \\ -4\gamma_c \delta_{\alpha, x_0} < 1|_1 & z - B + 4\gamma_c \delta_{\alpha, x_0} < l|_2 \end{pmatrix} \begin{pmatrix} q \\ u \end{pmatrix} = \begin{pmatrix} w_1 \\ w_2 \end{pmatrix}$$

and confer this latter expression with formulae of rank two perturbations. We let

$$\begin{pmatrix} A_{11} - f_{11} < l_1|_1 & -f_{12} < l_2|_2 \\ -f_{21} < l_1|_1 & A_{22} - f_{22} < l_2|_2 \end{pmatrix} := \begin{pmatrix} z + \Omega^2 & -\Omega^2 < l|_2 \\ -4\gamma_c\delta_{\alpha,x_0} < 1|_1 & z - B + 4\gamma_c\delta_{\alpha,x_0} < l|_2 \end{pmatrix}$$
 and next take

and next take

$$\begin{array}{ll} A_{11} := z + \Omega^2 \,, & < l_1|_1 := < 1|_1 \,, & A_{22} := z - B \,, & < l_2|_2 := < l|_2 \\ f_{11} := 0 \, & f_{12} := \Omega^2 \\ f_{21} := 4 \gamma_c \delta_{\alpha, x_0} \, & f_{22} := -4 \gamma_c \delta_{\alpha, x_0} \end{array}$$

After substituting these values of  $A_{\cdots}$  and  $f_{\cdots}$  in

$$\begin{pmatrix} 1 - \langle l_1 | A_{11}^{-1} f_{11} \rangle & - \langle l_1 | A_{11}^{-1} f_{12} \rangle \\ - \langle l_2 | A_{22}^{-1} f_{21} \rangle & 1 - \langle l_2 | A_{22}^{-1} f_{22} \rangle \end{pmatrix} \begin{pmatrix} c_1 \\ c_2 \end{pmatrix} = \begin{pmatrix} \langle l_1 | A_{11}^{-1} w_1 \rangle \\ \langle l_2 | A_{22}^{-1} w_2 \rangle \end{pmatrix}$$

we obtain

$$\begin{pmatrix} 1 - \langle 1|_1(z + \Omega^2)^{-1}0 \rangle & - \langle 1|_1(z + \Omega^2)^{-1}(\Omega^2) \rangle \\ - \langle l|_2(z - B)^{-1}(4\gamma_c\delta_{\alpha,x_0}) \rangle & 1 - \langle l|_2(z - B)^{-1}(-4\gamma_c\delta_{\alpha,x_0}) \rangle \end{pmatrix} \begin{pmatrix} c_1 \\ c_2 \end{pmatrix} \\ = \begin{pmatrix} \langle 1|_1(z + \Omega^2)^{-1}w_1 \rangle \\ \langle l|_2(z - B)^{-1}w_2 \rangle \end{pmatrix}$$

Then

$$\begin{pmatrix} 1 & -\frac{\Omega^2}{z+\Omega^2} \\ -4\gamma_c < l|(z-B)^{-1}\delta_{\alpha,x_0} > & 1+4\gamma_c < l|(z-B)^{-1}\delta_{\alpha,x_0} > \end{pmatrix} \begin{pmatrix} c_1 \\ c_2 \end{pmatrix} \\ = \begin{pmatrix} \frac{w_1}{z+\Omega^2} \\ < l|(z-B)^{-1}w_2 > \end{pmatrix}$$

Now let us choose

$$B := c^2 \Delta$$

In this case one can show that for any complex z such that  $z\not\in(-\infty,0]$ 

$$(z - B)^{-1} \equiv (z - c^2 \Delta))^{-1}$$

exists and is an integral operator which can be described as following (see e.g. Appendix A) :

$$((z-c^{2}\Delta))^{-1}F)(x) = \int_{-\infty}^{+\infty} \frac{i}{2kc^{2}} e^{ik|x-x'|} F(x')dx \text{ where } k^{2} = -z/c^{2}, \quad Im \, k > 0$$

If we choose in addition

$$\delta_{\alpha,x_0}(x) := \delta(x - x_0), \quad < l|F> := F(x_0)$$

then we obtain:

$$< l|(z-B)^{-1}\delta_{\alpha,x_0}> = \frac{i}{2kc^2}$$

and equation

$$\begin{pmatrix} 1 & -\frac{\Omega^2}{z+\Omega^2} \\ -4\gamma_c < l|(z-B)^{-1}\delta_{\alpha,x_0} > & 1+4\gamma_c < l|(z-B)^{-1}\delta_{\alpha,x_0} > \end{pmatrix} \begin{pmatrix} c_1 \\ c_2 \end{pmatrix} \\ = \begin{pmatrix} \frac{w_1}{z+\Omega^2} \\ < l|(z-B)^{-1}w_2 > \end{pmatrix}$$

becomes as following:

$$\begin{pmatrix} 1 & -\frac{\Omega^2}{z+\Omega^2} \\ -4\gamma_c \frac{i}{2kc^2} & 1+4\gamma_c \frac{i}{2kc^2} \end{pmatrix} \begin{pmatrix} c_1 \\ c_2 \end{pmatrix} = \begin{pmatrix} \frac{w_1}{z+\Omega^2} \\ \int_{-\infty}^{+\infty} \frac{i}{2kc^2} e^{ik|x_0-x'|} w_2(x')dx \end{pmatrix}$$

$$\begin{pmatrix} c_1 \\ c_2 \end{pmatrix} = \frac{1}{determinant} \begin{pmatrix} 1+4\gamma_c \frac{i}{2kc^2} & \frac{\Omega^2}{z+\Omega^2} \\ 4\gamma_c \frac{i}{2kc^2} & 1 \end{pmatrix} \begin{pmatrix} \frac{w_1}{z+\Omega^2} \\ \int_{-\infty}^{+\infty} \frac{i}{2kc^2} e^{ik|x_0-x'|} w_2(x')dx \end{pmatrix}$$

$$=\frac{1}{determinant}\left(\begin{array}{c} (1+4\gamma_c\frac{i}{2kc^2})\frac{w_1}{z+\Omega^2} + \frac{\Omega^2}{z+\Omega^2} \int_{-\infty}^{+\infty} \frac{i}{2kc^2} e^{ik|x_0-x'|}w_2(x')dx \\ 4\gamma_c\frac{i}{2kc^2}\frac{w_1}{z+\Omega^2} + \int_{-\infty}^{+\infty} \frac{i}{2kc^2} e^{ik|x_0-x'|}w_2(x')dx \end{array}\right)$$

where

$$k^2 = -z/c^2, \quad Im \, k > 0$$

and

$$determinant := 1 + 4\gamma_c \frac{i}{2kc^2} - \frac{\Omega^2}{z + \Omega^2} 4\gamma_c \frac{i}{2kc^2}$$
$$= 1 + 2\gamma_c \frac{i}{kc^2} - \frac{\Omega^2}{z + \Omega^2} 2\gamma_c \frac{i}{kc^2}$$

i.e.,

$$determinant = \frac{kc^{2}(z+\Omega^{2}) + 2\gamma_{c}i(z+\Omega^{2}) - 2\gamma_{c}i\Omega^{2}}{kc^{2}(z+\Omega^{2})} = \frac{kc^{2}(z+\Omega^{2}) + 2\gamma_{c}iz}{kc^{2}(z+\Omega^{2})}$$

An Interaction of An Oscillator with ... . II: Resolvents formulae

$$= \frac{kc^{2}(-k^{2}c^{2} + \Omega^{2}) - 2\gamma_{c}ik^{2}c^{2}}{kc^{2}(-k^{2}c^{2} + \Omega^{2})} = \frac{-k^{2}c^{2} + \Omega^{2} - 2i\gamma_{c}k}{-k^{2}c^{2} + \Omega^{2}}$$
$$= -\frac{k^{2} + \frac{2i\gamma_{c}}{c^{2}}k - \frac{\Omega^{2}}{c^{2}}}{-k^{2} + \frac{\Omega^{2}}{c^{2}}} = \frac{(k + \frac{i\gamma_{c}}{c^{2}})^{2} - \frac{\Omega^{2}}{c^{2}} + (\frac{\gamma_{c}}{c^{2}})^{2}}{k^{2} - \frac{\Omega^{2}}{c^{2}}}$$

We observe that determinant vanishes at two values of k:

$$k_{1,2} = -\frac{i\gamma_c}{c^2} \pm \sqrt{\frac{\Omega^2}{c^2} - (\frac{\gamma_c}{c^2})^2} \equiv -\frac{i\gamma_c}{c^2} \mp i\sqrt{-\frac{\Omega^2}{c^2} + (\frac{\gamma_c}{c^2})^2} \,.$$

But these values are forbidden to k because of the condition Im k > 0 and of course because of the condition  $\gamma_c > 0$  !!!

We temporarily defer discussing 'what does it mean'. In the next section, we will specially return to this question, for the time being, we turn to calculating the resolvent associated with the Friedrichs model, i.e., with the equation array

$$z \begin{pmatrix} q_0 \\ \phi \end{pmatrix} - \begin{pmatrix} -\Omega^2 & \gamma_1 < l|_2 \\ 4\gamma_{2,c}\delta_{\alpha,x_0} < 1|_1 & B \end{pmatrix} \begin{pmatrix} q_0 \\ \phi \end{pmatrix} = \begin{pmatrix} w_1 \\ w_2 \end{pmatrix}$$

We take

$$\begin{array}{ll} A_{11} := z + \Omega^2, & < l_1|_1 := < 1|_1, & A_{22} := z - B, & < l_2|_2 := < l|_2 \\ f_{11} := 0 & & f_{12} := \gamma_1 \\ f_{21} := 4\gamma_{2,c}\delta_{\alpha,x_0} & & f_{22} := 0 \end{array}$$

and after substituting these values of  $A_{\cdots}$  and  $f_{\cdots}$  in

$$\begin{pmatrix} 1 - \langle l_1 | A_{11}^{-1} f_{11} \rangle & - \langle l_1 | A_{11}^{-1} f_{12} \rangle \\ - \langle l_2 | A_{22}^{-1} f_{21} \rangle & 1 - \langle l_2 | A_{22}^{-1} f_{22} \rangle \end{pmatrix} \begin{pmatrix} c_1 \\ c_2 \end{pmatrix} = \begin{pmatrix} \langle l_1 | A_{11}^{-1} w_1 \rangle \\ \langle l_2 | A_{22}^{-1} w_2 \rangle \end{pmatrix}$$

we obtain

$$\begin{pmatrix} 1 - \langle 1|_1(z + \Omega^2)^{-1}_0 \rangle & - \langle 1|_1(z + \Omega^2)^{-1}_1(\gamma_1) \rangle \\ - \langle l|_2(z - B)^{-1}_1(4\gamma_{2,c}\delta_{\alpha,x_0}) \rangle & 1 - \langle l|_2(z - B)^{-1}_0 \rangle \end{pmatrix} \begin{pmatrix} c_1 \\ c_2 \end{pmatrix} \\ = \begin{pmatrix} \langle 1|_1(z + \Omega^2)^{-1}_1w_1 \rangle \\ \langle l|_2(z - B)^{-1}_1w_2 \rangle \end{pmatrix}$$

i.e.,

$$\begin{pmatrix} 1 & -\frac{\gamma_1}{z+\Omega^2} \\ -4\gamma_{2,c} < l|(z-B)^{-1}\delta_{\alpha,x_0} > & 1 \end{pmatrix} \begin{pmatrix} c_1 \\ c_2 \end{pmatrix}$$
$$= \begin{pmatrix} \frac{w_1}{z+\Omega^2} \\ < l|(z-B)^{-1}w_2 > \end{pmatrix}$$

Now let us take again

$$B := c^2 \Delta$$
  
$$\delta_{\alpha, x_0}(x) := \delta(x - x_0), \quad \langle l | F \rangle := F(x_0)$$

Since in this case for any complex z such that  $z\not\in (-\infty,0]$  the operator

$$(z-B)^{-1} \equiv (z-c^2\Delta))^{-1}$$

exists and since

$$< l|(z-B)^{-1}\delta_{\alpha,x_0}> = \frac{i}{2kc^2}$$
 where  $k^2 = -z/c^2$ ,  $Im \, k > 0$ 

the recent equation to  $c_1,c_2$  becomes as following:

$$\begin{pmatrix} 1 & -\frac{\gamma_1}{z+\Omega^2} \\ -4\gamma_{2,c}\frac{i}{2kc^2} & 1 \end{pmatrix} \begin{pmatrix} c_1 \\ c_2 \end{pmatrix} = \begin{pmatrix} \frac{w_1}{z+\Omega^2} \\ < l|_2(z-B)^{-1}w_2 > \end{pmatrix}$$

$$\begin{pmatrix} c_1 \\ c_2 \end{pmatrix} = \frac{1}{determinant_F} \begin{pmatrix} 1 & \frac{\gamma_1}{z + \Omega^2} \\ 4\gamma_{2,c}\frac{i}{2kc^2} & 1 \end{pmatrix} \begin{pmatrix} \frac{w_1}{z + \Omega^2} \\ \int_{-\infty}^{+\infty} \frac{i}{2kc^2} e^{ik|x_0 - x'|} w_2(x') dx \end{pmatrix}$$

$$= \frac{1}{determinant_F} \begin{pmatrix} \frac{w_1}{z+\Omega^2} + \frac{\gamma_1}{z+\Omega^2} \int_{-\infty}^{+\infty} \frac{i}{2kc^2} e^{ik|x_0-x'|} w_2(x') dx \\ 4\gamma_{2,c} \frac{i}{2kc^2} \frac{w_1}{z+\Omega^2} + \int_{-\infty}^{+\infty} \frac{i}{2kc^2} e^{ik|x_0-x'|} w_2(x') dx \end{pmatrix}$$

where

$$k^2 = -z/c^2, \quad Im \, k > 0$$

and

$$determinant_{F} := 1 + \frac{\gamma_{1}}{z + \Omega^{2}} 4\gamma_{2,c} \frac{i}{2kc^{2}}$$
$$= 1 + \frac{\gamma_{1}}{-k^{2}c^{2} + \Omega^{2}} \gamma_{2,c} \frac{i}{kc^{2}}$$
$$\frac{(-k^{2}c^{2} + \Omega^{2})kc^{2} + i\gamma_{1}\gamma_{2,c}}{(-k^{2}c^{2} + \Omega^{2})kc^{2}}$$

i.e.,

$$determinant_F = \frac{(-k^2 + \Omega^2/c^2)k + i\gamma_1\gamma_{2,c}/c^4}{(-k^2 + \Omega^2/c^2)k}$$

We conclude this section with two remarks.

**Remark 1.** We have considered the operator

$$\left( \begin{array}{cc} z + \Omega^2 & -\Omega^2 < l|_2 \\ -4\gamma_c \delta_{\alpha,x_0} < 1|_1 & z - B + 4\gamma_c \delta_{\alpha,x_0} < l|_2 \end{array} \right)$$

as a perturbation of the operator

$$\left(\begin{array}{cc} z+\Omega^2 & 0\\ 0 & z-B \end{array}\right)\,.$$

The perturbation, i.e. the operator

$$\left( \begin{array}{cc} 0 & -\Omega^2 < l|_2 \\ -4\gamma_c \delta_{\alpha,x_0} < 1|_1 & 4\gamma_c \delta_{\alpha,x_0} < l|_2 \end{array} \right)$$

is a rank-two operator, we have already mentioned it. If we initially take

$$\left(\begin{array}{cc}z&0\\0&z-B\end{array}\right)$$

as an unperturbated operator, then the perturbation becomes

$$\begin{pmatrix} \Omega^2 & -\Omega^2 < l|_2 \\ -4\gamma_c \delta_{\alpha,x_0} < 1|_1 & 4\gamma_c \delta_{\alpha,x_0} < l|_2 \end{pmatrix},$$

and this operator is a rank-one one.

Actually,

$$\begin{pmatrix} \Omega^2 & -\Omega^2 < l|_2 \\ -4\gamma_c \delta_{\alpha,x_0} < 1|_1 & 4\gamma_c \delta_{\alpha,x_0} < l|_2 \end{pmatrix} = \begin{pmatrix} \Omega^2 \\ -4\gamma_c \delta_{\alpha,x_0} \end{pmatrix} < 1 \oplus -l|$$
$$= \begin{pmatrix} \Omega^2 \\ -4\gamma_c \delta_{\alpha,x_0} \end{pmatrix} \begin{pmatrix} <1|_1 \oplus - < l|_2 \end{pmatrix}.$$

We have used here the fact that, as an operator  $\mathbf{C} \to \mathbf{C}$ ,  $-\Omega^2 < 1|_1$  coincides with the operator of multiplication by  $-\Omega^2$ :

$$-\Omega^2 < 1|_1 q > = -\Omega^2 q, \, q \in \mathbf{C}$$

-in other words,

$$-\Omega^2 < 1|_1 = -\Omega^2 \cdot = -\Omega^2 I_{\mathbf{C}} = \Omega^2 \,.$$

 $Of\ course,\ we\ have\ identified\ the\ operator\ multiplication\ by\ a\ number\ with\ this\ number\ itself:$ 

the operator multiplication by a number  $\equiv$  this number itself.

#### S.A. Chorošavin

### Remark 2. As for an explicit and detailed expression of

$$\left( \begin{array}{cc} z + \Omega^2 & -\Omega^2 < l|_2 \\ -4\gamma_c \delta_{\alpha, x_0} < 1|_1 & z - B + 4\gamma_c \delta_{\alpha, x_0} < l|_2 \end{array} \right)^{-1},$$

 $we\ have$ 

$$\begin{pmatrix} q \\ u \end{pmatrix} = \begin{pmatrix} A_{11}^{-1}w_1 \\ A_{22}^{-1}w_2 \end{pmatrix} + \begin{pmatrix} A_{11}^{-1}f_{11} & A_{11}^{-1}f_{12} \\ A_{22}^{-1}f_{21} & A_{22}^{-1}f_{22} \end{pmatrix} \begin{pmatrix} c_1 \\ c_2 \end{pmatrix}$$

$$= \begin{pmatrix} (z+\Omega^2)^{-1}w_1 \\ (z-B)^{-1}w_2 \end{pmatrix} + \begin{pmatrix} (z+\Omega^2)^{-1}\cdot 0 & (z+\Omega^2)^{-1}\Omega^2 \\ (z-B)^{-1}4\gamma_c\delta_{\alpha,x_0} & -(z-B)^{-1}4\gamma_c\delta_{\alpha,x_0} \end{pmatrix} \begin{pmatrix} c_1 \\ c_2 \end{pmatrix}$$

$$= \begin{pmatrix} (z+\Omega^2)^{-1}w_1 \\ (z-B)^{-1}w_2 \end{pmatrix} + \begin{pmatrix} 0 & (z+\Omega^2)^{-1}\Omega^2 \\ (z-B)^{-1}4\gamma_c\delta_{\alpha,x_0} & -(z-B)^{-1}4\gamma_c\delta_{\alpha,x_0} \end{pmatrix} \begin{pmatrix} c_1 \\ c_2 \end{pmatrix} ,$$

$$\begin{pmatrix} c_1 \\ c_2 \end{pmatrix} \cdot det$$

$$= \begin{pmatrix} \left(1 + 4\gamma_c < l|(z-B)^{-1}\delta_{\alpha,x_0} > \right) & \frac{\Omega^2}{z+\Omega^2} \\ 4\gamma_c < l|(z-B)^{-1}\delta_{\alpha,x_0} > & 1 \end{pmatrix} \begin{pmatrix} \frac{w_1}{z+\Omega^2} \\ < l|(z-B)^{-1}w_2 > \end{pmatrix}$$

$$= \begin{pmatrix} \left(1 + 4\gamma_c < l|(z-B)^{-1}\delta_{\alpha,x_0} > \right) \frac{w_1}{z+\Omega^2} + \frac{\Omega^2}{z+\Omega^2} < l|(z-B)^{-1}w_2 > \\ 4\gamma_c < l|(z-B)^{-1}\delta_{\alpha,x_0} > \frac{w_1}{z+\Omega^2} + < l|(z-B)^{-1}w_2 > \end{pmatrix}$$

where

$$det = 1 + 4\gamma_c < l|(z - B)^{-1}\delta_{\alpha, x_0} > -\frac{\Omega^2}{z + \Omega^2} 4\gamma_c < l|(z - B)^{-1}\delta_{\alpha, x_0} > .$$

Notice, by the way,

$$det \cdot (z + \Omega^2) = (z + \Omega^2) + 4\gamma_c z < l | (z - B)^{-1} \delta_{\alpha, x_0} >$$
  
=  $(1 + 4\gamma_c < l | (z - B)^{-1} \delta_{\alpha, x_0} >) z + \Omega^2,$ 

$$\begin{pmatrix} c_1 \\ c_2 \end{pmatrix} det \cdot (z + \Omega^2)$$

$$= \begin{pmatrix} \left(1 + 4\gamma_c < l|(z - B)^{-1}\delta_{\alpha, x_0} > \right)w_1 + \Omega^2 < l|(z - B)^{-1}w_2 > \\ \\ 4\gamma_c < l|(z - B)^{-1}\delta_{\alpha, x_0} > w_1 + (z + \Omega^2) < l|(z - B)^{-1}w_2 > \end{pmatrix}.$$

#### 3 Resonances, Resolvents, and the Second Sheet

In the previous section we have observed that the *determinant* vanishes at two values of k:

$$k_{1,2} = -\frac{i\gamma_c}{c^2} \pm \sqrt{\frac{\Omega^2}{c^2} - (\frac{\gamma_c}{c^2})^2} .$$
$$\equiv -\frac{i\gamma_c}{c^2} \mp i\sqrt{-\frac{\Omega^2}{c^2} + (\frac{\gamma_c}{c^2})^2} .$$

The circumstance having excited an interest is that these values are forbidden (!!!) to k because of the condition  $Im k > 0^{5}$ . In this section, we return specially to this fact, and we are now discussing 'what does it mean'.

Firstly, we fix on

$$det_{-1,+}(k) := ((z + \Omega^2)determinant)^{-1} \equiv ((-k^2c^2 + \Omega^2)determinant)^{-1}$$

as a function of k, —we need rather this latter quantity than the determinant . We have:

$$det_{-1}(k,+) = -\frac{1/c^2}{k^2 + \frac{2i\gamma_c}{c^2}k - \frac{\Omega^2}{c^2}} = \frac{1/c^2}{(k + \frac{i\gamma_c}{c^2})^2 - \frac{\Omega^2}{c^2} + (\frac{\gamma_c}{c^2})^2},$$
  
(where  $k^2 = -z/c^2$ ,  $Im k > 0$ ).

The first factor which calls attention to itself is that the written form of  $det_{-1}(k)$  in itself need not the relation  $k^2 = -z/c^2$ , Im k > 0 being fulfilled, and provokes to introduce the formal extension of  $det_{-1}(k, +)$ ,  $Det_{-1}(k)$  say, defined by

$$Det_{-1}(k) := -\frac{1/c^2}{k^2 + \frac{2i\gamma_c}{c^2}k - \frac{\Omega^2}{c^2}}, \qquad k^2 + \frac{2i\gamma_c}{c^2}k - \frac{\Omega^2}{c^2} \neq 0.$$

This extension  $Det_{-1}(k)$  is an analytic function of k, and at real k

$$|Det_{-1}(k)|^2 = |det_{-1,+}(k+i0))|^2 = \frac{1/c^4}{(k^2 - \frac{\Omega^2}{c^2})^2 + (\frac{2\gamma_c}{c^2}k)^2}, \quad (k \in \mathbf{R}).$$

For a moment, let us renormalise  $\Omega$  and  $\gamma_c$  so that

$$\frac{\Omega}{c} \longrightarrow \Omega \,, \, \frac{\gamma_c}{c^2} \longrightarrow \gamma \,,$$

or choose the measure units so that c = 1. Then we have:

$$det_{-1,+}(k) = -\frac{1}{k^2 + 2i\gamma k - \Omega^2} = \frac{1}{(k+i\gamma)^2 - \Omega^2 + \gamma^2},$$
  
(where  $k^2 = -z$ ,  $Im k > 0$ ),

$$|det_{-1}(k+i0))|^2 = \frac{1}{(k^2 - \Omega^2)^2 + (2\gamma k)^2}, \quad (k \in \mathbf{R}).$$

 $<sup>^5 {\</sup>rm and}$  of course because of the condition  $\gamma_c > 0$ 

We had already seen a similar expression. We had seen it in the previous paper.

In the previous paper  $^6,$  in subsection 1.4, where just c=1, we had seen: given an incident wave of the kind

$$A_{mp}\sin(k(x+t))$$
 where  $k \in \mathbf{R}$ ,  $A_{mp} \in \mathbf{R}$ ,

then

$$q(t) = \frac{\Omega^2 A_{mp}}{(-k^2 + \Omega^2)^2 + (2\gamma k)^2} \left( (-k^2 + \Omega^2) \sin(kt) - 2\gamma k \cos(kt) \right) + const_1 + const_1 + \frac{2\gamma}{\Omega^2} \frac{\partial q(t)}{\partial t} \Big|_{t=0} + e^{-\gamma t} \left( const_1 \cos(\Omega_\gamma t) + const_2 \sin(\Omega_\gamma t) \right);$$

we can write this relation as follows:

$$q(t) = \frac{\Omega^2 A_{mp}}{\sqrt{(-k^2 + \Omega^2)^2 + (2\gamma k)^2}} \sin(kt + \phi_k) + const_1 + \frac{2\gamma}{\Omega^2} \frac{\partial q(t)}{\partial t}\Big|_{t=0} + e^{-\gamma t} \Big(const_1 \cos(\Omega_{\gamma} t) + const_2 \sin(\Omega_{\gamma} t)\Big)$$

where  $\phi_k$  is such that

$$\cos \phi_k = \frac{-k^2 + \Omega^2}{\sqrt{(-k^2 + \Omega^2)^2 + (2\gamma k)^2}}, \ \sin \phi_k = -\frac{2\gamma k}{\sqrt{(-k^2 + \Omega^2)^2 + (2\gamma k)^2}}$$

We had also seen that the field u(t, x) is given by

$$u(t,x) = \left\{ \begin{array}{cc} Q(t-|x|) - A_{mp}\sin(k(t-|x|)) &, & \text{if } 0 \le t - |x| \\ 0 &, & \text{if } t - |x| < 0 \le t \end{array} \right\} + A_{mp}\sin(k(x+t)),$$

where

$$Q(t) - A_{mp}\sin(kt) = \frac{-2\gamma A_{mp}k}{(-k^2 + \Omega^2)^2 + (2\gamma k)^2} \left(2\gamma k\sin(kt) + (-k^2 + \Omega^2)\cos(kt)\right) + \frac{2\gamma}{\Omega^2} \frac{\partial q(t)}{\partial t}\Big|_{t=0} + e^{-\gamma t} \left(const_3\cos(\Omega_{\gamma}t) + const_4\sin(\Omega_{\gamma}t)\right)$$

and, thus, at points of the left real half-line the field is given by the relationships:

$$u(t,x) = \left\{ \begin{array}{cc} Q(t+x) &, & \text{if } x < 0, 0 \le t+x \\ 0 &, & \text{if } x < 0, t+x < 0 \le t \end{array} \right\}$$

$$\begin{aligned} Q(t) &- \frac{2\gamma}{\Omega^2} \frac{\partial q(t)}{\partial t} \Big|_{t=0} - e^{-\gamma t} \Big( const_3 \cos(\Omega_\gamma t) + const_4 \sin(\Omega_\gamma t) \Big) \\ &= \Big( \frac{-(2\gamma k)^2 A_{mp}}{(-k^2 + \Omega^2)^2 + (2\gamma k)^2} + A \Big) \sin(kt) + \frac{-2\gamma A_{mp}k(-k^2 + \Omega^2)}{(-k^2 + \Omega^2)^2 + (2\gamma k)^2} \cos(kt) \\ &= \frac{(-k^2 + \Omega^2)^2 A_{mp}}{(-k^2 + \Omega^2)^2 + (2\gamma k)^2} \sin(kt) + \frac{-2\gamma A_{mp}k(-k^2 + \Omega^2)}{(-k^2 + \Omega^2)^2 + (2\gamma k)^2} \cos(kt) \\ &= \frac{(-k^2 + \Omega^2) A_{mp}}{(-k^2 + \Omega^2)^2 + (2\gamma k)^2} \left( (-k^2 + \Omega^2) \sin(k(t)) - 2\gamma k \cos(kt) \right) \\ &= \frac{(-k^2 + \Omega^2) A_{mp}}{\sqrt{(-k^2 + \Omega^2)^2 + (2\gamma k)^2}} \sin(kt + \phi_k) \,. \end{aligned}$$

 $^{6}$ we mean [Ch03-1]

In addition,

$$q(t) - Q(t) - e^{-\gamma t} \Big( (const_1 - const_3) \cos(\Omega_{\gamma} t) + (const_2 - const_4) \sin(\Omega_{\gamma} t) \Big) \\ = \frac{k^2 A_{mp}}{(-k^2 + \Omega^2)^2 + (2\gamma k)^2} \left( (-k^2 + \Omega^2) \sin(k(t)) - 2\gamma k \cos(kt) \right) \\ = \frac{k^2 A_{mp}}{\sqrt{(-k^2 + \Omega^2)^2 + (2\gamma k)^2}} \sin(kt + \phi_k) \,.$$

Now, let us concentrate ourselves upon the expressions of the amplitudes of the harmonic parts of q(t), Q(t) and q(t)-Q(t), —expressions as functions of k,— i.e., let us concentrate upon oscillator amplitude, transmitted wave one, and a 'deformation' amplitude. These amplitudes are exactly

$$\begin{aligned} amplitude_q(k) &= \frac{\Omega^2 A_{mp}}{\sqrt{(-k^2 + \Omega^2)^2 + (2\gamma k)^2}}, \\ amplitude_Q(k) &= \frac{(-k^2 + \Omega^2)A_{mp}}{\sqrt{(-k^2 + \Omega^2)^2 + (2\gamma k)^2}}, \\ amplitude_{qQ}(k) &= \frac{k^2 A_{mp}}{\sqrt{(-k^2 + \Omega^2)^2 + (2\gamma k)^2}}; \end{aligned}$$

we will also need  $amplitude_q(k)^2$  ,  $amplitude_Q(k)^2$  , and  $amplitude_{qQ}(k)^2$  :

$$\begin{split} amplitude_q(k)^2 &= \frac{\Omega^4 A_{mp}^2}{(-k^2 + \Omega^2)^2 + (2\gamma k)^2} \,, \\ amplitude_Q(k)^2 &= \frac{(-k^2 + \Omega^2)^2 A_{mp}^2}{(-k^2 + \Omega^2)^2 + (2\gamma k)^2} \,, \\ amplitude_{qQ}(k)^2 &= \frac{(k^2)^2 A_{mp}^2}{(-k^2 + \Omega^2)^2 + (2\gamma k)^2} \,. \end{split}$$

After standard transformations—

$$(-k^{2} + \Omega^{2})^{2} + (2\gamma k)^{2} = k^{4} - 2(\Omega^{2} - 2\gamma^{2})k^{2} + \Omega^{4} = (k^{2} - (\Omega^{2} - 2\gamma^{2}))^{2} + 4\gamma^{2}(\Omega^{2} - \gamma^{2})$$

—we also write  $amplitude_q(k)^2$  and  $amplitude_{qQ}(k)^2$  as follows:

$$amplitude_q(k)^2 = \frac{\Omega^4 A_{mp}^2}{(-k^2 + \Omega^2)^2 + (2\gamma k)^2} = \frac{\Omega^4 A_{mp}^2}{(k^2 - (\Omega^2 - 2\gamma^2))^2 + 4\gamma^2(\Omega^2 - \gamma^2)},$$
$$amplitude_{qQ}(k)^2 = \frac{\Omega^4 A_{mp}^2}{1 - 2(\Omega^2 - 2\gamma^2)k^{-2} + \Omega^4 k^{-4}}.$$

 $\overline{7}$ 

We emphasise: in transforming no special assumption has been made. But now recall, we have been discussing incident wave where k is a wave-number, so, we presume that  $k \in \mathbf{R}$ .

Thus, since  $k \in \mathbf{R}$ , we had seen: there is a value of a system parameter, —this parameter is k, the wave-number of the incident wave, —such that the oscillator

<sup>7</sup>In addition, we have

$$amplitude_Q(k)^2 = \frac{A_{mp}^2}{1 - (2\gamma)^2/(-k^2 + \Omega^2) + (2\gamma)^2 \Omega^2/(-k^2 + \Omega^2)^2} \,.$$

amplitude becomes **maximal**. This phenomenon is called a **resonance**, in the proper, usual sense of the word. Temporarily, we call this phenomenon **resonance** of the first kind.

We had also seen: there is a value of a system parameter, —this parameter is k again, the wave-number of the incident wave, —such that the incident harmonic wave is **completely reflected**. This phenomenon is a kind of **resonance**, we will temporarily call it **resonance of the second kind**.

We have finally seen: there is a value of k such that amplitude of q(t) - Q(t) becomes maximal. This phenomenon is also a kind of **resonance**, we will temporarily call it **resonance of the third kind**.

Currently, the first kind resonant values of k are exactly

$$k_{1st.res} = \pm \sqrt{\Omega^2 - 2\gamma^2}$$

the second kind ones are

$$k_{2nd.res} = \pm \Omega$$

and the third kind resonant values of k are

$$k_{3rd.res} = \pm \sqrt{\frac{\Omega^4}{\Omega^2 - 2\gamma^2}}.$$

Another standard of transforming the denominator in the expressions of  $amplitude_q(k)^2$  and  $amplitude_Q(k)^2$  is this:

$$(-k^{2} + \Omega^{2})^{2} + (2\gamma k)^{2} = (k^{2} + 2i\gamma k - \Omega^{2})(k^{2} - 2i\gamma k - \Omega^{2}).$$

Thus, we observe that

$$amplitude_{q}(k)^{2} = \frac{\Omega^{4}A_{mp}^{2}}{(-k^{2} + \Omega^{2})^{2} + (2\gamma k)^{2}} = \frac{\Omega^{4}A_{mp}^{2}}{(k^{2} + 2i\gamma k - \Omega^{2})(k^{2} - 2i\gamma k - \Omega^{2})}$$

$$amplitude_{Q}(k)^{2} = \frac{(-k^{2} + \Omega^{2})^{2}A_{mp}^{2}}{(-k^{2} + \Omega^{2})^{2} + (2\gamma k)^{2}} = \frac{(-k^{2} + \Omega^{2})^{2}A_{mp}^{2}}{(k^{2} + 2i\gamma k - \Omega^{2})(k^{2} - 2i\gamma k - \Omega^{2})}$$

$$amplitude_{qQ}(k)^{2} = \frac{(k^{2})^{2}A_{mp}^{2}}{(-k^{2} + \Omega^{2})^{2} + (2\gamma k)^{2}} = \frac{(k^{2})^{2}A_{mp}^{2}}{(k^{2} + 2i\gamma k - \Omega^{2})(k^{2} - 2i\gamma k - \Omega^{2})}$$

and recall that

$$det_{-1,+}(k) = -\frac{1}{k^2 + 2i\gamma k - \Omega^2} = \frac{1}{(k+i\gamma)^2 - \Omega^2 + \gamma^2},$$
  
( where  $k^2 = -z$ ,  $Im k > 0$ ),

$$|det_{-1}(k+i0))|^2 = \frac{1}{(k^2 - \Omega^2)^2 + (2\gamma k)^2} = \frac{1}{(k^2 + 2i\gamma k - \Omega^2)(k^2 - 2i\gamma k - \Omega^2)} \quad (k \in \mathbf{R}) = \frac{1}{(k^2 - \Omega^2)^2 + (2\gamma k)^2} = \frac{1}{(k^2 - \Omega^2)^2 + (2\gamma k)^2} = \frac{1}{(k^2 - \Omega^2)(k^2 - 2i\gamma k - \Omega^2)} \quad (k \in \mathbf{R}) = \frac{1}{(k^2 - \Omega^2)^2 + (2\gamma k)^2} = \frac{1}{(k^2 - \Omega^2)(k^2 - 2i\gamma k - \Omega^2)}$$

First we focus on expressions of  $amplitude_q(k)^2$  and  $det_{-1}(k)$  . In the first moment we tend to write

$$amplitude_q(k)^2 = \Omega^4 A_{mp}^2 det_{-1,+}(k) det_{-1,+}(-k) + \frac{1}{2} det_{-1,+}(-k) + \frac{1}{2$$

We may not do it. There are at least two reasons, both of them are connected with the domains.

If Im k > 0 then Im(-k) < 0. Hence, if  $det_{-1,+}(k)$  is defined, then  $det_{-1,+}(-k)$ is not.

Secondly, if we deal with  $det_{-1,+}(k)$ , then Im k > 0; if we deal with  $amplitude_{q}(k)^{2}$  then  $k \in \mathbf{R}$ , i.e., Im k = 0; so, we must deal here with two 'different' k-s.

It is no surprise: k as a parameter of  $amplitude_q(k)^2$  means wave-number of incident wave, whereas k as a parameter of  $det_{-1}(k, +)$  is a term in describing of a resolvent of an operator.

In other words, k in amplitude<sub>q</sub>(k)<sup>2</sup> and k in  $det_{-1}(k, +)$  are of different natures. Nevertheless, the relation

$$amplitude_{q}(k)^{2} = \Omega^{4} A_{mp}^{2} |det_{-1,+}(k+i0))|^{2}$$

is mathematically correct, and we see: in order to find resonant values of the amplitude we have to find positions of extremal values of a function, which is absolute values function of boundery values function of an analytic function, analytic in the open upper half-plane (Im k > 0).

Let us discuss the recent idea in terms of analytic functions. We see, the written form of  $amplitude_q(k)^2$ ,  $amplitude_Q(k)^2$  and  $amplitude_{qQ}(k)^2$  in itself need not the relation  $k^2 = -z/c^2$ , Im k > 0 being fulfilled, and provokes to introduce the formal extensions of these quantities,  $Amplitude_q(k)^2$ ,  $Amplitude_Q(k)^2$  and  $Amplitude_{qQ}(k)^2$  say, defined by

$$\begin{split} Amplitude_q(k)^2 &:= \frac{\Omega^4 A_{mp}^2}{(-k^2 + \Omega^2)^2 + (2\gamma k)^2} = \frac{\Omega^4 A_{mp}^2}{(k^2 + 2i\gamma k - \Omega^2)(k^2 - 2i\gamma k - \Omega^2)} \\ Amplitude_Q(k)^2 &:= \frac{(-k^2 + \Omega^2)^2 A_{mp}^2}{(-k^2 + \Omega^2)^2 + (2\gamma k)^2} = \frac{(-k^2 + \Omega^2)^2 A_{mp}^2}{(k^2 + 2i\gamma k - \Omega^2)(k^2 - 2i\gamma k - \Omega^2)} \\ Amplitude_{qQ}(k)^2 &:= \frac{(k^2)^2 A_{mp}^2}{(-k^2 + \Omega^2)^2 + (2\gamma k)^2} = \frac{(k^2)^2 A_{mp}^2}{(k^2 + 2i\gamma k - \Omega^2)(k^2 - 2i\gamma k - \Omega^2)} \\ \text{wherever} \\ & (-k^2 + \Omega^2)^2 + (2\gamma k)^2 \neq 0 \,. \end{split}$$

These extensions are analytic functions of k, with poles at

$$\begin{aligned} k_{1,2}^- &= -i\gamma \pm \sqrt{\Omega^2 - \gamma^2} \equiv k_{1,2} \,, \\ k_{1,2}^+ &= +i\gamma \mp \sqrt{\Omega^2 - \gamma^2} \equiv -k_{1,2}^- \,, \end{aligned}$$

and  $Amplitude_q(k)^2$  is decreasing as  $|k| \to \infty$ . Thus we expect, the maximal value of  $amplitude_q(k)^2$  is somewhere near the poles of  $Amplitude_q(k)^2$ , at least at small  $\gamma$ .

...We have yet one analytic extension defined on C except for some poles. It is  $Det_{-1}(k)$ , introduced in the beginning of this section:

$$Det_{-1}(k) := -\frac{1}{k^2 + 2i\gamma k - \Omega^2}, \qquad k^2 + 2i\gamma k - \Omega^2 \neq 0.$$

The expressions of Amplitude-s and  $Det_{-1}(k)$  suggest writing

$$\begin{split} Amplitude_q(k)^2 &= \Omega^4 A_{mp}^2 \, Det_{-1}(k) \, Det_{-1}(-k) \,, \\ Amplitude_Q(k)^2 &= (-k^2 + \Omega^2)^2 A_{mp}^2 \, Det_{-1}(k) \, Det_{-1}(-k) \,, \end{split}$$

$$Amplitude_{qQ}(k)^2 = (k^2)^2 A_{mp}^2 Det_{-1}(k) Det_{-1}(-k) \,.$$

These relations are mathamatically correct. Therefore, let us discuss *amplitude*-s connecting them with  $Det_{-1}(k)$ .

We notice that  $Det_{-1}(k)$  is analytic extension of  $det_{-1,+}(k)$ , thus  $Det_{-1}(k)$  as well as  $det_{-1,+}(k)$ , is related rather to the resolvent, R(z), resolvent as a function <sup>8</sup> of parameter  $z, z \in \mathbb{C}^{9}$ . But the 'natural' parameter of resolvent is z, not k. Thus we would like to reformulate  $Det_{-1}(k)$  as a function of z.

Let us resume what we have. The situation with resolvent and resonances discussed in this section looks generally like this:

#### **Observation 1.**

1

(1) The resolvent is related to the function

$$det_{-1,+}(k) := -\frac{1}{k^2 + 2i\gamma k - \Omega^2} = \frac{1}{(k+i\gamma)^2 - \Omega^2 + \gamma^2},$$
  
( where  $Im \, k > 0$ );

(1a) the domain of  $det_{-1,+}(k)$  is

$$\{k|Im\,k>0\}$$

and is one-to-one to the set

$$\{z|z=-k^2; Im\,k>0\} \equiv \mathbf{C} \setminus (-\infty, 0];$$

(2) The amplitude-s are related to the function

$$det_{-1,0}(k) := -\frac{1}{k^2 + 2i\gamma k - \Omega^2} = \frac{1}{(k + i\gamma)^2 - \Omega^2 + \gamma^2},$$
  
( where  $Im k = 0$ );

(2a) the domain of  $det_{-1,0}(k)$  is

$$\{k | Im \, k = 0\} \equiv \{k | Im \, k = 0, \, Re \, k > 0\} \cup \{k | \, k = 0\} \cup \{k | Im \, k = 0, \, Re \, k < 0\},\$$

with the possible exception of the poles  $^{10}$  . Note,

$$\{z|z=-k^2; Im \, k=0 \,, \, Re \, k>0\} \equiv (-\infty,0) \equiv \{z|z=-k^2; Im \, k=0 \,, \, Re \, k<0\}$$

(3) The resonances are related to the poles of the function

$$det_{-1,-,0}(k) := -\frac{1}{k^2 + 2i\gamma k - \Omega^2} = \frac{1}{(k+i\gamma)^2 - \Omega^2 + \gamma^2},$$
  
( where  $Im \ k \le 0, \ k^2 + 2i\gamma k - \Omega^2 \ne 0$ );

(3a) the poles,  $k_{pol}$ , of  $det_{-1,-,0}(k)$ , and resonant values are connected by:

$$k_{pol} = -i\gamma \pm i\sqrt{-\Omega^2 + \gamma^2} \,,$$

<sup>&</sup>lt;sup>8</sup>operator-valued function

<sup>&</sup>lt;sup>9</sup>we mean usual notations in usual writing  $R(z) = (z - A)^{-1}$ 

 $<sup>^{10}{\</sup>rm they}$  exist, iff  $\gamma=0$ 

An Interaction of An Oscillator with ... . II: Resolvents formulae

$$k_{1st.res} = \pm \sqrt{\Omega^2 - 2\gamma^2},$$
  

$$k_{2nd.res} = \pm \Omega,$$
  

$$k_{3rd.res} = \pm \sqrt{\frac{\Omega^4}{\Omega^2 - 2\gamma^2}}.$$

(3b) The domain of  $det_{-1,-,0}(k)$  is

$$\{k|Im \, k \le 0\} = \{k|Im \, k < 0\} \cup \{k|Im \, k = 0\},\$$

with the exception of the poles;

the set

$$\{k|Im\,k<0\}$$

is one-to-one to the set

$$\{z|z=-k^2; Im\,k<0\} \equiv \mathbf{C} \setminus (-\infty,0];$$

(4) All these functions are consistent: there is an analytic function,—

$$Det_{-1}(k) = -\frac{1}{k^2 + 2i\gamma k - \Omega^2}, \qquad k^2 + 2i\gamma k - \Omega^2 \neq 0,$$

- such that:

(4a) every  $det_{-1,\#}$  is a restricton of  $Det_{-1}(k)$ ;

(4b) the union of the domains of  $det_{-1,\#}$  is the domain of  $Det_{-1}(k)$ .

Now then, we have no problem with reformulating  $det_{-1,+}(k)$  as a function of z: we do it by defining

$$\begin{aligned} d_{-1,+}(z) &:= \quad det_{-1}(k(z)) = -\frac{1}{k(z)^2 + 2i\gamma k(z) - \Omega^2} = \frac{1}{(k(z) + i\gamma)^2 - \Omega^2 + \gamma^2} \,, \\ &\quad \text{if } z \not\in (-\infty, 0] \,, \end{aligned}$$

where k(z) is defined as the unique solution to

$$k^2 = -z$$
,  $Im k > 0$ .

We need now reformulate  $Det_{-1}(k)$  which is an extension of  $det_{-1}(k)$ . In other words, we have to extend  $d_{-1,+}(z)$ , but 'into WHAT?'

Of course, we can extend  $d_{-1}(z)$  onto

$$z \in \mathbf{C}, \, z = -k^2, \, k^2 + 2i\gamma k - \Omega^2 \neq 0.$$

We can do it, by defining, e.g.,

$$\begin{array}{rcl} d_{-1(1)}(z) &:= & -\frac{1}{k_+(z)^2 + 2i\gamma k_+(z) - \Omega^2} = \frac{1}{(k_+(z) + i\gamma)^2 - \Omega^2 + \gamma^2} \,, \\ & & (k_+(z)^2 + 2i\gamma k_+(z) - \Omega^2 \neq 0) \end{array}$$

where  $k_+(z)$  is defined as the unique solution to

$$k_+^2 = -z$$
,  $Im k_+ > 0$ ,

if

$$z \notin \left(-\infty, 0\right],$$

and to

$$k_{+}^{2} = -z, Re k_{+} \ge 0,$$
  
 $z \in (-\infty, 0],$ 

but further, there is 'no place' for the 'rest of  $Det_{-1}$  ', i.e., for  $det_{-1,-,0}$  .

With  $det_{-1,-,0}$ , we connect another function of z:

$$d_{-1(2)}(z) := -\frac{1}{k_{-}(z)^{2} + 2i\gamma k_{-}(z) - \Omega^{2}} = \frac{1}{(k_{-}(z) + i\gamma)^{2} - \Omega^{2} + \gamma^{2}},$$
$$(k_{-}(z)^{2} + 2i\gamma k_{-}(z) - \Omega^{2} \neq 0)$$

where  $k_{-}(z)$  is defined as the unique solution to

$$k_{-}^2 = -z$$
,  $Im k_{-} < 0$ ,

if

$$z \not\in (-\infty, 0],$$

 $k_{-}^2 = -z$ ,  $Re k_{-} < 0$ ,

and to

if

$$z \in (-\infty, 0]$$
.

Thus we are not able to reformulate  $Det_{-1}(k)$  in terms of ONE function of z. . We can only reformulate  $Det_{-1}(k)$  in terms of TWO functions of ONE variable:  $d_{-1(1)}(z)$  and  $d_{-1(2)}(z)$ .

Nevertheless, physicists prefer to say, there is ONE analytic function  $D_{-1}(z)$  defined on 'TWO exemplars' of **C**, so that

(1) on the first 'exemplar' of **C**,  $D_{-1}(z) = d_{-1(1)}(z)$ ; this first 'exemplar' of **C** is called **first** or **physical sheet** of F(z);

(2) on the second 'exemplar' of  $\mathbf{C}$ ,  $D_{-1}(z) = d_{-1(2)}(z)$ ; this second 'exemplar' of  $\mathbf{C}$  is called **second** or **unphysical sheet** of F(z).

One can say also:

 $d_{-1(2)}(z)$  is a part of an **extended**  $d_{-1(1)}(z)$ , the part which is defined on the second 'exemplar' of **C**, and this second 'exemplar' of **C** is named the **second sheet** associated with R(z).

—In these words,  $z_{res}$  is a **resonant pole**, if  $z_{res}$  is a pole placed on the second sheet of the resolvent.

From the formal mathematical standpoint, the notion we have just introduced can be explained as following: 'two exemplars' of  $\mathbf{C}$  can be defined as

$$(\mathbf{C} \times \{1\}) \cup (\mathbf{C} \times \{2\});$$

the 'first' sheet is identified with  $\mathbf{C} \times \{1\}$ , and the second one is identified with  $\mathbf{C} \times \{2\}$ ; finally put

$$\begin{split} D_{-1}(z\times\{1\}) &:= d_{-1(1)}(z)\,,\\ D_{-1}(z\times\{2\}) &:= d_{-1(2)}(z)\,. \end{split}$$

22

if

The constructions of such kind are basic objects of the theory of **Riemann** surfaces, but we will not go into details.

There are reasons for it, and one of them is that the concept of Riemann surface has been elaborated so as to avoid introducing principal distinctions between the sheets, whereas we would like to contradistinguish them,—we need two or even three objects,—one object connected immediately with resolvent and the other(s) one(s) done with amplitudes and resonant poles. In other words, we prefer constructions like that we have displayed as **Observation 1**.

Another reason is that we need rather R(z) and  $R(-k^2)$  than  $det_{\#}$ , but the formers are not usual scalar functions. They are operator-valued function, thus, we must give exact definitions suited to the case. First we must define what we mean by 'element of  $R(-k^2)$  is an analytic function', what we mean by 'analytic extension', and then many other things. Otherwise confusion and false conclusions will occur!!

How to find proper mathematical definitions, does not form the subject of this paper.

0

## Appendix A.

## Free Green's Function.

Here we reproduce a well-known proof that

$$(z - c^2 \Delta)^{-1}(x, x') = \frac{i}{2kc^2} e^{ik|x - x'|}$$
, where  $k^2 = -z/c^2$ ,  $Im \, k > 0$ 

It can be deduced from p-representation of  $(z - c^2 \Delta)^{-1}$ :

$$(z - c^{2}\Delta)^{-1}(x, x') = \frac{1}{2\pi} \int_{-\infty}^{+\infty} \frac{e^{ip(x-x')}}{z + c^{2}p^{2}} dp.$$

We have:

$$\begin{split} (z - c^2 \Delta)^{-1}(x, x') &= \frac{1}{2\pi} \int_{-\infty}^{\infty} \frac{e^{ip(x-x')}}{z + c^2 p^2} dp \\ &= \frac{1}{c^2} \frac{1}{2\pi} \int_{-\infty}^{\infty} \frac{e^{ip(x-x')}}{p^2 + z/c^2} dp \\ &= \frac{i}{c^2} \frac{1}{2\pi i} \int_{-\infty}^{\infty} \frac{e^{ip(x-x')}}{p^2 - k^2} dp \\ &= \frac{i}{c^2} \frac{1}{2\pi i} \int_{-\infty}^{\infty} \frac{e^{ip(x-x')}}{p + k} \frac{1}{p - k} dp \\ &= \frac{i}{c^2} \frac{e^{ik|x-x'|}}{2k}, \\ &= \frac{i}{2kc^2} e^{ik|x-x'|}, \text{ where } \quad k^2 = -z/c^2, \quad Im \, k > \end{split}$$

### References

- [AK] S. ALBEVERIO AND P. KURASOV, Singular Perturbations of Differential Operators. Solvable Scrödinger Type Operators, London Mathematical Society: Lecture Note Series. 271, 1999. CAMBRIDGE UNIVERSITY PRESS
- [BF] F.A. BEREZIN, L.D. FADDEEV, Remark on the Schrödinger equation with singular potential, Dokl. Akad. Nauk. SSSR, 137 (1961) 1011-1014 (in Russian).
- [Do] W. DONOGHUE, On the perturbation of spectra, Comm. Pure App. Math. 18 (1965) 559-579
- [Fog] S.R. FOGUEL, Finite Dimensional Perturbations In Banach Spaces, American Jornal of Mathematics, Volume 82, Issue 2 (Apr., 1960), 260-270
- [Fr] K.O. FRIEDRICHS, *Perturbation of Spectra in Hilbert Space*, American Mathematical Society, Providence, (1965)
- [Gan] F.R. GANTMAHER, *The theory of matrices.* 4-th ed., Moscow, Nauka (1988)(Russian); English translation: Chelsea publishing company, New York (1959).
- [Jack] J.D. JACKSON: Classical electrodynamics. John Wiley & Sons, Inc. New York-London, 1962. see 'Radiation Reaction, Abraham Lorentz Equation, Braking Radiation' and all that.
- [RS1] M. REED, B. SIMON, Methods of Modern Mathematical Physics, vol 1, Functional analysis, - N.Y.: Academic Press, 1972.
- [RS2] M. REED, B. SIMON, Methods of Modern Mathematical Physics, vol 2, Fourier analysis, Self-Adjointness, - N.Y.: Academic Press, 1975.
- [RS3] M. REED, B. SIMON, Methods of Modern Mathematical Physics, vol 3, Scattering Theory, - N.Y.: Academic Press, 1979.
- [RS4] M. REED, B. SIMON, Methods of Modern Mathematical Physics, vol 4, Analysis of Operators, - N.Y.: Academic Press, 1978.

#### **Electronic Print:**

#### Mathematical Physics Preprint Archive

[DerFr] mp\_arc 02-275

Derezinski J., Fruboes R. Renormalization of the Friedrichs Hamiltonian (16K, LATeX 2e)

http://www.ma.utexas.edu/mp\_arc-bin/mpa?yn=02-275 http://mpej.unige.ch/mp\_arc-bin/mpa?yn=02-275 http://www.maia.ub.es/mp\_arc-bin/mpa?yn=02-275

#### [Der] mp\_arc 02-300

Jan Derezinski

Van Hove Hamiltonians—exactly solvable models of the infrared and ultraviolet problem.

(62K, LATeX 2e)

http://www.ma.utexas.edu/mp\_arc-bin/mpa?yn=02-300 http://mpej.unige.ch/mp\_arc-bin/mpa?yn=02-300 http://www.maia.ub.es/mp\_arc-bin/mpa?yn=02-300

[03-33] mp\_arc 03-33

Sergej A. Choroszavin ( sergius@pve.vsu.ru )

An Interaction of An Oscillator with An One-Dimensional Scalar Field. Simple Exactly Solvable Models based on Finite Rank Perturbations Methods. I: D'Alembert-Kirchhoff-like formulae

(88K, LaTeX 2.09) Jan 29, 03

http://www.ma.utexas.edu/mp\_arc-bin/mpa?yn=03-33 http://mpej.unige.ch/mp\_arc-bin/mpa?yn=03-33 http://www.maia.ub.es/mp\_arc-bin/mpa?yn=03-33 http://kleine.mat.uniroma3.it/mp\_arc-bin/mpa?yn=03-33

#### arXiv.org e-Print archive (LANL E-Print)

[AMN] Paper: physics/0001009 From: adolfo@lafexSu1.lafex.cbpf.br (Adolfo Malbouisson) Date: Wed, 5 Jan 2000 19:20:08 GMT (15kb)

> Title: An Exact Approach to the Oscillator Radiation Process in an Arbitrarily Large Cavity Authors: N.P. Andion, A.P.C. Malbouisson and A. Mattos Neto Comments: 27 pages Subj-class: Atomic Physics; Mathematical Physics http://arXiv.org/abs/physics/0001009

[BKZ] Paper: math-ph/0210051 From: Volker Bach <vbach@mathematik.uni-mainz.de> Date: Wed, 30 Oct 2002 00:06:05 GMT (31kb)

> Title: Mathematical analysis of the photoelectric effect Authors: Volker Bach, Frederic Klopp and Heribert Zenk Comments: See also http://www.intlpress.com/ATMP Subj-class: Mathematical Physics Journal-ref: Adv. Theor. Math. Phys. 5 (2001) 969-999 http://arXiv.org/abs/math-ph/0210051

[Ch03-1] Paper: math.DS/0301167 From: "Sergej A. Choroszavin" <sergius@pve.vsu.ru> Date: Thu, 16 Jan 2003 04:34:16 GMT (18kb) Date (revised v2): Fri, 17 Jan 2003 03:07:58 GMT (18kb) Date (revised v3): Mon, 27 Jan 2003 19:00:54 GMT (18kb)

Title: An Interaction of An Oscillator with An One-Dimensional Scalar Field. Simple Exactly Solvable Models based on Finite Rank Perturbations

26

Methods. I: D'Alembert-Kirchhoff-like formulae Author: Sergej A. Choroszavin Comments: Latex 2.09 Subj-class: Dynamical Systems; Mathematical Physics http://arXiv.org/abs/math.DS/0301167

This paper is the same as in Mathematical Physics Preprint Archive, PAPER: 03-33

[MPB] Paper: hep-th/9207033 From: physth@ulb.ac.be Date: Fri, 10 Jul 92 16:52:54 +0200 (20kb)

> Title: On the Problem of the Uniformly Accelerated Oscillator Authors: S. Massar, R. Parentani, R. Brout Comments: 14 pages (+postscript figures attached), ULB-TH-03/92 http://arXiv.org/abs/hep-th/9207033