

# Singularities, Lax degeneracies and Maslov indices of the periodic Toda chain

JA Foxman<sup>†</sup> and JM Robbins<sup>\*,‡</sup>

<sup>†</sup> School of Mathematics

University of Bristol, University Walk, Bristol BS8 1TW, UK

<sup>‡</sup> The Mathematical Sciences Research Institute

1000 Centennial Drive, #5070, Berkeley, CA 94720-5070, USA

December 15, 2017

## Abstract

The  $n$ -particle periodic Toda chain is a well known example of an integrable but nonseparable Hamiltonian system in  $\mathbb{R}^{2n}$ . We show that  $\Sigma_k$ , the  $k$ -fold singularities of the Toda chain, ie points where there exist  $k$  independent linear relations amongst the gradients of the integrals of motion, coincide with points where there are  $k$  (doubly) degenerate eigenvalues of representatives  $L$  and  $\bar{L}$  of the two inequivalent classes of Lax matrices (corresponding to degenerate periodic or antiperiodic solutions of the associated second-order difference equation). The singularities are shown to be nondegenerate, so that  $\Sigma_k$  is a codimension- $2k$  symplectic submanifold.  $\Sigma_k$  is shown to be of elliptic type, and the frequencies of transverse oscillations under Hamiltonians which fix  $\Sigma_k$  are computed in terms of spectral data of the Lax matrices.

If  $\mu(C)$  is the (even) Maslov index of a closed curve  $C$  in the regular component of  $\mathbb{R}^{2n}$ , then  $(-1)^{\mu(C)/2}$  is given by the product of the holonomies (equal to  $\pm 1$ ) of the even- (or odd-) indexed eigenvector bundles of  $L$  and  $\bar{L}$ .

---

\*E-mail address: j.robbs@bristol.ac.uk

# 1 Introduction

The Toda chain is a canonical example of a nonseparable but integrable Hamiltonian system. It consists of  $n$  particles on the line with exponential interactions between consecutively labeled particles. In the periodic Toda chain, the  $n$ th particle is coupled to the first. The Hamiltonian is

$$H = \sum_{j=1}^n \frac{1}{2}(p_j^2 + b_j^2), \quad (1.1)$$

where

$$b_j = e^{(q_j - q_{j+1})/2}, \quad (1.2)$$

and  $q_{n+1} \equiv q_1$ .

The Toda lattice was introduced in [24]. Its integrability was established by Hénon [19] and Flaschka [13] using the method of Lax pairs. There now exists an extensive literature on the problem (see, eg, [14]). Properties of eigenvectors of Lax matrices over their associated spectral curve were studied in classical papers by Adler and van Moerbeke [1, 2, 3] and van Moerbeke and Mumford [26]. Audin [6] has used these results to analyse the topology of the set of regular values of the integrals of motion. A recent account is given by Babelon, Bernard and Talon [7].

In the last 20 years there has been much interest in the topology of integrable finite-dimensional Hamiltonian systems. The generic local structure and dynamics is given by the Liouville-Arnold theorem [5], according to which neighbourhoods of phase space are foliated into invariant Lagrangian submanifolds diffeomorphic to  $\mathbb{R}^{n-k} \times \mathbb{T}^k$ , where  $\mathbb{T}^k$  is the  $k$ -torus, and the dynamics is linearised by action-angle coordinates. This local behaviour breaks down at critical points of the energy-momentum map, which comprise invariant sets of lower dimension. A Morse theory for integrable Hamiltonian systems, wherein the global topology is described in terms of these critical sets, has been extensively developed by Fomenko [15], Eliasson [12], Vey [27] and Tien Zung [28], among others.

Our interest here is in the phase space topology of the Toda chain. The paper is organised as follows. In Section 2.1, we obtain Lax generators for the flows generated by a complete set of  $n$  integrals  $F : \mathbb{R}^{2n} \rightarrow \mathbb{R}^n$ . These Lax generators are constructed from representatives  $L$  and  $\bar{L}$  of the two inequivalent classes of symmetric Lax matrices for the periodic Toda chain (these classes correspond to periodic and antiperiodic solutions of the associated second-order difference equation). Our formulation is related to the classical treatment of van Moerbeke and Mumford of general periodic finite difference operators [26], but our approach is self-contained and quite elementary.

These results are used to establish, in Section 3, a one-to-one correspondence between singularities of the Toda flow (critical points of  $F$ ) and eigenvalue degeneracies of the Lax matrices  $L$  and  $\bar{L}$ . More precisely, the corank of  $dF$  is equal to the number of (doubly) degenerate eigenvalues of  $L$  and  $\bar{L}$ . Similar ideas relating singularities to points of degeneracy on the spectral curve are discussed by Audin [6].

In Section 4 we determine the local structure of the corank- $k$  singularities  $\Sigma_k$ . These are shown to be codimension- $2k$  symplectic submanifolds of elliptic type composed of  $\binom{n-1}{k}$  components disconnected from each other. The eigenvalues of the linearised integrable flows which fix them are computed in terms of the spectral data of  $L$  and  $\bar{L}$ . There are singularities of corank between 1 and  $(n-1)$ , and  $\Sigma_k$  is contained in the closure of  $\Sigma_j$  for  $k > j$ . For  $n = 3$ , related results on the singularities and critical values for the Toda chain and its algebraic generalisations have been obtained by Polyakova [22] following the programme of Fomenko [15].

The Maslov index is a topological invariant of Lagrangian tori of integrable systems which appears in the semiclassical (EBK) quantisation conditions [21, 20]. In a companion paper [16], we show that the Maslov index is determined by corank-one singularities. In Section 5, it is shown that the (even) Maslov index of a closed curve in the set of regular points of  $F$  is determined, modulo 4, by the product of the even- (or odd-) indexed holonomies of the eigenvector bundles of  $L$  and  $\bar{L}$ . We note that, because the Toda chain is nonseparable, application of the semiclassical quantisation rules is not straightforward.

## 2 Higher Lax flows

The Lax formulation of the equations of motion for the Toda chain was obtained by Flaschka [13]. Let  $L$  be the  $n \times n$  symmetric matrix given by

$$L = \begin{pmatrix} p_1 & b_1 & 0 & \cdots & 0 & b_n \\ b_1 & p_2 & b_2 & 0 & \cdots & 0 \\ \vdots & & & & & \vdots \\ 0 & \cdots & 0 & b_{n-2} & p_{n-1} & b_{n-1} \\ b_n & 0 & \cdots & 0 & b_{n-1} & p_n \end{pmatrix}, \quad (2.1)$$

and  $M_{(2)}$  the  $n \times n$  antisymmetric matrix given by

$$M_{(2)} = \frac{1}{2} \begin{pmatrix} 0 & b_1 & 0 & \cdots & 0 & -b_n \\ -b_1 & 0 & b_2 & 0 & \cdots & 0 \\ \vdots & & & & & \vdots \\ 0 & \cdots & 0 & -b_{n-2} & 0 & b_{n-1} \\ b_n & 0 & \cdots & 0 & -b_{n-1} & 0 \end{pmatrix}. \quad (2.2)$$

(To simplify notion, in particular in Section 4, we will sometimes write  $M$  instead of  $M_{(2)}$ .) It is straightforward to verify that Hamilton's equations for the Hamiltonian (1.1) imply the Lax equation

$$\dot{L} := \{L, H\} = [L, M_{(2)}]. \quad (2.3)$$

Conversely, (2.3) along with the (independent) equation  $\sum_{r=1}^n p_r = \sum_{r=1}^n \dot{q}_r$  imply Hamilton's equations. Thus, Hamilton's equations and the Lax equation are essentially equivalent. It is convenient to express  $L$  and  $M_{(2)}$  in index form,

$$L_{rs} = p_r \delta_{rs} + b_r \delta_{r+1,s} + b_s \delta_{r,s+1}, \quad (2.4a)$$

$$M_{(2)rs} = b_r \delta_{r+1,s} - b_s \delta_{r,s+1}. \quad (2.4b)$$

For convenience, here and elsewhere, matrices and vectors are regarded as being periodic in their indices, with period  $n$ . Thus  $L_{r+n,s} = L_{r,s+n} = L_{rs}$ . Similarly, we take  $\delta_{rs}$  to be one if  $r = s \pmod n$  and to be zero otherwise.

As is well known, the Lax equation implies that the eigenvalues of  $L$ , as well as functions of them, are constants of the motion. In particular, the  $n$  functions

$$F_j = \frac{1}{j} \text{Tr } L^j, \quad 1 \leq j \leq n, \quad (2.5)$$

are conserved. We note that

$$F_1 = p_1 + \cdots + p_n \quad (2.6)$$

is the centre-of-mass momentum, while  $F_2$  is the Hamiltonian.

In this section we construct Lax equations for the Hamiltonian flows generated by each of the  $F_j$ 's. That is, we find antisymmetric matrices  $M_{(j)}$  such that

$$\{L, F_j\} = [L, M_{(j)}] \quad (2.7)$$

We may take  $M_{(1)} = 0$  (since  $F_1$  generates uniform translations and  $L$  is translation-invariant), while  $M_{(2)}$  is given by (2.1). The formulation presented here is related to that of van Moerbeke and Mumford [26], who give

nilpotent (triangular) Lax generators for the higher flows of general periodic finite difference operators. Before giving expressions for  $M_{(j)}$  for  $j > 2$ , we recall that the higher Lax equations (2.7) already imply that the functions  $F_j$  are in involution. Indeed,

$$\{F_{j+1}, F_k\} = \text{Tr} (L^j \{L, F_k\}) = \text{Tr} (L^j [L, M_{(k)}]) = 0, \quad (2.8)$$

where the last equality follows from the cyclicity of the trace. We recall, too, that integrability does not follow immediately from (2.8); one also needs to show that the  $F_j$ 's are functionally independent (here, functional independence is implied by Theorem 4.1 below).

We note that  $H$  is invariant under the  $n$  substitutions

$$b_j \mapsto -b_j. \quad (2.9)$$

It follows that the Lax equation (2.3) is similarly invariant under (2.9). It is easily seen that an even number of such substitutions can be generated by conjugations  $L \mapsto SLS^{-1}$ ,  $M_{(2)} \mapsto SM_{(2)}S^{-1}$  for  $S$  a diagonal matrix of  $\pm 1$ 's. The Lax equation (2.3) is trivially invariant under such conjugations, as are the  $F_j$ 's. There are, therefore, two inequivalent classes of Lax pairs with respect to (2.9). These are characterised by whether the number of negative  $b_j$ 's is even or odd. A spectral characterisation of these even and odd classes is provided by the difference equation

$$b_r v_{r+1} + p_r v_r + b_{r-1} v_{r-1} = \lambda v_r. \quad (2.10)$$

Eigenvalues  $\lambda$  (with eigenvectors  $\mathbf{v}$ ) of even  $L$ 's correspond to periodic solutions  $v_{r+n} = v_r$  of (2.10), whereas eigenvalues of odd  $L$ 's correspond to antiperiodic solutions  $v_{r+n} = -v_r$ . For definiteness and convenience, we take  $L$ , as given in (2.1), to be the even representative, and  $\bar{L}$  to be given by replacing  $b_n$  by  $-b_n$ , ie

$$\bar{L}_{rs} = p_r \delta_{rs} + \sigma_r b_r \delta_{r+1,s} + \sigma_s b_s \delta_{r,s+1}, \quad (2.11)$$

where

$$\sigma_r = \begin{cases} -1, & r = 0 \pmod n, \\ 1, & \text{otherwise,} \end{cases} \quad (2.12)$$

to be the odd representative.

We say that an  $n \times n$  symmetric matrix  $A$  is *off-banded of width  $j$* , or  *$j$ -off-banded*, if it has precisely  $(n - j)$  consecutive zero diagonals on or above

main diagonal (thus,  $j$  is the number of diagonals above these zero diagonals, the first of which does not vanish). Equivalently,  $A$  is  $j$ -off-banded if

$$A_{r,r+d} = 0, \quad 1 \leq r \leq n, \quad 0 \leq d < \min(n - j - 1, n - r), \quad (2.13a)$$

$$\sum_{r=1}^j |A_{r,r+n-j}| \neq 0. \quad (2.13b)$$

**Proposition 2.1.** *For  $1 \leq j \leq n$ ,  $L^j - \bar{L}^j$  is  $j$ -off-banded. Moreover, its elements on the first nonzero diagonal are given by*

$$\left( L^j - \bar{L}^j \right)_{r,r+j} = 2b_{r-1}b_{r-2} \cdots b_{r-j}, \quad j < n, \quad (2.14a)$$

$$= 4, \quad j = n, \quad (2.14b)$$

where  $1 \leq r \leq j$ .

*Proof.* Fix  $1 \leq r \leq n$ , and take  $d$  such that  $0 \leq d \leq \min(n - j - 1, n - r)$ . From (2.11) and (2.12) it is clear that  $(L^j - \bar{L}^j)_{r,r+d}$  is given by twice the sum of terms of  $(L^j)_{r,r+d}$  which are of odd degree in  $b_n$ . The terms of  $(L^j)_{r,r+d}$  are products  $T(t_0, \dots, t_j)$  of the form

$$T(t_0, \dots, t_j) = L_{t_0 t_1} L_{t_1 t_2} \cdots L_{t_{j-1} t_j}, \quad (2.15)$$

where  $t_0 = r$  and  $t_j = r + d$ .  $L_{1n}$  and  $L_{n1}$ , if they appear, contribute the only factors of  $b_n$  to  $T(t_0, \dots, t_j)$ .

Since  $L$  is banded modulo  $n$  (cf (2.4a)),  $T(t_0, \dots, t_j)$  vanishes unless each pair of consecutive indices  $t_k$  and  $t_{k+1}$  in (2.15) differ by 0 or  $\pm 1$  modulo  $n$ . Let us call the factor  $L_{t_k t_{k+1}}$  a *right step* if  $t_{k+1} = t_k + 1 \pmod{n}$ , and a *left step* if  $t_{k+1} = t_k - 1 \pmod{n}$  (diagonal factors, for which  $t_{k+1} = t_k \pmod{n}$ , are neither left nor right steps). Let  $u$  denote the number of right steps minus the number of left steps in  $T(t_0, \dots, t_j)$ . Then

$$u = d \pmod{n}. \quad (2.16)$$

Since  $-j \leq u \leq j$  and, by assumption,  $1 \leq j \leq n$  and  $0 \leq d < n$ , it follows that either  $u = d$  or  $u = d - n$ .

First, suppose that  $u = d - n$ . We show that  $T(t_0, \dots, t_j)$  is of odd degree in  $b_n$ . This is certainly true for terms which contain no right steps. Such terms are products of diagonal elements of  $L$  (which do not contribute factors of  $b_n$ ) and the left-step-only product

$$L_{r,r-1} L_{r-1,r-2} \cdots L_{r+(n-d)+1,r+(n-d)}. \quad (2.17)$$

$b_n$  appears just once in (2.17), in the factor  $L_{10} := L_{1n}$ . A general term  $T(t_0, \dots, t_j)$  with  $u = d - n$  is a product of diagonal elements, the left-step-only product (2.17), and palindromic products of off-diagonal elements, ie, products of the form

$$L_{k,k+1}L_{k+1,k+2} \cdots L_{k+p-1,k+p} \cdot L_{k+p,k+p-1} \cdots L_{k+1,k+2}L_{k,k+1} \quad (2.18)$$

in which every factor appears twice.

Next, suppose that  $d = u$ . A similar argument implies that  $T(t_0, \dots, t_j)$  is of even degree in  $b_n$ . In this case, we note that the right-step-only product  $L_{r,r+1}L_{r+1,r+2} \cdots L_{r+d-1,r+d}$  contains no factors of  $L_{1n}$  or  $L_{n1}$ , and that, in general,  $T(t_0, \dots, t_j)$  is a product of diagonal elements, the right-step-only product, and palindromic products (2.18).

Thus,  $T(t_0, \dots, t_j)$  is of odd degree in  $b_n$  if and only if  $u = d - n$ , or, equivalently,  $d = n - |u|$ . Since  $|u| \leq j$ , this condition can be satisfied only if  $d \geq n - j$ . Thus, if  $d < n - j$ , then  $(L^j - \bar{L}^j)_{r,r+d} = 0$ , in accord with (2.13a).

To establish (2.13b), we verify (2.14), which implies that  $L^j - \bar{L}^j$  has nonzero elements on the  $(n - j)$ th diagonal above the main diagonal. Let  $d = n - j$ . For  $T(t_0, \dots, t_j)$  to be of odd degree in  $b_n$ , we require that  $u = j$ . For  $j < n$ , there is only one such term, namely the left-step-only product  $L_{r,r-1}L_{r-1,r-2} \cdots L_{r-j+1,r-j} = b_{r-1}b_{r-2} \cdots b_{r-j}$ . (2.14a) follows. For  $j = n$ , there are two nonzero terms with  $u = n$ . The first is the left-step-only product  $L_{r,r-1}L_{r-1,r-2} \cdots L_{r-n+1,r-n} = b_{r-1}b_{r-2} \cdots b_{r-n} = 1$ . The second is the right-step-only product  $L_{r,r+1}L_{r+1,r+2} \cdots L_{r+n-1,r+n} = b_{r+1}b_{r+2} \cdots b_r = 1$ . (2.14b) follows.  $\square$

From Proposition 2.1 it follows immediately that

$$\text{Tr } L^j = \begin{cases} \text{Tr } \bar{L}^j, & 1 \leq j < n, \\ \text{Tr } \bar{L}^n + 4n, & j = n. \end{cases} \quad (2.19)$$

Also, we note that if a linear combination of off-banded matrices vanishes and the matrices all have different widths, then the coefficient of each matrix vanishes (argue inductively, starting with the matrix of greatest width). Therefore, Proposition 2.1 also implies the following:

**Corollary 2.1.** *If  $\sum_{j=2}^n c_j(L^{j-1} - \bar{L}^{j-1}) = 0$ , then  $c_2 = \cdots = c_n = 0$ .*

We introduce the following notation: given an  $n \times n$  matrix  $A$ , let  $A_+$  denote its strictly upper triangular part, ie the matrix elements  $A_{r,r+d}$  with  $1 \leq r \leq n$  and  $1 \leq d \leq n - r$ . The generators of the Lax flows for the  $F_j$ 's are given by the following:

**Proposition 2.2.** Let  $M_{(j)}$  and  $\overline{M}_{(j)}$  be the antisymmetric matrices given by

$$M_{(j)+} = \frac{1}{2}(\overline{L}^{j-1})_+, \quad (2.20a)$$

$$\overline{M}_{(j)+} = \frac{1}{2}(L^{j-1})_+ \quad (2.20b)$$

for  $1 \leq j \leq n$ . Then

$$\{L, F_j\} = [L, M_{(j)}], \quad (2.21a)$$

$$\{\overline{L}, F_j\} = [\overline{L}, \overline{M}_{(j)}]. \quad (2.21b)$$

We note that  $M_{(1)}$  and  $\overline{M}_{(1)}$  both vanish (consistent with the fact that  $L$  is invariant under uniform translations) while for  $j = 2$ , (2.20a) agrees with (2.1).

*Proof.* First, we note that (2.19) implies that under the substitution  $b_n \mapsto -b_n$ ,  $F_j$  changes by at most a constant. Therefore, the two sets of Lax equations (2.21a) and (2.21b) are related by this substitution, and it suffices to verify just one of them. For definiteness we consider (2.21b).

Since both sides are symmetric matrices, it suffices to verify (2.21b) for elements on or above the main diagonal. The elements of the left-hand side are given by

$$\{\overline{L}_{rs}, F_j\} = \frac{1}{j} \text{Tr} \{ \overline{L}_{rs}, L^j \} = \text{Tr} ( \{ \overline{L}_{rs}, L \} L^{j-1} ). \quad (2.22)$$

From (2.4a), (2.11) and

$$\{b_r, p_s\} = \frac{1}{2} b_r (\delta_{rs} - \delta_{r+1,s}), \quad (2.23)$$

a straightforward calculation gives, for  $1 \leq r \leq s \leq n$ , that

$$\{\overline{L}_{rs}, F_j\} = \begin{cases} b_{r-1} (L^{j-1})_{r-1,r} - b_r (L^{j-1})_{r,r+1}, & r = s, \\ \frac{1}{2} b_r ((L^{j-1})_{rr} - (L^{j-1})_{r+1,r+1}), & r+1 = s, \\ \frac{1}{2} b_n ((L^{j-1})_{11} - (L^{j-1})_{nn}), & r = 1, s = n, \\ 0, & \text{otherwise.} \end{cases} \quad (2.24)$$

Next we evaluate the right-hand side of (2.21b), ie  $[\overline{L}, \overline{M}_{(j)}]$ . From (2.11),

$$[\overline{L}, \overline{M}_{(j)}]_{rs} = p_r \overline{M}_{(j)rs} + \sigma_{r-1} b_{r-1} \overline{M}_{(j)r-1,s} + \sigma_r b_r \overline{M}_{(j)r+1,s} + (r \leftrightarrow s). \quad (2.25)$$

For  $r = s$ , (2.25) yields  $b_{r-1} L^{j-1}_{r-1,r} - b_r L^{j-1}_{r+1,r}$ , in agreement with (2.24) (note that  $\sigma_{r-1} \overline{M}_{(j)r-1,r} = \frac{1}{2} L^{j-1}_{r-1,r}$ , and similarly,  $\sigma_r \overline{M}_{(j)r+1,r} = -\frac{1}{2} L^{j-1}_{r+1,r}$ ).

To evaluate the off-diagonal elements  $1 \leq r < s \leq n$  in (2.24), we will make use of the trivial identity  $[\mathbf{L}, \frac{1}{2}\mathbf{L}^{j-1}] = 0$ , or

$$[\mathbf{L}, \frac{1}{2}\mathbf{L}^{j-1}]_{rs} = \frac{1}{2}p_r(\mathbf{L}^{j-1})_{rs} + \frac{1}{2}b_{r-1}(\mathbf{L}^{j-1})_{r-1,s} + \frac{1}{2}b_r(\mathbf{L}^{j-1})_{r+1,s} - (r \leftrightarrow s) = 0. \quad (2.26)$$

Referring to the terms in (2.25), we use (2.20b) to express  $M_j$  in terms of  $\mathbf{L}^{j-1}$ , as follows:

$$\begin{aligned} p_r \overline{M}_{(j)rs} + (r \leftrightarrow s) &= p_r \left( \frac{1}{2} \mathbf{L}^{j-1} \right)_{rs} - (r \leftrightarrow s), \\ \sigma_{r-1} b_{r-1} \overline{M}_{(j)r-1,s} + (r \leftrightarrow s) &= b_{r-1} \left( \frac{1}{2} \mathbf{L}^{j-1} \right)_{r-1,s} - (r \leftrightarrow s) \\ &\quad - \delta_{r1} \delta_{sn} b_n \left( \frac{1}{2} \mathbf{L}^{j-1} \right)_{nn} + \delta_{r+1,s} b_r \left( \frac{1}{2} \mathbf{L}^{j-1} \right)_{rr}, \\ \sigma_r b_r \overline{M}_{(j)r+1,s} + (r \leftrightarrow s) &= b_r \left( \frac{1}{2} \mathbf{L}^{j-1} \right)_{r+1,s} - (r \leftrightarrow s) \\ &\quad + \delta_{r1} \delta_{sn} b_n \left( \frac{1}{2} \mathbf{L}^{j-1} \right)_{11} - \delta_{r+1,s} b_r \left( \frac{1}{2} \mathbf{L}^{j-1} \right)_{r+1,r+1}. \end{aligned} \quad (2.27)$$

Substituting the preceding into (2.25) and using the identity (2.26), we get that

$$\begin{aligned} [\overline{\mathbf{L}}, \overline{M}_{(j)}]_{rs} &= [\mathbf{L}, \frac{1}{2}\mathbf{L}^{j-1}]_{rs} + \frac{1}{2} \delta_{r+1,s} b_r ((\mathbf{L}^{j-1})_{rr} - (\mathbf{L}^{j-1})_{r+1,r+1}) + \\ &\quad \frac{1}{2} b_n \delta_{r1} \delta_{sn} ((\mathbf{L}^{j-1})_{11} - (\mathbf{L}^{j-1})_{nn}). \end{aligned} \quad (2.28)$$

As the first term vanishes, this agrees with (2.24).  $\square$

### 3 Singularities and eigenvalue degeneracies

The singularities of an integrable system  $F : \mathbb{R}^{2n} \rightarrow \mathbb{R}^n$  are the critical points of  $F$ . Here we show that singularities of the Toda chain coincide with eigenvalue degeneracies of the Lax matrices  $\mathbf{L}$  and  $\overline{\mathbf{L}}$ . Let  $\Sigma$  denote the set of singularities of the Toda chain, and let  $\Sigma_k \subset \Sigma$  denote the subset in which there are precisely  $k$  linear relations amongst the  $dF_j$ 's, ie

$$\Sigma_k = \{(\mathbf{q}, \mathbf{p}) \mid \text{corank } dF = k\}. \quad (3.1)$$

We observe that eigenvalues of  $\mathbf{L}$  and  $\overline{\mathbf{L}}$  are at most two-fold degenerate, since the associated eigenvectors are solutions of the second-order linear difference equation (2.10), which for a given value of  $\lambda$  has at most two linearly independent solutions. Let  $\nu$  and  $\overline{\nu}$  denote the number of doubly degenerate eigenvalues of  $\mathbf{L}$  and  $\overline{\mathbf{L}}$  respectively.

**Theorem 3.1.**

$$\text{corank } dF = \nu + \overline{\nu}. \quad (3.2)$$

*Proof.* Let  $\mathbf{z}_* \in \mathbb{R}^{2n}$ . For convenience, let  $F_*$  denote  $F(\mathbf{z}_*)$ ,  $L_*$  denote  $L(\mathbf{z}_*)$ , etc. Let  $V_* \subset \mathbb{R}^n$  denote the space of linear relations amongst the  $dF_{j*}$ 's, ie

$$V_* = \left\{ \mathbf{c} \in \mathbb{R}^n \mid \sum_{j=1}^n c_j dF_{j*} = 0 \right\}. \quad (3.3)$$

so that  $\text{corank } dF_* = \dim V_*$ . We show that

$$\dim V_* = \nu_* + \bar{\nu}_*. \quad (3.4)$$

We first show that  $\dim V_* \leq \nu_* + \bar{\nu}_*$ . Let  $\mathcal{A}_*$  denote the real antisymmetric commutant of  $L_*$ ; that is,  $\mathcal{A}_*$  consists of all  $n$ -dimensional real antisymmetric matrices which commute with  $L_*$ . For  $\mathbf{c} \in V_*$ , Proposition 2.2 and Eq. (3.3) imply that

$$\left[ L_*, \sum_{j=1}^n c_j M_{(j)*} \right] = \left\{ L, \sum_{j=1}^n c_j F_j \right\}_* = 0, \quad (3.5)$$

so that  $\sum_{j=1}^n c_j M_{(j)*} \in \mathcal{A}_*$ . Similarly, letting  $\bar{\mathcal{A}}_*$  denote the real antisymmetric commutant of  $\bar{L}$ , we have that  $\sum_{j=1}^n c_j \bar{M}_{(j)*} \in \bar{\mathcal{A}}_*$ . Regarding  $\mathcal{A}_*$  and  $\bar{\mathcal{A}}_*$  as real vector spaces, we define a linear map from  $V_*$  to  $\mathcal{A}_* \oplus \bar{\mathcal{A}}_*$  according to

$$\mathbf{c} \mapsto \left( \sum_{j=1}^n c_j M_{(j)*} \right) \oplus \left( \sum_{j=1}^n c_j \bar{M}_{(j)*} \right). \quad (3.6)$$

This map is 1-1, for if  $\sum_{j=1}^n c_j M_{(j)*} = \sum_{j=1}^n c_j \bar{M}_{(j)*} = 0$ , then from (2.20) and Corollary 2.1 it follows that

$$c_2 = \cdots = c_n = 0. \quad (3.7)$$

But (3.7) and (3.3) imply that  $c_1 dF_1 = 0$ . Since  $dF_1 \neq 0$  (cf (2.6)), we must have  $c_1 = 0$ , and therefore  $\mathbf{c} = 0$ . Thus, (3.6) is 1-1, and

$$\dim V_* \leq \dim \mathcal{A}_* + \dim \bar{\mathcal{A}}_*. \quad (3.8)$$

To compute  $\dim \mathcal{A}_*$ , we note that  $\mathcal{A}_*$  is the direct sum of the spaces of antisymmetric linear maps on the eigenspaces of  $L_*$  (endowed with the standard inner product from  $\mathbb{R}^n$ ). In general, the space of antisymmetric linear maps on a  $k$ -dimensional inner product space is of dimension  $k(k-1)/2$ . Since  $L_*$  has  $\nu_*$  two-dimensional eigenspaces and  $(n - 2\nu_*)$  one-dimensional eigenspaces, it follows that  $\dim \mathcal{A}_* = \nu_*$ . Similarly,  $\dim \bar{\mathcal{A}}_* = \bar{\nu}_*$ . Substitution into (3.8) yields

$$\dim V_* \leq \nu_* + \bar{\nu}_*. \quad (3.9)$$

Next, we show that  $\nu_* + \bar{\nu}_* \leq \dim V_*$ . Let  $\mathcal{T}_*$  denote the set of polynomials which annihilate  $L_*$ . Elements of  $\mathcal{T}_*$  are divisible by the minimum polynomial of  $L_*$ , which we denote by  $P_*(x)$ .  $P_*(x)$  is of degree  $n - \nu_*$ . Regarding  $\mathcal{T}_*$  as a vector space, we let  $\mathcal{T}_*^n$  denote the subspace of polynomials of degree at most  $n$ , ie

$$\mathcal{T}_*^n = \{R(x) | R(L_*) = 0, \deg R \leq n\}. \quad (3.10)$$

Clearly  $\dim \mathcal{T}_*^n = \nu_*$  (elements of  $\mathcal{T}_*^n$  are products of  $P_*(x)$  with arbitrary polynomials of degree at most  $\nu_*$ ). Suppose  $\sum_{j=1}^n c_j x^{j-1} \in \mathcal{T}_*^n$ . Then

$$\sum_{j=1}^n c_j L_*^{j-1} = 0 \implies \text{Tr} \left( \sum_{j=1}^n c_j L_*^{j-1} dL_* \right) = 0 \implies \sum_{j=1}^n c_j dF_{j*} = 0. \quad (3.11)$$

Therefore,

$$\sum_{j=1}^n c_j x^{j-1} \mapsto \mathbf{c} \quad (3.12)$$

defines a linear map from  $\mathcal{T}_*^n$  to  $V_*$ . Clearly the map (3.12) is 1-1. Similarly, let  $\bar{\mathcal{T}}_*^n$  denote the  $\bar{\nu}_*$ -dimensional space of polynomials of degree at most  $n$  which annihilate  $\bar{L}_*$ , ie

$$\bar{\mathcal{T}}_*^n = \{\bar{R}(x) | \bar{R}(\bar{L}_*) = 0, \deg \bar{R} \leq n\}. \quad (3.13)$$

Arguing as above, we see that (3.12) also defines a 1-1 linear map from  $\bar{\mathcal{T}}_*^n$  to  $V_*$ .

Regarded as polynomial subspaces,  $\mathcal{T}_*^n$  and  $\bar{\mathcal{T}}_*^n$  are transverse. For if  $\sum_{j=1}^n c_j x^{j-1}$  belongs to both, then  $\sum_{j=1}^n c_j L_*^{j-1} = \sum_{j=1}^n c_j \bar{L}_*^{j-1} = 0$ . Corollary 2.1 implies that  $c_2 = \dots = c_n = 0$ , which in turn implies that  $c_1 = 0$ . Therefore, (3.12) defines a 1-1 map from  $\mathcal{T}_*^n \oplus \bar{\mathcal{T}}_*^n$  to  $V_*$ , and

$$\nu_* + \bar{\nu}_* = \dim(\mathcal{T}_*^n \oplus \bar{\mathcal{T}}_*^n) \leq \dim V_*, \quad (3.14)$$

as required.  $\square$

Since we actually have an equality in (3.14), we deduce the following:

**Corollary 3.1.** *Given  $\mathcal{T}_*^n$ ,  $\bar{\mathcal{T}}_*^n$  and  $V_*$  as above,*

$$\sum_{j=1}^n c_j x^{j-1} \mapsto \mathbf{c} \quad (3.15)$$

*is an isomorphism from  $\mathcal{T}_*^n \oplus \bar{\mathcal{T}}_*^n$  to  $V_*$ .*

## 4 Structure of singular sets

The Toda Hamiltonian has no corank- $n$  singularities, since  $dF_1 \neq 0$ . The singularities of corank  $(n - 1)$  are relative equilibria, as is shown in the following:

**Proposition 4.1.** *Let  $\Omega_{n-1} = \{(\mathbf{q}, \mathbf{p}) \mid q_1 = \cdots = q_n, p_1 = \cdots = p_n\}$  denote the set of points  $(\mathbf{q}, \mathbf{p})$  for which the components of  $\mathbf{q}$  are all the same and the components of  $\mathbf{p}$  are all the same. Then*

$$\Sigma_{n-1} = \Omega_{n-1}. \quad (4.1)$$

*Proof.* First we show that  $\Sigma_{n-1} \subset \Omega_{n-1}$ . Let  $(\mathbf{q}, \mathbf{p}) \in \Sigma_{n-1}$ . Since  $dF_1 \neq 0$ ,  $dH(\mathbf{q}, \mathbf{p})$  must be proportional to  $dF_1$ . But

$$dH = \sum_{j=1}^n p_j dp_j + \sum_{j=1}^n (b_j^2 - b_{j-1}^2) dq_j. \quad (4.2)$$

For  $dH$  to be proportional to  $dF_1$ , we must have that the  $p_j$ 's are all the same and the  $b_j$ 's (which are positive) are all the same. The latter implies that  $q_{j+1} - q_j$  is a constant independent of  $j$ , and periodicity, ie  $q_{n+1} = q_1$ , then implies that the  $q_j$ 's are all the same.

Next we show that  $\Omega_{n-1} \subset \Sigma_{n-1}$ . For  $(\mathbf{q}, \mathbf{p}) \in \Omega_{n-1}$ , the difference equation (2.10) simplifies to

$$v_{r-1} + v_{r+1} = (\lambda - p)v_r, \quad (4.3)$$

where  $p$  is the common value of the components of  $\mathbf{p}$ . Periodic and antiperiodic solutions of (4.3) are given by

$$u_{(r)j}^{\pm} = \exp(\pm \pi i j r / n), \quad \lambda_r = p + 2 \cos(\pi r / n), \quad 0 \leq r \leq n, \quad r \text{ even}, \quad (4.4)$$

$$\bar{u}_{(s)j}^{\pm} = \exp(\pm \pi i j s / n), \quad \bar{\lambda}_s = p + 2 \cos(\pi s / n), \quad 0 < s \leq n, \quad s \text{ odd}. \quad (4.5)$$

The  $\lambda_r$ 's are doubly degenerate except for  $r$  equal to 0 and (if  $n$  is even)  $n/2$ , while the  $\bar{\lambda}_s$ 's are all doubly degenerate. Thus,

$$\nu = [\frac{1}{2}(n - 1)], \quad \bar{\nu} = [\frac{1}{2}n], \quad (4.6)$$

where  $[x]$  denotes the integer part of  $x$ . In general,  $\nu + \bar{\nu} = n - 1$ , so Theorem 3.1 implies that  $(\mathbf{q}, \mathbf{p}) \in \Sigma_{n-1}$ .  $\square$

The explicit expressions (4.4) and (4.5) which hold in  $\Sigma_{n-1}$  allows us to deduce the following general result:

**Proposition 4.2.** *Let  $\lambda_1 \geq \dots \geq \lambda_n$  and  $\bar{\lambda}_1 \geq \dots \geq \bar{\lambda}_n$  denote the eigenvalues of  $L$  and  $\bar{L}$  in descending order. Then  $\lambda_r > \bar{\lambda}_r$  for  $r$  odd,  $\bar{\lambda}_r > \lambda_r$  for  $r$  even, and the allowed degeneracies are  $\lambda_r = \lambda_{r+1}$  for  $r$  even and  $\bar{\lambda}_r = \bar{\lambda}_{r+1}$  for  $r$  odd.*

*Proof.* From (4.4) and (4.5), these degeneracies are simultaneously realised at the points of  $\Sigma_{n-1}$ . Indeed, for points in  $\Sigma_{n-1}$ , we have that

$$\lambda_1 > \bar{\lambda}_1 = \bar{\lambda}_2 > \lambda_2 = \lambda_3 > \dots \bar{\lambda}_{2j-1} = \bar{\lambda}_{2j} > \lambda_{2j} = \lambda_{2j+1} > \dots \quad (4.7)$$

(the form of the end of the sequence depends on whether  $n$  is even or odd).

For any  $\mathbf{z} \in \mathbb{R}^{2n}$ , (2.19) along with Newton's identities imply that the characteristic polynomials of  $L$  and  $\bar{L}$  differ by a constant:

$$\det(x - L(\mathbf{z})) = \det(x - \bar{L}(\mathbf{z})) + 4. \quad (4.8)$$

Therefore, in general, eigenvalues of  $L$  and  $\bar{L}$  cannot coincide. Since the eigenvalues depend continuously on  $\mathbf{z}$ , it follows that the inequalities in (4.7) hold not only in  $\Sigma_{n-1}$  but everywhere else. Therefore, in general, we have that

$$\lambda_1 > \bar{\lambda}_1 \geq \bar{\lambda}_2 > \lambda_2 \geq \lambda_3 > \dots \bar{\lambda}_{2j-1} \geq \bar{\lambda}_{2j} > \lambda_{2j} \geq \bar{\lambda}_{2j+1} > \dots \quad (4.9)$$

It follows that  $\lambda_{2j} = \lambda_{2j+1}$  and  $\bar{\lambda}_{2j-1} = \bar{\lambda}_{2j}$  are the only possible degeneracies.  $\square$

To determine the local structure of the singular set  $\Sigma$ , it is convenient to bring the Lax matrices  $L$  and  $\bar{L}$  to a canonical form. As above, let  $\mathbf{z}_* \in \Sigma_k$ . Let  $L_*$  denote  $L(\mathbf{z}_*)$ , and let other functions evaluated at  $\mathbf{z}_*$  be similarly denoted. Let  $\nu_*$  and  $\bar{\nu}_*$  denote the number of (doubly) degenerate eigenvalues of  $L_*$  and  $\bar{L}_*$  respectively. From Theorem 3.1,  $\nu_* + \bar{\nu}_* = k$ . Let  $\lambda_{r*}$ ,  $1 \leq r \leq n$  denote the eigenvalues of  $L_*$  with degenerate eigenvalues repeated, ordered so that  $\lambda_{1*} = \lambda_{2*}, \dots, \lambda_{2\nu_*-1*} = \lambda_{2\nu_*}$ . Let  $\mathbf{u}_r$  denote an orthonormal set of eigenvectors of  $L_*$ . Define  $\bar{\lambda}_{s*}$  and  $\bar{\mathbf{u}}_s$  similarly with respect to  $\bar{L}_*$ . Then for  $\mathbf{z}$  in some neighbourhood of  $\mathbf{z}_*$ , there exists an orthogonal matrix  $R(\mathbf{z})$  depending smoothly on  $\mathbf{z}$ , with  $R_* = I$ , such that  $R^T(\mathbf{z})L(\mathbf{z})R(\mathbf{z})$  is block diagonal with respect to the  $\mathbf{u}_r$ -basis, with two-dimensional blocks for  $1 \leq r \leq 2\nu_*$  and diagonal for  $2\nu_* < r \leq n$ . That is, letting  $\Lambda(\mathbf{z})$  be the symmetric matrix with elements

$$\Lambda_{rt}(\mathbf{z}) = \mathbf{u}_r \cdot R^T(\mathbf{z})L(\mathbf{z})R(\mathbf{z}) \cdot \mathbf{u}_t, \quad (4.10)$$

we have that

$$\Lambda(\mathbf{z}) = \begin{pmatrix} * & * & & & & & \\ * & * & & & & & \\ & & \ddots & & & & \\ & & & * & * & & \\ & & & * & * & & \\ & & & & & * & \\ & & & & & & \ddots \\ & & & & & & & * \end{pmatrix}, \quad (4.11)$$

where omitted entries are zeros. At  $\mathbf{z}_*$ ,

$$\Lambda_* = \begin{pmatrix} \lambda_{1*} & 0 & & & & & \\ 0 & \lambda_{1*} & & & & & \\ & & \ddots & & & & \\ & & & \lambda_{r*} & 0 & & \\ & & & 0 & \lambda_{r*} & & \\ & & & & & \lambda_{r+1*} & \\ & & & & & & \ddots \\ & & & & & & & \lambda_{n-2\nu*} \end{pmatrix}. \quad (4.12)$$

Similarly, in some neighbourhood of  $\mathbf{z}_*$ , there exists an orthogonal matrix  $\bar{\mathbf{R}}(\mathbf{z})$  such that  $\bar{\mathbf{R}}^T(\mathbf{z})\bar{\mathbf{L}}(\mathbf{z})\bar{\mathbf{R}}(\mathbf{z})$  is block diagonal in the  $\bar{\mathbf{u}}_{s*}$  basis, with two-dimensional blocks for  $1 \leq s \leq 2\bar{\nu}_*$  and diagonal for  $2\nu_* < s \leq n$ . Define  $\bar{\Lambda}(\mathbf{z})$  in analogy with (4.10), ie

$$\bar{\Lambda}_{su}(\mathbf{z}) = \bar{\mathbf{u}}_s \cdot \bar{\mathbf{R}}^T(\mathbf{z})\bar{\mathbf{L}}(\mathbf{z})\bar{\mathbf{R}}(\mathbf{z}) \cdot \bar{\mathbf{u}}_u. \quad (4.13)$$

Let

$$\begin{aligned} \xi_r &= \frac{1}{2}(\Lambda_{2r,2r} - \Lambda_{2r-1,2r-1}), & \eta_r &= \Lambda_{2r-1,2r}, & 1 \leq r \leq \nu, \\ \bar{\xi}_s &= \frac{1}{2}(\bar{\Lambda}_{2s,2s} - \bar{\Lambda}_{2s-1,2s-1}), & \bar{\eta}_s &= \bar{\Lambda}_{2s-1,2s}, & 1 \leq s \leq \bar{\nu}. \end{aligned} \quad (4.14)$$

Then, from (4.10) and (4.13),  $\Sigma_k$  is given locally by the vanishing of  $\xi_r$ ,  $\eta_r$  and  $\bar{\xi}_s$ ,  $\bar{\eta}_s$ , eg

$$\xi_r = \eta_r = 0, \quad 1 \leq r \leq \nu_*, \quad \bar{\xi}_s = \bar{\eta}_s = 0, \quad 1 \leq s \leq \bar{\nu}_* \quad (4.15)$$

in some neighbourhood of  $\mathbf{z}_*$ .

The next result, Proposition 4.3, is a general expression for Poisson brackets of spectral components of the Lax matrices evaluated at  $\mathbf{z}_*$ . It is used in Proposition 4.4 to show that the Poisson brackets amongst  $\xi_r$ ,  $\eta_s$ ,  $\bar{\xi}_s$  and  $\bar{\eta}_s$

evaluated at  $\mathbf{z}_*$  are, up to normalisation, of canonical form. For the statement of Proposition 4.3, it is convenient to introduce Lax matrices in which the  $b_j$ 's are allowed to have arbitrary signs. Let  $\epsilon$  be an  $n$ -tuple of signs, and let  $L^\epsilon$  and  $M^\epsilon$  be the matrices obtained by replacing  $b_j$  with  $\epsilon_j b_j$  in the expressions (2.4) for  $L$  and  $M$ . Thus, for  $\epsilon = (1, \dots, 1)$ ,  $L^\epsilon = L$ , while for  $\epsilon = (1, \dots, 1, -1)$ ,  $L^\epsilon = \bar{L}$ . We regard  $\epsilon$  as  $n$ -periodic in its index.

**Proposition 4.3.** *Let  $\mathbf{u}$  and  $\mathbf{v}$  denote eigenvectors of  $L_*^\epsilon$  with the same eigenvalue  $\lambda$ , ie*

$$L_*^\epsilon \cdot \mathbf{u} = \lambda \mathbf{u}, \quad L_*^\epsilon \cdot \mathbf{v} = \lambda \mathbf{v}, \quad (4.16)$$

where  $\mathbf{u}$  and  $\mathbf{v}$  need not be linearly independent. Similarly, let  $\mathbf{w}$  and  $\mathbf{x}$  denote eigenvectors of  $L^\sigma$  with the same eigenvalue  $\mu$ ,

$$L_*^\sigma \cdot \mathbf{w} = \mu \mathbf{w}, \quad L_*^\sigma \cdot \mathbf{x} = \mu \mathbf{x}. \quad (4.17)$$

Let

$$\begin{aligned} L_{uv}^\epsilon &= \mathbf{u} \cdot L^\epsilon \cdot \mathbf{v}, \\ L_{wx}^\sigma &= \mathbf{w} \cdot L^\sigma \cdot \mathbf{x}. \end{aligned} \quad (4.18)$$

Then if  $\lambda = \mu$  and  $\prod_{m=1}^n \epsilon_m \sigma_m = 1$ ,

$$\{L_{uv}^\epsilon, L_{wx}^\sigma\}_* = \frac{1}{n} ((\mathbf{v} \cdot \mathbf{D} \cdot \mathbf{x})(\mathbf{u} \cdot M_*^\epsilon \mathbf{D} \cdot \mathbf{w}) + (\mathbf{u} \cdot \mathbf{D} \cdot \mathbf{w})(\mathbf{v} \cdot M_*^\epsilon \mathbf{D} \cdot \mathbf{x})), \quad (4.19)$$

where  $\mathbf{D}$  is the diagonal matrix with diagonal elements

$$d_m = \epsilon_1 \sigma_1 \cdots \epsilon_{m-1} \sigma_{m-1}. \quad (4.20)$$

Otherwise, ie if  $\lambda \neq \mu$  or  $\prod_{m=1}^n \epsilon_m \sigma_m = -1$ , then

$$\{L_{uv}^\epsilon, L_{wx}^\sigma\}_* = 0. \quad (4.21)$$

We note that  $\prod_{m=1}^n \epsilon_m \sigma_m = 1$  if and only if  $L^\epsilon$  and  $L^\sigma$  are conjugate. In fact, we shall only use the cases where  $L^\epsilon$  and  $L^\sigma$  are (independently) either  $L$  or  $\bar{L}$ .

*Proof.* Using (2.4) and (2.23), it is straightforward to show that

$$\begin{aligned} \{L_{uv}^\epsilon, L_{wx}^\sigma\} &= \frac{1}{2} \sum_{m=1}^n v_m x_m (A_m(\mathbf{u}, \mathbf{w}) + \epsilon_{m-1} \sigma_{m-1} A_{m-1}(\mathbf{u}, \mathbf{w})) + \\ &\quad \frac{1}{2} \sum_{m=1}^n u_m w_m (A_m(\mathbf{v}, \mathbf{x}) + \epsilon_{m-1} \sigma_{m-1} A_{m-1}(\mathbf{x}, \mathbf{v})), \end{aligned} \quad (4.22)$$

where

$$\begin{aligned} A_m(\mathbf{u}, \mathbf{w}) &= b_m(\epsilon_m u_{m+1} w_m - \sigma_m u_m w_{m+1}), \\ A_m(\mathbf{v}, \mathbf{x}) &= b_m(\epsilon_m v_{m+1} x_m - \sigma_m v_m x_{m+1}). \end{aligned} \quad (4.23)$$

Indeed, (4.22) holds independently of the eigenvector equations (4.16) and (4.17). The eigenvector equations imply additionally that, at  $\mathbf{z}_*$ ,  $A_{m*}(\mathbf{u}, \mathbf{w})$  and  $A_{m*}(\mathbf{v}, \mathbf{x})$  satisfy a Wronskian-like first-order difference equation. Indeed, (4.16) and (4.17) yield second-order difference equations for  $\mathbf{u}$  and  $\mathbf{w}$  (cf (2.10)),

$$(\epsilon_m b_m u_{m+1} + p_m u_m + \epsilon_{m-1} b_{m-1} u_{m-1})_* = \lambda u_m, \quad (4.24a)$$

$$(\sigma_m b_m w_{m+1} + p_m w_m + \sigma_{m-1} b_{m-1} w_{m-1})_* = \mu w_m. \quad (4.24b)$$

Multiplying (4.24a) by  $w_m$  and (4.24b) by  $u_m$ , and subtracting, we get

$$A_{m*}(\mathbf{u}, \mathbf{w}) - \epsilon_{m-1} \sigma_{m-1} A_{m-1*}(\mathbf{u}, \mathbf{w}) = (\lambda - \mu) u_m w_m. \quad (4.25a)$$

Similarly,

$$A_{m*}(\mathbf{v}, \mathbf{x}) - \epsilon_{m-1} \sigma_{m-1} A_{m-1*}(\mathbf{v}, \mathbf{x}) = (\lambda - \mu) v_m x_m. \quad (4.25b)$$

Suppose that  $\lambda \neq \mu$ . From (4.25), we have that

$$u_m w_m = \frac{1}{\Delta} (A_m(\mathbf{u}, \mathbf{w}) - \epsilon_{m-1} \sigma_{m-1} A_{m-1}(\mathbf{u}, \mathbf{w}))_*, \quad (4.26a)$$

$$v_m x_m = \frac{1}{\Delta} (A_m(\mathbf{v}, \mathbf{x}) - \epsilon_{m-1} \sigma_{m-1} A_{m-1}(\mathbf{v}, \mathbf{x}))_*, \quad (4.26b)$$

where  $\Delta = \lambda - \mu$ . Substituting into (4.22), we get that

$$\begin{aligned} \{L_{uv}^\epsilon, L_{wx}^\sigma\}_* &= \frac{1}{2\Delta} \sum_{m=1}^n \left[ \right. \\ & (A_m(\mathbf{v}, \mathbf{x}) - \epsilon_{m-1} \sigma_{m-1} A_{m-1}(\mathbf{v}, \mathbf{x})) \times (A_m(\mathbf{u}, \mathbf{w}) + \epsilon_{m-1} \sigma_{m-1} A_{m-1}(\mathbf{u}, \mathbf{w})) + \\ & \left. (A_m(\mathbf{u}, \mathbf{w}) - \epsilon_{m-1} \sigma_{m-1} A_{m-1}(\mathbf{u}, \mathbf{w})) \times (A_m(\mathbf{v}, \mathbf{x}) + \epsilon_{m-1} \sigma_{m-1} A_{m-1}(\mathbf{v}, \mathbf{x})) \right]_* \\ &= \frac{1}{\Delta} \sum_{m=1}^n \left[ A_m(\mathbf{u}, \mathbf{w}) A_m(\mathbf{v}, \mathbf{x}) - A_{m-1}(\mathbf{u}, \mathbf{w}) A_{m-1}(\mathbf{v}, \mathbf{x}) \right]_* = 0, \end{aligned} \quad (4.27)$$

as required by (4.21).

Next, suppose that  $\lambda = \mu$ . From (4.25),

$$A_{m*}(\mathbf{u}, \mathbf{w}) = \epsilon_{m-1} \sigma_{m-1} A_{m-1*}(\mathbf{u}, \mathbf{w}), \quad (4.28a)$$

$$A_{m*}(\mathbf{v}, \mathbf{x}) = \epsilon_{m-1} \sigma_{m-1} A_{m-1*}(\mathbf{v}, \mathbf{x}). \quad (4.28b)$$

Substituting into (4.22), we get that

$$\{L_{uv}^\epsilon, L_{wx}^\sigma\}_* = \sum_{m=1}^n v_m x_m A_{m*}(\mathbf{u}, \mathbf{w}) + \sum_{m=1}^n u_m w_m A_{m*}(\mathbf{v}, \mathbf{x}). \quad (4.29)$$

Iterating the relations (4.28)  $n$  times, we get that  $A_{m*}(\mathbf{u}, \mathbf{w}) = \epsilon_1 \sigma_1 \cdots \epsilon_n \sigma_n A_{m*}(\mathbf{u}, \mathbf{w})$  and similarly for  $A_{m*}(\mathbf{v}, \mathbf{x})$ . Therefore, if  $\epsilon_1 \sigma_1 \cdots \epsilon_n \sigma_n = -1$ ,  $A_{m*}(\mathbf{u}, \mathbf{w}) = A_{m*}(\mathbf{v}, \mathbf{x}) = 0$ , so that  $\{L_{uv}^\epsilon, L_{wx}^\sigma\}_* = 0$ , as in (4.21).

Now suppose that  $\epsilon_1 \sigma_1 \cdots \epsilon_n \sigma_n = 1$ . In this case, (4.28a) implies that

$$d_m A_{m*}(\mathbf{u}, \mathbf{w}) = d_{m-1} A_{m-1*}(\mathbf{u}, \mathbf{w}), \quad (4.30)$$

where  $d_m$  is given by (4.20), ie,  $d_m A_{m*}(\mathbf{u}, \mathbf{w})$  is independent of  $m$ . Thus, the first term in (4.29) can be expressed as

$$\begin{aligned} \sum_{m=1}^n v_m x_m A_{m*}(\mathbf{u}, \mathbf{w}) &= \left( \sum_{m=1}^n d_m v_m x_m \right) \left( \frac{1}{n} \sum_{m=1}^n d_m A_{m*}(\mathbf{u}, \mathbf{w}) \right) \\ &= (\mathbf{v} \cdot \mathbf{D} \cdot \mathbf{x}) \left( \frac{1}{n} \sum_{m=1}^n d_m A_{m*}(\mathbf{u}, \mathbf{w}) \right). \end{aligned} \quad (4.31)$$

The sum in the last term in (4.31) can be expressed as

$$\begin{aligned} \sum_{m=1}^n d_m A_{m*}(\mathbf{u}, \mathbf{w}) &= \sum_{m=1}^n d_m b_{m*} (\epsilon_m u_{m+1} w_m - \sigma_m u_m w_{m+1}) = \\ &= \sum_{m=1}^n \epsilon_m b_{m*} (u_{m+1} (\mathbf{D} \cdot \mathbf{w})_m - u_m (\mathbf{D} \cdot \mathbf{w})_{m+1}) = \mathbf{u} \cdot \mathbf{M}_*^\epsilon \mathbf{D} \cdot \mathbf{w}. \end{aligned} \quad (4.32)$$

Substituting (4.31) and (4.32) into the first term of (4.29) and making analogous substitutions for the second term, we get

$$\{L_{uv}^\epsilon, L_{wx}^\sigma\}_* = \frac{1}{n} (\mathbf{v} \cdot \mathbf{D} \cdot \mathbf{x}) (\mathbf{u} \cdot \mathbf{M}_*^\epsilon \mathbf{D} \cdot \mathbf{w}) + \frac{1}{n} (\mathbf{u} \cdot \mathbf{D} \cdot \mathbf{w}) (\mathbf{v} \cdot \mathbf{M}_*^\sigma \mathbf{D} \cdot \mathbf{x}), \quad (4.33)$$

as in (4.19).  $\square$

**Proposition 4.4.** *Let  $\xi_r, \eta_r, \bar{\xi}_s$  and  $\bar{\eta}_s$  be given by (4.14). Then*

$$\{\xi_r, \bar{\xi}_s\}_* = \{\xi_r, \bar{\eta}_s\}_* = \{\eta_r, \bar{\xi}_s\}_* = \{\eta_r, \bar{\eta}_s\}_* = 0, \quad (4.34a)$$

$$\{\xi_r, \xi_t\}_* = \{\eta_r, \eta_t\}_* = \{\bar{\xi}_s, \bar{\xi}_u\}_* = \{\bar{\eta}_s, \bar{\eta}_u\}_* = 0, \quad (4.34b)$$

$$\{\xi_r, \eta_t\}_* = \frac{1}{n} \mathbf{u}_{2r} \cdot \mathbf{M}_* \cdot \mathbf{u}_{2r-1} \delta_{rt}, \quad \{\bar{\xi}_s, \bar{\eta}_u\}_* = \frac{1}{n} \bar{\mathbf{u}}_{2s} \cdot \bar{\mathbf{M}}_* \cdot \bar{\mathbf{u}}_{2s-1} \delta_{su}. \quad (4.34c)$$

Moreover,  $\mathbf{u}_{2r} \cdot \mathbf{M}_* \cdot \mathbf{u}_{2r-1}$  and  $\bar{\mathbf{u}}_{2s} \cdot \bar{\mathbf{M}}_* \cdot \bar{\mathbf{u}}_{2s-1}$  do not vanish.

*Proof.* We treat a representative case for each set of relations in (4.34). For (4.34a), we consider  $\{\eta_r, \bar{\xi}_s\}_*$ . From (4.14) and (4.10),

$$\begin{aligned} d\eta_{r*} &= d(\mathbf{u}_{2r-1} \cdot \mathbf{R}^T \mathbf{L} \mathbf{R} \cdot \mathbf{u}_{2r})_* = \mathbf{u}_{2r-1} \cdot [\mathbf{L}, d\mathbf{R}]_* \cdot \mathbf{u}_{2r} + \mathbf{u}_{2r-1} \cdot d\mathbf{L}_* \cdot \mathbf{u}_{2r} \\ &= \mathbf{u}_{2r-1} \cdot d\mathbf{L}_* \cdot \mathbf{u}_{2r}, \end{aligned} \quad (4.35)$$

where we have used the fact that  $\mathbf{R}_* = \mathbf{I}$ ,  $d\mathbf{R}_*$  is antisymmetric (since  $\mathbf{R}$  is orthogonal) and, in the last equality, that  $\mathbf{u}_{2r-1}$  and  $\mathbf{u}_r$  are eigenvectors of  $\mathbf{L}_*$  with the same eigenvalue. Similarly, from (4.14) and (4.13),

$$d\bar{\xi}_{s*} = \frac{1}{2} \bar{\mathbf{u}}_{2s} \cdot d\bar{\mathbf{L}}_* \cdot \bar{\mathbf{u}}_{2s} - \frac{1}{2} \bar{\mathbf{u}}_{2s-1} \cdot d\bar{\mathbf{L}}_* \cdot \bar{\mathbf{u}}_{2s-1}. \quad (4.36)$$

Together, (4.35) and (4.36) imply that

$$\{\eta_r, \bar{\xi}_s\}_* = \{\mathbf{u}_{2r-1} \cdot \mathbf{L} \cdot \mathbf{u}_{2r}, \frac{1}{2} \bar{\mathbf{u}}_{2s} \cdot \bar{\mathbf{L}} \cdot \bar{\mathbf{u}}_{2s} - \frac{1}{2} \bar{\mathbf{u}}_{2s-1} \cdot \bar{\mathbf{L}} \cdot \bar{\mathbf{u}}_{2s-1}\}_*. \quad (4.37)$$

Proposition 4.3 then implies, with  $\epsilon = (1, \dots, 1)$  and  $\sigma = (1, \dots, -1)$ , that

$$\{\eta_r, \bar{\xi}_s\}_* = 0, \quad (4.38)$$

since  $\prod_{m=1}^n \epsilon_m \sigma_m = -1$ . The remaining relations in (4.34a) are obtained similarly.

For (4.34b), we consider  $\{\eta_r, \eta_t\}_*$ . If  $r = t$  the bracket obviously vanishes, so we take  $r \neq t$ . From (4.35) it follows that

$$\{\eta_r, \eta_t\}_* = \{\mathbf{u}_{2r-1} \cdot \mathbf{L} \cdot \mathbf{u}_{2r}, \mathbf{u}_{2t-1} \cdot \mathbf{L} \cdot \mathbf{u}_{2t}\}_*. \quad (4.39)$$

Since  $\lambda_r \neq \lambda_t$ , Proposition 4.3 implies that  $\{\eta_r, \eta_t\}_* = 0$ . The remaining relations in (4.34b) are obtained similarly.

For (4.34c), we consider  $\{\xi_r, \eta_t\}_*$ . From (4.35) and (4.36) (or rather, its analog for  $d\xi_r$ ), we get that

$$\{\xi_r, \eta_t\}_* = \frac{1}{2} \{\mathbf{u}_{2r} \cdot \mathbf{L} \cdot \mathbf{u}_{2r}, \mathbf{u}_{2t-1} \cdot \mathbf{L} \cdot \mathbf{u}_{2t}\}_* - \frac{1}{2} \{\mathbf{u}_{2r-1} \cdot \mathbf{L} \cdot \mathbf{u}_{2r-1}, \mathbf{u}_{2t-1} \cdot \mathbf{L} \cdot \mathbf{u}_{2t}\}_*. \quad (4.40)$$

Proposition 4.3 then implies, with  $\epsilon = \sigma = (1, \dots, 1)$ , that (4.40) vanishes if  $r \neq t$  (as, in this case,  $\lambda_r \neq \lambda_t$ ). On the other hand, for  $r = t$ , Proposition 4.3, with  $\mathbf{D} = \mathbf{I}$ , and  $\mathbf{u}_{2r-1}, \mathbf{u}_{2r}$  orthonormal gives that

$$\{\xi_r, \eta_r\}_* = \frac{1}{2n} \mathbf{u}_{2r} \cdot \mathbf{M}_* \cdot \mathbf{u}_{2r-1} - \frac{1}{2n} \mathbf{u}_{2r-1} \cdot \mathbf{M}_* \cdot \mathbf{u}_{2r} = \frac{1}{n} \mathbf{u}_{2r-1} \cdot \mathbf{M}_* \cdot \mathbf{u}_{2r}. \quad (4.41)$$

The quantity  $\mathbf{u}_{2r} \cdot \mathbf{M}_* \cdot \mathbf{u}_{2r-1}$  which appears in (4.41) does not vanish. This is because, from (4.30) and (4.32) (with  $d_m = 1$ ), we have that  $\mathbf{u}_{2r} \cdot \mathbf{M}_* \cdot \mathbf{u}_{2r-1}$  is equal to  $nb_m(u_{2r-1, m+1} u_{2r, m} - u_{2r-1, m} u_{2r, m+1})$  for all  $m$ .  $\mathbf{u}_{2r} \cdot \mathbf{M}_* \cdot \mathbf{u}_{2r-1} = 0$  would imply that  $\mathbf{u}_{2r-1}$  and  $\mathbf{u}_{2r}$  are proportional, contradicting orthonormality. The result for  $\{\bar{\xi}_s, \bar{\eta}_u\}_*$  follows similarly.  $\square$

**Theorem 4.1.**  $\Sigma_k$  is a codimension- $2k$  symplectic submanifold. It consists of  $\binom{n-1}{k}$  components which are disconnected from each other.  $\Sigma_k$  is contained in the closure of  $\Sigma_j$  for all  $j < k$ .

*Proof.* Let  $\mathbf{z}_* \in \Sigma_k$  and let  $\nu_*, \bar{\nu}_*$  denote the number of degenerate eigenvalues of  $L_*$  and  $\bar{L}_*$ , respectively, so that, by Theorem 3.1,  $\nu_* + \bar{\nu}_* = k$ . Let  $\xi_r, \eta_r$ ,  $1 \leq r \leq \nu_*$ , and  $\bar{\xi}_s, \bar{\eta}_s$ ,  $1 \leq s \leq \bar{\nu}_*$ , be given by (4.14) in some neighbourhood of  $\mathbf{z}_*$ . In this neighbourhood,  $\Sigma_k$  is given by

$$\xi_r = \eta_r = \bar{\xi}_s = \bar{\eta}_s = 0. \quad (4.42)$$

Proposition 4.4 implies that the derivatives of  $\xi_r, \eta_r, \bar{\xi}_s$  and  $\bar{\eta}_s$  are linearly independent at  $\mathbf{z}_*$ . It follows from the implicit function theorem that  $\Sigma_k$  is codimension- $2k$  submanifold.

Next, we show that the tangent space of  $\Sigma_k \subset \mathbb{R}^{2n}$  at  $\mathbf{z}_*$  is symplectic. Let  $\mathbf{X}_* \in T_{\mathbf{z}_*}\Sigma_k$ .  $\mathbf{X}_*$  may be extended to a vector field  $\mathbf{X}$  defined in a neighbourhood of  $\mathbf{z}_*$  whose restriction to  $\Sigma_k$  is tangent to  $\Sigma_k$ . Then (4.42) implies, for example, that  $(\mathbf{X} \cdot d\xi_r)_* = 0$ . But  $(\mathbf{X} \cdot d\xi_r)_*$  is just the symplectic inner product of  $\mathbf{X}_*$  with the Hamiltonian vector field generated by  $\xi_r$  at  $\mathbf{z}_*$ . Let  $E_*$  denote the subspace of  $T_*\mathbb{R}^{2n}$  spanned by the Hamiltonian vector fields generated by  $\xi_r, \eta_r$ , and  $\bar{\xi}_s, \bar{\eta}_s$ . Arguing as above, we may conclude that  $\mathbf{X}_*$ , and therefore  $T_*\Sigma_k$ , is skew-orthogonal to  $E_*$ . Proposition 4.4 implies that  $E_*$  is symplectic of dimension  $2k$ . Since  $\text{codim } T_{\mathbf{z}_*}\Sigma_k = 2k$ , it follows that  $T_{\mathbf{z}_*}\Sigma_k$  is the skew-orthogonal complement of  $E_*$ . The skew-orthogonal complement of a symplectic subspace is itself symplectic, so  $T_{\mathbf{z}_*}\Sigma_k$  is symplectic, and therefore  $\Sigma_k$  is a symplectic submanifold.

$\Sigma_k$  can be partitioned into distinct components according to the particular  $k$  pairs of eigenvalues which are degenerate. From Proposition 4.2 there are  $(n-1)$  possible degeneracies ( $[\frac{1}{2}(n-1)]$  from  $L$  and  $[\frac{1}{2}n]$  from  $\bar{L}$ ), so the maximum number of components is  $\binom{n-1}{k}$ . Every choice of  $k$  pairs can be realised by setting, in a neighbourhood of a point of  $\Sigma_{n-1}$ , the appropriate pairs of coordinates  $(\xi_r, \eta_r), (\bar{\xi}_s, \bar{\eta}_s)$  to be nonzero, and the remaining pairs to be zero. Thus  $\Sigma_k$  has precisely  $\binom{n-1}{k}$  components. Along any path connecting points in different components, the number of eigenvalue degeneracies must change, so the path must leave  $\Sigma_k$ . Therefore, the components are disconnected from each other.

Any neighbourhood of  $\mathbf{z}_* \in \Sigma_k$  contains points of  $\Sigma_{j < k}$ . Such points are obtained by setting  $k-j$  of the  $k$  coordinate pairs  $(\xi_r, \eta_r)$  and  $(\bar{\xi}_s, \bar{\eta}_s)$  to be nonzero. Therefore,  $\Sigma_k$  is contained in the closure of  $\Sigma_{j < k}$ .  $\square$

We consider next the transverse stability of  $\Sigma_k$ . Let  $\mathbf{z}_* \in \Sigma_k$ . In analogy

with (4.14), we define the local functions

$$\begin{aligned}\tau_r &= \frac{1}{2}(\Lambda_{2r,2r} + \Lambda_{2r-1,2r-1}), & 1 \leq r \leq \nu, \\ \bar{\tau}_s &= \frac{1}{2}(\bar{\Lambda}_{2s,2s} + \bar{\Lambda}_{2s-1,2s-1}), & 1 \leq s \leq \bar{\nu},\end{aligned}\quad (4.43)$$

with

$$\begin{aligned}\tau_{r*} &= \frac{1}{2}\mathbf{u}_{2r} \cdot d\mathbf{L}_* \cdot \mathbf{u}_{2r} + \frac{1}{2}\mathbf{u}_{2r-1} \cdot d\mathbf{L}_* \cdot \mathbf{u}_{2r-1}, & 1 \leq r \leq \nu, \\ \bar{\tau}_{s*} &= \frac{1}{2}\bar{\mathbf{u}}_{2s} \cdot d\bar{\mathbf{L}}_* \cdot \bar{\mathbf{u}}_{2s} + \frac{1}{2}\bar{\mathbf{u}}_{2s-1} \cdot d\bar{\mathbf{L}}_* \cdot \bar{\mathbf{u}}_{2s-1}, & 1 \leq s \leq \bar{\nu}.\end{aligned}\quad (4.44)$$

With arguments similar to those of Proposition 4.4, one can show that the Poisson brackets of the  $\tau_r$ 's and the  $\bar{\tau}_s$ 's with  $\xi_r$ ,  $\eta_r$ ,  $\bar{\xi}_s$  and  $\bar{\eta}_s$  all vanish at  $\mathbf{z}_*$ . Likewise, the Poisson brackets of the  $\tau_r$ 's and  $\bar{\tau}_s$ 's amongst themselves vanish at  $\mathbf{z}_*$ . We record this briefly as

$$\{\tau, \chi\}_* = \{\tau, \tau.\}_* = 0, \quad (4.45)$$

where  $\chi.$  denotes  $\xi_r$ ,  $\eta_r$ ,  $\bar{\xi}_s$  or  $\bar{\eta}_s$ , and  $\tau, \tau.$  denotes  $\tau_r$  or  $\bar{\tau}_s$ .

Let

$$\begin{aligned}T_r(x) &= \frac{\det(\mathbf{L}_* - x\mathbf{I})}{\lambda_{r*} - x} = \sum_{j=1}^n c_{rj}x^{j-1}, & 1 \leq r \leq \nu, \\ \bar{T}_{s*}(x) &= \frac{\det(\bar{\mathbf{L}}_* - x\mathbf{I})}{\bar{\lambda}_{s*} - x} = \sum_{j=1}^n \bar{c}_{sj}x^{j-1}, & 1 \leq s \leq \bar{\nu}.\end{aligned}\quad (4.46)$$

Clearly  $T_r(\mathbf{L}_*) = 0$ , and moreover, the  $T_r$ 's are linearly independent (since, if  $\sum_{t=1}^{\nu} a_t T_t(\lambda_{r*}) = 0$ , then  $\sum_{t=1}^{\nu} a_t T'_t(\lambda_{r*}) = 0$ , and the fact that  $T'_t(\lambda_{r*}) = 0$  for  $r \neq t$  implies that  $a_r = 0$  for each  $r$ ). Therefore, the  $T_r$ 's constitute a basis for  $\mathcal{T}_*^n$ , the space of polynomials of degree at most  $n$  that annihilate  $\mathbf{L}_*$  (cf (3.10)). Likewise, the  $\bar{T}_s$ 's constitute a basis for  $\bar{\mathcal{T}}_*^n$ , the space of polynomials of degree at most  $n$  that annihilate  $\bar{\mathbf{L}}_*$  (cf (3.13)). It follows from Corollary 3.1 that the vectors  $\mathbf{c}_r = (c_{r1}, \dots, c_{rn})$  and  $\bar{\mathbf{c}}_s = (\bar{c}_{s1}, \dots, \bar{c}_{sn})$  constitute a basis for  $V_*$ , the space of linear relations amongst the  $dF_{j*}$ 's. Therefore, letting

$$G_r = \sum_{j=1}^n c_{rj}F_j, \quad \bar{G}_s = \sum_{j=1}^n \bar{c}_{sj}F_j, \quad (4.47)$$

we have that the Hamiltonian flows generated by  $G_r$  and  $\bar{G}_s$  have  $\mathbf{z}_*$  as a fixed point. The stability of these flows at  $\mathbf{z}_*$  is given by the following:

**Theorem 4.2.**

$$\begin{aligned} G''_{r*} &= 2T'_r(\lambda_{r*}) (d\xi_r \otimes d\xi_r + d\eta_r \otimes d\eta_r + d\tau_r \otimes d\tau_r)_*, \\ \overline{G}''_{s*} &= 2\overline{T}'_s(\overline{\lambda}_{s*}) (d\overline{\xi}_s \otimes d\overline{\xi}_s + d\overline{\eta}_s \otimes d\overline{\eta}_s + d\overline{\tau}_s \otimes d\overline{\tau}_s)_*. \end{aligned} \quad (4.48)$$

Thus, the linearised  $G_r$ -flow about  $\mathbf{z}_*$  produces elliptic oscillations in the  $(\xi_r, \eta_r)$  plane with frequencies

$$\omega_r = 2nT'(\lambda_{r*})/(\mathbf{u}_{2r} \cdot \mathbf{M}_* \cdot \mathbf{u}_{2r-1}). \quad (4.49)$$

Similarly, the linearised  $\overline{G}_s$ -flow about  $\mathbf{z}_*$  produces elliptic oscillations in the  $(\overline{\xi}_s, \overline{\eta}_s)$  plane with frequencies

$$\overline{\omega}_s = 2n\overline{T}'(\overline{\lambda}_{s*})/(\overline{\mathbf{u}}_{2s} \cdot \overline{\mathbf{M}}_* \cdot \overline{\mathbf{u}}_{2s-1}). \quad (4.50)$$

*Proof.* We carry out the calculations for  $G_r$ ; those for  $\overline{G}_s$  are similar. From (4.47) and (2.5), we have that

$$G''_{r*} = \sum_{j=1}^n \text{Tr}(c_{jr} L_*^{j-1} L_*'') + \sum_{j=1}^n c_{jr} \text{Tr}(dL^{j-1} \otimes dL)_*. \quad (4.51)$$

As  $T_r(L_*) = 0$ , the first term vanishes. As for the second term,

$$\begin{aligned} \sum_{j=1}^n c_{jr} \text{Tr}(dL^{j-1} \otimes dL)_* &= \sum_{j=1}^n \sum_{k=0}^{j-2} c_{jr} \text{Tr}(L_*^k dL_* L_*^{j-k-2} \otimes dL_*) \\ &= \sum_{a=1}^{n-\nu} \sum_{b=1}^{n-\nu} \sum_{j=1}^n \sum_{k=0}^{j-2} c_{jr} \lambda_{a*}^k \lambda_{b*}^{j-k-2} \text{Tr}(\rho_{a*} dL_* \otimes \rho_{b*} dL_*), \end{aligned} \quad (4.52)$$

where we have introduced the spectral resolution of  $L_*$ ,

$$L_* = \sum_{a=1}^{n-\nu} \lambda_{a*} \rho_{a*}, \quad \rho_{a*} = \frac{\prod_{r \neq a} (L_* - \lambda_{r*})}{\prod_{r \neq a} (\lambda_{r*} - \lambda_{a*})}. \quad (4.53)$$

$\rho_{a*}$  is the symmetric projector onto the  $\lambda_{a*}$ -eigenspace of  $L_{a*}$ , so that  $\rho_{a*} \rho_{b*} = \delta_{ab} \rho_{a*}$ . The sums over  $j$  and  $k$  in (4.52) are readily performed to give

$$\sum_{j=1}^n \sum_{k=0}^{j-2} c_{jr} \lambda_{a*}^k \lambda_{b*}^{j-k-2} = \begin{cases} (T_r(\lambda_{a*}) - T_r(\lambda_{b*})) / (\lambda_{a*} - \lambda_{b*}), & a \neq b, \\ T'_r(\lambda_{a*}), & a = b. \end{cases} \quad (4.54)$$

We have that  $T_r(\lambda_{a*}) = 0$  for all  $a$ , while  $T'_r(\lambda_{a*})$  vanishes unless  $a = r$ . Thus, the sums over  $a$  and  $b$  in (4.52) collapse to the single term  $a = b = r$ . Substituting into (4.51), we get

$$G''_{r*} = T'_r(\lambda_{r*}) \text{Tr}(\rho_{r*} dL_* \otimes \rho_{r*} dL_*). \quad (4.55)$$

The projector  $\rho_{r*}$  may be written as the diadic  $\mathbf{u}_{2r-1} \mathbf{u}_{2r-1} + \mathbf{u}_{2r} \mathbf{u}_{2r}$ . Substituting into (4.55), we get

$$\begin{aligned} G''_{r*} = T'_r(\lambda_{r*}) [ & (\mathbf{u}_{2r} \cdot dL_* \cdot \mathbf{u}_{2r}) \otimes (\mathbf{u}_{2r} \cdot dL_* \cdot \mathbf{u}_{2r}) \\ & + 2(\mathbf{u}_{2r-1} \cdot dL_* \cdot \mathbf{u}_{2r}) \otimes (\mathbf{u}_{2r-1} \cdot dL_* \cdot \mathbf{u}_{2r}) \\ & + (\mathbf{u}_{2r-1} \cdot dL_* \cdot \mathbf{u}_{2r-1}) \otimes (\mathbf{u}_{2r-1} \cdot dL_* \cdot \mathbf{u}_{2r-1})]. \end{aligned} \quad (4.56)$$

Since (cf (4.35), (4.36), (4.44))

$$\begin{aligned} d\xi_{r*} + d\tau_{r*} &= \mathbf{u}_{2r} \cdot dL_* \cdot \mathbf{u}_{2r}, \\ d\xi_{r*} - d\tau_{r*} &= \mathbf{u}_{2r-1} \cdot dL_* \cdot \mathbf{u}_{2r-1}, \\ d\eta_{r*} &= \mathbf{u}_{2r-1} \cdot dL_* \cdot \mathbf{u}_{2r}, \end{aligned} \quad (4.57)$$

(4.56) yields the required result (4.48).

(4.48), along with the Poisson bracket relations of Proposition 4.4 and those recorded in (4.45), imply that the equations of motion for the linearised  $G_r$ -flow about  $\mathbf{z}_*$  transverse to  $\Sigma_k$  are given by

$$\dot{\xi}_r = \omega_r \eta_r, \quad \dot{\eta}_r = -\omega_r \xi_r, \quad (4.58)$$

while the remaining transverse coordinates  $\xi_{u \neq r}, \eta_{u \neq r}, \bar{\xi}_s, \bar{\eta}_s$  are fixed. Thus, the linearised transverse  $G_r$ -flow is elliptic in the  $(\xi_r, \eta_r)$ -plane with frequency  $\omega_r$ .  $\square$

## 5 Maslov indices and eigenvector monodromies.

Consider an integrable system in  $\mathbb{R}^{2n}$  with integrals of motion  $F : \mathbb{R}^{2n} \rightarrow \mathbb{R}^n$ . In the set of regular points  $R$  of  $F$ , the Hamiltonian vector fields generated by the  $F_j$ 's span a  $n$ -dimensional Lagrangian plane, which we denote  $\lambda(\mathbf{z})$ . As shown by Arnold [4], the space of  $n$ -dimensional Lagrangian planes,  $\Lambda(n)$ , has fundamental group  $\pi_1(\Lambda(n)) = \mathbb{Z}$ , and the Maslov index of a continuous, oriented closed curve  $C$  in  $R$  is the degree (winding number) of  $\lambda(C)$  in  $\Lambda(n)$ ,

$$\mu(C) = \text{wn } \lambda(C). \quad (5.1)$$

For angle contours  $C_j$  on an invariant torus, the Maslov index appears in the semiclassical EBK quantisation conditions for the associated action variables [21, 20],

$$I_j = (n_j + \frac{1}{4}\mu_j)\hbar. \quad (5.2)$$

In this case  $\mu(C)$  is always even (this is because the distribution of Lagrangian planes  $\lambda(\mathbf{z})$  over  $R$  is orientable).

In a companion paper [16], we show that under certain genericity conditions, the Maslov index of  $C$  can be expressed as a sum of contributions from the nondegenerate corank-one singularities it encloses. We briefly summarise the results. Let  $\Sigma$  denote the critical set of  $F$ , and let

$$\Sigma_k = \{\mathbf{z} \in \mathbb{R}^{2n} \mid \text{corank } dF(\mathbf{z}) = k\} \quad (5.3)$$

denote the set of critical points of  $F$  of corank  $k$ . Given  $\mathbf{z}_* \in \Sigma_1$ , let  $\mathbf{c} \in \mathbb{R}^n$  be a nontrivial solution of  $\sum_{j=1}^n c_j dF_{j*} = 0$  (here  $dF_* = dF(\mathbf{z}_*)$ ). Let  $K_* = \mathbf{J} \left( \sum_{j=1}^n c_j F''_{j*} \right)$ . We say that  $\mathbf{z}_*$  is *nondegenerate* if  $\text{Tr } K_*^2 \neq 0$ . Let  $\Delta$  denote the set of nondegenerate points in  $\Sigma_1$ ,

$$\Delta = \{\mathbf{z}_* \in \Sigma_1 \mid \text{Tr } K_*^2 \neq 0\}. \quad (5.4)$$

In general,  $\Delta$  is a codimension-2 symplectic submanifold. Let  $S : D^2 \rightarrow \mathbb{R}^{2n}$  denote a map of the oriented unit two-disk  $D^2$  into  $\mathbb{R}^{2n}$  smooth on the interior of  $D^2$  such that  $S|_{\partial D^2} = C$ . We assume that  $S$  is *transverse* to  $\Sigma$ . By this we mean that i) the only critical points of  $dF$  contained in the image of  $S$  belong to  $\Delta$ , ii)  $S^{-1}(\Delta)$  consists of a finite set of points  $e_1, \dots, e_r$  in the interior of  $D^2$ , and iii)  $dS(e_j)$  has full rank. (If  $\Sigma - \Sigma_1$  is composed of submanifolds of codimension three or more, such an  $S$  can always be found.) Let  $\mathbf{z}_j = S(e_j)$  denote the critical points of  $F$  in the image of  $S$ . Let  $E_{\mathbf{z}_j}$  denote the skew-orthogonal complement of  $T_{\mathbf{z}_j}\Delta$ .  $E_{\mathbf{z}_j}$  is a two-dimensional symplectic plane. Let  $P_j : T_{\mathbf{z}_j}\mathbb{R}^{2n} \rightarrow E_{\mathbf{z}_j}$  denote the projection onto  $E_{\mathbf{z}_j}$  with respect to the decomposition

$$T_{\mathbf{z}_j}\mathbb{R}^{2n} = T_{\mathbf{z}_j}\Delta \oplus E_{\mathbf{z}_j}. \quad (5.5)$$

Then the map  $P_j \circ dS : T_{e_j}D^2 \rightarrow E_{\mathbf{z}_j}$  is nonsingular. Let  $\sigma_j = \pm 1$  according to whether this map is orientation-preserving or reversing (the orientation on  $E_{\mathbf{z}_j}$  is given by the symplectic form). Then

$$\mu(C) = 2 \sum_j \sigma_j \cdot \text{sgn } \text{Tr } K_{\mathbf{z}_j}^2. \quad (5.6)$$

For the periodic Toda chain, all critical points are nondegenerate of elliptic type (cf Theorem 4.2), so that  $\text{Tr } K_{\mathbf{z}_j}^2 < 0$ . Also,  $\Sigma - \Sigma_1 = \cup_{k>1} \Sigma_k$

is composed of submanifolds of codimension four or more (Theorem 4.1). Thus, for  $C$  a continuous closed curve in the regular component of the Toda integrals of motion, we have that

$$\mu(C) = -2 \sum_j \sigma_j. \quad (5.7)$$

We can also associate to  $C$  the signs acquired by the normalised eigenvectors of  $L$  and  $\bar{L}$  on continuation around  $C$ . Explicitly, let the eigenvalues of  $L$  and  $\bar{L}$  be indexed in increasing order, so that  $\lambda_1 < \dots < \lambda_n$  and  $\bar{\lambda}_1 < \dots < \bar{\lambda}_n$  in  $R$ . Let  $\mathbf{z}(t)$ ,  $0 \leq t \leq 1$  denote a parameterisation of  $C$ , and let  $\mathbf{u}_r(t)$  denote a continuously varying, normalised real eigenvector of  $L(\mathbf{z}(t))$  with eigenvalue  $\lambda_r(\mathbf{z}(t))$ . Then

$$\mathbf{u}_r(1) = \gamma_r(C) \mathbf{u}_r(0), \quad (5.8)$$

where  $\gamma_r(C) = \pm 1$ . Defining  $\bar{\mathbf{u}}_s(t)$  analogously to be a continuously varying, normalised eigenvector of  $\bar{L}(\mathbf{z}(t))$  with eigenvalue  $\bar{\lambda}_s(\mathbf{z}(t))$ , we have that

$$\bar{\mathbf{u}}_s(1) = \bar{\gamma}_s(C) \bar{\mathbf{u}}_s(0), \quad (5.9)$$

where  $\bar{\gamma}_s(C) = \pm 1$ .  $\gamma_r$  and  $\bar{\gamma}_s$  may be regarded as the holonomies of the real eigenvector line bundles  $E_r$  and  $\bar{E}_s$  over  $R$ ,

$$\begin{aligned} E_r &= \{(\mathbf{z}, \mathbf{u}) \in R \times \mathbb{R}^n \mid (L(\mathbf{z}) - \lambda_r(\mathbf{z})) \cdot \mathbf{u} = 0\}, \\ \bar{E}_s &= \{(\mathbf{z}, \mathbf{u}) \in R \times \mathbb{R}^n \mid (\bar{L}(\mathbf{z}) - \bar{\lambda}_s(\mathbf{z})) \cdot \mathbf{u} = 0\}. \end{aligned} \quad (5.10)$$

They are examples of (real) geometric phases [8], though in this context (holonomies of eigenvectors of real symmetric matrices) have a long history (see, eg, [5] and [9]).

For the periodic Toda chain, the Maslov index and eigenvector holonomies are related by the following:

**Theorem 5.1.**

$$(-1)^{\mu/2} = \prod_{r \text{ even}} \gamma_r \prod_{s \text{ even}} \bar{\gamma}_s. \quad (5.11)$$

We note that the unrestricted product  $\prod_r \gamma_r$  is always +1, since this gives the holonomy of the (trivial) determinant bundle of  $R \times \mathbb{R}^n$ . Similarly for the unrestricted product  $\prod_s \bar{\gamma}_s$ . Thus, either of the products in (5.11) may be restricted to odd rather than even indices. (Alternatively, from Proposition 4.2 one can deduce that  $\gamma_r = \gamma_{r+1}$  for  $r$  even and  $\bar{\gamma}_r = \bar{\gamma}_{r+1}$  for  $r$  odd.)

*Proof.* Let  $C$  be a continuous closed curve in the regular set  $R$  of  $F$ , and let  $S$  be a transverse disk with boundary  $C$ . Let  $N$  denote the number of singular points in the image of  $S$ . From (5.7),

$$(-1)^{\mu(C)/2} = (-1)^N. \quad (5.12)$$

At the singular points  $\mathbf{z}_j \in \Sigma_1$ , there is precisely one doubly degenerate eigenvalue of either  $L(\mathbf{z}_j)$  or  $\bar{L}(\mathbf{z}_j)$ . For definiteness, suppose  $\lambda_r$  and  $\lambda_{r+1}$  are degenerate at  $\mathbf{z}_*$ . As in (4.10) and (4.11), in a neighbourhood of  $\mathbf{z}_*$ ,  $\bar{L}(\mathbf{z})$  is smoothly conjugate to a diagonal matrix, while  $L(\mathbf{z})$  is smoothly conjugate to a block diagonal matrix  $\Lambda(\mathbf{z})$  with a single two-dimensional block, with elements  $\Lambda_{ij}(\mathbf{z})$ ,  $i, j = 1, 2$ , and the rest diagonal. Let

$$\xi(\mathbf{z}) = \frac{1}{2}(\Lambda_{11}(\mathbf{z}) - \Lambda_{22}(\mathbf{z})), \quad \eta(\mathbf{z}) = \Lambda_{12}(\mathbf{z}). \quad (5.13)$$

As in Proposition 4.4,  $\xi$  and  $\eta$  are functionally independent near  $\mathbf{z}_*$ , and  $\xi(\mathbf{z}_*) = \eta(\mathbf{z}_*) = 0$ . Construct local coordinates  $(\xi, \eta, a_1, \dots, a_{2n-2})$  with origin at  $\mathbf{z}_*$ . Let  $C_*$  denote the closed curve parameterised by  $\xi(t) = \epsilon \cos 2\pi t$ ,  $\eta(t) = \epsilon \sin 2\pi t$ ,  $a_k = 0$ , with  $\epsilon$  chosen small enough so that  $C_*$  lies in the coordinate neighbourhood. Along  $C_*$ , let  $\Lambda^{(2)}(t)$  denote the two-dimensional block of  $\Lambda$ . Then

$$\Lambda^{(2)}(t) = \frac{1}{2}\tau(t)\mathbf{I} + \begin{pmatrix} -\cos 2\pi t & \sin 2\pi t \\ \sin 2\pi t & \cos 2\pi t \end{pmatrix}, \quad (5.14)$$

where  $\tau(t)$  is given by  $\Lambda_{22} + \Lambda_{11}$  along  $C_*$ . The eigenvectors  $(\cos \pi t \ \sin \pi t)^T$  and  $(\sin \pi t \ -\cos \pi t)^T$  of  $\Lambda^{(2)}(t)$  change sign around  $C_*$ . This corresponds to holonomies

$$\gamma_r(C_*) = \gamma_{r+1}(C_*) = -1 \quad (5.15)$$

in the associated eigenvectors  $\mathbf{u}_r$  and  $\mathbf{u}_{r+1}$  of  $L$ . The other holonomies, ie  $\gamma_{r \neq p}(C_*)$ ,  $\bar{\gamma}_s(C_*)$ , are trivially  $+1$ .

We introduce contours  $C_j$  analogously for all singularities  $\mathbf{z}_j$  in the image of  $S$  (replacing  $L$  by  $\bar{L}$  in the preceding as appropriate).  $C$  is homologous to an oriented sum of circuits  $C_j$ . Around each  $C_j$ , exactly one even-indexed eigenvector holonomy of either  $L$  or  $\bar{L}$  is  $-1$  (cf (5.15)), and the rest are  $+1$ . Thus,

$$\prod_{r, \text{even}} \gamma_r(C) \prod_{s, \text{even}} \gamma_s(C) = (-1)^N. \quad (5.16)$$

Comparing (5.16) and (5.12), we obtain the result (5.11).  $\square$

## 6 Discussion

$\Sigma_k$ , the space of codimension- $2k$  singularities of the Toda chain, is a symplectic submanifold corresponding to points where there are  $k$  doubly degenerate eigenvalues of the Lax matrices  $L$  and  $\bar{L}$ , for  $1 \leq k \leq n - 1$ . These submanifolds are of elliptic type, and we have calculated the frequencies of transverse oscillations under integrable flows that preserve them pointwise. The codimension-two singularities are sources for the Maslov index. The (even) Maslov index of a closed curve  $C$  is determined, modulo 4, by the product of the even- (or odd-) indexed eigenvector holonomies of  $L$  and  $\bar{L}$ . It would be interesting to relate higher Maslov classes [23, 25] to singularities of higher codimension, and to compute these higher Maslov classes explicitly for the Toda chain. In this context, it would also be interesting to study higher-order corrections [10, 11] to the semiclassical quantization conditions (5.2) for the quantum Toda chain [17, 18].

### Acknowledgments

We thank the referees for helpful remarks. JAF was supported by a grant from the EPSRC. JMR thanks the MSRI for hospitality and support while some of this work was carried out.

## References

- [1] M. Adler and P. van Moerbeke. Completely integrable systems, Euclidean lie algebras and curves. *Adv. Math.*, 38:267–317, 1980.
- [2] M. Adler and P. van Moerbeke. Linarization of Hamiltonian systems, Jacobi varieties and representation theory. *Adv. Math.*, 38:318–379, 1980.
- [3] M. Adler and P. van Moerbeke. The Toda lattice, Dynkin diagrams, singularities and Abelian varieties. *Adv. Math.*, 38:318–379, 1980.
- [4] V.I. Arnold. Characteristic class entering quantization conditions. *Functional Anal. Appl.*, 1:1–13, 1967.
- [5] V.I. Arnold. *Mathematical methods of classical mechanics*. Springer-Verlag, 2nd edition, 1989.
- [6] M. Audin. *Spinning Tops*. Cambridge University Press, 1996.
- [7] O. Babelon, D. Bernard, and M. Talon. *Introduction to classical integrable systems*. Cambridge University Press, 2003.

- [8] M.V. Berry. Quantal phase factors accompanying adiabatic changes. *Proc. Roy. Soc. London Ser. A*, 392:45–57, 1984.
- [9] M.V. Berry. Anticipations of the geometric phase. *Physics Today*, 43:34–40, 1990.
- [10] M. Cargo, A. Gracia-Saz, R.G. Littlejohn, M.W. Reinsch, and P. de M. Rios. Quantum normal forms, Moyal star product and Bohr-Sommerfeld approximation. *J. Phys. A*, 38:1977–2004, 2005.
- [11] Y. Colin de Verdière. Bohr-Sommerfeld rules to all orders. to be published in *Ann. Inst. Henri Poincaré*, year = 2004.
- [12] L.H. Eliasson. Normal form for Hamiltonian systems with Poisson commuting integrals – elliptic case. *Comm. Math. Helv.*, 65:4–35, 1990.
- [13] H. Flaschka. The toda lattice. i. existence of integrals. *Phys. Rev. B*, 9:1924–1925, 1974.
- [14] H. Flaschka. Integrable systems and torus actions. In P. Cartier O. Babelon and Y. Kosmann-Schwarzbach, editors, *Lectures on integrable systems*. World Scientific, 1994.
- [15] A. Fomenko. *Topological classification of integrable systems*. Number 6 in Advances in Soviet Mathematics. AMS, 1991.
- [16] J.A. Foxman and J.M. Robbins. The Maslov index and nondegenerate singularities of integrable systems. preprint, 2004.
- [17] M.C. Gutwiller. The quantum mechanical Toda lattice. *Ann. Physics*, 124, 1980.
- [18] M.C. Gutwiller. The quantum mechanical Toda lattice. ii. *Ann. Physics*, 133:304–331, 1981.
- [19] M. Hénon. Integrals of the Toda lattice. *Phys. Rev. B*, 9:1921–1923, 1974.
- [20] J.B. Keller. Corrected Bohr-Sommerfeld quantum conditions for non-separable systems. *Ann. Phys.*, 4:180–188, 1958.
- [21] V.P. Maslov and M. Fedoriuk. *Semiclassical approximation in quantum mechanics*. Reidel, 1981.

- [22] L.S. Polyakova. Topological invariants for some algebraic analogs of the toda lattice. In *Topological classification of integrable systems*, volume 6 of *Adv. Sov. Math.*, pages 185–207. 1991.
- [23] H. Suzuki. Residue classes of Lagrangian subbundles and Maslov classes. *Trans. Amer. Math. Soc.*, 347:189–202, 1995.
- [24] M. Toda. Waves in a nonlinear lattice. *Prog. Theor. Phys. Suppl.*, 45:174–200, 1970.
- [25] V.V. Trofimov. Generalized Maslov classes on the path space of a symplectic manifold. *Proc. Steklov Institute of Mathematics*, 205:157–179, 1995.
- [26] P. van Moerbeke and D. Mumford. The spectrum of difference operators and algebraic curves. *Acta Math.*, 143:93–154, 1979.
- [27] J. Vey. Sur certains systèmes dynamiques séparables. *Amer. J. Math.*, 100:591–614, 1978.
- [28] Nguyen Tien Zung. Symplectic topology of integrable Hamiltonian systems, I: Arnold-Liouville with singularities. *Compositio Math.*, 101:179–215, 1996.