

PERIODIC CELLULAR AUTOMATA AND BETHE ANSATZ

ATSUO KUNIBA AND AKIRA TAKENOUCI

ABSTRACT. We review and generalize the recent progress in a soliton cellular automaton known as the periodic box-ball system. It has the extended affine Weyl group symmetry and admits the commuting transfer matrix method and the Bethe ansatz at $q = 0$. Explicit formulas are proposed for the dynamical period and the number of states characterized by conserved quantities.

1. INTRODUCTION

In [13], a class of periodic soliton cellular automata is introduced associated with non-exceptional quantum affine algebras. The dynamical period and a state counting formula are proposed by the Bethe ansatz at $q = 0$ [11]. In this paper we review and generalize the results on the $A_n^{(1)}$ case, where the associated automaton is known as the periodic box-ball system [14, 19]. The box-ball system was originally introduced on the infinite lattice without boundary [18, 17]. Here is a collision of two solitons with amplitudes 3 and 1 interchanging internal degrees of freedom with a phase shift:

$$\begin{aligned} & \cdots 1114221111131111111111111111 \cdots \\ & \cdots 1111114221113111111111111111 \cdots \\ & \cdots 1111111114221311111111111111 \cdots \\ & \cdots 1111111111142321111111111111 \cdots \\ & \cdots 1111111111111214321111111111 \cdots \\ & \cdots 1111111111111121114321111111 \cdots \\ & \cdots 111111111111111211111432111 \cdots \end{aligned}$$

The system was identified [4, 3] with a solvable lattice model [1] at $q = 0$, which led to a direct formulation by the crystal base theory [5] and generalizations to the soliton cellular automata with quantum group symmetry [7, 6]. Here we develop the approach to the periodic case in [13] further by combining the commuting transfer matrix method [1] and the Bethe ansatz [2] at $q = 0$.

In section 2 we formulate the most general periodic automaton for $\mathfrak{g}_n = A^{(1)}$ in terms of the crystal theory. A commuting family of time evolutions $\{T_j^{(a)}\}$ is introduced as commuting transfer matrices under periodic boundary condition. The associated conserved quantities form an n -tuple of Young diagrams $m = (m^{(1)}, \dots, m^{(n)})$, which we call the *soliton content*.

In section 3 we invoke the Bethe ansatz at $q = 0$ [11] to study the Bethe eigenvalue $\Lambda_i^{(r)}$ relevant to $T_i^{(r)}$. The Bethe equation is linearized into the string center equation and the $\Lambda_i^{(r)}$ is shown to be a root of unity. We also recall an explicit weight multiplicity formula obtained by counting the off-diagonal solutions to the string center equation [11]. It is a version of the fermionic formula called the combinatorial completeness of the string hypothesis at $q = 0$. These results are parameterized with the number of strings, which we call the *string content*.

In section 4 two applications of the results in section 3 are presented under the identification of the soliton and the string contents. First we relate the root of unity in the Bethe eigenvalue $\Lambda_l^{(r)}$ to the dynamical period of the periodic $A_n^{(1)}$ automaton under the time evolution $T_l^{(r)}$. Second we connect each summand in the weight multiplicity formula [11] to the number of states characterized by conserved quantities.

In [13], similar results have been announced concerning the highest states. Our approach here is based on conserved quantities and covers a wider class of states without recourse to the combinatorial Bethe ansatz at $q = 1$ [10]. We expect parallel results in general \mathfrak{g}_n . In fact all the essential claims in this paper make sense also for $\mathfrak{g}_n = D_n^{(1)}$ and $E_{6,7,8}^{(1)}$. Our formulas (11) and (13) include the results in [19] proved by a different approach as the case $\mathfrak{g}_n = A_1^{(1)}$ with $B = (B^{1,1})^{\otimes L}$ and $l = \infty$. For the standard notation and facts in the crystal theory, we refer to [5, 9, 6].

2. PERIODIC $A_n^{(1)}$ AUTOMATON

Let $B^{a,j}$ ($1 \leq a \leq n, j \in \mathbb{Z}_{\geq 1}$) be the crystal [9] of the Kirillov-Reshetikhin module $W_j^{(a)}$ over $U_q(A_n^{(1)})$. Elements of $B^{a,j}$ are labeled with semistandard tableaux on an $a \times j$ rectangular Young diagram with letters $\{1, 2, \dots, n+1\}$. For example when $n = 2$, one has $B^{1,1} = \{1, 2, 3\}$, $B^{1,2} = \{11, 12, 13, 22, 23, 33\}$, $B^{2,2} = \left\{ \begin{array}{cc} 1 & 1 \\ 2 & 2 \end{array}, \begin{array}{cc} 1 & 1 \\ 2 & 3 \end{array}, \begin{array}{cc} 1 & 1 \\ 3 & 3 \end{array}, \begin{array}{cc} 1 & 2 \\ 2 & 3 \end{array}, \begin{array}{cc} 1 & 2 \\ 3 & 3 \end{array}, \begin{array}{cc} 2 & 2 \\ 3 & 3 \end{array} \right\}$ as sets. $\text{Aff}(B^{a,j}) = \{\zeta^{db} \mid b \in B^{a,j}, d \in \mathbb{Z}\}$ denotes the affine crystal. The combinatorial R is the isomorphism of affine crystals $\text{Aff}(B^{a,j}) \otimes \text{Aff}(B^{b,k}) \xrightarrow{\sim} \text{Aff}(B^{b,k}) \otimes \text{Aff}(B^{a,j})$ [16]. It has the form $R(\zeta^{db} \otimes \zeta^e c) = \zeta^{e+H} \tilde{c} \otimes \zeta^{d-H} \tilde{b}$, where $H = H(b \otimes c)$ is the energy function. We normalize it so as to attain the maximum at $H(u^{a,j} \otimes u^{b,k}) = 0$, where $u^{a,j} \in B^{a,j}$ denotes the classically highest element. We set $B = B^{r_1, l_1} \otimes B^{r_2, l_2} \otimes \dots \otimes B^{r_L, l_L}$ and write $\text{Aff}(B^{r_1, l_1}) \otimes \dots \otimes \text{Aff}(B^{r_L, l_L})$ simply as $\text{Aff}(B)$. An element of B is called a state. Given a state $p = b_1 \otimes \dots \otimes b_L \in B$, regard it as the element $\zeta^0 b_1 \otimes \dots \otimes \zeta^0 b_L \in \text{Aff}(B)$ and seek an element $v \in B^{r, l}$ such that $\zeta^0 v \otimes p \simeq (\zeta^{d_1} b'_1 \otimes \dots \otimes \zeta^{d_L} b'_L) \otimes \zeta^e v$ under the isomorphism $\text{Aff}(B^{r, l}) \otimes \text{Aff}(B) \simeq \text{Aff}(B) \otimes \text{Aff}(B^{r, l})$. If such a v exists and $\zeta^{d_1} b'_1 \otimes \dots \otimes \zeta^{d_L} b'_L$ is unique even if v is not unique, we say that p is (r, l) -evolvable and write $T_l^{(r)}(p) = b'_1 \otimes \dots \otimes b'_L \in B$ and $E_l^{(r)}(p) = e = -d_1 - \dots - d_L$. Otherwise we say that p is not (r, l) -evolvable or $T_l^{(r)}(p) = 0$. We formally set $T_l^{(r)}(0) = 0$. $E_l^{(r)} \in \mathbb{Z}_{\geq 0}$ holds under this normalization. Our $A_n^{(1)}$ automaton is a dynamical system on $B \sqcup \{0\}$ equipped with the family of time evolutions $\{T_l^{(r)} \mid 1 \leq r \leq n, l \in \mathbb{Z}_{\geq 1}\}$. $T_l^{(r)}$ is the $q = 0$ analogue of the transfer matrix in solvable vertex models. It is invertible and weight preserving on B . Using the Yang-Baxter equation of the combinatorial R , one can show (cf. [3, 6])

Theorem 2.1. *The commutativity $T_j^{(a)} T_k^{(b)}(p) = T_k^{(b)} T_j^{(a)}(p)$ is valid for any $(a, j), (b, k)$ and $p \in B$, where the both sides are either in B or 0. In the former case $E_j^{(a)}(T_k^{(b)}(p)) = E_j^{(a)}(p)$ and $E_k^{(b)}(T_j^{(a)}(p)) = E_k^{(b)}(p)$ hold.*

Thus, $\{E_j^{(a)} \mid 1 \leq a \leq n, j \in \mathbb{Z}_{\geq 1}\}$ is a family of conserved quantities.

Conjecture 2.1. *For any $1 \leq a \leq n$ and $p \in B$, there exists $i \geq 1$ such that $T_k^{(a)}(p) \neq 0$ if and only if $k \geq i$. The limit $\lim_{k \rightarrow \infty} T_k^{(a)}(p) \in B$ exists and $E_i^{(a)}(p) < E_{i+1}^{(a)}(p) < \dots < E_j^{(a)}(p) = E_{j+1}^{(a)}(p) = \dots$ holds for some $j \geq i$.*

Let S_0, S_1, \dots, S_n be the Weyl group operators [5] and pr be the promotion operator [16] acting on B component-wise. For instance for $A_3^{(1)}$, $\text{pr}\left(\begin{smallmatrix} 2 & 2 & 3 \\ 3 & 3 & 4 \end{smallmatrix} \otimes 1344\right) = \begin{smallmatrix} 1 & 3 & 3 \\ 4 & 4 & 4 \end{smallmatrix} \otimes 1124 \in B^{2,3} \otimes B^{1,4}$. They act on B as the extended affine Weyl group $\widetilde{W}(\mathfrak{g}_n) = \widetilde{W}(A_n^{(1)}) = \langle \text{pr}, S_0, S_1, \dots, S_n \rangle$.

Theorem 2.2. *If $T_j^{(a)}(p) \neq 0$, then for any $w \in \widetilde{W}(A_n^{(1)})$, the relations $wT_j^{(a)}(p) = T_j^{(a)}(w(p))$ and $E_j^{(a)}(w(p)) = E_j^{(a)}(p)$ are valid.*

A state $p \in B$ is called *evolvable* if it is (a, j) -evolvable for any (a, j) . In Conjecture 2.1, we expect that the convergent limit $T_\infty^{(a)}$ equals a translation in $\widetilde{W}(A_n^{(1)})$. Compared with T_l in [13], the family $\{T_j^{(a)}\}$ here is more general and enjoys a larger symmetry $\widetilde{W}(A_n^{(1)})$. Define the subset of B by

$$(1) \quad P(m) = \{p \in B \mid p : \text{evolvable}, E_j^{(a)}(p) = \sum_{k \geq 1} \min(j, k) m_k^{(a)}\}.$$

Pictorially, $m = (m^{(1)}, \dots, m^{(n)})$ is the n -tuple of Young diagrams and $E_k^{(a)}$ (resp. $m_k^{(a)}$) is the number of nodes in the first k columns (resp. number of length k rows) in $m^{(a)}$. We call m the *soliton content*.

Remark 2.1. $P(m)$ is $\widetilde{W}(A_n^{(1)})$ -invariant due to Theorem 2.2.

Given $p \in P(m)$, $T_j^{(a)}(p) \in P(m)$ is not necessarily valid. For instance $p = 112233 \in P(((22), (2))) \subset B = (B^{1,1})^{\otimes 6}$ but $T_1^{(2)}(p) = 213213$ is not evolvable since $(T_1^{(2)})^2(p) = 0$. On the other hand one can show

Proposition 2.1. *If $p \in P(m)$ and $(T_j^{(a)})^t(p) \neq 0$ for any t , then $(T_j^{(a)})^t(p) \in P(m)$ for any t .*

Let W, Λ_a, α_a be the Weyl group, the fundamental weights and the simple roots of A_n , respectively. We specify $p_j^{(a)} = p_j^{(a)}(m)$ by (6) and set

$$(2) \quad \lambda(m) = \sum_{a=1}^n p_\infty^{(a)} \Lambda_a, \quad H = \{(a, j) \mid 1 \leq a \leq n, j \in \mathbb{Z}_{\geq 1}, m_j^{(a)} > 0\}.$$

Conjecture 2.2. $P(m) \neq \emptyset$ if and only if $p_j^{(a)} \geq 0$ for all $(a, j) \in H$. $\{\text{wtp} \mid p \in P(m)\} = W\lambda(m)$.

The claim on the weights is consistent with the $\widetilde{W}(A_n^{(1)})$ -invariance of $P(m)$.

Here is an example of time evolutions in $B = B^{1,1} \otimes B^{1,1} \otimes B^{1,3} \otimes B^{1,1} \otimes B^{1,1} \otimes B^{1,1} \otimes B^{1,2}$ with $\widetilde{W}(A_3^{(1)})$ symmetry. The leftmost column is $p_0, T_2^{(1)}(p_0), T_1^{(2)}T_2^{(1)}(p_0)$ and $T_2^{(3)}T_1^{(2)}T_2^{(1)}(p_0)$ from the top to the bottom. At each time step, the states connected by the Weyl group actions S_0 and S_1 are shown, forming commutative diagrams. (\cdot signifies \otimes .) All these states belong to $P(((3111), (21), (1)))$.

$$\begin{array}{ccccc} 2 \cdot 1 \cdot 233 \cdot 4 \cdot 1 \cdot 2 \cdot 12 & \xrightarrow{S_0} & 2 \cdot 4 \cdot 233 \cdot 4 \cdot 1 \cdot 2 \cdot 24 & \xrightarrow{S_1} & 1 \cdot 4 \cdot 133 \cdot 4 \cdot 1 \cdot 2 \cdot 14 \\ 1 \cdot 2 \cdot 123 \cdot 3 \cdot 4 \cdot 1 \cdot 22 & & 4 \cdot 2 \cdot 234 \cdot 3 \cdot 4 \cdot 1 \cdot 22 & & 4 \cdot 1 \cdot 134 \cdot 3 \cdot 4 \cdot 1 \cdot 12 \\ 1 \cdot 2 \cdot 112 \cdot 3 \cdot 2 \cdot 3 \cdot 24 & & 1 \cdot 2 \cdot 244 \cdot 3 \cdot 2 \cdot 3 \cdot 24 & & 1 \cdot 2 \cdot 144 \cdot 3 \cdot 1 \cdot 3 \cdot 14 \\ 2 \cdot 3 \cdot 112 \cdot 4 \cdot 2 \cdot 3 \cdot 12 & & 2 \cdot 3 \cdot 244 \cdot 4 \cdot 2 \cdot 3 \cdot 12 & & 2 \cdot 3 \cdot 144 \cdot 4 \cdot 1 \cdot 3 \cdot 11 \end{array}$$

Remark 2.2. Let $R_j (1 \leq j \leq L-1)$ be the combinatorial R that exchanges the j -th and $(j+1)$ -st components in $B = B^{r_1, l_1} \otimes \cdots \otimes B^{r_L, l_L}$ and $\pi(b_1 \otimes \cdots \otimes b_L) = b_L \otimes b_1 \otimes \cdots \otimes b_{L-1}$. Together with $R_0 = \pi^{-1} R_1 \pi$, they act on $\cup_{s \in \mathfrak{S}_L} B^{r_{s_1}, l_{s_1}} \otimes \cdots \otimes B^{r_{s_L}, l_{s_L}}$ as the extended affine Weyl group $\widetilde{W}(A_{L-1}^{(1)}) = \langle \pi, R_0, \dots, R_{L-1} \rangle$. Theorem 2.2 is actually valid for $w \in \widetilde{W}(\mathfrak{g}_n = A_n^{(1)}) \times \widetilde{W}(A_{L-1}^{(1)})$ as in [8]. In the homogeneous case $(r_1, l_1) = \cdots = (r_L, l_L)$, the $\widetilde{W}(A_{L-1}^{(1)})$ symmetry shrinks down to the π -symmetry, which is the origin of the adjective “periodic”.

3. BETHE ANSATZ AT $q = 0$

Eigenvalues of row transfer matrices in trigonometric vertex models are given by the analytic Bethe ansatz [15, 12]. Let $Q_r(u) = \prod_k \sinh \pi(u - \sqrt{-1}u_k^{(r)})$ be Baxter’s Q -function where $\{u_j^{(a)}\}$ satisfy the Bethe equation eq.(2.1) in [11]. We set $q = e^{-2\pi\hbar}$ and $\zeta = e^{2\pi u}$. For the string solution ([11] Definition 2.3), the relevant quantity to our $T_l^{(r)}$ is the top term of the eigenvalue $\Lambda_l^{(r)}(u)$ (cf.[12] (2.12)):

$$(3) \quad \lim_{q \rightarrow 0} \frac{Q_r(u - l\hbar)}{Q_r(u + l\hbar)} = \zeta^{-E_l^{(r)}} \Lambda_l^{(r)}, \quad \Lambda_l^{(r)} := \prod_{j\alpha} (-z_{j\alpha}^{(r)})^{\min(j,l)}.$$

Here $z_{j\alpha}^{(a)}$ is the center of the α -th string having color a and length j . Denote by $m_j^{(a)}$ the number of such strings. We call the data $m = (m_j^{(a)})$ the *string content*. The product in (3) is taken over $j \in \mathbb{Z}_{\geq 1}$ and $1 \leq \alpha \leq m_j^{(r)}$. $E_l^{(r)}$ is given by the same expression as in (1) as the function of m . At $q = 0$ the Bethe equation becomes the string center equation ([11] (2.36)):

$$(4) \quad \prod_{(b,k) \in H} \prod_{\beta=1}^{m_k^{(b)}} (z_{k\beta}^{(b)})^{A_{aj\alpha, bk\beta}} = (-1)^{p_j^{(a)} + m_j^{(a)} + 1},$$

$$(5) \quad A_{aj\alpha, bk\beta} = \delta_{ab} \delta_{jk} \delta_{\alpha\beta} (p_j^{(a)} + m_j^{(a)}) + C_{ab} \min(j, k) - \delta_{ab} \delta_{jk},$$

$$(6) \quad p_j^{(a)} = \sum_{i=1}^L \min(j, l_i) \delta_{ar_i} - \sum_{(b,k) \in H} C_{ab} \min(j, k) m_k^{(b)},$$

where $(C_{ab})_{1 \leq a, b \leq n}$ is the Cartan matrix of A_n . To avoid a notational complexity we temporarily abbreviate the triple indices $aj\alpha$ to j , $bk\beta$ to k and accordingly $z_{k\beta}^{(b)}$ to z_k etc. Then (3) and (4) read

$$(7) \quad \Lambda_l^{(r)} = \prod_k (-z_k)^{\rho_k}, \quad \prod_k (-z_k)^{A_{j,k}} = (-1)^{s_j},$$

where ρ_k is given by $\rho_k = \delta_{br} \min(k, l)$ for k corresponding to $bk\beta$, and s_j is an integer. Note that $A_{j,k} = A_{k,j}$. Suppose that the $q = 0$ eigenvalue satisfies $(\Lambda_l^{(r)})^{\mathcal{P}_l^{(r)}} = \pm 1$ for generic solutions to the string center equation ¹. It means that there exist integers ξ_j such that $\sum_j \xi_j A_{j,k} = \mathcal{P}_l^{(r)} \rho_k$, or equivalently $\xi_j = \mathcal{P}_l^{(r)} \frac{\det A[j]}{\det A}$, where $A[j]$ denotes the matrix $A = (A_{j,k})$ with its j -th column replaced by ${}^t(\rho_1, \rho_2, \dots)$. In view of the condition $\forall \xi_j \in \mathbb{Z}$, the minimum integer allowed for $\mathcal{P}_l^{(r)}$ is $\mathcal{P}_l^{(r)} = \text{LCM}\left(1, \cup_k \frac{\det A}{\det A[k]}\right)$, where LCM stands for the least common multiple and \cup_k' means the union over those

¹ $\mathcal{P}_l^{(r)}$ here should not be confused with the symbol in (6).

k such that $A[k] \neq 0$. Back in the original indices, the determinants here can be simplified (cf. [11] (3.9)) to those of matrices indexed with H :

$$(8) \quad \mathcal{P}_l^{(r)} = \text{LCM}\left(1, \bigcup'_{(b,k) \in H} \frac{\det F}{\det F[b, k]}\right),$$

where the matrix $F = (F_{aj, bk})_{(a,j), (b,k) \in H}$ is defined by

$$(9) \quad F_{aj, bk} = \delta_{ab} \delta_{jk} p_j^{(a)} + C_{ab} \min(j, k) m_k^{(b)}.$$

The matrix $F[b, k]$ is obtained from F by replacing its (b, k) -th column as

$$(10) \quad F[b, k]_{aj, cm} = \begin{cases} F_{aj, cm} & (c, m) \neq (b, k), \\ \delta_{ar} \min(j, l) & (c, m) = (b, k). \end{cases}$$

The union in (8) is taken over those (b, k) such that $\det F[b, k] \neq 0$.

The LCM (8) can further be simplified for $A_1^{(1)}$, $r = 1, \forall r_i = \forall l_i = 1$. We write $p_j^{(1)}$ just as p_j and parameterize the set $H = \{j \in \mathbb{Z}_{\geq 1} \mid m_j^{(1)} > 0\}$ as $H = \{(0 <) J_1 < \dots < J_s\}$. Setting $i_k = \min(J_k, l)$ and $i_0 = 0$, one has

$$(11) \quad \mathcal{P}_l^{(1)} = \text{LCM}\left(1, \bigcup'_{k=0}^t \frac{p_{i_{k+1}} p_{i_k}}{(i_{k+1} - i_k) p_{i_s}}\right),$$

where $0 \leq t \leq s - 1$ is the maximum integer such that $i_{t+1} > i_t$.

Let us turn to another Bethe ansatz result, the character formula called combinatorial completeness of the string hypothesis at $q = 0$ [11]:

$$(12) \quad \prod_{i=1}^L \text{ch} B^{r_i, l_i} = \sum_m \Omega(m) e^{\lambda(m)},$$

$$(13) \quad \Omega(m) = \det F \prod_{(a,j) \in H} \frac{1}{m_j^{(a)}} \binom{p_j^{(a)} + m_j^{(a)} - 1}{m_j^{(a)} - 1} \in \mathbb{Z},$$

where $\binom{s}{t} = s(s-1)\dots(s-t+1)/t!$ and $\text{ch} B^{r, l}$ is the character of $B^{r, l}$. $\lambda(m)$, $p_j^{(a)}$ and F are defined by (2), (6) and (9). The sum in (12) extends over all $m_j^{(a)} \in \mathbb{Z}_{\geq 0}$ canceling out exactly leaving the character of B . (12) and (13) are the special cases of eq.(5.13) and eq.(4.1) in [11], respectively. $\Omega(m)$ (denoted by $R(\nu, N)$ therein) is the number of off-diagonal solutions to the string center equation with string content m . It is known ([11] Lemma 3.7) that $\Omega(m) \in \mathbb{Z}_{\geq 1}$ provided that $p_j^{(a)} \geq 0$ for all $(a, j) \in H$.

4. DYNAMICAL PERIOD AND STATE COUNTING

In (1), the soliton content $m = (m_j^{(a)})$ is introduced as the conserved quantity associated with the commuting transfer matrices. On the other hand, the $m_j^{(a)}$ in the string content $m = (m_j^{(a)})$ is the number of strings of color a and length j in the Bethe ansatz in section 3. From now on we identify them motivated by the factor $\zeta^{-E_i^{(r)}}$ in (3) and some investigation of Bethe vectors at $q = 0$. In view of Conjecture 2.2, the data of the form $m = (m_j^{(a)})$ is defined to be a *content* if and only if $p_j^{(a)} \geq 0$ for all $(a, j) \in H$. Thus $\det F > 0$ and $\lambda(m)$ in (2) is a dominant weight for any content m .

Conjecture 4.1. *If $p \in P(m)$ and $(T_l^{(r)})^t(p) \neq 0$ for any t , the dynamical period of p under $T_l^{(r)}$ (minimum positive integer t such that $(T_l^{(r)})^t(p) = p$) is equal to $\mathcal{P}_l^{(r)}$ (8) generically and its divisor otherwise.*

In the situation under consideration, the whole $T_l^{(r)}$ orbit of p belongs to $P(m)$ due to Proposition 2.1. Naturally we expect $(\Lambda_l^{(r)})^{\mathcal{P}_l^{(r)}} = 1$, which can indeed be verified for $A_1^{(1)}$. Conjecture 4.1 has been checked, for example in $A_3^{(1)}$ case, for $B = (B^{1,1})^{\otimes 3} \otimes B^{2,2}$ and $B^{2,1} \otimes B^{2,1} \otimes B^{3,1} \otimes B^{3,2}$.

Let us present more evidence of Conjecture 4.1. To save the space, $\cdot = \otimes$ is dropped when $B = (B^{1,1})^{\otimes L}$. In each table, the period under $T_l^{(r)}$ with maximum l is equal to that under $T_\infty^{(r)}$.

$A_1^{(1)}$, state = 1221121122221, content = ((321))

(r, l)	LCM of				= period
(1,1)	1,	13,	13,	13	13
(1,2)	1,	$\frac{91}{3}$,	$\frac{91}{16}$,	$\frac{91}{16}$	91
(1,3)	1,	91,	$\frac{273}{16}$,	$\frac{273}{107}$	273

$A_1^{(1)}$, state = 122 · 112 · 12 · 1222 · 2 · 11111 · 1122 · 111, content = ((4321))

(r, l)	LCM of				= period
(1,1)	1,	2			2
(1,2)	1,	7,	$\frac{7}{2}$,	21, 42	42
(1,3)	1,	14,	7,	$\frac{21}{4}$, $\frac{21}{2}$	42
(1,4)	1,	21,	$\frac{21}{2}$,	$\frac{63}{8}$, $\frac{126}{29}$	126

$A_3^{(1)}$, state = 134 · 34 · 1 · 134 · 23 · 1 · 13, content = ((432), (31), (1))

(r, l)	LCM of							= period
(1,1)	1,	$\frac{380}{39}$,	$\frac{95}{6}$,	$\frac{95}{6}$,	$\frac{380}{31}$,	$\frac{380}{27}$,	$\frac{380}{29}$	380
(1,2)	1,	$\frac{190}{39}$,	$\frac{95}{12}$,	$\frac{95}{12}$,	$\frac{190}{31}$,	$\frac{190}{27}$,	$\frac{190}{29}$	190
(2,1)	1,	$\frac{190}{13}$,	$\frac{95}{4}$,	$\frac{95}{4}$,	$\frac{190}{137}$,	$\frac{190}{9}$,	$\frac{190}{73}$	190
(2,2)	1,	$\frac{76}{5}$,	$\frac{38}{3}$,	$\frac{38}{3}$,	$\frac{76}{41}$,	$\frac{76}{21}$,	$\frac{76}{31}$	76
(2,3)	1,	$\frac{95}{6}$,	$\frac{95}{11}$,	$\frac{95}{11}$,	$\frac{95}{34}$,	$\frac{95}{48}$,	$\frac{95}{41}$	95
(3,1)	1,	$\frac{380}{13}$,	$\frac{95}{2}$,	$\frac{95}{2}$,	$\frac{380}{137}$,	$\frac{380}{9}$,	$\frac{380}{263}$	380

$A_3^{(1)}$, state = 233 · $\frac{1}{2}$ $\frac{2}{3}$ $\frac{3}{4}$ · $\frac{1}{3}$ $\frac{1}{4}$ · 1, content = ((3), (3), (2))

(r, l)	LCM of			= period
(1,1)	1,	$\frac{11}{2}$,	11, 22	22
(1,2)	1,	$\frac{11}{4}$,	$\frac{11}{2}$,	11
(1,3)	1,	$\frac{11}{6}$,	$\frac{11}{3}$,	$\frac{22}{3}$
(2,1)	1,	11,	$\frac{33}{7}$,	$\frac{66}{7}$
(2,2)	1,	$\frac{11}{2}$,	$\frac{33}{14}$,	$\frac{33}{7}$
(2,3)	1,	$\frac{11}{3}$,	$\frac{11}{7}$,	$\frac{22}{7}$
(3,2)	1,	11,	$\frac{33}{7}$,	$\frac{33}{20}$

Here, $\text{content} = ((3111), (44), (2))$ for example means that $m_1^{(1)} = 3, m_3^{(1)} = m_2^{(3)} = 1, m_4^{(2)} = 2$ and the other $m_j^{(a)}$'s are 0.

Let us turn to another application of the Bethe ansatz results (12) and (13). We introduce $\mathcal{T}(P(m)) = \bigcup_{a=1}^n \bigcup_{j \geq 1} \{T_j^{(a)}(p) \mid p \in P(m)\}$, which is the subset of B consisting of all kinds of one step time evolutions of $P(m)$. Under Conjecture 2.1, any state $p \in P(m)$ is (a, j) -evolvable for j sufficiently large. Thus from Proposition 2.1, p is expressed as $p = (T_j^{(a)})^k(p)$ for some k , showing that $\mathcal{T}(P(m)) \supseteq P(m)$. In general $\mathcal{T}(P(m))$ can contain non-evolvable states which do not belong to $P(m)$.

Conjecture 4.2. *For any content m such that $\mathcal{T}(P(m)) = P(m)$, the following relation holds:*

$$(14) \quad \Omega(m) = \frac{|P(m)|}{|W\lambda(m)|}.$$

In view of Remark 2.1 and Conjecture 2.2, the right hand side is the number of states in the periodic $A_n^{(1)}$ automaton having the content m and a fixed weight. Thus it is equal to $\#\{p \in P(m) \mid \text{wtp} = \lambda(m)\}$. In case $\mathcal{T}(P(m)) \not\supseteq P(m)$, we expect that $|P(m)|/|W\lambda(m)|$ is a divisor of $\Omega(m)$.

Let us present two examples of Conjecture 4.2. In the periodic $A_3^{(1)}$ automaton with $B = B^{1,2} \otimes B^{1,1} \otimes B^{1,2} \otimes B^{1,1}$, there are 1600 states among which 824 are evolvable. They are classified according to the contents m in the following table.

m	$\lambda(m)$	$ W\lambda(m) $	$ P(m) $	$\Omega(m)$
$(\emptyset, \emptyset, \emptyset)$	$(6, 0, 0, 0)$	4	4	1
$((1), \emptyset, \emptyset)$	$(5, 1, 0, 0)$	12	48	4
$((11), \emptyset, \emptyset)$	$(4, 2, 0, 0)$	12	24	2
$((2), \emptyset, \emptyset)$	$(4, 2, 0, 0)$	12	72	6
$((21), \emptyset, \emptyset)$	$(3, 3, 0, 0)$	6	24	4
$((3), \emptyset, \emptyset)$	$(3, 3, 0, 0)$	6	36	6
$((11), (1), \emptyset)$	$(4, 1, 1, 0)$	12	96	8
$((22), (2), \emptyset)^*$	$(2, 2, 2, 0)$	4	24	12
$((21), (1), \emptyset)$	$(3, 2, 1, 0)$	24	432	18
$((111), (11), (1))$	$(3, 1, 1, 1)$	4	16	4
$((211), (11), (1))$	$(2, 2, 1, 1)$	6	48	8

In the second column, $(\lambda_1, \lambda_2, \lambda_3, \lambda_4)$ means $\lambda(m) = (\lambda_1 - \lambda_2)\Lambda_1 + (\lambda_2 - \lambda_3)\Lambda_2 + (\lambda_3 - \lambda_4)\Lambda_3$. In the last two cases, the subsets of $P(m)$ having the dominant weight $\lambda(m)$ are given by

$$\begin{aligned} & \{11 \cdot 2 \cdot 13 \cdot 4, 12 \cdot 3 \cdot 14 \cdot 1, 13 \cdot 4 \cdot 11 \cdot 2, 14 \cdot 1 \cdot 12 \cdot 3\} \text{ for } m = ((111), (11), (1)), \\ & \{11 \cdot 2 \cdot 23 \cdot 4, 12 \cdot 2 \cdot 13 \cdot 4, 12 \cdot 3 \cdot 24 \cdot 1, 13 \cdot 4 \cdot 12 \cdot 2, \\ & \quad 14 \cdot 1 \cdot 22 \cdot 3, 22 \cdot 3 \cdot 14 \cdot 1, 23 \cdot 4 \cdot 11 \cdot 2, 24 \cdot 1 \cdot 12 \cdot 3\} \text{ for } m = ((211), (11), (1)). \end{aligned}$$

In the case of $B = B^{2,1} \otimes B^{2,1} \otimes B^{2,2}$, there are 720 states among which 518 are evolvable.

m	$\lambda(m)$	$ W\lambda(m) $	$ P(m) $	$\Omega(m)$
$(\emptyset, \emptyset, \emptyset)$	$(4, 4, 0, 0)$	6	6	1
$(\emptyset, (1), \emptyset)$	$(4, 3, 1, 0)$	24	72	3
$(\emptyset, (11), (1))$	$(4, 2, 1, 1)$	12	36	3
$(\emptyset, (2), \emptyset)$	$(4, 2, 2, 0)$	12	48	4
$((1), (11), \emptyset)$	$(3, 3, 2, 0)$	12	36	3
$((1), (11), (1))$	$(3, 3, 1, 1)$	6	72	12
$((1), (21), (1))$	$(3, 2, 2, 1)$	12	240	20
$((2), (22), (2))^*$	$(2, 2, 2, 2)$	1	8	32

The assumption $\mathcal{T}(P(m)) = P(m)$ of the conjecture is valid for all the contents except $((22), (2), \emptyset)$ and $((2), (22), (2))$ marked with $*$.

REFERENCES

- [1] R. J. Baxter, *Exactly solved models in statistical mechanics*, Academic Press, London (1982).
- [2] H. A. Bethe, *Z. Physik* **71** 205 (1931).
- [3] K. Fukuda, M. Okado, Y. Yamada, *Int. J. Mod. Phys. A* **15** 1379 (2000).
- [4] G. Hatayama, K. Hikami, R. Inoue, A. Kuniba, T. Takagi and T. Tokihiro, *J. Math. Phys.* **42** 274 (2001).
- [5] M. Kashiwara, *Duke Math. J.* **71** 839 (1993).
- [6] G. Hatayama, A. Kuniba, M. Okado, T. Takagi and Y. Yamada, *Contemporary Math.* **297** 151 (2002).
- [7] G. Hatayama, A. Kuniba, and T. Takagi, *Nucl. Phys. B* **577**[PM] 619 (2000), *J. Stat. Phys.* **102** 843 (2001), *J. Phys. A: Math. Gen.* **34** 10697 (2001).
- [8] K. Kajiwara, M. Noumi and Y. Yamada, *Lett. Math. Phys.* **60** 211 (2002).
- [9] S-J. Kang, M. Kashiwara, K. C. Misra, T. Miwa, T. Nakashima and A. Nakayashiki, *Duke Math. J.* **68** 499 (1992).
- [10] S.V.Kerov, A.N.Kirillov and N.Yu.Reshetikhin, *Zap. Nauch. Semin. LOMI.* **155** 50 (1986).
- [11] A. Kuniba and T. Nakanishi, *J. Alg.* **251** 577 (2002).
- [12] A. Kuniba and J. Suzuki, *Commun. Math. Phys.* **173** 225 (1995).
- [13] A. Kuniba and A. Takenouchi, Bethe ansatz at $q = 0$ and periodic box-ball systems, preprint (nlin.SI/0509001).
- [14] J. Mada, M. Idzumi and T. Tokihiro, *J. Math. Phys.* **46** 022701 (2005).
- [15] N. Yu. Reshetikhin, *Sov. Phys. JETP* **57** 691 (1983).
- [16] M. Shimoazono, *J. Algebraic Combin.* **15** 151 (2002).
- [17] D. Takahashi, *Proceedings of the International Symposium on Nonlinear Theory and Its Applications (NOLTA '93)* 555 (1993).
- [18] D. Takahashi and J. Satsuma, *J. Phys. Soc. Jpn.* **59** 3514 (1990).
- [19] D. Yoshihara, F. Yura and T. Tokihiro, *J. Phys. A: Math. Gen.* **36** 99 (2003).

INSTITUTE OF PHYSICS, UNIVERSITY OF TOKYO, TOKYO 153-8902, JAPAN
E-mail address: atsuo@gokutan.c.u-tokyo.ac.jp

INSTITUTE OF PHYSICS, UNIVERSITY OF TOKYO, TOKYO 153-8902, JAPAN
E-mail address: takenouchi@gokutan.c.u-tokyo.ac.jp