

Bessel functions of integer order in terms of hyperbolic functions

V. Bârsan

IFIN, Bucharest-Magurele, Romania

S. Cojocaru

Dipartimento Di Fisica "E. R. Caianiello"

Universita degli Studi di Salerno, Via S. Allende

8408, Baronissi (Sa), Italy and

Institute of Applied Physics, Chişinău 2028, Moldova

February 8, 2020

Abstract

Using Jacobi's identity we derive a simple expression for the Bessel functions of integer order in terms of combinations of powers and hyperbolic functions of the same argument.

In this article we propose a simple formula expressing the modified Bessel functions of integer order, I_n , in terms of powers and hyperbolic functions of the same argument. It can be easily adapted for the Bessel functions J_n .

The starting point is the generalization of the Jacobi identity ([1] p.22) used in the calculation of lattice sums [2]:

$$\frac{1}{N} \sum_{m=0}^{N-1} \exp \left[\frac{z}{2} \left(w e^{i \frac{2\pi m}{N}} + \frac{1}{w} e^{-i \frac{2\pi m}{N}} \right) \right] = \sum_{k=-\infty}^{\infty} w^{kN} \cdot I_{kN}(z). \quad (1)$$

For $w = 1$ and $N = 2$, (1) gives a well-known formula:

$$\cosh z = \sum_{k=-\infty}^{\infty} I_{2k}(z) = I_0(z) + 2[I_2(z) + I_4(z) + \dots], \quad (2)$$

(see for instance [3], eq.9.6.39)

It is interesting to exploit (1) at $w = 1$, for larger values of N . For $N = 4$,

$$\frac{1}{2} (1 + \cosh z) = \cosh^2 \frac{z}{2} = I_0(z) + 2 \sum_{k=1}^{\infty} I_{4k}(z), \quad (3)$$

and for $N = 8$:

$$\frac{1}{4}(1 + \cosh z + 2 \cosh \frac{z}{\sqrt{2}}) = I_0(z) + 2 \sum_{k=1}^{\infty} I_{8k}(z). \quad (4)$$

It is easy to see that, if the l.h.s. of these equations contains p hyperbolic cosines, it provides an exact expression for the series of $I_0(z)$, cut off at the z^{4p} term. The generalization of (4) for $N = 4p$, with p - an arbitrary integer, is indeed:

$$\frac{1}{2p} \left[1 + \cosh z + 2 \cosh \left(z \cos \frac{\pi}{2p} \right) + \dots + 2 \cosh \left(z \cos (p-1) \frac{\pi}{2p} \right) \right] = I_0(z) + 2 \sum_{k=1}^{\infty} I_{4pk}(z). \quad (5)$$

Because

$$I_n(-iz) = i^{-n} J_n(z). \quad (6)$$

our result (5) can be written as:

$$\begin{aligned} \frac{1}{2p} \left[1 + \cos z + 2 \cos \left(z \cos \frac{\pi}{2p} \right) + \dots + 2 \cos \left(z \cos (p-1) \frac{\pi}{2p} \right) \right] = \\ = J_0(z) + 2 \sum_{k=1}^{\infty} J_{4pk}(z). \end{aligned} \quad (7)$$

It is easy to obtain formulae similar to (5) for any modified Bessel function of integer order. Let us introduce the notations:

$$c_1 = \cos \frac{\pi}{2p}, \quad \dots \quad c_{p-1} = \cos \frac{p-1}{2p} \pi. \quad (8)$$

and let us define the functions:

$$S_q(z) = \sinh z + 2c_1^q \sinh(c_1 z) + \dots + 2c_p^q \sinh(c_p z) \quad (9)$$

$$C_q(z) = \cosh z + 2c_1^q \cosh(c_1 z) + \dots + 2c_p^q \cosh(c_p z) \quad (10)$$

We have:

$$\frac{dS_q(z)}{dz} = C_{q+1}(z); \quad \frac{dC_q(z)}{dz} = S_{q+1}(z) \quad (11)$$

Using the formula ([4] 8.486.6):

$$\left(\frac{1}{z} \frac{d}{dz} \right)^n I_0(z) = \frac{I_n(z)}{z^n}, \quad (12)$$

and the notation:

$$T_n(z) = \left(\frac{1}{z} \frac{d}{dz} \right)^n C_0(z), \quad (13)$$

we get:

$$\frac{1}{2p} z^n T_n = I_n(z) + 2z^n \left(\frac{1}{z} \frac{d}{dz} \right)^n \sum_{k=1}^{\infty} I_{4pk}(z). \quad (14)$$

For $n = 1, \dots, 4$:

$$T_1(z) = z^{-1} S_1(z), \quad T_2 = -z^{-3} S_1(z) + z^{-2} C_2(z), \quad (15)$$

$$T_3(z) = 3z^{-5} S_1(z) - 3z^{-4} C_2(z) + z^{-3} S_3(z), \quad (16)$$

$$T_4(z) = -15z^{-7} S_1(z) + 15z^{-6} C_2(z) - 6z^{-5} S_3(z) + z^{-4} C_4(z). \quad (17)$$

The general expressions are:

$$T_{2n} = z^{-2n} \left[\alpha_1^{(2n)} z^{-2n+1} S_1 + \alpha_2^{(2n)} z^{-2n+2} C_2 + \dots + \alpha_{2n}^{(2n)} C_{2n} \right], \quad (18)$$

$$T_{2n+1} = z^{-2n-1} \left[\alpha_1^{(2n+1)} z^{-2n} S_1 + \alpha_2^{(2n+1)} z^{-2n+1} C_2 + \dots + \alpha_{2n+1}^{(2n+1)} S_{2n+1} \right], \quad (19)$$

We get:

$$\alpha_1^{(n)} = -\alpha_2^{(n)} = (-1)^{n+1} (2n-3)!!, \quad \alpha_{n-1}^{(n)} = -\frac{(n-1)n}{2}, \quad \alpha_n^{(n)} = 1. \quad (20)$$

and the following recurrence relations for the coefficients $\alpha_q^{(p)}$:

$$\alpha_{n-p}^{(n)} = \alpha_{n-p-1}^{(n-1)} - (n+p-2) \alpha_{n-p}^{(n-1)}, \quad 2 \leq p \leq n-3. \quad (21)$$

Other general expressions of the coefficients are:

$$\alpha_3^{(n)} = (-1)^{n+1} (n-2) (2n-5)!! \quad (22)$$

$$\alpha_4^{(n)} = (-1)^n (n-3)! \left\{ \frac{2^{n-4}}{0!} 1!! + \frac{2^{n-3}}{1!} 3!! + \dots + \frac{2^0}{(n-4)!} (2n-7)!! \right\}. \quad (23)$$

Ignoring the series in the r.h.s. of (14), we get approximate expression for I_n . Let us give here these expressions for the value $p = 2$ and $n = 0, 1, 2, 3$:

$$I_0^{(ap)}(z) = \frac{1}{4} \cdot \left(1 + \cosh z + 2 \cosh \frac{z}{\sqrt{2}} \right), \quad (24)$$

$$I_1^{(ap)}(z) = \frac{1}{4} \cdot \left(\sinh z + \sqrt{2} \sinh \frac{z}{\sqrt{2}} \right), \quad (25)$$

$$I_2^{(ap)}(z) = \frac{1}{4} \cdot \left(-\frac{1}{z} \left(\sinh z + \sqrt{2} \sinh \frac{z}{\sqrt{2}} \right) + \cosh z + \cosh \frac{z}{\sqrt{2}} \right), \quad (26)$$

$$I_3^{(ap)}(z) = \frac{1}{4} \left[\frac{3}{z^2} \left(\sinh z + \sqrt{2} \sinh \frac{z}{\sqrt{2}} \right) - \frac{3}{z} \left(\cosh z + \cosh \frac{z}{\sqrt{2}} \right) - \frac{3}{z} \left(\cosh z + \cosh \frac{z}{\sqrt{2}} \right) + \left(\sinh z + \frac{1}{\sqrt{2}} \sinh \frac{z}{\sqrt{2}} \right) \right]. \quad (27)$$

According to the Table 1, even for this very small value of p , the approximation provided by these functions, for "moderate" values of the argument ($z \lesssim 4$), is very good.

Table 1

z	1	2	3	4
$\frac{I_0^{(ap)} - I_0}{I_0}$	1.6×10^{-7}	2.4×10^{-5}	3.3×10^{-4}	1.7×10^{-3}
$\frac{I_1^{(ap)} - I_1}{I_1}$	2.3×10^{-6}	1.4×10^{-4}	1.2×10^{-3}	4.4×10^{-3}
$\frac{I_2^{(ap)} - I_2}{I_2}$	7.1×10^{-5}	10^{-3}	4.5×10^{-3}	1.2×10^{-2}
$\frac{I_3^{(ap)} - I_3}{I_3}$	1.8×10^{-3}	7.3×10^{-3}	1.7×10^{-2}	3×10^{-2}

It is visible that the precision of the approximation decreases with the order n of the Bessel function I_n . We can increase it arbitrarily, by increasing the value of p .

The extension of these results to Bessel functions of real argument is trivial, using the formula (6) and:

$$S_q(-iz) = -i \left[\sin z + 2c_1^q \sin(c_1 z) + \dots + 2c_p^q \sin(c_p z) \right], \quad (28)$$

$$C_q(-iz) = \cos z + 2c_1^q \cos(c_1 z) + \dots + 2c_p^q \cos(c_p z). \quad (29)$$

In conclusion, we have proposed a controlled analytic approximation for Bessel functions of integer order. The first $4p - n$ terms of the series representation of I_n is generated exactly by the first $4p - n$ terms of the elementary functions in the l.h.s. of eq.(14). So, our formulae can be used, for instance, to find the series expansions of the powers of Bessel functions, a subject discussed recently by Bender et al [5]

The results presented in this paper can be applied to a large variety of problems, mainly with cylindrical symmetry, involving Bessel functions at "moderate" arguments. They may provide also a useful "visualization" of J_n and I_n

in terms of elementary functions. The method cannot be used for asymptotic problems.

References

- [1] G.N.Watson 1948 A Treatise on the Theory of Bessel Functions (Cambridge University Press)
- [2] Cojocaru S 2005 *Int. J. Mod. Phys.* **20** 593
- [3] Abramowitz M and Stegun I A 1972 Handbook of Mathematical Functions (New York: Dover Publications)
- [4] Gradshteyn I S and Ryzhik I M 1980 Tables of Integrals, Series and Products (New York, London: Academic Press)
- [5] Bender C M, Brody D C and Meister B K 2003 *J.Math.Phys.* **44** 309