

A new integral representation for the Riemann Zeta function

Sandeep Tyagi, Christian Holm

Frankfurt Institute for Advanced Studies, Max-von-Laue-Straße 1, and

D-60438 Frankfurt am Main, Germany

Abstract

A new integral representation for the Riemann zeta function is derived. This new representation of $\zeta(s)$ covers the important region of the complex plane where $0 < \text{Re}[s] < 1$. Using this new representation, we obtain new functional identities for the Riemann zeta function.

I. INTRODUCTION

The Riemann zeta function has an illustrious history of 150 years¹. It arises in many branches of mathematics and, and is also intimately related to the distribution of prime numbers. The most facinating thing related to the Riemann zeta function is the Riemann's hypothesis which states that all non-trivial zeros of the Riemann zeta function $\zeta(s)$ fall on the strip $\text{Re}[s] = 1/2$. This hypothesis has so far not been proven despite the efforts of some of the most brilliant scientists in mathematics such as Riemann, Hilbert, Hardy and Polya¹. There are a number of integral representations of the Riemann zeta function. A few of them cover the important region of $0 \leq \text{Re}[s] \leq 1$. In this paper, we report a new integral representation.

The Riemann zeta function is defined as

$$\zeta(s) = \sum_{n=1}^{\infty} \frac{1}{n^s}$$

However, this definition is valid only in the region $\text{Re}[s] > 1$. By analytical continuation, the definition can be extended to the whole complex plane. The new formula valid for the whole complex plane is

$$\zeta(s) = -\frac{1}{1-2^{(1-s)}} \sum_{n=1}^{\infty} \frac{(-1)^n}{n^s}. \quad (1.1)$$

The Riemann zeta function has just a single simple pole in the whole complex plane which lies at $s = 1$. The $\zeta(s)$ has a number of integral representations; a comprehensive list can be obtained from the mathworld website². Our aim in this paper is to derive a new integral representation and obtain new functional identity using it.

II. A NEW REPRESENTATION

We obtain this representation by rewriting a general term of Eq. (1.1) in an integral form and then taking the sum over n inside the integral sign. We start with Eq. (1.1). We will first work with a special case of $s = 1/2+ib$. In the end, we will use analytical continuation to obtain the result for any $0 < \text{Re}[s] < 2$. For $s = 1/2 + ib$ we have from Eq. (1.1),

$$\zeta(1/2 + ib) = -\frac{1}{1-2^{(1/2-ib)}} \sum_{n=1}^{\infty} (-1)^n \frac{n^{-ib}}{\sqrt{n}}. \quad (2.1)$$

At this point, we cannot proceed any further as the summation over n cannot be performed in a straightforward way. We have to re-express the term n^{-ib}/\sqrt{n} in Eq. (2.1) so that the summation over n can be carried out. To this end, we would like to have an integral involving the dummy index x and the parameters n and b such that this integral is equal to n^{-ib}/\sqrt{n} apart from some factors independent of n . To achieve such an integral, we can try the complex residue theorem. It dictates that the denominator should have a form of $x^2 + n$ or $x^4 + n^2$ because then the residues would give rise to terms having \sqrt{n} . One can go ahead and try it with $x^2 + n$, but that leads to difficulties in the series summation later on. So, let us try with $x^4 + n^2$, which does lead to a sensible series. To get rid of some other unwanted terms that would arise out of the presence of residues, we consider the integral

$$\int_{-\infty}^{+\infty} \frac{x^2 x^{-2ib}}{x^4 + n^2} dx. \quad (2.2)$$

This integral has been chosen such that when x^2 is substituted by \sqrt{n} , then the x^2 in the numerator will be cancelled by a term arising in the denominator and the x^{-2ib} would give rise to n^{-ib} . Finally a term of \sqrt{n} would arise as a result of the other residue. The integral in Eq. (2.3) can be easily evaluated using the complex residue theorem by choosing an appropriate contour. Even though we have assumed b to be real when writing the integral in Eq. (2.3), it turns out that the integral is defined for all b in the range of $-3/2 < \text{Im}[b] < 1/2$. The integral can be performed to yield

$$\int_{-\infty}^{+\infty} \frac{x^2 x^{-2ib}}{x^4 + n^2} dx = \frac{\pi n^{-ib}}{2 \sqrt{n}} \csc\left(\frac{\pi}{2}(1/2 + ib)\right) \quad -3/2 < \text{Im}[b] < 1/2. \quad (2.3)$$

Combining Eq. (2.1) and (2.3) we obtain

$$\zeta(1/2 + ib) = -\frac{2 \sin\left(\frac{\pi}{2}(1/2 + ib)\right)}{\pi (1 - 2^{(1/2-ib)})} \sum_{n=1}^{\infty} (-1)^n \int_{-\infty}^{+\infty} \frac{x^2 x^{-2ib}}{x^4 + n^2} dx. \quad (2.4)$$

The transformation above has led to a form where the summation over n can now be performed using well known results. We can interchange the summation and integration and use the summation formula,

$$\sum_{n=1}^{\infty} \frac{(-1)^n}{x^4 + n^2} = \frac{\pi x^2 \csc(\pi x^2) - 1}{2x^4}, \quad (2.5)$$

to write

$$\zeta(1/2 + ib) = \frac{2}{(1 - 2^{(1/2-ib)})} \sin\left(\frac{\pi}{2}(1/2 + ib)\right) \int_0^{+\infty} \frac{x^{-2ib}}{\pi x^2} (1 - \pi x^2 \csc(\pi x^2)) dx. \quad (2.6)$$

Note that the integral is symmetrical and for this reason we have written

$$\int_{-\infty}^{+\infty} \dots dx = 2 \int_0^{+\infty} \dots dx.$$

Eq. (2.6) is our desired integral representation. It is defined only for $\text{Re}[s] = 1/2$, although the integral in Eq. (2.2) is valid for $-3/2 < \text{Im}[b] < 1/2$. Writing $b = b' + iq$ where $-3/2 < q < 1/2$ and b' is real, we can express

$$1/2 + ib = (1/2 - q) + ib'.$$

Thus the real part of $s = 1/2 + ib$ actually changes between 0 and 2 as q changes varies from $-3/2$ and $1/2$. Substituting $s = 1/2 + ib$ in Eq. (2.6) we obtain

$$\zeta(s) = \frac{2}{(1 - 2^{1-s})} \sin\left(\frac{\pi s}{2}\right) \int_0^{\infty} \frac{x^{-2s}}{\pi x} (1 - \pi x^2 \csc[\pi x^2]) dx \quad 0 < \text{Re}[s] < 2. \quad (2.7)$$

Changing variable $\pi x^2 = y$ we can write

$$\zeta(s) = \frac{\pi^{s-1}}{(1 - 2^{1-s})} \sin\left(\frac{\pi s}{2}\right) \int_0^{\infty} y^{-s} \left(\frac{1}{y} - \frac{1}{\sinh(y)}\right) dy \quad 0 < \text{Re}[s] < 2. \quad (2.8)$$

Note that Eqs. (2.7) and (2.8) are valid for $0 < \text{Re}[s] < 2$. However, the Riemann zeta function has a simple pole on $s = 1$. Thus, these equations make sense only when $\text{Im}[s] \neq 0$ for $\text{Re}[s] = 1$.

There are two formulas in the literature that resemble our formula in Eq. (2.8). The first one is

$$\zeta(s) = \frac{2^{s-1}}{(1 - 2^{1-s}) \Gamma(s)} \int_0^{\infty} y^{s-1} \frac{\exp(-y)}{\cosh(y)} dy \quad \text{Re}[s] > 0 \quad (2.9)$$

$$= \frac{2^{s-1}}{(1 - 2^{1-s}) \Gamma(s)} \int_0^{\infty} \frac{y^{s-1}}{\exp(y) + 1} dy \quad \text{Re}[s] > 0, \quad (2.10)$$

where Γ stands for the Gamma function. The second formula is due to Ramanujan¹ and it is given by

$$\zeta(s) = \frac{\sin(\pi s)}{\pi} \int_0^{\infty} y^{-s} \left(\log(y) - \frac{\Gamma'(y)}{\Gamma(y)}\right) dy \quad 0 \leq \text{Re}[s] < 1. \quad (2.11)$$

There is another way to view 2.8. The $\sinh(y)$ in denominator will lead to poles at integers with residues $(-1)^n$. The presence of $1/y$ is basically to remove the pole at $n = 0$. Thus 2.8 can be derived using residue theorem as well.

III. NEW FUNCTIONAL IDENTITY

So far there have been only a few functional forms and identities satisfied by the Riemann zeta. One of the most important functional relation is given by

$$\Gamma\left(\frac{s}{2} - 1\right) \pi^{-\frac{s}{2}} \zeta(s) = \Gamma\left(\frac{1-s}{2} - 1\right) \pi^{-\frac{1-s}{2}} \zeta(1-s). \quad (3.1)$$

Our representation leads to other functional identities involving the Riemann zeta and Gamma functions. To derive a new functional representation we use the integral formula

$$\int_0^\infty \frac{y^k}{\sinh(y)} dy = (2 - 2^{-k}) \Gamma(1+k) \zeta(1+k) \quad \text{where } k > 0. \quad (3.2)$$

We can transform our general representation given in Eq. (2.8) into this form by using the series expansion

$$\sinh(y) - y = \frac{y^3}{3!} + \frac{y^5}{5!} + \dots \quad (3.3)$$

This allows us to write Eq. (2.8) as

$$\begin{aligned} \zeta(s) &= \frac{\pi^{s-1}}{(1-2^{1-s})} \sin\left(\frac{\pi s}{2}\right) \int_0^\infty y^{-s-1} \left(\frac{\sinh(y) - y}{\sinh(y)}\right) dy \\ &= \frac{\pi^{s-1}}{(1-2^{1-s})} \sin\left(\frac{\pi s}{2}\right) \sum_{n=1}^\infty \frac{1}{(2n+1)!} \int_0^\infty y^{-s-1} \left(\frac{y^{(2n+1)}}{\sinh(y)}\right) dy. \end{aligned} \quad (3.4)$$

Inserting the integral formula of Eq. (3.2) leads to the following new functional representation of the Riemann zeta function:

$$\zeta(s) = \frac{\pi^{s-1}}{(1-2^{1-s})} \sin\left(\frac{\pi s}{2}\right) \sum_{n=1}^\infty \frac{(2-2^{-2n+s})}{(2n+1)!} \Gamma(2n-s+1) \zeta(2n-s+1). \quad (3.5)$$

A part of the series in Eq. (3.5) can be summed up analytically:

$$\begin{aligned} \sum_{n=1}^\infty \frac{(2-2^{-2n+s})}{(2n+1)!} \Gamma(2n-s+1) &= [-1 + 3^s + s(2^s - 2)] \Gamma(-s) \\ &\quad + \sqrt{\pi} \frac{\Gamma(1-s) \Gamma(s)}{\Gamma\left(1 + \frac{s}{2}\right) \Gamma\left(\frac{1+s}{2}\right)}. \end{aligned} \quad (3.6)$$

Using the above relation we can obtain the following functional relation

$$\begin{aligned} \zeta(s) &= \frac{\pi^{s-1}}{(1-2^{1-s})} \sin\left(\frac{\pi s}{2}\right) \left(\sum_{n=1}^\infty \frac{(2-2^{-2n+s})}{(2n+1)!} \Gamma(2n-s+1) [\zeta(2n-s+1) - 1] \right. \\ &\quad \left. + [-1 + 3^s + s(2^s - 2)] \Gamma(-s) + \sqrt{\pi} \frac{\Gamma(1-s) \Gamma(s)}{\Gamma\left(1 + \frac{s}{2}\right) \Gamma\left(\frac{1+s}{2}\right)} \right). \end{aligned} \quad (3.7)$$

Identity in Eq. (3.7) appears to be new.

IV. CONCLUSION

In this work we have derived an integral representation for the Riemann zeta function. Using this representation, we are able to obtain a new identity for the Riemann zeta function. This representation allows us to express the Riemann zeta function in terms of the Riemann zeta and Gamma functions. Note that similar identities have been discovered by Milgram³.

¹ H. M. Edwards, *Riemann zeta function*, New York, Dover 2001

² <http://functions.wolfram.com/ZetaFunctionsandPolylogarithms/Zeta/07/01/01/>

³ M. S. Milgram, Integral and Series Representations of Riemann's Zeta function, Dirichelet's Eta Function and a Medley of Related Results, <https://arxiv.org/abs/1208.3429>