## Five-Dimensional Tangent Vectors in Space-Time

Alexander Krasulin

Institute for Nuclear Research of the Russian Academy of Sciences 60th October Anniversary Prospect, 7a, 117312 Moscow, Russia

## Abstract

This article is a summary of a series of papers where I examine a special kind of geometric objects that can be defined in space-time — five-dimensional tangent vectors. Similar objects exist in any other differentiable manifold, and their dimension is one unit greater than that of the manifold. Like ordinary tangent vectors, the considered five-dimensional vectors and the tensors constructed out of them can be used for describing certain local quantities and in this capacity find direct application in physics. For example, such familiar physical quantities as the stress-energy and angular momentum tensors prove to be parts of a single five-tensor. In this paper I describe several different mathematical definitions of five-dimensional tangent vectors, discuss their basic algebraic and differential properties, and speak about their possible application in the theory of gravity and in gauge theories.

1. Adding a dimension to tangent vectors in spacetime is not a new idea in physics. A well-known example is the Kaluza–Klein model [1] and the models that succeeded it, where the extra dimension of tangent vectors results from adding a dimension to the space-time manifold itself. Another example are the theories of gravity formulated as Yang–Mills gauge theories of the de Sitter group [2] and similar models, where the additional dimension is assigned not to the tangent vectors themselves, but to the internal vector space where the vierbein field takes its values. Unlike all these constructions, for introducing the five-dimensional vectors I consider in this paper one does not need to change the space-time manifold in any way nor to endow it with any additional structure. The vectors I am going to discuss here, which I will call *five-dimensional tangent vectors* or simply *five-vectors*, should be viewed as another type of geometric objects that can be defined in space-time and which are more suited for describing certain kinds of geometric and physical quantities than ordinary tangent vectors and tensors.

A hint to the existence of five-dimensional tangent vectors can be found in spinors. For the type of 4-spinors commonly used in physics, the symmetry group of the corresponding Clifford algebra is SO(3,2). Accordingly, there exist five constituents of the Clifford algebra (five matrices)  $\Gamma_A$ , where A runs 0, 1, 2, 3, and 5, that all transform alike under Dirac and charge conjugation:

$$\bar{\Gamma}_A = \Gamma_A \text{ and } \Gamma_A^c = \Gamma_A,$$
(1)

and that satisfy the following anticommutation relations:

$$\Gamma_A \Gamma_B + \Gamma_B \Gamma_A = -2 \eta_{AB}, \qquad (2)$$

where  $\eta_{AB} \equiv \text{diag}(+1, -1, -1, -1, +1)$ . It is evident that one can obtain a new set of five constituents satisfying the same conjugation and anticommutation relations by applying an arbitrary O(3,2) transformation to the original set. Moreover, any two sets of constituents that satisfy relations (1) and (2) prove to be connected by an O(3,2) transformation. For an appropriate choice of the constituent set, the standard  $\gamma$ -matrices (the ones identified with the components of the basis four-dimensional tangent vectors) are expressed in terms of  $\Gamma_A$ 's as

$$\gamma_{\mu} = \frac{i}{2} (\Gamma_{\mu} \Gamma_5 - \Gamma_5 \Gamma_{\mu}), \qquad (3)$$

where  $\mu = 0, 1, 2, \text{ or } 3$ .

These observations may give one the idea to consider a new type of vectors that make up a real fivedimensional vector space endowed with a symmetric nondegenerate inner product with the signature (+---+) or (-+++-). Considering the relation that exists between multiplication in a Clifford algebra and exterior multiplication of multivectors and forms, on the grounds of equation (3) one may further suppose that there should exist a certain correspondence between four-dimensional tangent vectors and part of the bivectors constructed from elements of the mentioned five-dimensional vector space. It is apparent that these latter bivectors should be of the form  $\mathbf{u} \wedge \mathbf{e}$ , where  $\mathbf{u}$  is arbitrary and  $\mathbf{e}$  belongs to a fixed one-dimensional subspace whose elements have a norm squared of such a sign that the inner product induced on the subspace of all such bivectors is of Lorentz type.

Basing on these assumptions one can make a formal study of the basic algebraic and differential properties of five-dimensional tangent vectors, as it is done in part I of the long version [3]. Though not really necessary, this formal analysis may serve as a guide in developing a more sophisticated theory of five-vectors basing on the principles of differential geometry, as it is done in part II of the long version [4]. Within this latter theory five-dimensional tangent vectors are introduced either as equivalence classes of parametrized curves or, more rigorously, as a particular kind of differential-algebraic operators that act upon scalar functions. The first of these representations is obtained as follows.

Consider a set of all smooth parametrized curves that go through an arbitrary space-time point Q. In an evident way, for any such curve  $\mathcal{A}$  one can evaluate the derivative of any smooth scalar function f defined in the vicinity of Q, and I will denote this derivative as  $\partial_{\mathcal{A}} f|_Q$ . Let us now focus our attention on the behaviour of the curves in the infinitesimal vicinity of Q. From that point of view, the considered set can be divided into classes of equivalent curves that coincide in direction or in direction and parametrization. One can consider three degrees to which two given curves,  $\mathcal{A}$  and  $\mathcal{B}$ , may coincide:

i. The two curves come out of Q in the same direction. A more precise formulation is the following: there exists a real positive number a such that for any scalar function f

$$\partial_{\mathcal{A}} f|_Q = a \cdot \partial_{\mathcal{B}} f|_Q. \tag{4}$$

ii. The two curves come out of Q in the same direction and in the vicinity of Q their parameters change with equal rates. More precisely: for any scalar function f

$$\partial_{\mathcal{A}} f|_Q = \partial_{\mathcal{B}} f|_Q. \tag{5}$$

iii. The two curves come out of Q in the same direction; their parameters,  $\lambda_{\mathcal{A}}$  and  $\lambda_{\mathcal{B}}$ , change with equal rates in the vicinity of Q; and the values of these parameters at Q are the same. This means that

$$\lambda_{\mathcal{A}}(Q) = \lambda_{\mathcal{B}}(Q) \tag{6a}$$

and for any scalar function f

$$\partial_{\mathcal{A}} f|_Q = \partial_{\mathcal{B}} f|_Q. \tag{6b}$$

It is evident that relations (4), (5) and (6) are all equivalence relations on the considred set of curves, and for each of them one can introduce the corresponding quotient set—the set whose elements are classes of equivalent curves.

Relation (4) is of no interest to us and I will not consider it any further.

Let us denote the elements of the quotient set corresponding to relation (5) with capital boldface Roman letters: **A**, **B**, **C**, etc. According to relation (5), the derivative of any scalar function f at Q is the same for all the curves belonging to a given class **A**, so it makes sense to introduce the notation  $\partial_{\mathbf{A}} f|_Q$ . In a natural way, one can define the addition of two equivalence classes **A** and **B** and the product of an equivalence class **A** and a real number k:  $\mathbf{A} + \mathbf{B}$  and  $k\mathbf{A}$  are such equivalence classes that for any scalar function f

$$\partial_{\mathbf{A}+\mathbf{B}}f|_Q = \partial_{\mathbf{A}}f|_Q + \partial_{\mathbf{B}}f|_Q$$

and

$$\partial_{k\mathbf{A}}f|_Q = k \cdot \partial_{\mathbf{A}}f|_Q.$$

With thus defined addition and multiplication by a real number, the set of all equivalence classes corresponding to relation (5) becomes a vector space, which I will denote as  $V_4$ . It is evident that its elements can be identified with ordinary (four-dimensional) tangent vectors, and in the following I will refer to them as to four-vectors.

In a similar manner one can deal with the quotient set associated with relation (6). Let us denote its elements with lower-case boldface Roman letters: **a**, **b**, **c**, etc. As in the case of four-vectors, one can introduce the notation  $\partial_{\mathbf{a}} f|_Q$  for the common value of the derivatives of any scalar function f along all the curves belonging to a given equivalence class **a**. Likewise, the common value of the parameters of all these curves at Q will be denoted as  $\lambda_{\mathbf{a}}(Q)$ . One can then give the following definition to the sum of two equivalence classes **a** and **b** and to the product of an equivalence class **a** and **a** real number k:  $\mathbf{a} + \mathbf{b}$  and  $k\mathbf{a}$  are such equivalence classes that

$$\begin{split} \lambda_{\mathbf{a}+\mathbf{b}}(Q) &= \lambda_{\mathbf{a}}(Q) + \lambda_{\mathbf{b}}(Q) \\ \lambda_{k\mathbf{a}}(Q) &= k \cdot \lambda_{\mathbf{a}}(Q) \end{split}$$

and for any scalar function  $\boldsymbol{f}$ 

$$\partial_{\mathbf{a}+\mathbf{b}} f|_Q = \partial_{\mathbf{a}} f|_Q + \partial_{\mathbf{b}} f|_Q$$
$$\partial_{k\mathbf{a}} f|_Q = k \cdot \partial_{\mathbf{a}} f|_Q.$$

These two operations turn the quotient set associated with relation (6) into a vector space whose dimension is evidently five and which I will denote as  $V_5$ . By examining the properties of these five-dimensional vectors more closely, one can show that there indeed exists a natural isomorphism (actually, two of them) between the space of ordinary tangent vectors and the subspace of all bivectors of the form  $\mathbf{u} \wedge \mathbf{e}$ , where  $\mathbf{e}$  is an element of a certain distinguished one-dimensional subspace in  $V_5$ .

A still more rigorous way of introducing fivedimensional tangent vectors is similar to how one introduces ordinary tangent vectors in modern differential geometry, i.e. by identifying the fields of the latter with a particular kind of operators that act upon the scalar functions from a set  $\Im$  which determines the topological and differential properties of the manifold. Each five-vector field **u** is defined as a map

 $\mathbf{u}:\Im\to\Im$ 

that satisfies the following three requirements:

$$\mathbf{u}[k] = v \cdot k \text{ for any constant } k \in \mathfrak{I},$$
  
where  $v \in \mathfrak{I}$  is characteristic of  $\mathbf{u}$ ,  
$$\mathbf{u}[f+g] = \mathbf{u}[f] + \mathbf{u}[g] \text{ for any } f, g \in \mathfrak{I},$$
  
$$\mathbf{u}[fg] = \mathbf{u}[f] \cdot g + f \cdot \mathbf{u}[g] - \mathbf{u}[1]fg \text{ for}$$
  
any  $f, g \in \mathfrak{I},$  where 1 is the constant  
unity function. (7)

Similar to the case of four-vector fields, one can prove a theorem that in any local coordinate system each such map can be presented as the following differential-algebraic operator:

$$\mathbf{u} = u^{\alpha} (\partial/\partial x^{\alpha}) + u^5 \cdot \mathbf{1},\tag{8}$$

where  $\partial/\partial x^{\alpha}$  are derivatives along coordinate lines, **1** is the identity operator, and  $u^A$  are scalar functions from  $\Im$ . As in the case of ordinary tangent vectors, tangent five-vectors at a given point Q can be defined as equivalence classes of the above maps with respect to the equivalence relation

$$\mathbf{u} \equiv \mathbf{v} \Leftrightarrow \mathbf{u}[f](Q) = \mathbf{v}[f](Q)$$
 for any  $f \in \Im$ ,

and it is a simple matter to show that at every point there exists a natural isomorphism between the fivedimensional tangent vectors defined this way and the five-dimensional tangent vectors defined as elements of the quotient set associated with relation (6).

2. Let us now briefly discuss the basic algebraic properties of five-dimensional tangent vectors. To be definite, I will assume that the latter are defined as equivalence classes of parametrized curves, as it has been described above.

As any other vector space,  $V_5$  is completely isotropic with respect to its two composition laws and has no distinguished direction nor any other distinguished subspace of nonzero dimension. However, one *can* distinguish two subspaces in  $V_5$  by associating them with certain classes of parametrized curves.

Consider all those curves at Q for which  $\partial f|_Q = 0$ for any scalar function f. It is evident that all of them belong to the same equivalence class with respect to relation (5) and that this class is the zero vector in  $V_4$ . With respect to relation (6), the considered curves belong to equivalence classes that make up a onedimensional subspace in  $V_5$ , which will be denoted as  $\mathcal{E}$ . In the language of operators this means that  $\mathcal{E}$  is made up by all those five-vectors which at the considered point are represented by purely algebraic operators.

Another distinguished subspace in  $V_5$  can be obtained by considering all those curves for which  $\lambda(Q) = 0$ . The four-vectors corresponding to these curves are all the vectors of  $V_4$ . The corresponding five-vectors make up a four-dimensional subspace in  $V_5$ , which will be denoted as  $\mathcal{Z}$ . It is apparent that this subspace is made up by all those fivevectors which at the considered point are represented by purely differential operators and that  $V_5$  is the direct sum of  $\mathcal{E}$  and  $\mathcal{Z}$ .

From the definition of four- and five-vectors it follows that there exists a set-theoretic relation between  $V_4$  and  $V_5$ : the former is the quotient set corresponding to the following equivalence relation on  $V_5$ :

$$\mathbf{a} \equiv \mathbf{b} \Leftrightarrow \partial_{\mathbf{a}} f|_Q = \partial_{\mathbf{b}} f|_Q$$
 for any scalar function  $f$ .

Denoting this relation as R, one has  $V_4 = V_5/R$ . The fact that  $\mathbf{A}$  is the equivalence class of  $\mathbf{a}$  will be denoted as  $\mathbf{a} \in \mathbf{A}$ . From the definition of the symbols  $\partial_{\mathbf{a}}$  and  $\partial_{\mathbf{A}}$  it follows that  $\mathbf{a} \in \mathbf{A}$  iff  $\partial_{\mathbf{a}} = \partial_{\mathbf{A}}$ . It is easy to see that R can be reformulated as follows:

$$\mathbf{a} \equiv \mathbf{b} \pmod{R} \Leftrightarrow \mathbf{a} = \mathbf{b} + \mathbf{e}, \text{ where } \mathbf{e} \in \mathcal{E}.$$

The latter condition is equivalent to **a** and **b** having equal components in the four-dimensional subspace  $\mathcal{Z}$ . This means that there exists a one-to-one correspondence between five-vectors from  $\mathcal{Z}$  and fourvectors and that this correspondence is a homomorphism.

A typical five-vector basis will be denoted as  $\mathbf{e}_A$ , where A (as all capital latin indices) runs 0, 1, 2, 3, and 5. One can choose a basis in  $V_5$  arbitrarily, but it is more convenient to select the fifth basis vector belonging to  $\mathcal{E}$ . Such bases will be called *standard* and will be used in all calculations.<sup>1</sup>

<sup>&</sup>lt;sup>1</sup>As one can see, the basis vector and vector components related to the fifth dimension are labled with the index 5 rather than 4. This corresponds to the index convention used for

The basis in  $V_4$  can be chosen arbitrarily and independently of the basis in  $V_5$ . It is more convenient though to associate it with the five-vector basis. A natural choice is to take  $\mathbf{E}_{\alpha}$  to be the equivalence classes of the basis five-vectors  $\mathbf{e}_{\alpha}$  (the equivalence class of  $\mathbf{e}_5$  is the zero four-vector). I will refer to this basis as to the one *associated* with the basis  $\mathbf{e}_A$  in  $V_5$ .

It is also convenient to introduce the notions of a *regular* basis and of a *coordinate* five-vector basis. By definition, the former is a standard five-vector basis whose first four elements belong to  $\mathcal{Z}$  and the fifth basis vector is normalized in some particular way. A coordinate five-vector basis is one for which the associated four-vector basis is a coordinate basis in the usual sense.

If  $\mathbf{e}_A$  and  $\mathbf{e}'_A$  are two standard bases in  $V_5$  and  $\mathbf{e}'_A = \mathbf{e}_B L^B_A$ , then  $L^B_A$  can be shown to satisfy the condition

$$L_5^{\alpha} = 0$$
 for all  $\alpha$ .

The corresponding equivalence classes are related as  $\mathbf{E}'_{\alpha} = \mathbf{E}_{\beta} L^{\beta}_{\alpha}$ .

As for ordinary tangent vectors, for five-vectors the Riemannian metric of space-time fixes a certain symmetric inner product, which I will denote as g, too. This inner product is such that for any two fivevectors **u** and **v** 

$$g(\mathbf{u}, \mathbf{v}) = g(\mathbf{U}, \mathbf{V}),$$

where **U** and **V** are such four-vectors that  $\mathbf{u} \in \mathbf{U}$ and  $\mathbf{v} \in \mathbf{V}$ . From the latter equation it follows that g is degenerate on  $V_5$ . It is a simple matter to see that the subspace of all degenerate five-vectors for it coincides with  $\mathcal{E}$  and that g is nondegenerate within any subspace complementary to  $\mathcal{E}$ .

It is not difficult to construct from g a *non*degenerate inner product on  $V_5$ . For that one should consider another natural measure that exists for five-vectors: to each five-vector  $\mathbf{u}$  one can put into correspondence the value of the relevant curve parameter,  $\lambda_{\mathbf{u}}$ . If one then interprets this latter number as the length of vector  $\mathbf{u}$ , one will obtain another inner product, which for any  $\mathbf{u}$  and  $\mathbf{v}$  can be shown to equal  $\lambda_{\mathbf{u}} \cdot \lambda_{\mathbf{v}}$ . Consequently, the subspace of all degenerate vectors for this latter inner product coincides with  $\mathcal{Z}$  and it is nondegenerate within any (one-dimensional) subspace complementary to  $\mathcal{Z}$ .

One should now notice that the subspaces of degenerate vectors for the two considered inner products are complementary to each other, which means that their sum will be a nondegenerate inner product on  $V_5$ . The only problem in constructing such a sum is that for any **u** and **v** the quantities  $g(\mathbf{u}, \mathbf{v})$  and  $\lambda_{\mathbf{u}}\lambda_{\mathbf{v}}$  are of *different* dimension. Therefore, to construct a nondegenerate inner product for five-vectors, one needs some dimensional constant,  $\xi$ , which would play a role similar to that of the speed of light: it would establish a relation between different units used to measure the same quantity. The resulting inner product, measured in the same units as g, will be

$$h(\mathbf{u}, \mathbf{v}) \equiv g(\mathbf{u}, \mathbf{v}) + \xi \,\lambda_{\mathbf{u}} \lambda_{\mathbf{v}}.\tag{9}$$

Having selected the constant  $\xi$  in the latter formula somehow, one is then able to fix the length of the fifth basis vector in a regular five-vector basis by requiring that  $h(\mathbf{e}_5, \mathbf{e}_5) = \operatorname{sign} \xi$ . In the following, such a regular five-vector basis will be called *normalized*.

As in the case of any other type of vectors, one can consider linear forms corresponding to five-vectors. Such forms will be denoted with lower-case boldface Roman letters with a tilde:  $\tilde{\mathbf{a}}, \tilde{\mathbf{b}}, \tilde{\mathbf{c}}$ , etc., and their space will be denoted as  $\tilde{V}_5$ . To distinguish a *p*form associated with five-vectors from a *p*-form associated with four-vectors the former will be called a *five-vector p*-form and the latter a *four-vector p*-form.

Five-vector 1-forms have all the properties common to linear forms in general. In addition, they have several specific features which are due to their association with five-vectors.

The existence of two distinguished subspaces in  $V_5$ results in the existence of two distinguished subspaces in  $\tilde{V}_5$ . The first of these subspaces is made up by all those 1-forms from  $\tilde{V}_5$  whose contraction with any five-vector from  $\mathcal{E}$  is zero. It is evident that this subspace is four-dimensional, and I will denote it as  $\tilde{\mathcal{Z}}$ . The other distinguished subspace is made up by all those 1-forms that have a zero contraction with any five-vector from  $\mathcal{Z}$ . This subspace is one-dimensional, and I will denote it as  $\tilde{\mathcal{E}}$ . It is evident that  $\tilde{V}_5$  is the direct sum of  $\tilde{\mathcal{Z}}$  and  $\tilde{\mathcal{E}}$ .

If  $\mathbf{e}_A$  is a standard five-vector basis and  $\widetilde{\mathbf{o}}^A$  is the corresponding dual basis of five-vector 1-forms, then  $\widetilde{\mathbf{o}}^{\alpha} \in \widetilde{\mathcal{Z}}$  for all  $\alpha$ . The fifth basis 1-form will not necessarily be an element of  $\widetilde{\mathcal{E}}$ : this will be the case only if all  $\mathbf{e}_{\alpha} \in \mathcal{Z}$ . The same conclusions follow from the transformation formulae for the dual basis of 1-forms, corresponding to the transformation  $\mathbf{e}'_A = \mathbf{e}_B L^B_A$  from one standard five-vector basis to another: since in this case  $(L^{-1})^{\alpha}_{5} = 0$ , one has

$$\widetilde{\mathbf{o}}'^{\,\alpha} = (L^{-1})^{\alpha}_{\,B} \, \widetilde{\mathbf{o}}^{B} = (L^{-1})^{\alpha}_{\,\beta} \, \widetilde{\mathbf{o}}^{\beta}$$
$$\widetilde{\mathbf{o}}'^{\,5} = (L^{-1})^{5}_{\,5} \, \widetilde{\mathbf{o}}^{5} + (L^{-1})^{5}_{\,\beta} \, \widetilde{\mathbf{o}}^{\beta}.$$

but

 $<sup>\</sup>gamma$ -matrices, where the notation  $\gamma_4$  is reserved for the timelike  $\gamma$ -matrix in the Pauli metric:  $\gamma_4 = i\gamma_0$ . This also better suits the words "fifth dimension", and accentuates the fact that this direction in  $V_5$  is distinguished as being the one that corresponds to the one-dimensional subspace  $\mathcal{E}$ .

The fact that  $\mathcal{Z}$  is isomorphic to  $V_4$  enables one to establish a natural isomorphism between  $\widetilde{\mathcal{Z}}$  and the space of four-vector 1-forms, which will be denoted as  $\widetilde{V}_4$ . Namely, to each five-vector 1-form  $\widetilde{\mathbf{w}}$  from  $\widetilde{\mathcal{Z}}$ one can put into correspondence such a four-vector 1-form  $\widetilde{\mathbf{W}}$  that for any five-vector  $\mathbf{u} \in \mathcal{Z}$  one would have  $\langle \widetilde{\mathbf{w}}, \mathbf{u} \rangle = \langle \widetilde{\mathbf{W}}, \mathbf{U} \rangle$  where  $\mathbf{u} \in \mathbf{U}$ . It is evident that this isomorphism can be extended to a map of  $\widetilde{V}_5$  onto  $\widetilde{V}_4$ , which will be a homomorphism but will not be a one-to-one correspondence.

As in the case of any other vector space, each inner product on  $V_5$  defines a certain correspondence between five-vectors and five-vector 1-forms. Since one has two inner products on  $V_5 - g$  and h, there are two such correspondences, which will be denoted as  $\vartheta_g$  and  $\vartheta_h$ , respectively. By definition,  $\vartheta_g(\mathbf{u})$  is such a five-vector 1-form that

$$\langle \vartheta_q(\mathbf{u}), \mathbf{v} \rangle = g(\mathbf{u}, \mathbf{v}) \text{ for any } \mathbf{v} \in V_5.$$

The definition of the 1-form  $\vartheta_h(\mathbf{u})$  is similar. It is evident that both  $\vartheta_g$  and  $\vartheta_h$  are linear maps of  $V_5$ into  $\widetilde{V}_5$ . If  $u^A$  are components of some five-vector **u** in a certain five-vector basis, then the components of  $\vartheta_g(\mathbf{u})$  and  $\vartheta_h(\mathbf{u})$  in the corresponding dual basis of 1-forms are  $g_{AB}u^B$  and  $h_{AB}u^B$ , respectively. Since the matrix  $h_{AB}$  is nondegenerate, this means that  $\vartheta_h$  is a one-to-one correspondence and is a map of  $V_5$  onto  $V_5$ . It is also easy to see that  $\vartheta_h(\mathcal{Z}) = \mathcal{Z}$ and  $\vartheta_h(\mathcal{E}) = \widetilde{\mathcal{E}}$ . By contrast,  $\vartheta_q$  is neither a oneto-one correspondence nor a surjection. It is evident that  $\vartheta_q(\mathbf{u}) = \vartheta_q(\mathbf{u}^{\mathcal{Z}}) = \vartheta_h(\mathbf{u}^{\mathcal{Z}})$ , so  $\vartheta_q(\mathcal{Z}) = \widetilde{\mathcal{Z}}$ , but  $\vartheta_q(\mathcal{E}) = \{\mathbf{0}\}$ . Consequently, one can use  $g_{AB}$  only to lower five-vector indices. Raising indices with  $q_{AB}$ is possible only if one confines oneself to five-vectors from  $\mathcal{Z}$  and to 1-forms from  $\widetilde{\mathcal{Z}}$ .

3. Let us now turn to the differential properties of five-vectors. As for any other type of vector-like objects considered in space-time, one can speak of parallel transport of five-vectors from one space-time point to another. One can then define the covariant derivative of five-vector fields; introduce the connection coefficients corresponding to a given five-vector basis; construct the corresponding curvature tensor; etc. In doing all this one does not have to use in any way the fact that five-vectors are associated with space-time by their definition.

One should expect that the origin of five-vectors manifests itself in that the rules of their parallel transport are related in some way to similar rules for fourvectors and, possibly, to the Riemannian geometry of space-time. It is obvious that this relation cannot be derived from the algebraic properties of five-vectors, and to obtain it one has to make some new assumptions about five-vectors, which ought to be regarded as part of their definition.

Let us first consider the relation between the rules of parallel transport for four- and five-vectors. The simplest and the most natural form of this relation is obtained by postulating that parallel transport preserves the set-theoretic relation between four- and five-vectors considered above. A more precise formulation of this statement is the following:

If four-vector **U** is the equivalence class of five-vector **u**, then the transported **U** is the equivalence class of the transported **u**. (10)

This assumption is quite natural considering that  $\mathbf{u} \in \mathbf{U}$  means that  $\mathbf{u}$  and  $\mathbf{U}$  correspond to the same direction in the manifold. It has two consequences, which can be conveniently expressed in terms of connection coefficients. Let us define the latter for five-vectors as

$$\nabla_{\mu} \mathbf{e}_A = \mathbf{e}_B G^B_{A\mu}$$

where  $\nabla_{\mu} \equiv \nabla_{\mathbf{E}_{\mu}}$  denotes the covariant derivative in the direction of the basis four-vector  $\mathbf{E}_{\mu}$ . The connection coefficients for four-vectors will be denoted in the usual way:

$$\nabla_{\mu} \mathbf{E}_{\alpha} = \mathbf{E}_{\beta} \Gamma^{\beta}_{\alpha \mu}.$$

Let us consider the parallel transport of vectors from an arbitrary point Q to a nearby point Q'.

If two five-vectors at Q belong to the same equivalence class, then according to assumption (10), the transported five-vectors should also be equivalent. Since parallel transport is a linear operation, this means that vectors from  $\mathcal{E}_{\text{at }Q}$  are transported into vectors from  $\mathcal{E}_{\text{at }Q'}$ . Consequently, in *any* standard five-vector basis,

$$G^{\alpha}_{5\mu} = 0.$$
 (11)

Let  $\mathbf{e}_A$  be an arbitrary standard five-vector basis and let  $\mathbf{E}_{\alpha}$  be the associated basis of four-vectors. If  $\mathbf{E}_{\alpha}(Q)$  are transported into vectors  $\mathbf{E}_{\beta}(Q')C_{\alpha}^{\beta}$ , then according to assumption (10),  $\mathbf{e}_{\alpha}(Q)$  should be transported into vectors  $\mathbf{e}_{\beta}(Q')C_{\alpha}^{\beta} + \mathbf{e}_{5}(Q')C_{\alpha}^{5}$ , where the coefficients  $C_{\alpha}^{\beta}$  are the same in both cases. This means that in the selected bases,

$$G^{\alpha}_{\ \beta\mu} = \Gamma^{\alpha}_{\ \beta\mu}.\tag{12}$$

It is evident that assumption (10) tells one nothing about  $G^5_{\alpha\mu}$  and  $G^5_{5\mu}$ . To get an idea of what these coefficients can be like, one may consider a particular case where the connection for five-vector fields is such that there exists a certain local symmetry which can be formulated as the following principle:

For any set of scalar, five-vector and five-tensor fields defined in the vicinity of any point Q in space-time, by means of a certain procedure one can construct a set of fields in the vicinity of any other point Q', such that at Q'these new fields (which will be called *equivalent*) satisfy the same algebraic and first-order differential relations that the original fields satisfy at Q. (13)

The procedure by means of which the equivalent fields are constructed can be formulated as follows:

i. Introduce at Q a system of local Lorentz coordinates  $x^{\alpha}.$ 

Introduce the corresponding regular coordinate five-vector basis  $\mathbf{e}_A$ .

Introduce the corresponding bases for all other five-tensors.

ii. Each scalar field f in the vicinity of Q will then determine and be determined by one real coordinate function f(x).

Each five-vector field  $\mathbf{u}$  in the vicinity of Q will determine and be determined by five real coordinate functions  $u^A(x)$  (= components of  $\mathbf{u}$  in the basis  $\mathbf{e}_A$ ).

Each five-tensor field  $\mathbf{T}$  in the vicinity of Q will determine and be determined by an appropriate number of real coordinate functions  $T_{DE...F}^{AB...C}(x)$ (= components of  $\mathbf{T}$  in the relevant tensor basis corresponding to  $\mathbf{e}_A$ ).

- iii. Introduce at Q' a system of local Lorentz coordinates  $x'^{\alpha}$  such that  $x'^{\alpha}(Q') = x^{\alpha}(Q)$ . Introduce the corresponding regular coordinate five-vector basis  $\mathbf{e}'_A$ . Introduce the corresponding bases for all other five-tensors.
- iv. Then the equivalent scalar, five-vector and fivetensor fields in the vicinity of Q' will be determined in coordinates  $x'^{\alpha}$  and in the corresponding bases by the same functions  $f(\cdot), u^{A}(\cdot), \ldots, T_{DE...F}^{AB...C}(\cdot)$  that determine the original fields in the vicinity of Q in coordinates  $x^{\alpha}$  and in the corresponding bases.

It is not difficult to show that from the above symmetry principle follows that in any normalized regular basis

$$G^5_{5\mu} = 0$$
 and  $G^5_{\alpha\mu} = -\kappa g_{\alpha\mu}$ , (14)

where  $\kappa$  is a certain constant of dimension  $(length)^{-1}$ , which is not fixed by symmetry considerations. The latter formula suggests that at  $\kappa \neq 0$  it may be convenient to change the normalization of the fifth basis vector in such a way that one would have  $G^5_{\ \alpha\mu} = -g_{\alpha\mu}$ . In the following, such a regular basis will be called *active*.

One should also observe that there is no sense in talking about five-vectors if  $\kappa = 0$ , for it is impossible to distinguish a five-vector with such rules of parallel transport from a pair consisting of a four-vector and a scalar. Indeed,  $V_5$  is isomorphic to the direct sum of  $V_4$  and the space of scalars (regarded as one-dimensional vectors), and it is apparent that at  $\kappa = 0$  this isomorphism is preserved by parallel transport. Considering this, in the following I will always assume that  $\kappa \neq 0$ .

4. Let us now discuss in more detail the case of flat space-time. Supposing that the symmetry principle holds, from formulae (11), (12) and (14) one finds that for any active regular basis  $\mathbf{e}_A$  associated with a system of global Lorentz coordinates one has

$$G^{\alpha}_{\ \beta\mu} = G^{\alpha}_{\ 5\mu} = G^{5}_{\ 5\mu} = 0 \text{ and } G^{5}_{\ \beta\mu} = -\eta_{\beta\mu}, \quad (15)$$

where  $\eta_{\beta\mu} \equiv \text{diag}(+1, -1, -1, -1)$ . Such a set of five-vector basis fields will be called an *O*-basis ('O' stands for '*orthonormal*'). As one can see, the fields  $\mathbf{e}_A$  are not self-parallel. This is a distinctive feature of the considered five-vector connection, with respect to which the inner product *h* (regarded as a five-tensor) is *not* covariantly constant. The latter fact results in that the requirements of orthonormality and selfparallelism become conflicting in the sense that one can have either orthonormality or self-parallelism but not both at the same time.

Let us now construct a self-parallel five-vector basis,  $\mathbf{p}_A$ , that would coincide with  $\mathbf{e}_A$  at the origin of the considered Lorentz coordinate system. Being self-parallel, each  $\mathbf{p}_A$  should satisfy the equation  $\nabla_{\mu}\mathbf{p}_A = 0$ . Hence, if  $\mathbf{p}_A = \mathbf{e}_B N_A^B$ , one should have

$$\partial_{\mu}N^A_B(x) + G^A_{C\mu}N^C_B(x) = 0,$$

where  $G_{B\mu}^{A}$  are given by equations (15). Considering also that  $\mathbf{p}_{A}$  and  $\mathbf{e}_{A}$  should coincide at x = 0, one finds that

$$\begin{split} N_5^5(x) &= 1, \; N_\beta^\alpha(x) = \delta_\beta^\alpha, \\ N_5^\alpha(x) &= 0, \; N_\alpha^\beta(x) = x_\alpha, \end{split}$$

where  $x_{\alpha} \equiv \eta_{\alpha\beta} x^{\beta}$  are the corresponding covariant Lorentz coordinates. Thus,  $\mathbf{p}_A$  are expressed in terms of  $\mathbf{e}_A$  as follows:

$$\mathbf{p}_{\alpha}(x) = \mathbf{e}_{\alpha}(x) + x_{\alpha}\mathbf{e}_{5}(x)$$
 and  $\mathbf{p}_{5}(x) = \mathbf{e}_{5}(x)$ . (16)

The set  $\mathbf{p}_A$  will be called a *P*-basis ('*P*' stands for '*parallel*') associated with the considered system of Lorentz coordinates. Simple calculations show that

$$h(\mathbf{p}_{\alpha}, \mathbf{p}_{\beta}) = \eta_{\alpha\beta} + \kappa^2 x_{\alpha} x_{\beta}$$
  

$$h(\mathbf{p}_{\alpha}, \mathbf{p}_5) = h(\mathbf{p}_5, \mathbf{p}_{\alpha}) = \kappa^2 x_{\alpha},$$
(17)

so  $\mathbf{p}_A$  are orthogonal only at the origin.

Let us now derive the transformation formulae for the components of five-vectors and of other fivetensors corresponding to the transformation from one system of Lorentz coordinates to another. It is not difficult to show that under the transformation

$$x^{\mu} \to x^{\prime \mu} = \Lambda^{\mu}_{\nu} x^{\nu} + a^{\mu} \tag{18}$$

the elements of the O-basis transforms as

$$\mathbf{e}_{\alpha}' = \mathbf{e}_{\beta} \left( \Lambda^{-1} \right)_{\alpha}^{\beta} \quad \text{and} \quad \mathbf{e}_{5}' = \mathbf{e}_{5}, \tag{19}$$

whence one obtains the following transformation laws for the corresponding components of five-vectors and five-vector 1-forms:

$$\left\{ \begin{array}{ll} v^{\prime\alpha} = \Lambda^{\alpha}_{\ \beta} \, v^{\beta} \\ v^{\prime 5} = v^5 \end{array} \right. \ \, \text{and} \ \, \left\{ \begin{array}{ll} w^{\prime}_{\alpha} = w_{\beta} (\Lambda^{-1})^{\beta}_{\ \alpha} \\ w^{\prime}_{5} = w_{5}. \end{array} \right.$$

As one can see, the first four components of any fivevector or five-vector 1-form in the O-basis transform exactly as components of a four-vector or a fourvector 1-form, while the fifth component behaves as scalar.

By using equations (16) and the obvious formula for transformation of covariant Lorentz coordinates, one can easily find that under transformation (18) the elements of the P-basis transforms as

$$\mathbf{p}_{\alpha}' = \mathbf{p}_{\beta} \left( \Lambda^{-1} \right)_{\alpha}^{\beta} + a_{\alpha} \mathbf{p}_{5} \text{ and } \mathbf{p}_{5}' = \mathbf{p}_{5}, \qquad (20)$$

where  $a_{\alpha} = \eta_{\alpha\beta} a^{\beta}$ . From the latter formulae one obtains the following transformation laws for the components of five-vectors and five-vector 1-forms in the *P*-basis:

$$\begin{cases} v^{\prime\alpha} = \Lambda^{\alpha}_{\ \beta} v^{\beta} \\ v^{\prime 5} = v^{5} - a_{\alpha} \Lambda^{\alpha}_{\ \beta} v^{\beta} \end{cases}$$
(21a)

and

<

$$\begin{cases} w'_{\alpha} = w_{\beta} (\Lambda^{-1})^{\beta}_{\alpha} + a_{\alpha} w_5 \\ w'_5 = w_5. \end{cases}$$
(21b)

As one can see, these components transform nontrivially under space-time translations, and now one is able to understand why.

A global *P*-basis can exist only in flat space-time, where the parallel transport of five-vectors is independent of the path along which it is made. A *P*-basis can be constructed by choosing an orthonormal fivevector basis (with the fifth basis vector normalized as in an active regular basis) at one point and transporting it parallelly to all other points in space-time. Since (at  $\kappa \neq 0$ ) the inner product h is not conserved by parallel transport, the *P*-basis cannot be orthonormal at every point. Actually, the rules of parallel transport for five-vectors are such that  $\mathbf{p}_A$  will be orthogonal only at the origin. Moreover, as one can see from formulae (17), at each point the inner product matrix  $h_{AB} \equiv h(\mathbf{p}_A, \mathbf{p}_B)$  has its own value, different from the values it has at all other points. This means that having a *P*-basis, one is able to distinguish points without using any coordinates. In fact, if need be, one can recover the relevant Lorentz coordinates by simply calculating the inner product of  $\mathbf{p}_{\alpha}$  and  $\mathbf{p}_{5}$ . Thus, the *P*-basis is a structure which is rigitly connected to space-time points and to one of the Lorentz coordinate systems. When the latter is changed, the *P*-basis changes too.

5. One may now ask the following question: are there any geometric or physical quantities which are described by five-vectors or by other nontrivial fivetensors (by the ones not reducible to a four-tensor)? This brings us to another question: how can one discover a five-vector or a five-tensor? One possible answer to this question is the same as to a similar question for four-vectors: one has to find several quantities that under Lorentz transformations and translations in flat space-time transform as components of a five-vector or of some other five-tensor. Since one is talking about components, one has to specify the basis in which they are evaluated. This is a simple matter if the definition of the quantities one considers involves only scalars and components of four-tensors in a Lorentz basis: since in either case  $\nabla_{\mu} = \partial_{\mu}$ , the same should be true for the quantities defined, and considering that in this basis  $g_{\mu\nu} = \eta_{\mu\nu}$ , one concludes that the five-tensor components should correspond to a *P*-basis and consequently should transform according to formulae (21).

The simplest example of quantities that transform as components of a nontrivial five-tensor are covariant Lorentz coordinates. Indeed, under Lorentz transformations and translations the five quantities  $x_A$ , where  $x_5 \equiv 1$ , transform as components of a five-vector 1form. Consequently, if  $\tilde{\mathbf{q}}^A$  is the basis of five-vector 1-forms dual to the *P*-basis associated with the selected Lorentz coordinates, the 1-form  $\tilde{\mathbf{x}}$  constructed according to the formula

$$\widetilde{\mathbf{x}}(x) \equiv x_{\alpha} \widetilde{\mathbf{q}}^{\alpha}(x) + \widetilde{\mathbf{q}}^{5}(x), \qquad (22)$$

will be the same no matter which system of Lorentz

coordinates is used.

From equations (20) one can easily obtain the formulae that relate the basis  $\tilde{\mathbf{q}}^A$  to the basis of fivevector 1-forms  $\tilde{\mathbf{o}}^A$  dual to the *O*-basis corresponding to the same coordinates:

$$\widetilde{\mathbf{q}}^{\alpha}(x) = \widetilde{\mathbf{o}}^{\alpha}(x) \text{ and } \widetilde{\mathbf{q}}^{5}(x) = \widetilde{\mathbf{o}}^{5}(x) - x_{\alpha}\widetilde{\mathbf{o}}^{\alpha}(x).$$
 (23)

Substituting these relations into definition (22), one obtains the following expression for the 1-form  $\tilde{\mathbf{x}}$  in the basis  $\tilde{\mathbf{o}}^A$ :

$$\widetilde{\mathbf{x}}(x) = x_{\alpha} \widetilde{\mathbf{o}}^{\alpha}(x) + \widetilde{\mathbf{o}}^{5}(x) - x_{\alpha} \widetilde{\mathbf{o}}^{\alpha}(x) = \widetilde{\mathbf{o}}^{5}(x),$$

from which one can clearly see that  $\tilde{\mathbf{x}}$  is indeed independent of the choice of the coordinate system.

Another example of geometric quantities that transform as components of a nontrivial five-tensor are parameters of Poincare transformations. Let us recall that the symmetry properties of flat space-time can be formulated as a principle similar to the one presented above, only now instead of local Lorentz coordinates one should speak of global Lorentz coordinate systems. It is evident that in this case the latter are used only as a tool for constructing the equivalent fields. By itself, the replacement of a given set of fields with an equivalent set, which is nothing but an active field transformation, is an invariant procedure and can be considered without referring to any coordinates. However, depending on how the latter are selected, a given field transformation will correspond to different coordinate transformations. Let us now find how the parameters of these coordinate transformations change as one passes from one system of Lorentz coordinates to another.

The idea of the following calculation is very simple. One selects some set of fields and a system of Lorentz coordinates, and by means of an arbitrary Poincare transformation constructs the equivalent set of fields. One then considers another system of Lorentz coordinates and determines the precise Poincare transformation that one has to make in these new coordinates to obtain the same set of equivalent fields. Finally, one expresses the parameters of this second Poincare transformation in terms of the parameters of the first one.

As a set of fields it is convenient to choose the covariant coordinates associated with the selected Lorentz coordinate system  $x^{\alpha}$ , i.e. the four scalar fields  $\varphi_{(\alpha)}$  such that

$$\varphi_{(\alpha)}(Q) = \eta_{\alpha\beta} x^{\beta}(Q)$$

at every point Q. Let us consider an arbitrary Poincare transformation that corresponds to the coordinate transformation

$$x_{\alpha} \to y_{\alpha} = x_{\beta} L^{\beta}_{\ \alpha} + b_{\alpha}. \tag{24}$$

The equivalent fields obtained by this transformation are

$$\varphi_{(\alpha)}^{\text{equiv}} = y_{\alpha} = x_{\beta} L^{\beta}_{\ \alpha} + b_{\alpha}.$$

Let us now consider another system of Lorentz coordinates:

$$x^{\prime\alpha} = \Lambda^{\alpha}_{\ \beta} x^{\beta} + a^{\alpha}.$$

In these new coordinates the original fields acquire the form

$$\varphi_{(\alpha)} = (x'_{\beta} - a_{\beta})\Lambda^{\beta}_{\alpha},$$

and the equivalent fields are

$$\varphi_{(\alpha)}^{\text{equiv}} = (x_{\gamma}' - a_{\gamma})\Lambda_{\beta}^{\gamma}L_{\alpha}^{\beta} + b_{\alpha}.$$

One should now present the right-hand side of the latter equation as

$$\varphi_{(\alpha)}^{\text{equiv}} = (y_{\beta}' - a_{\beta})\Lambda_{\alpha}^{\beta},$$

where

$$y'_{\alpha} \equiv x'_{\beta} L'^{\beta}_{\alpha} + b'_{\alpha},$$

and then express  $L^{\prime\beta}_{\alpha}$  and  $b^{\prime}_{\alpha}$  in terms of  $L^{\beta}_{\alpha}$  and  $b_{\alpha}$ . Straightforward calculations give

$$L^{\prime\alpha}_{\ \beta} = \Lambda^{\alpha}_{\ \sigma} L^{\sigma}_{\ \tau} (\Lambda^{-1})^{\tau}_{\ \beta} b^{\prime}_{\ \beta} = b_{\tau} (\Lambda^{-1})^{\tau}_{\ \beta} + a_{\beta} - a_{\rho} \Lambda^{\rho}_{\ \sigma} L^{\sigma}_{\ \tau} (\Lambda^{-1})^{\tau}_{\ \beta},$$
(25)

which shows that the quantities  $\mathcal{T}^A_B$  defined as

$$\mathcal{T}^{\alpha}_{\beta} = L^{\alpha}_{\beta}, \ \mathcal{T}^{5}_{\beta} = b_{\beta}, \ \mathcal{T}^{\alpha}_{5} = 0, \ \text{ and } \ \mathcal{T}^{5}_{5} = 1,$$

transform as components of a five-tensor of rank (1,1).

It is also interesting to find the transformation formulae for the parameters of infinitesimal Poicare transformations. In this case the matrix  $L^{\alpha}_{\beta}$  in equation (24) can be presented as

$$L^{\alpha}_{\ \beta} = \delta^{\alpha}_{\ \beta} + \frac{1}{2} (\delta^{\alpha}_{\nu} \eta_{\beta\mu} - \delta^{\alpha}_{\mu} \eta_{\beta\nu}) \, \omega^{\mu\nu},$$

where  $\omega^{\mu\nu} = -\omega^{\nu\mu}$ , and both  $\omega^{\mu\nu}$  and  $b_{\alpha}$  are infinitesimals. From formulae (25) one obtains

$$\omega'^{\mu\nu} = \Lambda^{\mu}_{\alpha}\Lambda^{\nu}_{\beta}\omega^{\alpha\beta} \quad \text{and} \quad b'^{\mu} = \Lambda^{\mu}_{\nu}(b^{\nu} - a_{\alpha}\Lambda^{\alpha}_{\beta}\omega^{\nu\beta}),$$

which shows that the quantities  $\mathcal{R}^{AB}$  defined as

$$\mathcal{R}^{\mu\nu} = \omega^{\mu\nu}, \ \mathcal{R}^{\mu5} = -\mathcal{R}^{5\mu} = b^{\mu}, \ \text{ and } \ \mathcal{R}^{55} = 0,$$

transform as components of an antisymmetric fivetensor of rank (2,0). Tensors  $\mathcal{T}_B^A$  and  $\mathcal{R}^{AB}$  are discussed in more detail in part III of the long version [5].

Let us now consider an example of physical quantities that transform as components of a five-tensor: the canonical stress-energy and angular momentum  $\mathcal{M}$  in the O-basis. Using relations (23), one has tensors,  $\Theta^{\mu}_{\alpha}$  and  $M^{\mu}_{\alpha\beta}$ .

Let us begin by writing out the formulae that express the components of these two tensors in one Lorentz coordinate system in terms of their components in another Lorentz coordinate system. If the two coordinate systems are related as in equation (18), then

$$\Theta_{\alpha}^{\prime\mu} = \Lambda_{\nu}^{\mu} \Theta_{\beta}^{\nu} (\Lambda^{-1})_{\alpha}^{\beta}, 
M_{\alpha\beta}^{\prime\mu} = x_{\alpha}^{\prime} \Theta_{\beta}^{\prime\mu} - x_{\beta}^{\prime} \Theta_{\alpha}^{\prime\mu} + \Sigma_{\alpha\beta}^{\prime\mu} 
= \Lambda_{\nu}^{\mu} M_{\sigma\tau}^{\nu} (\Lambda^{-1})_{\alpha}^{\sigma} (\Lambda^{-1})_{\beta}^{\tau} 
+ a_{\alpha} \Lambda_{\nu}^{\mu} \Theta_{\tau}^{\nu} (\Lambda^{-1})_{\alpha}^{\tau},$$
(26)

where  $\Sigma^{\mu}_{\alpha\beta}$  is the spin angular momentum tensor.

With respect to their lower indices,  $\Theta^{\mu}_{\alpha}$  and  $M^{\mu}_{\alpha\beta}$ are traditionally regarded as components of fourtensors, and the fact that under space-time translations  $M^{\mu}_{\alpha\beta}$  acquires additional terms proportional to  $\Theta^{\mu}_{\alpha}$  is interpreted as a consequence of one actually making a switch from one quantity-the angular momentum relative to the point  $x^{\mu} = 0$ , to another quantity-the angular momentum relative to the point  $x'^{\mu} = 0$ . Five-dimensional tangent vectors enable one to give this fact a different interpretation, which in several ways is more attractive.

One should notice that equations (26) coincide exactly with the transformation formulae for components in the P-basis of a tensor—let us denote it as  $\mathcal{M}$ —that has one (upper) four-vector index and two (lower) five-vector indices and whose components are related to  $\Theta^{\mu}_{\alpha}$  and  $M^{\mu}_{\alpha\beta}$  as follows:

$$\mathcal{M}^{\mu}_{\alpha\beta} = M^{\mu}_{\alpha\beta}, \quad \mathcal{M}^{\mu}_{5\alpha} = \Theta^{\mu}_{\alpha} \\ \mathcal{M}^{\mu}_{\alpha5} = -\Theta^{\mu}_{\alpha}, \quad \mathcal{M}^{\mu}_{55} = 0.$$
(27)

This coincidence means that  $\Theta^{\mu}_{\alpha}$  and  $M^{\mu}_{\alpha\beta}$  can be regarded as components of a single five-tensor. Since by definition  $M^{\mu}_{\alpha\beta} = -M^{\mu}_{\beta\alpha}$ , this tensor is antisymmetric in its lower (five-vector) indices.

Such an interpretation of  $\Theta^{\mu}_{\alpha}$  and  $M^{\mu}_{\alpha\beta}$  implies that there exists a single local physical quantity: the stress–energy–angular momentum tensor  $\mathcal{M}$ . The belief that there are many different angular momenta should now be regarded as merely a wrong impression created by interpreting  $\Theta^{\mu}_{\alpha}$  and  $M^{\mu}_{\alpha\beta}$  as four-tensors: in reality, all these angular momenta are simply the components of  $\mathcal{M}$  in different five-vector bases.

There is now no difficulty in defining the angular momentum density in curved space-time. To see how this can be done, let us evaluate the components of

$$\mathcal{M} = (x_{\alpha}\Theta^{\mu}_{\beta} - x_{\beta}\Theta^{\mu}_{\alpha} + \Sigma^{\mu}_{\alpha\beta}) \tilde{\mathbf{q}}^{\alpha} \otimes \tilde{\mathbf{q}}^{\beta} \otimes \mathbf{E}_{\mu} + (\Theta^{\mu}_{\beta}) \tilde{\mathbf{q}}^{5} \otimes \tilde{\mathbf{q}}^{\beta} \otimes \mathbf{E}_{\mu} + (-\Theta^{\mu}_{\alpha}) \tilde{\mathbf{q}}^{\alpha} \otimes \tilde{\mathbf{q}}^{5} \otimes \mathbf{E}_{\mu} + (\Theta^{\mu}_{\beta}) \tilde{\mathbf{o}}^{5} \otimes \tilde{\mathbf{o}}^{\beta} \otimes \mathbf{E}_{\mu} + (-\Theta^{\mu}_{\alpha}) \tilde{\mathbf{o}}^{\alpha} \otimes \tilde{\mathbf{o}}^{5} \otimes \mathbf{E}_{\mu}.$$

Thus, in the O-basis  $\mathcal{M}^{\mu}_{\alpha\beta}$  coincide with the components of the spin angular momentum tensor. In the case of flat space-time one gives preference to the Pbasis, since in it  $\nabla_{\mu} = \partial_{\mu}$ , and, accordingly, the  $\mathcal{M}^{\mu}_{\alpha\beta}$ components acquire additional terms proportional to covariant Lorentz coordinates and to the components  $\mathcal{M}^{\mu}_{\alpha 5}$  and  $\mathcal{M}^{\mu}_{5\beta}$ . In the case of curved space-time, where a global self-parallel basis does not exist, it is more convenient to use a regular basis and have  $\mathcal{M}^{\mu}_{\alpha\beta} = \Sigma^{\mu}_{\alpha\beta}.$ 

Let us now recall that canonical  $\Theta^{\mu}_{\alpha}$  and  $M^{\mu}_{\alpha\beta}$ are defined as Noether currents corresponding to Poincare transformations and as such satisfy the following "conservation laws":

$$\begin{split} \partial_{\mu}\Theta^{\mu}_{\alpha} &= 0\\ \partial_{\mu}M^{\mu}_{\alpha\beta} &= \eta_{\alpha\mu}\Theta^{\mu}_{\beta} - \eta_{\beta\mu}\Theta^{\mu}_{\alpha} + \partial_{\mu}\Sigma^{\mu}_{\alpha\beta} = 0. \end{split}$$

One can now replace these two four-tensor equations with a single covariant five-tensor equation:

$$\mathcal{M}^{\mu}_{\alpha\beta;\mu} = 0, \qquad (28)$$

where it has been taken into account that in the Pbasis all five-vector connection coefficients are zero. It is interesting to see how equation (28) works in the O-basis. One has

$$\begin{aligned} \mathcal{M}^{\mu}_{5\alpha;\,\mu} &= \partial_{\mu}\mathcal{M}^{\mu}_{5\alpha} - \mathcal{M}^{\mu}_{A\alpha}G^{A}_{5\mu} - \mathcal{M}^{\mu}_{5A}G^{A}_{\alpha\mu} \\ &= \partial_{\mu}\Theta^{\mu}_{\alpha} - \mathcal{M}^{\mu}_{55}G^{5}_{\alpha\mu} = \partial_{\mu}\Theta^{\mu}_{\alpha} = 0 \end{aligned}$$

and

$$\begin{aligned} \mathcal{M}^{\mu}_{\alpha\beta;\,\mu} &= \partial_{\mu}\mathcal{M}^{\mu}_{\alpha\beta} - \mathcal{M}^{\mu}_{A\beta}G^{A}_{\alpha\mu} - \mathcal{M}^{\mu}_{\alphaA}G^{A}_{\beta\mu} \\ &= \partial_{\mu}\Sigma^{\mu}_{\alpha\beta} - \Theta^{\mu}_{\beta}G^{5}_{\alpha\mu} + \Theta^{\mu}_{\alpha}G^{5}_{\beta\mu} \\ &= \partial_{\mu}\Sigma^{\mu}_{\alpha\beta} + \Theta^{\mu}_{\beta}\eta_{\alpha\mu} - \Theta^{\mu}_{\alpha}\eta_{\beta\mu} = 0. \end{aligned}$$

Thus, one obtains the same conservation laws for  $\Theta^{\mu}_{\alpha}$ and  $M^{\mu}_{\alpha\beta}$ , only now the terms proportional to  $\Theta^{\mu}_{\alpha}$ in the second equation come from connection coefficients.

6. The fact that five-dimensional tangent vectors and the tensors associated with them enable one to give a coordinate-independent description to finite and infinitesimal Poincare transformations and to describe as a single local object such quantities

as the stress-energy and angular momentum tensors. should be thought of only as a reason for considering five-vectors in the first place and for making an exploratory study of their basic properties. If this were all there is to it, i.e. if five-vectors only enabled one to present certain geometric quantities and the relations between them in a mathematically more attractive form, such vectors would hardly be of particular interest both to physicists, who typically do not care much for fancy mathematics unless it enables them to formulate new physical concepts, and to mathematicians, who would consider five-vectors as merely a particular combination of already known mathematical constructions. A more important reason why the concept of a five-dimensional tangent vector is worth considering is that it enables one to extend the notion of the affine connection on a manifold and of the connections which physicists call gauge fields, and thereby at no cost at all, i.e. without changing the manifold in any way and without introducing new gauge groups, to obtain new geometric properties of space-time in the form of a new kind of torsion and a new kind of gauge fields.

Before discussing these applications of five-vectors in more detail, let me say a few words about the fivevector generalization of exterior differential calculus, which is considered in detail in part IV of the long version [6]. This latter generalization is more a technical necessity—a necessity in replacing ordinary tangent vectors with five-vectors in all the formulae related to integration of differential forms and to exterior differentiation of the latter. Apart from allowing one to present certain relations in a more elegant form, for scalar-valued forms this generalization is equivalent to ordinary exterior calculus, which was to be expected since five-vectors in this case are used only for characterizing the infinitesimal elements of integration volumes, and the latter are not changed in any way themselves and are not endowed with any new additional structure.

Let us now discuss the five-vector generalization of the covariant derivative. Above, the latter has been introduced for five-vector fields, which is equivalent to introducing a map

$$\nabla: \, \mathcal{D} \times \mathcal{F} \to \mathcal{F}, \tag{29}$$

where  $\mathcal{F}$  is the set of all five-vector fields and  $\mathcal{D}$  is the set of all four-vector fields (derivations). Considering the way five-vectors are related to four-vectors, one can regard the structure defined on space-time by this map as an extension of the structure defined on it by ordinary (four-vector) affine connection.

The next step is to replace the operator  $\nabla_{\mathbf{U}}$  with the operator  $\nabla_{\mathbf{u}}$  defined by the equation

$$\nabla_{\mathbf{u}} = \nabla_{\mathbf{U}} \text{ for } \mathbf{u} \in \mathbf{U}.$$

It is obvious that  $\nabla_{\mathbf{u}}$  is absolutely equivalent to  $\nabla_{\mathbf{U}}$ , however, unlike the latter, it formally depends on a *five*-vector. From the above definition it follows that  $\nabla_{\mathbf{u}} = \nabla_{(\mathbf{u}^{\mathcal{Z}})}$  for any  $\mathbf{u}$ , so the replacement of  $\nabla_{\mathbf{U}}$ with  $\nabla_{\mathbf{u}}$  is equivalent to replacing map (29) with a map

$$\nabla: \, \mathcal{F}_{\mathcal{Z}} \times \mathcal{F} \to \mathcal{F}, \tag{30}$$

where  $\mathcal{F}_{\mathcal{Z}}$  is the subset of all five-vector fields from  $\mathcal{F}$  represented by purely differential operators. It now seems natural to make one more step in generalizing the concept of affine connection to five-vectors and consider a map

$$\square: \mathcal{F} \times \mathcal{F} \to \mathcal{F},$$

which I will call the *five-vector affine connection*. The image of a pair of fields  $(\mathbf{u}, \mathbf{v})$  with respect to  $\Box$  will be denoted as  $\Box_{\mathbf{u}}\mathbf{v}$  and will be called the *five-vector covariant derivative* of field  $\mathbf{v}$  in the direction of field  $\mathbf{u}$ . To give  $\Box$  a formal definition, one should formulate certain requirements that should be satisfied by  $\Box$ , similar to the requirements one usually imposes on  $\nabla$ . Two such requirements are quite obvious:

$$\Box_{(f\mathbf{u}+g\mathbf{v})}\mathbf{w} = f \cdot \Box_{\mathbf{u}}\mathbf{w} + g \cdot \Box_{\mathbf{v}}\mathbf{w}, \qquad (31a)$$
$$\Box_{\mathbf{u}}(\mathbf{v}+\mathbf{w}) = \Box_{\mathbf{u}}\mathbf{v} + \Box_{\mathbf{u}}\mathbf{w} \qquad (31b)$$

for any scalar functions f and g and any five-vector fields  $\mathbf{u}$ ,  $\mathbf{v}$ , and  $\mathbf{w}$ . To make a rational choice of the analog of the requirement on  $\nabla$  that expresses the Leibniz rule in application to the product of a fourvector field and a scalar function, one should first formulate explicitly the condition that the structure defined on space-time by  $\square$  is an extension of the structure defined on it by  $\nabla$ . The latter statement apparently means that the restriction of  $\square$  to  $\mathcal{F}_{\mathbb{Z}} \times \mathcal{F}$  should coincide with map (30), which in its turn means that

$$\square_{(\mathbf{u}^{\mathcal{Z}})} = \nabla_{\mathbf{u}} \tag{32}$$

for any five-vector field **u**. Together with requirement (31a), the latter equation yields

$$\Box_{\mathbf{u}} = \nabla_{\mathbf{u}} + \lambda_{\mathbf{u}} \, \Box_{\mathbf{i}},\tag{33}$$

where  $\mathbf{i}$  is the five-vector from  $\mathcal{E}$  that corresponds to the unity value of the parameter:  $\lambda_{\mathbf{i}} = 1$ . Since the elements of  $\mathcal{E}$  do not correspond to any direction in space-time, it will be assumed that  $\Box_{\mathbf{i}}$  is a *purely algebraic* operator, so that for any five-vector field  $\mathbf{v}$ and any function f,

$$\Box_{\mathbf{i}}(f\mathbf{v}) = f \cdot \Box_{\mathbf{i}} \mathbf{v}. \tag{34}$$

From the latter equation and formula (33) one obtains the relation

$$\Box_{\mathbf{u}}(f\mathbf{v}) = \partial_{\mathbf{u}}f \cdot \mathbf{v} + f \cdot \Box_{\mathbf{u}}\mathbf{v}, \qquad (31c)$$

which is the desired analog of the chain rule for  $\Box$ .

Let us now define the action of  $\Box$  on scalar functions. Considering what has been said above, it seems reasonable to think that the action of  $\Box_{\mathbf{u}}$  on an arbitrary scalar function f should produce a sum of the derivative  $\partial_{\mathbf{u}} f$  and a term of the form  $a\lambda_{\mathbf{u}} f$ , where ais a constant. One should now notice that if one adds to  $\Box$  a term proportional to  $\lambda_{\mathbf{u}} \mathbf{1}$ , one will obtain an operator that will still satisfy requirements (31), but whose action on scalar functions will be different. In particular, one can select this additional term in such a way that the action of the resulting operator on fwould yield  $\partial_{\mathbf{u}} f$ . In the following, the notation  $\Box_{\mathbf{u}}$ will refer to this particular choice of the five-vector covariant derivative operator, and so

$$\Box_{\mathbf{u}}f = \partial_{\mathbf{u}}f.$$

From the latter equation and equation (31c) it is seen that the action of  $\Box$  on the product of two scalar functions and on the product of a scalar function and a five-vector field obeys the Leibniz rule. One may assume that the same rule holds for the contraction and tensor product. This will enable one to define the action of  $\Box$  on an arbitrary five-vector 1-form field  $\tilde{s}$ according to the formula

$$< \Box_{\mathbf{u}} \widetilde{\mathbf{s}}, \mathbf{v} > = \partial_{\mathbf{u}} < \widetilde{\mathbf{s}}, \mathbf{v} > - < \widetilde{\mathbf{s}}, \Box_{\mathbf{u}} \mathbf{v} > 0$$

for any five-vector field  $\mathbf{v}$ , and, by induction, on the fields of all other five-tensors according to the formula

$$\Box_{\mathbf{u}}(\mathbf{m}\otimes\mathbf{n}) = \Box_{\mathbf{u}}\mathbf{m}\otimes\mathbf{n} + \mathbf{m}\otimes\Box_{\mathbf{u}}\mathbf{n},$$

where  $\mathbf{m}$  and  $\mathbf{n}$  are any two five-tensor fields.

There is one more constraint that should be imposed on  $\Box$ , which enables one to define in a natural way the action of  $\Box$  on four-vector fields. Namely, one should require that

$$\mathbf{v} \equiv \mathbf{w} \pmod{R} \implies \mathbf{\Box}_{\mathbf{u}} \mathbf{v} \equiv \mathbf{\Box}_{\mathbf{u}} \mathbf{w} \pmod{R}, \quad (35)$$

where R is the equivalence relation on  $V_5$  considered earlier. The derivative  $\Box_{\mathbf{u}} \mathbf{V}$  of an arbitrary fourvector field  $\mathbf{V}$  can then be defined as the equivalence class with respect to R of all the fields of the form  $\Box_{\mathbf{u}} \mathbf{v}$  with  $\mathbf{v} \in \mathbf{V}$ .

Let us now introduce the analogs of connection coefficients for  $\Box$ . For a given set of five-vector basis fields  $\mathbf{e}_A$ , it is natural to define the latter according to the equation

$$\Box_A \mathbf{e}_B = \mathbf{e}_C H^C_{BA}$$

where  $\Box_A \equiv \Box_{\mathbf{e}_A}$ . The quantities  $H^A_{BC}$  will be called *five-vector connection coefficients*. If  $\mathbf{e}_A$  is a regular basis, then from equation (32) it follows that

$$H^A_{\ B\mu} = G^A_{\ B\mu},$$

where  $G^{A}_{B\mu}$  are the connection coefficients associated with  $\nabla$ . Furthermore, from condition (35) it follows that in any standard five-vector basis

$$H^{\alpha}_{5B} = 0 \tag{36}$$

at all  $\alpha$  and B. In the usual way one can obtain the transformation formula for five-vector connection coefficients corresponding to the transformation  $\mathbf{e}'_A = \mathbf{e}_B L^B_A$ :

$$H_{BC}^{\prime A} = (L^{-1})_{D}^{A} H_{EF}^{D} L_{B}^{E} L_{C}^{F} + (L^{-1})_{D}^{A} (\partial_{F} L_{B}^{D}) L_{C}^{F}.$$

If both bases are standard, one will have

$$H'^{A}_{B5} = (L^{-1})^{A}_{D} H^{D}_{E5} L^{E}_{B} L^{5}_{5},$$

so the coefficients  $H_{B5}^A$  transform as components of a five-tensor and therefore cannot be nullified at a given space-time point by an appropriate choice of the five-vector basis fields.

The interpretation of the five-vector covariant derivative is discussed in detail in part V of the long version [7]. In particular, it is shown that  $\Box$  can be regarded as the operator of a derivative calculated by using certain rules of parallel transport for the vectors and tensors which are the values of the differentiated fields, but the properties of this transport will differ from the usual ones in that the derivative along a parametrized curve whose tangent four-vector is  $\mathbf{U} + \mathbf{V}$  in general will no longer equal the sum of the derivative along a curve whose tangent four-vector is  $\mathbf{U}$  and the derivative along a curve whose tangent four-vector is referred to paper [7].

The derivative  $\Box$  can also be defined for the fields whose values are some abstract vectors or tensors that have no direct relation to the space-time manifold. In the following such vectors and tensors will be referred to as *nonspacetime* vectors and tensors.

Let us consider a set  $\mathcal{V}$  of fields whose values are some *n*-dimensional nonspacetime vectors, which I will denote with small capital Roman letters with an arrow:  $\vec{A}, \vec{B}, \vec{C}$ , etc. Defining an ordinary covariant derivative for such fields is equivalent to fixing a map

$$\nabla: \mathcal{D} \times \mathcal{V} \to \mathcal{V},$$

or an equivalent map

$$\nabla: \, \mathcal{F}_{\mathcal{Z}} \times \mathcal{V} \to \mathcal{V}. \tag{37}$$

If  $\vec{\mathbf{E}}_i$  (i = 1, ..., n) is some set of basis fields in  $\boldsymbol{\mathcal{V}}$ , then the corresponding connection coefficients, which I will refer to as gauge fields, are defined by the equation

$$\nabla_{\mu} \vec{\mathbf{E}}_i = \vec{\mathbf{E}}_j A^j_{\ i\mu}.$$

In a similar manner one can formally define the fivevector covariant derivative for the fields from  $\mathcal{V}$ . This is equivalent to fixing a map

$$\Box: \mathcal{F} \times \mathcal{V} \to \mathcal{V}, \tag{38}$$

which will be regarded as an extension of map (37), so in this case, too, the operators  $\Box$  and  $\nabla$  will be related as in equation (32). In addition to this, map (38) should satisfy three requirements similar to requirements (31) for five-vector fields, which I will not present here.

The connection coefficients corresponding to derivative (38), which I will call *five-vector gauge fields*, are defined by the equation

$$\square_A \vec{\mathbf{E}}_i = \vec{\mathbf{E}}_j B^j_{\ iA}.\tag{39}$$

It is apparent that in any regular five-vector basis

$$B^i_{\ j\mu} = A^i_{\ j\mu}$$

for any i, j, and  $\mu$ . In the usual manner one can obtain the formula for transformation of five-vector gauge fields under the transformation  $\vec{\mathbf{E}}'_i = \vec{\mathbf{E}}_j L^j_i$  of the basis fields in  $\mathcal{V}$ :

$$B'_{jA}^{i} = (L^{-1})^{i}_{\ k} B^{k}_{\ lA} L^{l}_{\ j} + (L^{-1})^{i}_{\ k} \partial_{A} L^{k}_{\ j}.$$

From this formula it follows that in any standard five-vector basis

$$B'_{j5}^{i} = (L^{-1})^{i}_{\ k} B^{k}_{\ l5} L^{l}_{\ j},$$

so the fields  $B^i_{j5}$  transform as components of a tensor of rank (1,1) over  $\mathcal{V}$ . This latter fact, together with the facts that  $B^i_{j5}$  are Lorentz scalars and that in the equations of motion for matter fields they will appear at the place where the mass parameter usually stands, may suggest that some of these new gauge fields can effectively play the role of Higgs fields. A more detailed discussion of five-vector gauge fields can be found in Ref. [7].

7. Before turning to the next item, it is necessary to say a few words about the properties of five-vector bivectors. It is a simple matter to see that any such bivector  $\boldsymbol{\mathcal{A}}$  can be uniquely presented as a sum of two terms: (i) a bivector made only of five-vectors from  $\boldsymbol{\mathcal{Z}}$  and (ii) a wedge product of a five-vector from  $\boldsymbol{\mathcal{E}}$ and some other five-vector. In the following, these two parts of  $\mathcal{A}$  will be referred to as its  $\mathcal{Z}$ - and  $\mathcal{E}$ components, respectively, and will be denoted as  $\mathcal{A}^{\mathcal{Z}}$ and  $\mathcal{A}^{\mathcal{E}}$ .

Since  $\mathcal{Z}$  endowed with the inner product h is isomorphic to  $V_4$ , to the  $\mathcal{Z}$ -component of  $\mathcal{A}$  one can put into correspondence a certain four-vector bivector. If  $\mathbf{e}_A$  is an active regular basis and  $\mathbf{E}_{\alpha}$  is the associated four-vector basis, then the components of this fourvector bivector in the basis  $\mathbf{E}_{\alpha} \otimes \mathbf{E}_{\beta}$  equal the components  $\mathcal{A}^{\alpha\beta}$  of  $\mathcal{A}$  in the basis  $\mathbf{e}_A \otimes \mathbf{e}_B$ . In a similar way, since the subspace of five-vector bivectors with the zero  $\mathcal{Z}$ -component is isomorphic to  $V_4$ , too, to the  $\mathcal{E}$ -component of  $\mathcal{A}$  one can put into correspondence a certain four-vector. For practical reasons, it is convenient to establish the isomorphism between the above two vector spaces by supposing that the former is endowed not with the inner product induced by h, but with the inner product that differs from the latter by the factor  $\xi^{-1}$ . In this case the components of the mentioned four-vector in the basis  $\mathbf{E}_{\alpha}$  introduced above will equal the components  $\mathcal{A}^{\alpha 5}$  of the bivector  $\boldsymbol{\mathcal{A}}$  in the basis  $\mathbf{e}_A \otimes \mathbf{e}_B$ .

Let us now introduce a new kind of derivative whose argument is a five-vector bivector and which, in view of this, will be called the *bivector* derivative. Let us first define it for scalar, four-vector and fourtensor fields in flat space-time. To this end, let us consider the group of global active Poincare tranformations of the indicated fields and distinguish in it some one-parameter family  $\mathcal{H}$  that inludes the identity transformation. Let us denote the parameter of this family as s and the image of an arbitrary field  $\mathcal{G}$  under a transformation from  $\mathcal{H}$  as  $\Pi_s{\mathcal{G}}$ . It is convenient to take that the identity transformation corresponds to s = 0.

For the selected one-parameter family  $\mathcal{H}$  and for any sufficiently smooth field  $\mathcal{G}$  from the indicated class of fields, one can define the derivative

$$\mathsf{D}_{\mathcal{H}}\mathcal{G} \equiv (d/ds)\mathbf{\Pi}_s\{\mathcal{G}\}|_{s=0},\tag{40}$$

which apparently is a field of the same type as  $\mathcal{G}$ . It is also apparent that for every type of fields, the operators  $D_{\mathcal{H}}$  corresponding to all possible one-parameter families  $\mathcal{H}$  make up a ten-dimensional real vector space, which is nothing but the representation of the Lie algebra of the Poincare group that corresponds to the considered type of fields.

Let us introduce in space-time some system of global Lorentz coordinates  $x^{\alpha}$  and select a basis in the space of operators  $D_{\mathcal{H}}$  consisting of the six operators  $M_{\mu\nu}$  that correspond to rotations in the planes  $x^{\mu}x^{\nu}$  ( $\mu < \nu$ ) and of the four operators  $P_{\mu}$  that correspond to translations along the coordinate axes. If one parametrizes the indicated transformations with the parameters  $\omega^{\alpha\beta}$  and  $b^{\alpha}$  introduced earlier, then for an arbitrary scalar function f one will have

$$\begin{aligned}
\mathsf{P}_{\mu}f(x) &= \partial_{\mu}f(x) \\
\mathsf{M}_{\mu\nu}f(x) &= x_{\nu}\partial_{\mu}f(x) - x_{\mu}\partial_{\nu}f(x);
\end{aligned} (41)$$

for an arbitrary four-vector field **U** one will have

where  $(M_{\mu\nu})^{\alpha}_{\ \beta} \equiv \delta^{\alpha}_{\ \nu} g_{\mu\beta} - \delta^{\alpha}_{\ \mu} g_{\nu\beta}$  and the components correspond to the Lorentz four-vector basis associated with the selected coordinates; and so on.

It is a simple matter to see that with transition to another system of Lorentz coordinates, the quantities  $\mathsf{M}_{\mu\nu}\mathcal{G}$  and  $\mathsf{P}_{\mu}\mathcal{G}$  transform respectively as the  $\mu\nu$ - and  $\mu$ 5-components of a five-vector 2-form in the *P*-basis. Consequently, the field

$$\mathsf{D}\mathcal{G} \equiv \mathsf{M}_{|\mu\nu|}\mathcal{G} \cdot \widetilde{\mathbf{q}}^{\mu} \wedge \widetilde{\mathbf{q}}^{\nu} + \mathsf{P}_{\mu}\mathcal{G} \cdot \widetilde{\mathbf{q}}^{\mu} \wedge \widetilde{\mathbf{q}}^{5}, \qquad (43)$$

where  $\widetilde{\mathbf{q}}^A$  is the basis of five-vector 1-forms dual to the *P*-basis  $\mathbf{p}_A$  associated with the selected Lorentz coordinate system, will be the same at any choice of the latter. From definition (43) it follows that at every point in space-time

$$\mathsf{P}_{\mu}\mathcal{G} = < \mathsf{D}\mathcal{G} \,, \, \mathbf{p}_{\mu} \wedge \mathbf{p}_{5} > \ \mathsf{M}_{\mu
u}\mathcal{G} = < \mathsf{D}\mathcal{G} \,, \, \mathbf{p}_{\mu} \wedge \mathbf{p}_{
u} >,$$

and basing on these relations one can regard  $\mathsf{P}_{\mu}\mathcal{G}$  and  $M_{\mu\nu}G$  as particular values of a derivative whose argument is a five-vector bivector. For an arbitrary Lorentz coordinate system one will have

$$\mathsf{P}_{\mu}\mathcal{G} = \mathsf{D}_{\mathbf{p}_{\mu}\wedge\mathbf{p}_{5}}\mathcal{G} \text{ and } \mathsf{M}_{\mu\nu}\mathcal{G} = \mathsf{D}_{\mathbf{p}_{\mu}\wedge\mathbf{p}_{\nu}}\mathcal{G},$$
 (44)

where  $\mathbf{p}_A$  is the *P*-basis associated with these coordinates. Comparing the latter formulae with formulae (41) at the origin, one can see that for any active regular basis  $\mathbf{e}_A$  and any scalar function f,

$$\mathsf{D}_{\mathbf{e}_{\mu}\wedge\mathbf{e}_{5}}f = \partial_{\mathbf{e}_{\mu}}f \text{ and } \mathsf{D}_{\mathbf{e}_{\mu}\wedge\mathbf{e}_{\nu}}f = 0.$$
 (45)

From these equations it follows that at the point with coordinates  $x^{\alpha}$ .

$$\mathsf{D}_{\mathbf{p}_{\mu}\wedge\mathbf{p}_{5}}f=\mathsf{D}_{\mathbf{p}_{\mu}^{\mathbb{Z}}\wedge\mathbf{p}_{5}}f=\mathsf{D}_{\mathbf{e}_{\mu}\wedge\mathbf{e}_{5}}f=\partial_{\mathbf{e}_{\mu}}f=\partial_{\mu}f$$

and

$$\begin{aligned} \mathsf{D}_{\mathbf{p}_{\mu}\wedge\mathbf{p}_{\nu}}f &= \mathsf{D}_{\mathbf{e}_{\mu}\wedge\mathbf{e}_{\nu}}f + x_{\nu}\mathsf{D}_{\mathbf{e}_{\mu}\wedge\mathbf{e}_{5}}f + x_{\mu}\mathsf{D}_{\mathbf{e}_{5}\wedge\mathbf{e}_{\nu}}f \\ &= x_{\nu}\partial_{\mu}f - x_{\mu}\partial_{\nu}f, \end{aligned}$$

eral case (in the latter two chains of equations and in sponds to the  $\mathcal{E}$ -component of  $\mathcal{A}$ .

equations (46) and (47) that follow,  $\mathbf{e}_A$  denotes the O-basis associated with the considered coordinates). Comparing formulae (44) with formulae (42) at the origin, one can see that for any Lorentz four-vector basis  $\mathbf{E}_{\alpha}$ ,

 $\mathbf{D}_{\mathbf{e}_{\mu}\wedge\mathbf{e}_{5}}\mathbf{E}_{\alpha} = \mathbf{0}$  and  $\mathbf{D}_{\mathbf{e}_{\mu}\wedge\mathbf{e}_{\nu}}\mathbf{E}_{\alpha} = \mathbf{E}_{\beta} (M_{\mu\nu})_{\alpha}^{\beta}$ . (46)

Consequently, for any such basis,

$$\mathsf{D}_{\mathbf{p}_A \wedge \mathbf{p}_B} \mathbf{E}_\alpha = \mathsf{D}_{\mathbf{e}_A \wedge \mathbf{e}_B} \mathbf{E}_\alpha \tag{47}$$

at all A and B. From the properties of Poincare transformations and from definition (40) it follows that for any scalar function f and any four-vector field  $\mathbf{V}$ ,

$$\mathsf{D}_{\mathbf{p}_A \wedge \mathbf{p}_B}(f\mathbf{V}) = \mathsf{D}_{\mathbf{p}_A \wedge \mathbf{p}_B} f \cdot \mathbf{V} + f \cdot \mathsf{D}_{\mathbf{p}_A \wedge \mathbf{p}_B} \mathbf{V}, \quad (48)$$

which together with equations (46) and (47) gives formulae (42) for an arbitrary four-vector field U. Similar formulae can be obtained for all other four-tensor fields.

One can now consider a more general derivative than  $\mathsf{D}_{\mathcal{H}}$  by allowing the one-parameter family  $\mathcal H$  to vary from point to point. Everywhere below, when speaking of the bivector derivative I will refer to this more general type of differentiation. According to the results obtained above, any such derivative can be uniquely fixed by specifying a certain field of fivevector bivectors. Therefore, by analogy with the covariant derivative, for any type of fields D can be formally regarded as a map that puts into correspondence to every pair consisting of a bivector field and a field of the considered type another field of that type. For example, the bivector derivative for fourvector fields can be viewed as a map

$$\mathcal{F} \wedge \mathcal{F} \times \mathcal{D} \to \mathcal{D},$$
 (49)

where  $\mathcal{F} \wedge \mathcal{F}$  is the set of all fields of five-vector bivectors. From the definition of the bivector derivative it follows that map (49) has the following formal properties: for any scalar functions f and q, any four-vector fields **U** and **V**, and any bivector fields  $\boldsymbol{\mathcal{A}}$  and  $\boldsymbol{\mathcal{B}}$ ,

$$\mathsf{D}_{(f\mathcal{A}+g\mathcal{B})}\mathbf{U} = f \cdot \mathsf{D}_{\mathcal{A}}\mathbf{U} + g \cdot \mathsf{D}_{\mathcal{B}}\mathbf{U}$$
(50a)

$$\mathsf{D}_{\mathcal{A}}(\mathsf{U}+\mathsf{V}) = \mathsf{D}_{\mathcal{A}}\mathsf{U} + \mathsf{D}_{\mathcal{A}}\mathsf{V}$$
(50b)

$$\mathsf{D}_{\mathcal{A}}(f\,\mathsf{U}) = \mathsf{D}_{\mathcal{A}}f\cdot\mathsf{U} + f\cdot\mathsf{D}_{\mathcal{A}}\mathsf{U}.$$
 (50c)

In the third equation, the action of D on the function f is determined by the rules:

which is in agreement with formulae (41) in the gen- where **A** denotes the four-vector field that corre-

The properties of D presented above are similar to the three main properties of the covariant derivative which are used for defining the latter formally. Using properties (50) for the same purpose is not very convenient, since to define the bivector derivative completely one has to supplement them with the formulae that determine the relation of D to the space-time metric, and usually from such relations one is already able to derive part of the properties expressed by equations (50). As an example, let us present the formulae that express the operator D in terms of the operator  $\nabla$  of the torsion-free *q*-conserving covariant derivative and of the linear local operator **M** defined below, both of which are completely determined by the metric. For an arbitrary four-vector field **U** one has:

$$\mathsf{D}_{\mathcal{A}^{\mathcal{E}}}\mathbf{U} = \nabla_{\mathbf{A}}\mathbf{U} \text{ and } \mathsf{D}_{\mathcal{A}^{\mathcal{E}}}\mathbf{U} = \widehat{\mathbf{M}}_{\mathbf{B}}\mathbf{U},$$
 (52)

where  $\mathbf{A}$ , as in definition (51), denotes the four-vector field corresponding to the  $\mathcal{E}$ -component of  $\mathcal{A}$ ,  $\mathbf{B}$  denotes the field of four-vector bivectors corresponding to the  $\mathcal{Z}$ -component of  $\mathcal{A}$ , and the operator  $\widehat{\mathbf{M}}$ , which depends linearly on its argument, has the following components in an arbitrary four-vector basis  $\mathbf{E}_{\alpha}$ :

$$\mathbf{\widehat{M}}_{\mathbf{E}_{\alpha}\wedge\mathbf{E}_{\beta}}\mathbf{E}_{\mu}=\mathbf{E}_{\nu}(M_{\alpha\beta})_{\mu}^{\nu}.$$

It is easy to see that properties (50b) and (50c) follow from formulae (52) and property (50a), and property (50a) itself follows from equations (52) and the following simpler property:

$$\mathsf{D}_{(\mathcal{A}+\mathcal{B})}\mathbf{U}=\mathsf{D}_{\mathcal{A}}\mathbf{U}+\mathsf{D}_{\mathcal{B}}\mathbf{U},$$

which is similar to the first equation in definition (51) and which, together with equations (52), can serve as a definition of the bivector derivative for four-vector fields.

For the bivector derivative one can define the analogs of connection coefficients. Namely, for any set of basis four-vector fields  $\mathbf{E}_{\alpha}$  and any set of basis five-vector fields  $\mathbf{e}_{A}$  one puts

$$\mathsf{D}_{AB}\mathbf{E}_{\mu} = \mathbf{E}_{\nu}\Gamma^{\nu}_{\ \mu AB},\tag{53}$$

where  $\mathsf{D}_{AB} \equiv \mathsf{D}_{\mathbf{e}_A \wedge \mathbf{e}_B}$ . According to equations (46), for any Lorentz four-vector basis and any standard five-vector basis associated with it, one has

$$\Gamma^{\mu}_{\nu\alpha5} = -\Gamma^{\mu}_{\nu5\alpha} = 0$$
  

$$\Gamma^{\mu}_{\nu\alpha\beta} = -\Gamma^{\mu}_{\nu\beta\alpha} = (M_{\alpha\beta})^{\mu}_{\nu}.$$
(54)

The bivector connection coefficients for all other bases can be found either by using the following transformation formula:

$$\Gamma^{\prime\mu}_{\nu AB} = (\Lambda^{-1})^{\mu}_{\sigma} \Gamma^{\sigma}_{\tau ST} \Lambda^{\tau}_{\nu} L^{S}_{A} L^{T}_{B} + (\Lambda^{-1})^{\mu}_{\sigma} (\mathsf{D}_{ST} \Lambda^{\sigma}_{\nu}) L^{S}_{A} L^{T}_{B},$$
(55)

which corresponds to the transformations  $\mathbf{E}'_{\alpha} = \mathbf{E}_{\beta} \Lambda^{\beta}_{\alpha}$  and  $\mathbf{e}'_{A} = \mathbf{e}_{B} L^{B}_{A}$  of the four- and five-vector basis fields, or by using formulae (52). In particular, for an arbitrary four-vector basis and the corresponding active regular five-vector basis one has

$$\Gamma^{\mu}_{\nu\alpha5} = \Gamma^{\mu}_{\nu\alpha} \text{ and } \Gamma^{\mu}_{\nu\alpha\beta} = (M_{\alpha\beta})^{\mu}_{\nu}, \qquad (56)$$

where  $\Gamma^{\mu}_{\nu\alpha}$  are ordinary four-vector connection coefficients associated with  $\nabla$ .

In order to define the bivector derivative for scalar, four-vector and four-tensor fields in the case of arbitrary Riemannian geometry, one may observe that in the case of flat space-time D is determined only by the metric, and since with respect to its metric properties any sufficiently smooth space-time manifold is locally flat, the bivector derivative in the general case can be defined by postulating that in local Lorentz coordinates it has the same form at any space-time geometry. For scalar fields this means that the bivector derivative of an arbitrary function f is given by formula (45), where  $\mathbf{e}_A$  is now an active regular basis at the considered point. For four-vector fields the above assertion means that the bivector derivative of the basis fields  $\mathbf{E}_{\alpha}$  corresponding to any system of local Lorentz coordinates at the considered point is given by formula (46), where  $\mathbf{e}_A$  is the associated active regular five-vector basis. Furthermore, one should assume that in the general case, too, the bivector derivative has the properties expressed by equations (50), which will enable one to define the derivative  $D_{\mathcal{A}} W$  of any four-vector field W along any five-vector bivector  $\boldsymbol{\mathcal{A}}$ , and that the bivector derivative of the contraction and tensor product obeys the Leibniz rule, which will enable one to define the action of operator D on all other four-tensor fields.

The bivector derivative in the general case can also be defined without referring to local Lorentz coordinates. Instead, one can postulate that as in the case of flat space-time, it is expressed according to formulae (51) and (52) for scalar fields in terms of the directional derivative and for four-vector fields in terms of the torsion-free g-conserving ordinary covariant derivative  $\nabla$  and of the local operator  $\widehat{\mathbf{M}}$  introduced above.

8. The bivector derivative defined above possesses one important property: at any five-vector affine connection  $\Box$  with respect to which the metric tensor g is covariantly constant, the five-vector covariant derivative of any scalar, four-vector or four-tensor field is expressed linearly in terms of its bivector derivatives. More precisely this property can be formulated as follows: at any given five-vector affine connection for which  $\Box g = 0$ , at each space-time point there exists such a linear map  $\sigma$  from the tangent space of five-vectors to the tangent space of five-vector bivectors that for any five-vector **u** at that point

$$\square_{\mathbf{u}} \mathcal{G} = \mathsf{D}_{\sigma(\mathbf{u})} \mathcal{G} \tag{57}$$

for any field  $\mathcal{G}$  from the considered class of fields.

Owing to the linearity of  $\sigma$ , the image  $\sigma(\mathbf{u})$  can be presented as a contraction of  $\mathbf{u}$  with a certain fivevector 1-form,  $\tilde{\mathbf{s}}$ , whose values are five-vector bivectors. In any active regular basis this 1-form has the components

$$s^{\alpha 5}{}_{A} = -s^{5\alpha}{}_{A} = \delta^{\alpha}{}_{A} \text{ and } s^{\alpha \beta}{}_{A} = S^{\alpha \beta}{}_{A}, \quad (58)$$

where  $S_{A}^{\alpha\beta}$  are the components of another five-vector 1-form,  $\mathbf{\tilde{S}}$ , whose values are *four*-vector bivectors and which can be regarded as a generalization of the so-called contorsion tensor. In terms of the derivatives  $\square$  and  $\nabla$  and operator  $\widehat{\mathbf{M}}$ , this latter 1-form can be defined as follows: for any four-vector field  $\mathbf{W}$  and any five-vector  $\mathbf{u}$ 

$$\left(\mathbf{\Box}_{\mathbf{u}} - \nabla_{\mathbf{u}}\right) \mathbf{W} = \widehat{\mathbf{M}}_{<\widetilde{\mathbf{S}},\mathbf{u}>} \mathbf{W}.$$
 (59)

From the fact that Poincare transformations conserve the inner product g for four-vectors it follows that the bivector derivative of the metric tensor is identically zero, which means that requiring equation (57) to hold is equivalent to requiring g to be covariantly constant. As one will see below, this fact can be used to specify another particular case of the connection for five-vector fields, which is more general than the one considered above, the one where there exist local symmetry (13).

To explain the role equation (57) will play, let us go back and see how one arrives at the symmetry principle (13). As is known, in general relativity two constraints are imposed on ordinary (four-vector) connection: (i) that with respect to it the metric tensor be covariantly constant and (ii) that  $\nabla$  be torsionfree. Let us now try to determine the corresponding connection for five-vector fields. If one simply generalizes the above two constraints on ordinary connection to the case of five-vector fields, i.e. if one requires that the inner product *h* regarded as a five-tensor be covariantly constant and that five-vector torsion be identically zero<sup>2</sup>, one will obtain a connection given

$$\nabla_{\mathbf{u}}\mathbf{v} - \nabla_{\mathbf{v}}\mathbf{u} - [\mathbf{u}, \mathbf{v}],$$

by equations (11), (12) and (14) with  $\kappa = 0$ , at which, as it has been pointed out above, talking about fivevectors just does not make sense. It is not difficult to show that to enable a five-vector from  $\mathcal{Z}$  to acquire a nonzero  $\mathcal{E}$ -component in the process of transport at the same type of connection for four-vector fields, one has to weaken the constraint  $\nabla h = 0$ , replacing it with the less stringent requirement  $\nabla q = 0$ , where g is regarded as a *five*-tensor. Together with the requirement of zero five-vector torsion (and condition (10), which is always assumed to be imposed) this weaker constraint will give one the desired connection for four-vector fields, but for the connection coefficients that determine the  $\mathcal{E}$ -component of a transported five-vector one will obtain only that in any regular basis  $G_{5\mu}^5 = G_{[\mu\nu]}^5 = 0$ , whereas the symmetric part of  $G^5_{\mu\nu}$  will be completely undetermined.

To fix the connection for five-vector fields in this particular case more precisely, one can use a trick which is often done in mathematics: one should replace the mentioned constraints on  $\nabla$  for fourvector fields with different but equivalent requirements whose generalization to the case of five-vector fields would determine not only  $G_{5\mu}^5$  and  $G_{[\mu\nu]}^5$ , but also the symmetric part of  $G_{\mu\nu}^5$  (at least up to normalization). The analog of symmetry principle (13) for four-vector fields is just this equivalent requirement.

Let us now consider a more general case of ordinary affine connection where the latter is constrained only by the requirement of covariant constancy of q, and find the corresponding connection  $\nabla$  for five-vector fields. If one imposes only condition (10) and the requirement  $\nabla g = 0$  for g regarded as a five-tensor, one will obtain a connection for which there hold equations (11) and (12), in which  $\Gamma^{\alpha}_{\beta\mu}$  are the standard connection coefficients for four-vector fields in the Riemann-Cartan geomentry. However, both  $G^5_{5\mu}$ and  $G^{5}_{\mu\nu}$  will now be completely arbitrary. This arbitrariness can be reduced by using the same trick as above: the constraint  $\nabla g = 0$  for four-vector fields should be replaced with an equivalent requirement whose generalization to the case of five-vector fields would fix the connection coefficients  $G_{5\mu}^5$  and  $G_{\mu\nu}^5$ to a greater extent. It turns out that to this end one can use equation (57) with  $\square$  replaced by  $\nabla$ . As a result, one will obtain a connection for five-vector fields which is completely fixed by space-time metric and by ordinary four-vector torsion or, equivalently, by the metric and by the components  $S^{\alpha\beta}_{\ \mu}$  of the 1-form  $\widetilde{\mathbf{S}}$ 

$$[\mathbf{u},\mathbf{v}]f = \mathbf{u}(\mathbf{v}f) - \mathbf{v}(\mathbf{u}f).$$

<sup>&</sup>lt;sup>2</sup>By analogy with its four-vector counter-part, five-vector torsion can be defined as a five-vector-valued five-vector 2-form whose contraction with any five-vector bivector  $\mathbf{u} \wedge \mathbf{v}$  equals (or is proportional to)

where  $[\mathbf{u},\mathbf{v}]$  is the commutator of the five-vector fields  $\mathbf{u}$  and  $\mathbf{v},$  which by definition is a five-vector field whose action on any

scalar function f is given by the formula

introduced above, as is readily seen from relation (32) and definition (59). Moreover, the same method can be used to specify a still more general case of the connection for five-vector fields, where instead of  $\nabla$  one has  $\Box$ , and the latter satisfies equation (57) in its original form.

It is evident that in order to generalize equation (57) to the case of five-vector fields one should first define for the latter the notion of the bivector derivative. As in the case of four-vector fields, in flat space-time this can be done according to formula (40), where  $\mathcal{G}$  is now an arbitrary five-vector field. From formula (19) one then obtains that for an arbitrary O-basis

$$G^{A}_{B\mu5} = 0 \text{ and } G^{A}_{B\mu\nu} = (M_{\mu\nu})^{A}_{B}, \qquad (60)$$

where  $(M_{KL})_B^A \equiv \delta_L^A g_{KB} - \delta_K^A g_{LB}$  (according to the definition of g for five-vectors, in any standard basis  $g_{\alpha 5} = g_{5\alpha} = g_{55} = 0$ ) and the bivectors connection coefficients are defined according to the formula

$$\mathsf{D}_{KL}\mathbf{e}_A = \mathbf{e}_B G^B_{\ AKL}$$

In the case of arbitrary Riemannian geometry the bivector derivative for five-vector fields can be defined by postulating that formulae (60) hold for any active regular basis associated with a system of local Lorentz coordinates at the considered point.

Though such a definition of the bivector derivative for five-vector fields is quite permissible, it is not difficult to see that in that case the relation between  $\square$  and  $\square$  expressed by equation (57) cannot exist. Indeed, the five-vector covariant derivative of an arbitrary field represented by a purely differential operator will in general have a nonzero  $\mathcal{E}$ -component, whereas the bivector derivative of any such field defined as described above will always have a zero  $\mathcal{E}$ component, as is readily seen from equations (60). In view of this, if one does wish that equation (57)could hold for five-vector fields as well, one should try to define the derivative D for the latter in some other way. To see how this can be done, let us first observe that since the five-vector covariant derivative  $\square$  possesses property (35), for the bivector derivative one should require that

$$\mathbf{v} \equiv \mathbf{w} \pmod{R} \implies \mathsf{D}_{\mathcal{A}} \mathbf{v} \equiv \mathsf{D}_{\mathcal{A}} \mathbf{w} \pmod{R}$$

for any bivector field  $\mathcal{A}$ . Furthermore, it seems reasonable to suppose that as in the case of four-vector fields, the bivector derivative of any five-vector field should be determined only by space-time metric, and since with respect to its metric properties flat space-time is *homogeneous* and *isotropic*, one should require

that in the case of the latter the bivector connection coefficients have the same form for any *O*-basis. It is not difficult to show that the most general form of the bivector connection coefficients for such a basis that satisfy the above two requirements is the following:

$$G^{A}_{B\mu5} \propto (M_{\mu5})^{A}_{B}$$
 and  $G^{A}_{B\mu\nu} = (M_{\mu\nu})^{A}_{B}$ 

It is apparent that equations (60) are a particular case of the latter formulae, where the proportionality factor in the first relation is zero. To find the value of this factor at which equation (57) could hold, one may consider the particular case of five-vector connection where  $\Box_5 = 0$  and where there exists local symmetry (13). This way for an arbitrary *O*-basis one finds that

$$G^{A}_{B\mu5} = -(M_{\mu5})^{A}_{B}$$
 and  $G^{A}_{B\mu\nu} = (M_{\mu\nu})^{A}_{B}$ . (61)

By using the obvious formula for transformation of bivector connection coefficients one can find that for an arbitrary active regular basis

$$G^{\alpha}_{\ \beta\mu5} = G^{\alpha}_{\ \beta\mu}, \ G^{5}_{\ \beta\mu5} = -g_{\beta\mu}, G^{A}_{\ 5\mu5} = 0, \ G^{\alpha}_{\ \beta\mu\nu} = (M_{\mu\nu})^{\alpha}_{\beta},$$
(62)

where  $G^{\alpha}_{\beta\mu}$  are the connection coefficients associated with the torsion-free *g*-conserving ordinary covariant derivative  $\nabla$  fixed for five-vector fields by equations (11), (12) and (14). From the latter formulae it follows that the operator  $D_{\mathcal{A}}$  defined this way can be presented as a sum of the operator  $\nabla$ , whose argument will be the five-vector from  $\mathcal{Z}$  that corresponds to the  $\mathcal{E}$ -component of  $\mathcal{A}$ , and of the local operator  $\widehat{\mathbf{M}}$ , whose components in any standard five-vector basis are  $(M_{KL})^A_B$  and whose argument will be the  $\mathcal{Z}$ component of  $\mathcal{A}$ . For an arbitrary field  $\mathbf{u}$  one will thus have

$$\mathsf{D}_{\mathcal{A}}\mathbf{u} = \nabla_{\mathbf{a}}\mathbf{u} + \widehat{\mathbf{M}}_{\mathcal{A}^{\mathcal{Z}}}\mathbf{u},\tag{63}$$

where **a** is the five-vector from  $\mathcal{Z}$  that corresponds to  $\mathcal{A}^{\mathcal{E}}$ . To define the bivector derivative for five-vector fields in the case of arbitrary Riemannian geometry one can either postulate that formulae (61) hold for any active regular basis associated with a system of local Lorentz coordinates at the considered point or postulate that relation (63) holds in curved space-time as well.

Requiring equation (57) to be valid for an arbitrary five-vector field  $\mathcal{G}$ , one obtains the following relation between the connection coefficients associated with  $\square$  and D:

$$H^A_{BC} = G^A_{BKL} \, s^{|KL|}_{C}.$$

By using formulae (58) and (62), for an arbitrary active regular basis one finds that

$$\begin{aligned}
H^{\alpha}_{\ \beta\mu} &= G^{\alpha}_{\ \beta\mu} - s^{\alpha}_{\ \beta\mu}, H^{\alpha}_{\ \beta5} = -s^{\alpha}_{\ \beta5}, \\
H^{\alpha}_{\ 5\mu} &= H^{\alpha}_{\ 55} = 0, \quad H^{5}_{\ \beta\mu} = -g_{\beta\mu}, \\
H^{5}_{\ \beta5} &= H^{5}_{\ 5\mu} = H^{5}_{\ 55} = 0,
\end{aligned} \tag{64}$$

where  $s^{\alpha}_{\beta C} \equiv g_{\beta \omega} s^{\alpha \omega}_{C}$ . This particular connection for five-vector fields and the rules of parallel transport that correspond to it are discussed in more detail in part VI of the long version [8].

9. Let us now derive a possible set of field equations that would determine the geometry of space-time in the case of connection (64). To this end, let us first introduce the five-vectors analog of the curvature tensor, **R**. The latter can be defined in the usual manner: as a five-vector 2-form whose values are tensors of rank (1, 1) over  $V_5$  and whose contraction with any five-vector bivector  $\mathbf{u} \wedge \mathbf{v}$  equals

$$\Box_{\mathbf{u}}\Box_{\mathbf{v}} - \Box_{\mathbf{v}}\Box_{\mathbf{u}} - \Box_{[\mathbf{u},\mathbf{v}]}.$$

From the latter formula one can easily obtain a familiar expression for the components of  $\mathbf{R}$  in a fivevector basis for which all the commutators are zero, in terms of the corresponding five-vector connection coefficients:

$$\begin{aligned} R^A_{\ BCD} &= \partial_C H^A_{\ BD} - \partial_D H^A_{\ BC} \\ &+ H^A_{\ KC} H^K_{\ BD} - H^A_{\ KD} H^K_{\ BC}. \end{aligned}$$

For the connection that satisfies condition (57) one finds that in any active regular basis

$$R^{\alpha}_{5CD} = R^{5}_{5CD} = 0, \ R^{\alpha}_{\beta\mu\nu} = R^{(\nabla)\alpha}_{\ \beta\mu\nu}, R^{\alpha}_{\beta\mu5} = -\{\partial_{\mu}s^{\alpha}_{\beta5} + H^{\alpha}_{\omega\mu}s^{\omega}_{\beta5} + s^{\alpha}_{\omega5}H^{\omega}_{\beta\mu}\}, \ (65) R^{5}_{\beta CD} = -2g_{\beta\omega}s^{\omega}_{[CD]},$$

where  $R^{(\nabla) \alpha}_{\ \beta\mu\nu}$  are the components of the Riemann tensor corresponding to the ordinary covariant derivative  $\nabla$  related to  $\Box$  according to equation (32), in the associated four-vector basis. From the fact that g is covariantly constant it follows that

$$g_{\alpha\omega}R^{\omega}_{\ \beta CD} + g_{\beta\omega}R^{\omega}_{\ \alpha CD} = 0$$

This property of  $\mathbf{R}$  and the property of the latter expressed by the first double equation in (65) enable one to associate with it a certain five-vector 2-form,  $\mathbf{K}$ , whose values are five-vector bivectors and whose components are related to those of  $\mathbf{R}$  as follows:

$$K^{A\beta}_{\ CD} = -K^{\beta A}_{\ CD} = g^{\beta \omega} R^A_{\ \omega CD},$$

where  $g^{\beta\omega}$  is the inverse of the 4×4 matrix  $g_{\beta\omega}$ . From formulae (65) one finds that in any active regular basis

$$\begin{split} K^{\alpha 5}{}_{CD} &= -K^{5\alpha}{}_{CD} = 2s^{\alpha}{}_{[CD]}, \\ K^{\alpha \beta}{}_{\mu 5} &= -\left\{ \partial_{\mu}s^{\alpha \beta}{}_{5} + H^{\alpha}{}_{\omega \mu}s^{\omega \beta}{}_{5} + H^{\beta}{}_{\omega \mu}s^{\alpha \omega}{}_{5} \right\}, \quad (66) \\ K^{\alpha \beta}{}_{\mu \nu} &= g^{\beta \omega}R^{(\nabla)}{}_{\omega \mu \nu}^{\alpha}. \end{split}$$

Let us now consider a situation where one has several matter fields,  $\vec{u}_{\ell}$ , whose values can be vectors or

tensors of any nature (the index  $\ell$  lables the fields, not their components) and where the Lagrangian L that describes these fields is a function of the values of the fields themselves and of their five-vector covariant derivatives. As in ordinary theory, from the requirement of local isotropy and homogeneity of space-time one can derive certain relations, from which, by using the equations of motion for the considered fields, one can then derive equations that can be interpreted as a conservation law for a certain tensor quantity whose components in the limit of flat space-time coincide with the five-vector analogs of the Noether currents associated with the symmetry under global Poincare transformations. As is shown in part VI of the long version [8], this tensor quantity, which I will denote as  $\mathcal{M}$ , can be chosen to have the following components in an active regular basis:

$$\mathcal{M}^{A}_{\mu 5} = -\mathcal{M}^{A}_{5\mu} = \delta^{A}_{\mu} \mathbf{L} - \sum_{\ell} \frac{\partial \mathbf{L}}{\partial (\mathbf{\Box}_{A} \vec{\mathbf{u}}_{\ell})} \mathbf{\Box}_{\mu} \vec{\mathbf{u}}_{\ell} \quad (67)$$

and

$$\mathcal{M}^{A}_{\mu\nu} = -\sum_{\ell} \frac{\partial \mathbf{L}}{\partial (\mathbf{\Box}_{A} \vec{\mathrm{u}}_{\ell})} \mathsf{D}_{\mu\nu} \vec{\mathrm{u}}_{\ell} \,. \tag{68}$$

The corresponding conservation laws are

$$(\overset{*}{\Box}_{A}\mathcal{M})_{\mu 5}^{A} = \mathcal{M}_{ST}^{A} K_{\mu A}^{|ST|}, (\overset{*}{\Box}_{A}\mathcal{M})_{\mu \nu}^{A} = 0, \quad (69)$$

where, as usual, the vertical bars around the indices mean that summation extends only over K < L and the operator  $\square_A \equiv \square_A + 2 s^K_{[AK]}$  is the direct generalization of the corresponding four-vector operator  $\nabla_{\alpha} \equiv \nabla_{\alpha} - 2 T_{\alpha\omega}{}^{\omega}$ , where  $T_{\alpha\beta}{}^{\mu}$  are the components of four-vector torsion. By analogy with the usual terminology, the expressions in the left-hand sides of equations (69) will be called *modified* divergences.

It is apparent that the components  $\mathcal{M}^{\mu}_{5\alpha} = -\mathcal{M}^{\mu}_{\alpha5}$ and  $\mathcal{M}^{\mu}_{\alpha\beta}$  coincide with the components of the canonical stress-energy and angular momentum tensors, respectively, as is stated by equations (27). In addition to these one has two new quantities:

$$\mathcal{M}_{\mu 5}^{5} = -\mathcal{M}_{5\mu}^{5} = -\sum_{\ell} \frac{\partial \mathbf{L}}{\partial (\mathbf{\Box}_{5} \vec{\mathrm{u}}_{\ell})} \, \mathbf{\Box}_{\mu} \vec{\mathrm{u}}_{\ell}$$

and

$${\cal M}^5_{\mu
u} = -\sum_\ell \, rac{\partial {f L}}{\partial ({f \Box}_5 ec {f U}_\ell)} \, {\sf D}_{\mu
u} ec {f U}_\ell \, ,$$

whose geometric interpretation will be discussed elsewhere. Let us only observe that the first of these quantities has no effect on the conservation laws for  $\mathcal{M}$ , since in the left-hand side of the first equation in (69) it appears only in the term

$$\mathcal{M}_{\sigma 5}^{5} K^{\sigma 5}{}_{\mu 5} = 2 \, \mathcal{M}_{\sigma 5}^{5} s_{\, [\mu 5]}^{\sigma} = \mathcal{M}_{\sigma 5}^{5} s_{\, \mu 5}^{\sigma} \,,$$

and in the right-hand side of the same equation, only and finally in the term

$$( \stackrel{*}{\Box}_{5} \mathcal{M} )^{5}_{\mu 5} = \mathcal{M}^{5}_{\mu 5;5} + (H^{K}_{K5} - H^{K}_{5K}) \mathcal{M}^{5}_{\mu 5} = -\mathcal{M}^{5}_{\sigma 5} H^{\sigma}_{\mu 5} = \mathcal{M}^{5}_{\sigma 5} s^{\sigma}_{\mu 5} .$$

Consequently, its contributions cancel out.

We are now ready to discuss the possible field equations for connection (64). Let us first observe that  $\Box$ can be regarded as a composite structure consisting of an ordinary affine connection  $\nabla$ , which is related to  $\square$  by equation (32), and of another structure, which in the case we are now considering can be fixed by a field of four-vector bivectors whose components in any four-vector basis are proportional to the components  $s_{5}^{\alpha\beta}$  of the 1-form  $\tilde{\mathbf{s}}$  introduced above, in the associated regular five-vector basis. Let us now recall that the Einstein and Kibble–Sciama equations can be obtained from the action principle if the Lagrangian describing the geometry of space-time is taken (in our notations) to be (-1/2k)R, where k is Newton's gravitational constant times  $8\pi c^{-4}$  and R is the curvature scalar constructed out of the fourvector Riemann tensor, and the varied parameters are the components of the metric tensor and the components of the four-vector torsion tensor. Let us suppose that the gravitaional equations in the case of five-vector affine connection can be obtained in a similar way. By virtue of equations (66) and owing to the antisymmetry of the quantities  $s_{5}^{\alpha\beta}$  in their upper indices, one has

$$\begin{array}{rcl} K^{AB}_{\phantom{AB}AB} & = & 2K^{\alpha 5}_{\phantom{\alpha 5}\alpha 5} + K^{\alpha \beta}_{\phantom{\alpha \beta}\alpha \beta} \\ & = & 4s^{\alpha}_{\phantom{\alpha 5}\alpha 5} + g^{\beta \omega} R^{(\nabla) \, \alpha}_{\phantom{(\nabla) \alpha}\omega \alpha \beta} \; = \; R \, , \end{array}$$

and since the components  $R^{(\nabla) \alpha}_{\beta \mu \nu}$  are independent of  $s^{\alpha\beta}_{5}$ , to obtain a full system of equations from the action principle in the case of five-vector connection (64), to the Lagrangian (-1/2k)R one should add some additional term, which I will denote as  $\mathbf{L}_{add}$ . Thus.

$$\mathbf{L}_{\text{geom}} = (-1/2k)R + \mathbf{L}_{\text{add}}$$

As varied parameters let us choose  $g_{\mu\nu}$  and  $T_{\alpha\beta}^{\ \mu} =$  $-s^{\mu}_{\ \ \ \ \alpha\beta}$ , and also the six quantities  $s^{\alpha\beta}_{\ \ 5}$ . By direct calculation one obtains the following equations:

$$G^{\{\mu\nu\}} - (\stackrel{*}{\nabla}_{\omega} T^{(\mathrm{mod})})^{\mu\omega\nu} - (\stackrel{*}{\nabla}_{\omega} T^{(\mathrm{mod})})^{\nu\omega\mu} + kg^{\mu\nu} \mathbf{L}_{\mathrm{add}} + 2k(\delta \mathbf{L}_{\mathrm{add}}/\delta g_{\mu\nu}) = k\Theta^{\{\mu\nu\}} + \frac{1}{2}k(\stackrel{*}{\nabla}_{\omega}\Sigma)^{\mu\omega\nu} + \frac{1}{2}k(\stackrel{*}{\nabla}_{\omega}\Sigma)^{\nu\omega\mu} - kg_{\sigma\tau}s^{\sigma\{\mu}{}_{5}\mathcal{M}^{\nu\}\tau 5},$$
(70)

then

('

$$T^{(\text{mod}) \mu\lambda\nu} - T^{(\text{mod}) \nu\lambda\mu} + T^{(\text{mod}) \mu\nu\lambda} - kg^{\lambda\omega} (\delta \mathbf{L}_{\text{add}} / \delta T_{\mu\nu}^{\omega})$$
(71)  
$$= -\frac{1}{2} k (\Sigma^{\mu\lambda\nu} - \Sigma^{\nu\lambda\mu} + \Sigma^{\mu\nu\lambda}),$$

$$\delta \mathbf{L}_{\mathrm{add}} / \delta s^{\alpha\beta}{}_{5} = \frac{1}{2} \mathcal{M}^{5}_{\alpha\beta} , \qquad (72)$$

where  $G^{\mu\nu}$  are the components of the four-vector Einstein tensor,  $T^{(\text{mod}) \mu \omega \nu} = g^{\mu \sigma} g^{\omega \tau} (T_{\sigma \tau}^{\ \nu} + \delta^{\nu}_{\sigma} T_{\tau \rho}^{\ \rho} \delta^{\nu}_{\tau} T_{\sigma\rho}{}^{\rho}$ ,  $\mathcal{M}^{\mu\nu5} = g^{\mu\sigma} g^{\nu\tau} \mathcal{M}^{5}_{\sigma\tau}$ , the derivative  $\stackrel{*}{\nabla}_{\omega}$ acts on  $T^{(\text{mod})}$  and  $\Sigma$  as on four-tensors, and the components of the stress-energy and angular momentum tensors,  $\Theta^{\mu\nu}$  and  $\Sigma^{\mu\nu\alpha}$  are identified as in the convensional theory of gravity with spin and torsion (see, for example, review [9]). One should observe that none of the components  $\mathcal{M}^{5}_{\mu 5}$  act as a source.

Let us now try to select  $\mathbf{L}_{\mathrm{add}}$  in such a way that the field equations resulting from equations (70)-(72)in which the role of the source is played by  $\mathcal{M}^{\alpha}_{\mu 5}$  and  $\mathcal{M}^{\alpha}_{\mu\nu}$  would differ as little as possible from the Einstein and Kibble–Sciama equations, respectively. In the latter case this can be achieved quite easily: one has only to require that  $\mathbf{L}_{add}$  be *independent* of  $T_{\mu\nu}^{\alpha}$ . Equation (71) will then give

$$T^{(\text{mod})\ \alpha\beta\mu} = -\frac{1}{2}k\Sigma^{\alpha\beta\mu},\tag{73}$$

which is exactly the Kibble-Sciama equation that relates four-vector torsion to spin.<sup>3</sup> Substituting this value of  $T^{(mod)}$  into equation (70), one obtains

$$G^{\{\mu\nu\}} + kg^{\mu\nu}\mathbf{L}_{add} + 2k(\delta\mathbf{L}_{add}/\delta g_{\mu\nu}) = k\Theta^{\{\mu\nu\}} - kg_{\sigma\tau}s^{\sigma\{\mu}{}_{5}\mathcal{M}^{\nu\}\tau 5}.$$
(74)

It is impossible in general to get rid of the second term in the left-hand side of this equation, and as we will see below, there is no need to. One can, however, try to select  $\mathbf{L}_{add}$  in such a way that the last term in the left-hand side would calcel out with the last term in the right-hand side. This requirement gives one the second condition on  $\mathbf{L}_{add}$ :

$$\delta \mathbf{L}_{\mathrm{add}} / \delta g_{\mu\nu} = -\frac{1}{2} g_{\sigma\tau} s^{\sigma\{\mu}{}_{5} \mathcal{M}^{\nu\}\tau 5}, \qquad (75)$$

and equation (74) then acquires the form

$$G^{\{\mu\nu\}} + kg^{\mu\nu}\mathbf{L}_{add} = k\Theta^{\{\mu\nu\}}.$$

<sup>&</sup>lt;sup>3</sup>Some authors hide the factor  $-\frac{1}{2}$  by defining the fourvector torsion tensor with a different sign and by choosing a different normalization for the spin angular momentum. The simplest way to compare the definitions of these quantities adopted in a particular paper with ours is to evaluate the proportionality factor between  $\Sigma^{\mu}_{\ \alpha\beta;\mu} - 2 T_{\mu\omega}^{\ \omega} \Sigma^{\mu}_{\ \alpha\beta}$  and  $g_{\beta\mu} \Theta^{\mu}_{\ \alpha} - g_{\alpha\mu} \Theta^{\mu}_{\ \beta}$ (in our case it is unity) and the proportionality factor between  $T_{\alpha\beta}^{\ \mu}$  and  $\Gamma^{\alpha}_{[\mu\nu]}$  (in our case the latter is unity, too, provided the definition of the four-vector connection coefficients is the same as ours). The sign and normalization of the stress-energy tensor is fixed by the condition that  $\Theta_0^0$  be the energy density of matter.

As one can see, the symmetric parts of  $G^{\mu\nu}$  and differentiating (80) and using the covariant constancy  $k\Theta^{\mu\nu}$  are no longer equal to each other. However, one can try to choose  $\mathbf{L}_{add}$  in such a way that the antisymmetric parts of these tensors would coincide:

$$G^{[\mu\nu]} = k\Theta^{[\mu\nu]}. \tag{76}$$

If one succeeds, then after adding the latter two equations one will obtain

$$G^{\mu\nu} + kg^{\mu\nu}\mathbf{L}_{add} = k\Theta^{\mu\nu}.$$
 (77)

To derive from requirement (76) a constraint on  $\mathbf{L}_{add}$ , let us recall the differential identity that relates the modified four-dimensional divergence of  $T^{(mod)}$  to the antisymmetric part of the Einstein tensor:

$$\left(\nabla_{\alpha}^{*} T^{(\mathrm{mod})}\right)_{\mu\nu}^{\alpha} = G_{[\mu\nu]}.$$

Combining this identity with equation (76) and using (69) and (73), one finds that

$$\mathcal{M}^{\mu 5}_{\ \omega} s^{\omega}_{\ \nu 5} - s^{\mu}_{\ \omega 5} \mathcal{M}^{\omega 5}_{\ \nu} = 0, \tag{78}$$

meaning that the quantities  $s^{\mu}_{\ \nu 5}$  and  $\mathcal{M}^{\mu}_{\ \nu}{}^{5}$  regarded as matrices with respect to the indices  $\mu$  and  $\nu$  should commute with each other. Together with equation (72), the latter relation gives us one more constraint on  $\mathbf{L}_{add}$ .

Let us finally recall that in the case of arbitrary four-vector torsion the Einstein tensor satisfies the differential identity

$$\left(\mathop{\nabla}\limits_{\alpha}^{*} G\right)_{\mu}^{\alpha} = R^{\sigma\tau}_{\ \mu\alpha} T^{(\mathrm{mod})\,\alpha}_{\sigma\tau} - 2 T^{\sigma}_{\mu\alpha} G^{\alpha}_{\sigma}.$$

Combining the latter with equations (73) and (77)and using (66) and (69), one obtains the last condition on  $\mathbf{L}_{add}$ :

$$\partial_{\mu} \mathbf{L}_{\text{add}} = \frac{1}{2} \left\{ \partial_{\mu} s^{\sigma\tau}{}_{5} + H^{\sigma}{}_{\omega\mu} s^{\omega\tau}{}_{5} + H^{\tau}{}_{\omega\mu} s^{\sigma\omega}{}_{5} \right\} \mathcal{M}^{5}_{\sigma\tau}.$$
(79)

The simplest way to satisfy requirement (78) is to take  $s_{\sigma\tau5}$  proportional to  $\mathcal{M}_{\sigma\tau}^5$ . As one can see from equation (72), for that one should choose

$$\mathbf{L}_{\mathrm{add}} = a \cdot g_{\alpha\sigma} g_{\beta\tau} h^{55} s^{\alpha\beta}{}_5 s^{\sigma\tau}{}_5, \qquad (80)$$

where a is a certain constant and the factor  $h^{55}$  has been introduced so that the latter would not depend on the normalization of the fifth basis vector. Accordingly, one has

$$2a h^{55} s_{\sigma\tau 5} = \frac{1}{2} \mathcal{M}^{5}_{\sigma\tau}.$$
 (81)

It is a simple matter to check that at such  $\mathbf{L}_{add}$  conditions (79) and (75) are also satisfied. Indeed, by

of q, one obtains that

$$\partial_{\mu} \mathbf{L}_{\mathrm{add}} = 2a h^{55} s_{\sigma\tau 5} \cdot s^{\sigma\tau}_{5;\mu} = \frac{1}{2} \{ \partial_{\mu} s^{\sigma\tau}_{5} + H^{\sigma}_{\omega\mu} s^{\omega\tau}_{5} + H^{\tau}_{\omega\mu} s^{\sigma\omega}_{5} \} \mathcal{M}^{5}_{\sigma\tau} .$$

Similarly, by varying (80) with respect to  $g_{\mu\nu}$  and using (81), one obtains

$$\begin{split} \delta \mathbf{L}_{\text{add}} &= \delta g_{\mu\nu} \cdot 2a \, g_{\sigma\tau} h^{55} s^{\mu\sigma}_{5} s^{\nu\tau}_{5} \\ &= \delta g_{\mu\nu} \cdot \left\{ \frac{1}{2} \, g_{\sigma\tau} s^{\mu\sigma}_{5} \, \mathcal{M}^{\nu\tau5} \right\}, \end{split}$$

whence follows (75).

The dimension of the constant a can be easily established from formula (80). Since in the normalized regular basis  $h^{55}$  is dimensionless and the components  $s^{\alpha\beta}_{5}$  have the same dimension as  $s^{\alpha\beta}_{\mu}$ , the expression following a in formula (80) should have the same dimension as R, so  $a^{-1}$  should have the same dimension as k. In view of this, one may put  $a = (-1/2k) \rho$ , where  $\rho$  is some unknown dimensionless constant, whose value should be found experimentally. One will then have

$$\mathbf{L}_{\text{geom}} = (-1/2k) \left( R + \varrho \cdot g_{\alpha\sigma} g_{\beta\tau} h^{55} s^{\alpha\beta}{}_5 s^{\sigma\tau}{}_5 \right), \quad (82)$$

and the gravitational equations in the four-tensor notations will acquire the following form:

$$G_{\mu\nu} - g_{\mu\nu} \frac{1}{2} \varepsilon X_{\sigma\tau} X^{\sigma\tau} = k \Theta_{\mu\nu}$$
  

$$T_{\mu\nu}^{\ \alpha} + \delta^{\alpha}_{\mu} T_{\nu\sigma}^{\ \sigma} - \delta^{\alpha}_{\nu} T_{\mu\sigma}^{\ \sigma} = -\frac{1}{2} k \Sigma^{\alpha}_{\mu\nu} \qquad (83)$$
  

$$X_{\mu\nu} = -\frac{1}{2} \varepsilon^{-1} k \Xi_{\mu\nu} ,$$

where I have denoted  $X^{\mu\nu} \equiv s^{\mu\nu}_{5} \cdot |h^{55}|^{1/2}$ ,  $\Xi_{\mu\nu} \equiv \mathcal{M}^{5}_{\mu\nu} \cdot |h^{55}|^{-1/2}$ , and  $\varepsilon \equiv \rho \operatorname{sign} h^{55}$ .

10. The notion of the bivector derivative can be extended to the fields whose values are nonspacetime vectors or tensors. By doing so one obtains a more particular generalization of the traditional gauge field theory framework where the five-vector gauge fields introduced above are viewed as composite quantities constructed from more elementary connection coefficients—from those associated with the bivector derivative. This latter generalization is obtained by postulating that for the fields of nonspacetime vectors and tensors there exists a derivative whose argument is a five-vector bivector and that for any such field this derivative is related to its five-vector covariant derivative according to equation (57), where  $\sigma(\mathbf{u})$  is the same as it is for four-vector fields.

As before, let us consider a set  $\mathcal{V}$  of all sufficiently smooth fields whose values are some n-dimensional nonspacetime vectors. Defining the bivector derivative for such fields is equivalent to specifying a map

$$\mathsf{D}:\ \mathcal{F}\wedge\mathcal{F}\times\mathcal{V}\rightarrow\mathcal{V},$$

The latter should satisfy the usual requirements: for any scalar functions f and g, any bivector fields  $\mathcal{A}$ and  $\mathcal{B}$ , and any fields  $\vec{v}$  and  $\vec{v}$  from  $\mathcal{V}$ ,

$$\begin{split} \mathsf{D}_{(f\mathcal{A}+g\mathcal{B})} \vec{\mathrm{u}} &= f \cdot \mathsf{D}_{\mathcal{A}} \vec{\mathrm{u}} + g \cdot \mathsf{D}_{\mathcal{B}} \vec{\mathrm{u}} \\ \mathsf{D}_{\mathcal{A}} (\vec{\mathrm{u}} + \vec{\mathrm{v}}) &= \mathsf{D}_{\mathcal{A}} \vec{\mathrm{u}} + \mathsf{D}_{\mathcal{A}} \vec{\mathrm{v}} \\ \mathsf{D}_{\mathcal{A}} (f \vec{\mathrm{u}}) &= \mathsf{D}_{\mathcal{A}} f \cdot \vec{\mathrm{u}} + f \cdot \mathsf{D}_{\mathcal{A}} \vec{\mathrm{u}}, \end{split}$$

where the bivector derivative of the scalar field f is defined by equation (51).

If  $\vec{\mathbf{E}}_i$  (i = 1, ..., n) is some set of basis fields from  $\mathcal{V}$ , one can define for it the connection coefficients associated with the derivative D according to the formula

$$\mathsf{D}_{AB}\vec{\mathsf{E}}_i = \vec{\mathsf{E}}_j C^j_{\ iAB},$$

where, as before,  $\mathsf{D}_{AB} \equiv \mathsf{D}_{\mathbf{e}_A \wedge \mathbf{e}_B}$  and  $\mathbf{e}_A$  is the selected five-vector basis. These connection coefficients will be called *bivector* gauge fields. Under the transformation  $\mathbf{e}'_A = \mathbf{e}_B L^B_A$  of the five-vector basis these fields transform simply as

$$C'^{i}_{jAB} = C^{i}_{jST}L^{S}_{A}L^{T}_{B}.$$

Under the transformation  $\vec{\mathbf{E}}_i' = \vec{\mathbf{E}}_j \Lambda_i^j$  of the basis in  $\mathcal{V}$  they transform as

$$C'_{jAB}^{i} = (\Lambda^{-1})^{i}_{k} C^{k}_{lAB} \Lambda^{l}_{j} + (\Lambda^{-1})^{i}_{k} \mathsf{D}_{AB} \Lambda^{k}_{j},$$

so in any active regular basis one has

$$C'_{j\alpha5}^{i} = (\Lambda^{-1})^{i}_{k} C^{k}_{l\alpha5} \Lambda^{l}_{j} + (\Lambda^{-1})^{i}_{k} \partial_{\alpha} \Lambda^{k}_{j}$$

and

$$C'^{i}_{j\alpha\beta} = (\Lambda^{-1})^{i}_{k} C^{k}_{l\alpha\beta} \Lambda^{l}_{j}.$$

Thus, in such a basis the quantities  $C^i_{j\alpha5}$  transform as ordinary gauge fields, while the quantities  $C^i_{j\alpha\beta}$ transform as components of a tensor and cannot be nullified at a given space-time point by an appropriate choice of the basis in  $\mathcal{V}$ .

Let us now write down explicitly the relation between the derivatives  $\square$  and D for the considered type of fields. As it has been said above, for any field  $\vec{u}$ from  $\mathcal{V}$  one should have

$$\Box_{\mathbf{v}}\vec{\mathrm{U}} = \mathsf{D}_{\sigma(\mathbf{v})}\vec{\mathrm{U}}$$

at any **v**. For  $\vec{u} = \vec{e}_i$  and  $\mathbf{v} = \mathbf{e}_A$  one has

$$\vec{\mathbf{E}}_{i}B^{i}_{jA} = \boldsymbol{\Box}_{A}\vec{\mathbf{E}}_{j} = \boldsymbol{\mathsf{D}}_{\sigma(\mathbf{e}_{A})}\vec{\mathbf{E}}_{j}$$
$$= s^{|KL|}_{A}\boldsymbol{\mathsf{D}}_{KL}\vec{\mathbf{E}}_{j} = \vec{\mathbf{E}}_{i}C^{i}_{jKL}s^{|KL|}_{A}.$$

Consequently,

$$B^i_{\ jA} = C^i_{\ jKL} s^{|KL|}_{\ A}$$

so in any active regular basis one has

$$B^{i}_{j\alpha} = C^{i}_{j\alpha5} + C^{i}_{j\mu\nu}s^{|\mu\nu|}{}_{\alpha} \text{ and } B^{i}_{j5} = C^{i}_{j\mu\nu}s^{|\mu\nu|}{}_{5}$$

The latter formulae elucidate the meaning of the bivector gauge fields. Within the traditional gauge field theory scheme, the parallel transport of nonspacetime vectors is independent of torsion in the sense that there is no direct relation between the latter and the corresponding gauge fields associated with the covariant derivative. According to the scheme we are now discussing, the parallel transport of nonspacetime vectors *is* torsion-depend, which manifests itself in an additional rotation of transported vectors compared to the case where torsion is zero. Let us also note that the scheme with ordinary (four-vector) gauge fields can be viewed as a particular case of the one we are now considering, which corresponds to the situation where the fields  $C^{i}_{j\mu\nu}$  in any regular five-vector basis are all identically zero.

As in the case of four-vector and five-vector fields, the bivector derivative operator for the fields of nonspacetime vectors can be split into two parts:

$$\mathsf{D}_{\mathcal{A}} = \mathsf{D}_{\mathcal{A}^{\mathcal{E}}} + \mathsf{D}_{\mathcal{A}^{\mathcal{Z}}}.$$
(84)

The first operator in the right-hand side can be regarded as a function of the four-vector **A** that corresponds to the  $\mathcal{E}$ -component of the bivector  $\mathcal{A}$  (or as a function of the corresponding five-vector from  $\mathcal{Z}$ ), and it is a simple matter to show that when regarded this way, it has all the properties of an ordinary covariant derivative, which permits one to denote this operator as  $\nabla_{\mathbf{A}}$ . It is easy to see that in any four-vector basis the connection coefficients associated with  $\nabla$  equal  $C^i_{j\mu 5}$  provided that the latter are evaluated for the corresponding active regular five-vector basis.

In a similar manner, the second operator in the right-hand side of formula (84) can be viewed as a function of the four-vector bivector **B** that corresponds to  $\boldsymbol{\mathcal{A}}^{\mathcal{Z}}$ , and by analogy with the case of fourand five-vectors, I will denote it as  $\widehat{\mathbf{M}}_{\mathbf{B}}$ . Naturally, in the case of nonspacetime vectors the components of  $\widehat{\mathbf{M}}$  will no longer equal  $(M_{\mu\nu})^{\alpha}_{\beta}$  or  $(M_{\mu\nu})^{A}_{B}$ , but instead, in any four-vector basis one will have

$$(\widehat{\mathbf{M}}_{\alpha\beta})^{i}_{\ j} = C^{i}_{\ j\alpha\beta},$$

where the bivector gauge fields in the right-hand side are to be evaluated in the corresponding regular fivevector basis. The latter fact reflects the fundamental

difference between the case of four- and five-vectors and the case of nonspacetime vectors in relation to the bivector derivative: whereas for the former the operator  $\widehat{\mathbf{M}}$  is fixed and its components are constructed from the Lorentz-invariant quantities  $g_{\alpha\beta}$  and  $\delta^{\alpha}_{\beta}$ , for the latter the operator  $\widehat{\mathbf{M}}$  can be as arbitrary as is allowed by the constraints imposed on D and its components represent an independent element of the geometry associated with the considered type of nonspacetime vectors, just as within the traditional scheme this is done by ordinary gauge fields. Such a state of affairs has a certain logic to it. Since the components of the operator  $\widehat{\mathbf{M}}$  for four-vector fields are fixed, the additional rotation of such vectors in the process of their parallel transport compared to the case where torsion is zero but the Riemannian geometry is the same, is determined only by the quantities  $s^{\mu\nu}_{\ A}$ , and having found the latter this way, one can then make a similar comparison for the transport of considered nonspacetime vectors and determine the combinations  $C^{i}_{j\mu\nu}s^{|\mu\nu|}_{A}$ , from which, knowing the torsion, one can find the quantities  $C^{i}_{j\mu\nu}$  themselves.

The mathematics of bivector gauge fields is discussed in more detail in part VI of the long version [8]. Their physics will be examined more closely in a separate paper.

11. In conclusion of this paper let me say a few words about the nonspacetime analogs of five-vectors. The nonspacetime vectors I have been talking about so far—such as those that are used in physics for describing the internal symmetries of elementary particles, resemble ordinary tangent vectors in the sense that at each space-time point their vector space is endowed only with a nondegenerate inner product and has no other additional structure similar to the  $\mathcal{Z}$ - $\mathcal{E}$ splitting in the space of five-vectors. In accordance with this, on the parallel transport of such vectors one imposes no other constraints except for the requirements that it be linear and conserve the mentioned inner product, so at an appropriate choice of the relevant gauge fields, any given vector at the initial point can be transported into any vector of the same length at the final point, if this does not contradict the condition of the transport continuity. One may now ask the following question: can there be defined such nonspacetime vectors that would resemble five-vectors?

Let us try to imagine what properties such vectors should have. It goes without saying that at each space-time point they should make up a certain finitedimensional vector space, W, the dimension of which in the general case it is convenient to denote as n+1. Accordingly, in the following such vectors themselves will be referred to as (n + 1)-vectors, and will be denoted with lower-case Roman-type letters with an arrow:  $\vec{u}$ ,  $\vec{v}$ ,  $\vec{w}$ , etc. It is also natural to assume that the space of (n + 1)-vectors is endowed with a nondegenerate inner product, which I will denote as  $\eta$ . All this, however, applies to ordinary nonspacetime vectors as well. It seems reasonable to suppose that (n+1)-vectors should differ from the latter in that their space is "split" into two invariant subspaces, which I will denote as  $W^{\mathcal{Z}}$  and  $W^{\mathcal{E}}$ , the first one of dimension n, the other one-dimensional. The space W itself will be the direct sum of these two subspaces, and the components of an arbitrary (n+1)-vector in them will be referred to as its  $\mathcal{Z}$ - and  $\mathcal{E}$ -component, respectively.

Since as in the case of ordinary nonspacetime vectors, it is not supposed that (n+1)-vectors are associated with any manifold, the mentioned splitting will have a real meaning only if it manifests itself in some specific properties, basing on which one would be able to say that one is dealing with (n+1)-vectors and not with some type of ordinary nonspacetime vectors of dimension n + 1. It is apparent that if the space of (n+1)-vectors is not endowed with any additional structure, then the above specific properties can only be related to parallel transport. Basing on the analogy with five-vectors, one may assume that (n+1)-vectors from  $W^{\mathcal{E}}$  are transported into (n+1)vectors from  $W^{\mathcal{E}}$  and that (n+1)-vectors from  $W^{\mathcal{Z}}$ may acquire in the process of transport a nonzero  $\mathcal{E}$ -component. The first of these properties tells one that one is not dealing with ordinary nonspacetime vectors. The second property tells one that neither one is dealing with elements of the direct sum of two spaces of ordinary nonspacetime vectors (of dimension n and one). In addition to this, it will be assumed that parallel transport conserves the inner product

$$\theta(\vec{\mathbf{u}}, \vec{\mathbf{v}}) \equiv \eta(\vec{\mathbf{u}}^{\mathcal{Z}}, \vec{\mathbf{v}}^{\mathcal{Z}}),$$

which is the analog of the scalar product g for five-vectors.

In order to write down the indicated properties of (n + 1)-vectors in the form of equations, let us introduce the following notations. The set of all sufficiently smooth fields whose values are (n + 1)-vectors of the considered type will be denoted as  $\mathcal{W}$ . An arbitrary set of basis fields from  $\mathcal{W}$  will be denoted as  $\vec{e}_1, \ldots, \vec{e}_{n+1}$ . It will be taken that lower-case latin indices run 1 through n and that capital Greek indices run 1 through n+1. Often, instead of the value n+1 I will use the symbol &.

The basis in  $\mathcal{W}$  can be chosen arbitrarily. However, for practical reasons it is more convenient to select it in such a way that at each space-time point the (n+1)st basis vector belong to  $W^{\mathcal{E}}$ . Similarly to the case of five-vectors, such bases will be called standard. It is also useful to introduce the notion of a *regular* basis, whose first n elements belong to  $W^{\mathcal{Z}}$  and the (n+1)st element is normalized in some particular way. Since (n + 1)-vectors are not associated with any manifold, and therefore cannot be represented with differential-algebraic operators, and since, as one will see below, from the rules of their parallel transport one also cannot obtain any special normalization for the vectors from  $W^{\mathcal{E}}$ , the only condition that one can use for normalizing  $\vec{e}_{\&}$  is the requirement  $|\eta(\vec{e}_{\&},\vec{e}_{\&})| = 1$ , which is similar to the normalization condition for the fifth basis vector in a normalized regular five-vector basis.

The connection coefficients for an arbitrary set of basis fields  $\vec{e}_{\Theta}$  in  $\mathcal{W}$  are defined in the usual way:

$$\Box_A \vec{\mathbf{e}}_\Theta = \vec{\mathbf{e}}_\Xi C^\Xi_{\ \Theta A}.$$

The quantities  $C^{\Xi}_{\Theta A}$  will still be called five-vector gauge fields. From the assumptions made above about the parallel transport of (n + 1)-vectors it follows that for any standard basis

$$C^i_{\&A} = 0,$$
 (85)

which is the analog of constraint (36) on the connection coefficients for five-vector fields. Furthermore, if, for example, the considered (n + 1)-vectors are complex and their inner product  $\theta$  is Hermitian, there should hold the equation

$$\partial_A \theta_{ij} - \theta_{kj} (C^k_{\ iA})^* - \theta_{ik} C^k_{\ jA} = 0, \qquad (86)$$

similar to the usual constraint on the gauge fields associated with ordinary nonspacetime vectors.

From the assumptions made above it follows that parallel transport of (n + 1)-vectors preserves the following equivalence relation on W:

$$\vec{\mathbf{u}} \equiv \vec{\mathbf{v}} \iff \vec{\mathbf{u}} - \vec{\mathbf{v}} \in W^{\mathcal{E}}.$$

It is a simple matter to check that with regard to their properties, the elements of the quotient space  $W/W^{\mathcal{E}}$ are ordinary nonspacetime vectors, and that at each space-time point this quotient space, endowed with the inner product induced by the product  $\theta$  on W, is isomorphic to the subspace  $W^{\mathcal{Z}}$ . One should therefore expect that with each type of (n+1)-vectors there is associated a certain type of ordinary nonspacetime vectors, whose relation to the considered (n + 1)vectors is similar to the relation of four-vectors to five-vectors. For these associated vectors one can use all the notations and definitions that have been introduced earlier for ordinary nonspacetime vectors. In particular, if the gauge fields corresponding to them are defined by equation (39) and if the corresponding basis fields  $\vec{\mathbf{e}}_i$  are such that at each point  $\vec{\mathbf{e}}_i$  is the equivalence class of the basis (n + 1)-vector  $\vec{\mathbf{e}}_i$ , then by virtue of what has been said above there should hold the equation

$$C^i_{\ jA} = B^i_{\ jA},\tag{87}$$

which is the analog of relation (12) between the connection coefficients for four-vector and five-vector fields.

The formula for transformation of the fields  $C^{\Xi}_{\Theta A}$  as one passes to another set of basis fields in  $\mathcal{W}$  is the following:

$$C'^{\Theta}_{\Xi A} = (L^{-1})^{\Theta}_{\Delta} C^{\Delta}_{\Sigma A} L^{\Sigma}_{\Xi} + (L^{-1})^{\Theta}_{\Delta} \partial_A L^{\Delta}_{\Xi}, \quad (88)$$

where  $L^{\Xi}_{\Theta}$  is the basis transformation matrix. If both bases are standard, one has  $L^{i}_{\&} = (L^{-1})^{i}_{\&} = 0$ , and at  $\Theta = i$  and  $\Xi = \&$  obtains

$$\begin{split} C'{}^{i}{}_{\&A} &= (L^{-1})^{i}{}_{k}{}^{k}{}_{lA}{}^{l}{}_{\&} \\ &+ (L^{-1})^{i}{}_{k}{}^{k}{}_{\&A}{}^{k}{}_{\&} \\ &+ (L^{-1})^{i}{}_{k}{}^{k}{}_{\&A}{}^{k}{}_{\&} \\ \end{split}$$

which is actually a demonstration of the fact that from the validity of equation (85) in one standard basis follows its validity in any other such basis. In a similar manner, at  $\Theta = i$  and  $\Xi = j$  one has

$$C'{}^{i}{}_{jA} = (L^{-1})^{i}{}_{k} C^{k}{}_{lA} L^{l}{}_{j} + (L^{-1})^{i}{}_{k} \partial_{A} L^{k}{}_{j}$$

so the connection coefficients  $C^{i}_{jA}$  transform as gauge fields corresponding to ordinary nonspacetime vectors, which agrees with equation (87).

Let us now turn to the gauge fields that determine the  $\mathcal{E}$ -components of the transported (n+1)-vectors. The first question one has to ask is whether parallel transport conserves the length of the vectors from  $W^{\mathcal{E}}$ . Since (n+1)-vectors are not associated with any manifold, the only measure available for the vectors from  $W^{\mathcal{E}}$  is the scalar square constructed with the inner product  $\eta$ . As in the case of five-vectors, one may suppose that this scalar square does *not* change. In the case of real vectors this means that for any field of regular bases one should have  $C^{\&}_{\&A} = 0$ . In the case of complex vectors and Hermitian  $\eta$ , the fields  $C^{\&}_{\ \&A}$  for a regular basis do not have to vanish, and it is only necessary that they be imaginary. With transition to another basis in  $\mathcal{W}$ , but also a regular one, in the latter case one has  $L^{\&}_{\&} = e^{i\alpha}$ , so

$$C'^{\&}_{\&A} = (L^{-1})^{\&}_{\&} C^{\&}_{\&A} L^{\&}_{\&} + (L^{-1})^{\&}_{\&} \partial_A L^{\&}_{\&} \\ = C^{\&}_{\&A} + i \partial_A \alpha.$$

There is one more constraint that can be imposed on the parallel transport of (n+1)-vectors, which implicitly is very often imposed on the parallel transport of ordinary nonspacetime vectors. Namely, one can require that this transport conserve the Levi-Civita type tensor  $\epsilon$  associated with the considered (n + 1)vectors. In the case of real W this condition is equivalent to the conservation of the length of the (n + 1)vectors from  $W^{\mathcal{E}}$ . In the case of complex W this requirement can be shown to imply that in any basis where the components of  $\eta$  and  $\epsilon$  are constant, one should have  $C_{\Theta A}^{\Theta} = 0$ .

The gauge fields  $C^{\&}_{jA}$  are evidently the analogs of the five-vector connection coefficients  $H^{5}_{\mu A}$ . From formula (88) it follows that with transition to another basis in  $\mathcal{W}$  they transform as

$$\begin{split} C'{}^{\&}_{\ jA} &= (L^{-1}){}^{\&}_{\Xi} \, C^{\Xi}{}_{lA} \, L^{l}{}_{j} + (L^{-1}){}^{\&}_{\&} \, C^{\&}_{\ \&A} \, L^{\&}_{j} \\ &+ (L^{-1}){}^{\&}_{k} \, \partial_{A} L^{k}{}_{j} + (L^{-1}){}^{\&}_{\&} \, \partial_{A} L^{\&}_{j}. \end{split}$$

If both bases are regular, then  $(L^{-1})_{j}^{\&} = L_{j}^{\&} = 0$ , and one has

$$C'^{\&}_{jA} = (L^{-1})^{\&}_{\&} C^{\&}_{\ lA} L^{l}_{\ j}.$$
(89)

If, in addition, one has  $\vec{e}'_{\&} = \vec{e}_{\&}$ , then simply

$$C'_{jA}^{\&} = C_{lA}^{\&} L_{j}^{l}.$$

An essential difference between the gauge fields  $C_{jA}^{\&}$  and their five-vector counterparts is that for the former there does not exist a nonzero value that would be invariant under the transformations from the symmetry group of W. On the other hand, the value  $C_{jA}^{\&} = 0$ , which does not break this symmetry, has the unpleasant property that at it one cannot distinguish the considered (n + 1)-vectors from pairs made of an ordinary *n*-dimensional nonspacetime vector and a scalar. It is evident that at any nonzero  $C_{jA}^{\&}$  the inner product  $\eta$  is not conserved by parallel transport, and since neither the requirement of the covariant constancy of  $\theta$  nor a similar requirement for the *n*-plus-one-vector  $\epsilon$  tensor impose any constraints on  $C_{jA}^{\&}$ , the latter can be absolutely arbitrary.

Let us now examine in more detail the case of complex vectors for which the inner product  $\eta$  is Hermitian and is positively definite. At each space-time point, let us select the basis in W orthonormal and such that one would have  $\epsilon_{1...n\&} = 1$ . Condition (86) will then acquire the form

$$\theta_{kj} (C_{iA}^k)^* + \theta_{ik} C_{jA}^k = 0,$$

whence it follows that the quantities  $C_{ijA} \equiv \theta_{ik} C^{k}_{\ jA}$ are anti-Hermitian matrices with respect to the indices *i* and *j*. Since in the selected basis  $C_{ijA} = C^{i}_{\ jA}$ , one can write that

$$C^{i}_{jA} = (i/2) g (t_a)^{i}_{j} C^{a}_{A} + ig \sqrt{2n(n+1)} \delta^{i}_{j} C^{0}_{A},$$
(90)

where the index *a* runs 1 through  $n^2 - 1$ ; the matrices  $(t_a)_j^i$  are the usual (Hermitian) generators for the fundamental representation of SU(*n*), normalized by the condition  $\text{Tr}(t_a t_b) = 2\delta_{ab}$ ; the fields  $C_A^a$  and  $C_A^o$  are real; and *g* is a dimensionless constant, which together with the factors 1/2 and  $[2n(n + 1)]^{-1/2}$  is introduced for convenience. From the condition  $C_{\Theta A}^{\Theta} = 0$  it follows that

$$C^{\&}_{\&A} = -ig \left[n/2(n+1)\right]^{1/2} C^{0}_{A}.$$
 (91)

By using (90) and (91) one can write down the expression for the components of the five-vector covariant derivative of an arbitrary (n + 1)-vector field in the selected basis in the following way:

$$(\Box_{A}\vec{u})^{i} = \partial_{A}u^{i} + (i/2) g (t_{a})^{i}_{j} C^{a}_{A} u^{j} + ig [2n(n+1)]^{-1/2} C^{o}_{A} u^{i}, (\Box_{A}\vec{u})^{\&} = \partial_{A}u^{\&} - ig [n/2(n+1)]^{1/2} C^{o}_{A} u^{\&} + g X_{jA} u^{j},$$

$$(92)$$

where I have introduced the notation  $X_{jA} \equiv g^{-1}C_{jA}^{\&}$ . Similarly, the expression for the components of the five-vector covariant derivative of a field  $\tilde{v}$  whose values are elements of the space  $\widetilde{W}$  of linear forms on W can be written down as follows:

$$(\Box_A \widetilde{\mathbf{v}})_i = \partial_A v_i - (i/2) g v_j (t_a)_i^j C^a_A - ig [2n(n+1)]^{-1/2} v_i C^0_A - g v_\& X_{iA}, \qquad (93)$$

$$(\square_A \widetilde{\mathbf{v}})_{\&} = \partial_A v_{\&} + ig \left[ n/2(n+1) \right]^{1/2} v_{\&} C^0_A.$$

If one disregards the terms involving the fields  $X_{iA}$ . the expressions obtained will have such a form as if one was dealing with the gauge fields corresponding to ordinary nonspacetime vectors and the gauge group was  $SU(n) \times U(1)$ . With respect to SU(n) the sets of fields  $(u^1, \ldots, u^n)$  and  $(v_1, \ldots, v_n)$  transform according to the fundamental and anti-fundamental representations, respectively, and the fields  $u^{\&}$  and  $v_{\&}$  are singlets. With respect to the group U(1) the fields  $u^1, \ldots, u^n$  all have the charge  $g[2n(n+1)]^{-1/2}$ , the field  $u^{\&}$  has the charge  $-g[n/2(n+1)]^{1/2}$ , and the charges of the fields  $v_1, \ldots, v_k$  are opposite to those of  $u^1, \ldots, u^{\&}$ , which is in agreement with the fact that the field  $\tilde{v}$  can be obtained from some (n+1)vector field by conjugation (by the latter I mean the antilinear map from W to W fixed by the inner product  $\eta$ , which is the analog of the map  $\vartheta_h$  introduced above and which in the selected basis coincides with ordinary Hermitian conjugation).

Besides  $C_A^a$  and  $C_A^o$ , the above expressions for the derivatives involve the gauge fields  $X_{jA}$ , due to which the covariant differentiation of the considered (n+1)-

vector fields in general does not commute with conjugation, as it can be clearly seen by comparing formulae (92) and (93). To gain a better understanding of what this noncommutativity implies, let us recall how one assigns a representation to matter fields in ordinary gauge theory when introducing new gauge fields. To be definite, I will consider the case where the gauge group in question is unitary. As always, the starting point is the existence of several matter fields in the theory, say,  $\varphi^1, \ldots, \varphi^n$ , that enter the Lagrangian density in such a way that the latter is invariant under the replacement

$$\varphi^i \to \varphi'^{\,i} = L^i_{\,\,j} \,\varphi^j, \tag{94}$$

where  $L_j^i$  is an arbitrary constant unitary  $n \times n$  matrix. One then gauges this symmetry by introducing the corresponding gauge fields, and as a result obtains the following expression for the derivative of the set  $\varphi = (\varphi^1, \ldots, \varphi^n)$ :

$$(\nabla_{\alpha}\varphi)^{i} = \partial_{\alpha}\varphi^{i} + (i/2) g (t_{a})^{i}_{j} B^{a}_{\ \alpha} \varphi^{j} + ig' B^{o}_{\ \alpha} \varphi^{i},$$

where  $(t_a)_j^i$  are the same as in formula (90), and for simplicity I omit the connection coefficients corresponding to other possible degrees of freedom of  $\varphi$ . By presenting the transformation formula for the considered matter fields in the form (94) one thereby states that this set of fields transforms according to the *fundamental* representation of the gauge group (= these fields are components of a corresponding nonspacetime vector). Equally well, one can lable the fields with a *lower* index and, accordingly, write the rule for their transformation as

$$\varphi_i \to \varphi_i' = \varphi_j \, L^j_{\ i}.$$

By doing so one would state that the fields  $\varphi$  transform according to the *anti-fundamental* representation (= are components of a *linear form* associated with the relevant nonspacetime vectors), and the expression for the derivative would then acquire the form

$$(\nabla_{\alpha}\varphi)_{i} = \partial_{\alpha}\varphi_{i} - (i/2) g \varphi_{j} (t_{a})^{j}_{i} \widetilde{B}^{a}_{\alpha} - ig'\varphi_{i} \widetilde{B}^{0}_{\alpha},$$

where  $\varphi_i = \varphi^i$ ,  $\tilde{B}^0_{\ \alpha} = -B^0_{\ \alpha}$ ,  $\tilde{B}^a_{\ \alpha} = -\varepsilon^a_b B^b_{\ \alpha}$ , and the coefficients  $\varepsilon^a_b$  are determined by the equation  $(t_a)^i_j = (t_b)^j_i \varepsilon^b_a$ . The transition from the fields  $\varphi^i, B^0_{\ \alpha}, B^a_{\ \alpha}$  to the fields  $\varphi_i, \tilde{B}^0_{\ \alpha}, \tilde{B}^a_{\ \alpha}$  and vice versa is a part of the charge conjugation operation.

By making similar transformations in formulae (92)

and (93) one obtains

$$(\Box_{A}\vec{u})_{i} = \partial_{A}u_{i} - (i/2) g u_{j} (t_{a})_{i}^{j} C_{A}^{a} - ig [2n(n+1)]^{-1/2} u_{i} \widetilde{C}_{A}^{0},$$

$$(\Box_{A}\vec{u})_{\&} = \partial_{A}u_{\&} + ig [n/2(n+1)]^{1/2} u_{\&} \widetilde{C}_{A}^{0} + g u_{j} X_{A}^{j}$$
(95)

and

$$(\Box_A \widetilde{\mathbf{v}})^i = \partial_A v^i + (i/2) g (t_a)^i_j C^a_A v^j + ig [2n(n+1)]^{-1/2} \widetilde{C}^0_A v^i - g X^i_A v^{\&}, \qquad (96)$$

$$(\square_A \widetilde{\mathbf{v}})^{\&} = \partial_A v^{\&} - ig \left[ n/2(n+1) \right]^{1/2} v^{\&} \widetilde{C}^o_A,$$

where  $u_{\Theta} = u^{\Theta}$ ,  $v_{\Theta} = v^{\Theta}$ ,  $X_A^i = X_{iA}$ ,  $\tilde{C}_A^0 = -C_A^0$ ,  $\tilde{C}_A^a = -\varepsilon_b^a C_A^b$ , and the coefficients  $\varepsilon_b^a$  are the same as above. Comparing expressions (95) and (96) with expressions (93) and (92) respectively, one can see that at  $X_{iA} \neq 0$  they do not coincide. Consequently, the interaction with the fields  $X_{iA}$  is not C-invariant, and one should observe that in this case the charge asymmetry is implemented directly in the nonspacetime degrees of freedom of the fields.

## Acknowledgements

I would like to thank V. D. Laptev for supporting this work. I am grateful to V. A. Kuzmin for his interest and to V. A. Rubakov for a very helpful discussion and advice. I am indebted to A. M. Semikhatov of the Lebedev Physical Institute for a very stimulating and pleasant discussion and to S. F. Prokushkin of the same institute for consulting me on the Yang-Mills theories of the de Sitter group. I would also like to thank L. A. Alania, S. V. Aleshin, and A. A. Irmatov of the Mechanics and Mathematics Department of the Moscow State University for their help and advice.

## References

- Th. Kaluza, Sitzungsber. Preuss. Akad. Wiss. Berlin, Math.–Phys. K1. (1921) 966; O. Klein, Z. Phys. 46 (1927) 188.
- See e.g. K.S. Stelle and P.C. West, Phys. Rev. D 21 (1980) 1466.
- A.B. Krasulin, Five-Dimensional Tangent Vectors in Space-Time I: Introduction and Formal Theory, to be published.
- 4. A.B. Krasulin, Five-Dimensional Tangent Vectors in Space-Time II: Differential-Geometric Approach, to be published.

- 5. A.B. Krasulin, Five-Dimensional Tangent Vectors in Space-Time III: Some Applications, to be published.
- 6. A.B. Krasulin, Five-Dimensional Tangent Vectors in Space-Time IV: Generalization of Exterior Calculus, to be published.
- 7. A.B. Krasulin, Five-Dimensional Tangent Vectors in Space-Time V: Generalization of Covariant Derivative, to be published.
- 8. A.B. Krasulin, Five-Dimensional Tangent Vectors in Space-Time VI: Bivector Derivative and its Applications, to be published.
- W. Hehl et al., Reviews of Modern Physics 48 (1976) 393.