FROBENIUS-SCHUR FUNCTIONS: SUMMARY OF RESULTS

GRIGORI OLSHANSKI, AMITAI REGEV, AND ANATOLY VERSHIK

We introduce and study a family $\{FS_{\mu}\}$ of symmetric functions which we call the *Frobenius–Schur functions*. These are inhomogeneous functions indexed by partitions and such that FS_{μ} differs from the conventional Schur function s_{μ} in lower terms only.

Our interest in these new functions comes from the fact that they provide an explicit expression for $\dim \nu/\mu$ (dimension of a skew Young diagram ν/μ) in terms of the Frobenius coordinates of ν , see Theorem 2 below. The problem of studying $\dim \nu/\mu$ as a function of the Frobenius coordinates of ν originated in the asymptotic theory of the characters of the symmetric groups [VK]. We discuss this point below.

Our main result is a surprisingly simple determinantal expression of the Frobenius– Schur functions in terms of the conventional Schur functions, see Theorem 9.

Other results concerning the functions FS_{μ} are: certain generating series, the Giambelli formula, vanishing properties and interpolation, a combinatorial formula (representation in terms of tableaux), and a Sergeev–Pragacz–type formula.

We actually deal with a wider class of symmetric functions $s_{\mu;a}$ of the form s_{μ} + lower terms, which we call *multiparameter Schur functions*. Here $a = (a_i)_{i \in \mathbb{Z}}$ is an arbitrary doubly infinite sequence of parameters. The multiparameter Schur functions turn into the Frobenius–Schur functions when $(\ldots, a_0, a_1, a_2, a_3, \ldots) = (\ldots, -\frac{1}{2}, \frac{1}{2}, \frac{3}{2}, \frac{5}{2}, \ldots)$ and into the conventional Schur functions when $a_i \equiv 0$. All our results about the functions FS_{μ} can be stated for the functions $s_{\mu;a}$ without extra efforts. For this reason we decided to include this generalization (although at present we do not dispose of any applications of the functions $s_{\mu;a}$ with general parameters).

For proofs we refer to the detailed version of the paper, [ORV].

We are grateful to Alain Lascoux for fruitful criticism.

The authors were supported by Russian Foundation for Basic Research under grants 98-01-00303 and 99-01-00098 (G. O. and A. V., respectively) and by ISF grant 6629/1 (A. R.).

Preliminaries. We follow the notation of Macdonald [Ma1] in what concerns symmetric functions and Schur functions. Our base field is \mathbb{C} . Let Λ denote the algebra of symmetric functions. Recall that Λ is a graded algebra, isomorphic to the algebra of polynomials in the power sums p_1, p_2, \ldots A natural homogeneous basis in Λ is formed by the *Schur functions* s_{μ} , where the subscript μ is a partition (or, what is the same, a Young diagram).

The representation theory of the infinite symmetric group naturally leads to linear superalgebra and supersymmetric functions, see [VK], [W], [KV1], [KV2], [OI], [KOO]. A polynomial in two finite collections of indeterminates, x_1, \ldots, x_m and y_1, \ldots, y_n , is called *supersymmetric* if it is separately symmetric in the x's and in the y's, and, moreover, the following *cancellation property* holds: for any fixed i and j, specializing $x_i = -y_j = t$ we get a polynomial in m + n - 2 variables not depending on t.

Let \mathbb{C}_0^{∞} be the space of the infinite complex vectors with finitely many nonzero coordinates. A function f(x; y) on $\mathbb{C}_0^{\infty} \times \mathbb{C}_0^{\infty}$ is called a *supersymmetric function* if for any m and n, the specialization $x_{m+1} = x_{m+2} = \cdots = y_{n+1} = y_{n+2} = \cdots = 0$ turns f into a supersymmetric polynomial whose degree remains bounded as m and n go to infinity.

Specializing $\mathbf{p}_k \mapsto \sum_i x_i^k + (-1)^{k-1} \sum_j y_j^k$ we get an *isomorphism* between the algebra Λ and the algebra of supersymmetric functions, see [S], [P]. We agree to identify both algebras by means of this isomorphism. Given $f \in \Lambda$, we denote by f(x; y) the corresponding supersymmetric function on $\mathbb{C}_0^\infty \times \mathbb{C}_0^\infty$. In particular, we write $s_{\mu}(x; y)$ for the "supersymmetric Schur function".

If $x \in \mathbb{C}^m$, $y \in \mathbb{C}^n$, and f is a supersymmetric function, then we abbreviate $f(x;y) = f(x,0,0,\ldots;y,0,0,\ldots)$.

We will often use the *Frobenius notation* [Ma1, §I.1] for Young diagrams:

$$\nu = (p_1, \ldots, p_d \mid q_1, \ldots, q_d),$$

where $d = d(\nu)$ (the *depth* of ν) is the number of diagonal squares in ν , $p_i = \nu_i - i$ and $q_i = \nu'_i - i$ are the *Frobenius coordinates* of ν , and ν' stands for the transposed diagram. As usual, $|\nu|$ denotes the number of squares in ν .

The Frobenius–Schur functions: a motivation. Let S(n) denote the symmetric group of degree n. For a Young diagram ν let χ^{ν} denote the irreducible character of $S(|\nu|)$ indexed by ν and let dim $\nu = \chi^{\nu}(e)$ be the degree of χ^{ν} (i.e., the dimension of the corresponding representation), which coincides with the number of standard tableaux of shape ν . If σ is a partition of $|\nu|$ then χ^{ν}_{σ} denotes the value of χ^{ν} on the conjugacy class indexed by σ . Our starting point is the following result (see [VK], [W, III.6], [KO]):

Theorem 1. Fix a partition $\rho = (\rho_1, \ldots, \rho_l)$ of m. For any $n \ge m$ we agree to complete ρ to a partition of n by adding 1's. Let $\nu = (p_1, \ldots, p_d \mid q_1, \ldots, q_d)$ denote an arbitrary Young diagram with n boxes.

There exists an inhomogeneous symmetric function

$$\boldsymbol{p}_{\rho}^{\#} = \boldsymbol{p}_{\rho_1} \dots \boldsymbol{p}_{\rho_l} + \text{lower degree terms}$$

such that

$$\frac{\chi_{\rho\cup 1^{n-m}}^{\nu}}{\dim\nu} = \frac{\boldsymbol{p}_{\rho}^{\#}(p_1+\frac{1}{2},\ldots,p_d+\frac{1}{2};q_1+\frac{1}{2},\ldots,q_d+\frac{1}{2})}{n(n-1)\ldots(n-m+1)}$$

To avoid misunderstanding, let us explain once more the meaning of an expression like $p_{\rho}^{\#}(p_1 + \frac{1}{2}, \ldots, p_d + \frac{1}{2}; q_1 + \frac{1}{2}, \ldots, q_d + \frac{1}{2})$: here a symmetric function (specifically, $p_{\rho}^{\#}$) is first viewed as a supersymmetric function on $\mathbb{C}_0^{\infty} \times \mathbb{C}_0^{\infty}$ and then it is specialized at a finite vector from the space $\mathbb{C}^d \times \mathbb{C}^d$ canonically embedded into the larger space $\mathbb{C}_0^{\infty} \times \mathbb{C}_0^{\infty}$. It is worth noting that a symmetric function is uniquely determined by its superspecializations at the points of the form $(p_1 + \frac{1}{2}, \ldots, p_d + \frac{1}{2}; q_1 + \frac{1}{2}, \ldots, q_d + \frac{1}{2})$, so that $p_a^{\#}$ is defined uniquely.

Note also a discrepancy with the classical point of view; usually a character χ^{ν} is thought of as a function on the set of conjugacy classes whereas here the picture is reversed: we fix a conjugacy class and view χ^{ν} as a function of the index ν .

This theorem plays an important role in the following asymptotic problem considered in [VK]: describe all sequences $\{\nu\}$ of Young diagrams (where $|\nu| \to \infty$) such that the corresponding normalized irreducible characters $\chi^{\nu}/\dim\nu$ converge on the infinite symmetric group $S(\infty)$ (this group is defined as the union of the groups S(n)). It turns out that the limit functions $\lim_{\{\nu\}} \frac{\chi^{\nu}}{\dim\nu}$ are exactly the *characters* of

the group $S(\infty)$, discovered in [T].

One of the main conclusions of [VK] was that a solution to the asymptotic problem mentioned above can be described in terms of the limit behavior of the Frobenius coordinates of the diagrams ν . This explains why we find it so important to express χ^{ν}_{σ} through the Frobenius coordinates of ν and not the conventional row coordinates ν_1, ν_2, \ldots

Theorem 1 is equivalent to the following claim. Let $\dim(\mu, \nu)$ be the number of standard tableaux of skew shape ν/μ provided that $\mu \subseteq \nu$, and 0 otherwise.

Theorem 2. For any Young diagram μ there exists an inhomogeneous symmetric function

 $FS_{\mu} = s_{\mu} + \text{lower degree terms}$

such that for any diagram $\nu = (p_1, \ldots, p_d \mid q_1, \ldots, q_d)$ with $|\nu| = n \ge m = |\mu|$,

$$\frac{\dim(\mu,\nu)}{\dim\nu} = \frac{FS_{\mu}(p_1 + \frac{1}{2},\dots,p_d + \frac{1}{2};q_1 + \frac{1}{2},\dots,q_d + \frac{1}{2})}{n(n-1)\dots(n-m+1)}.$$

Again, the function in question is defined uniquely. We call the functions FS_{μ} the Frobenius-Schur functions, because their definition exploits Frobenius coordinates.

The expression $\frac{\dim(\mu,\nu)}{\dim\nu}$ has an important meaning: it is a conditional probability in a probabilistic interpretation of the characters of the group $S(\infty)$, which was suggested in [VK]. We will briefly explain this point, because it may help in understanding the role of the Frobenius–Schur functions.

Let Tab be the set of all infinite Young tableaux [VK], [KV2]. An element $\tau \in$ Tab is an infinite chain ($\tau_0 = \emptyset, \tau_1, \tau_2, \ldots$) of Young diagrams such that $|\tau_n| = n$ and τ_n is obtained from τ_{n-1} by adding a box. In other terms, τ is an infinite path in the Young graph. Given a probability measure P on Tab, we can speak about random paths. We say that P is a *central measure* if the probability that the random path has a fixed beginning (τ_0, \ldots, τ_n) depends only on the diagram τ_n (and not on the whole finite chain (τ_0, \ldots, τ_n)). It turns out that there is a natural one–to–one correspondence between the characters of $S(\infty)$ and the indecomposable central measures on Tab. Now we are in a position to explain what $\frac{\dim(\mu,\nu)}{\dim\nu}$ is: this is the probability that the random path at it passes through ν at "time" n. Note that this conditional probability does not depend on a central measure; it is rather a characteristic of the Young graph. For more detail, see [VK] and [KOr]

From quite a different point of view the Frobenius–Schur functions are considered in $[BO, \S 6]$.

The multiparameter Schur functions. Fix an arbitrary doubly infinite sequence $a = (a_i)_{i \in \mathbb{Z}}$ of complex parameters. We define elements $h_{1;a}, h_{2;a}, \ldots$ of Λ (which depend on the sequence a) by a generating series:

$$1 + \sum_{k=1}^{\infty} \frac{h_{k;a}}{(u-a_1)\dots(u-a_k)} = 1 + \sum_{k=1}^{\infty} \frac{h_k}{u^k},$$

where $h_1, h_2, \ldots \in \Lambda$ are the complete homogeneous symmetric functions, u is a formal variable, and the series above are understood as elements of the algebra $\Lambda[[\frac{1}{u}]]$. Equivalently,

$$h_{k;a} = \sum_{i=1}^{k} (-1)^{k-i} e_{k-i}(a_1, \dots, a_{k-1}) h_i,$$

where e_1, e_2, \ldots are the elementary symmetric functions. This implies, in particular, that $\{h_{k;a}\}_{k=1,2,\ldots}$ is a system of algebraically independent generators of Λ .

We agree that $h_{0;a} = 1$ and $h_{-1;a} = h_{-2;a} = \cdots = 0$. We also need the following notation: for $r \in \mathbb{Z}$, let $\tau^r a$ be the result of shifting a by r digits to the left: $(\tau^r a)_i = a_{i+r}$.

The multiparameter Schur function indexed by an arbitrary Young diagram μ is defined as follows:

$$s_{\mu;a} = \det[h_{\mu_i - i + j;\tau^{1-j}a}],$$

where the order of the determinant is any number greater or equal to $\ell(\mu)$, the number of rows in μ . Clearly, $h_{k;a} = s_{(k);a}$.

We have $s_{\mu;a} = s_{\mu} + \text{lower degree terms}$, which implies that the functions $s_{\mu;a}$ form a basis in Λ . When $a \equiv 0$, they reduce to s_{μ} .

Theorem 3. The Frobenius–Schur functions FS_{μ} coincide with the functions $s_{\mu;a}$ for the following special choice of the parameters: $a_i = i - \frac{1}{2}$.

Let us call $a_i = i - \frac{1}{2}$ the special parameter sequence.

Although we are mainly interested in Frobenius–Schur functions, all the results below are stated in a more general context of multiparameter Schur functions. To pass to the Frobenius–Schur case it suffices to substitute $a_i = i - \frac{1}{2}$.

Connections with earlier works. Some topics in the representation theory of infinite–dimensional classical groups and in the theory of Yangians naturally lead to the algebra of shifted symmetric functions. This algebra, denoted by Λ^* , is studied in detail in [OO]. This is a filtered algebra with associated graded algebra canonically isomorphic to Λ . In Λ^* , there is a natural basis formed by the shifted Schur functions. These functions are closely related to the factorial Schur polynomials discovered by Biedenharn and Louck [BL1], [BL2] (for more references, see [OO]). According to [KO], both Λ^* and Λ can be realized as certain algebras of functions on the set of partitions, which leads to an algebra isomorphism $\Lambda^* \to \Lambda$. It turns out that under this isomorphism, the shifted Schur functions turn into the Frobenius–Schur functions: this claim is equivalent to Theorem 8.1 in [OO] which expresses $\frac{\dim(\mu,\nu)}{\dim\nu}$ in terms of the shifted Schur functions evaluated at the row coordinates of ν (not in terms of the Frobenius coordinates!). A connection between shifted Schur functions and Schubert polynomials is explained in [L2], see also Lascoux's remark

in [OO, §15.7]. It is possible that the Schubert polynomial techniques [Ma3] can provide an alternative approach to the results of the present paper but we did not verify this.

Next, the functions $s_{\mu;a}$ can be obtained by an appropriate specialization of Lascoux's *multi–Schur functions* introduced in [L1] (see also [Ma3] and [L2]). We are grateful to Alain Lascoux for pointing out this fact. For more detail see [ORV].

Finally, the functions $s_{\mu;a}$ are closely connected with Molev's factorial supersymmetric Schur polynomials [Mo]. For more detail see [ORV]. At the early stage of the present work this connection helped us very much.

Duality. Consider the involutive automorphism $\omega : \Lambda \to \Lambda$, see [Ma1]. Recall that

$$\omega(\boldsymbol{p}_k) = (-1)^{k-1} \boldsymbol{p}_k, \quad \omega(h_k) = e_k, \quad \omega(e_k) = h_k, \quad \omega(s_\mu) = s_{\mu'}$$

In terms of supersymmetric functions, ω interchanges the arguments $x, y \in \mathbb{C}_0^{\infty}$.

Define the dual parameter sequence \hat{a} by $\hat{a}_i = -a_{-i+1}$. Thus, a can be written as $(\ldots, -\hat{a}_2, -\hat{a}_1, a_1, a_2, \ldots)$.

Theorem 4. We have $\omega(s_{\mu;a}) = s_{\mu';\hat{a}}$.

Set
$$e_{k;a} = s_{(1^k);a}$$
. By Theorem 4, $e_{k;a} = \omega(h_{k;\hat{a}})$ and we have a generating series

$$1 + \sum_{k=1}^{\infty} \frac{e_{k;a}}{(u - \hat{a}_1) \dots (u - \hat{a}_k)} = 1 + \sum_{k=1}^{\infty} \frac{e_k}{u^k}.$$

Or, equivalenty,

$$e_{k;a} = \sum_{i=1}^{k} (-1)^{k-i} e_{k-i}(\widehat{a}_1, \dots, \widehat{a}_{k-1}) e_i,$$

Further, $s_{\mu;a} = \det[e_{\mu'_i - i + j;\tau^{j-1}a}].$

These formulas can be immediately specialized for the case of the Frobenius– Schur functions. Here $(a_1, a_2, ...) = (\hat{a}_1, \hat{a}_2, ...) = (\frac{1}{2}, \frac{3}{2}, ...)$. Note that the special parameter sequence is selfdual, so that $\omega(FS_{\mu}) = FS_{\mu'}$, which is also evident from Theorem 2.

Hook functions. The definition of the multiparameter Schur functions has the form $s_{\mu;a} = \det[h_{\mu_i - i + j, j - 1}]$, where $\{h_{k,l}\}$ is a certain family of elements of Λ (depending of *a*). By a general result (see [Ma2], [Ma1, Ex. I.3.21]), it follows

Theorem 5 (Giambelli formula). Write μ in the Frobenius notation, $\mu = (p_1, \ldots, p_d \mid q_1, \ldots, q_d)$. Then $s_{\mu;a} = \det[s_{(p_i|q_j);a}]$.

The hook functions $s_{(p|q);a}$ can be determined from the following generating series. Set

$$H(u) = 1 + \sum_{k=1}^{\infty} \frac{h_k}{u^k}, \qquad E(v) = 1 + \sum_{k=1}^{\infty} \frac{e_k}{u^k}.$$

Theorem 6. We have

$$1 + (u+v)\sum_{p,q=0}^{\infty} \frac{s_{(p|q);a}}{(u-a_1)\dots(u-a_{p+1})(v-\hat{a}_1)\dots(v-\hat{a}_{q+1})} = H(u)E(v).$$

When $a \equiv 0$, this reduces to a known identity, see [Ma1, Ex. I.3.9]. For the Frobenius–Schur functions we get

$$1 + (u+v)\sum_{p,q=0}^{\infty} \frac{FS_{(p|q)}}{(u-\frac{1}{2})\dots(u-\frac{2p+1}{2})(v-\frac{1}{2})\dots(v-\frac{2q+1}{2})} = H(u)E(v).$$

Transition coefficients. Let $b = (b_i)_{i \in \mathbb{Z}}$ be another parameter sequence. By Theorem 6,

$$1 + (u+v) \sum_{p,q=0}^{\infty} \frac{s_{(p|q);a}}{(u-a_1)\dots(u-a_{p+1})(v-\hat{a}_1)\dots(v-\hat{a}_{q+1})}$$

= 1 + (u+v) $\sum_{p,q=0}^{\infty} \frac{s_{(p|q);b}}{(u-b_1)\dots(u-b_{p+1})(v-\hat{b}_1)\dots(v-\hat{b}_{q+1})}.$

It follows that

$$s_{(p \mid q);a} = \sum_{p'=0}^{p} \sum_{q'=0}^{q} c_{pp'}(a,b) c_{qq'}(\widehat{a},\widehat{b}) s_{(p' \mid q');b},$$

where

$$c_{pp'}(a,b) = h_{p-p'}(b_1,\ldots,b_{p'+1};-a_1,\ldots,-a_p), \qquad p \ge p' \ge 0.$$

Note that $c_{pp}(a, b) = 1$.

An important particular case is $b \equiv 0$. Then

$$c_{pp'}(a,0) = h_{p-p'}(0,\ldots,0;-a_1,\ldots,-a_p) = (-1)^{p-p'}e_{p-p'}(a_1,\ldots,a_p).$$

Together with the Giambelli formula (Theorem 5) these formulas provide a simple determinant expression for the transition coefficients between the bases $\{s_{\mu;a}\}$ and $\{s_{\mu;b}\}$.

Theorem 7. Let $a = (a_i)_{i \in \mathbb{Z}}$ and $b = (b_i)_{i \in \mathbb{Z}}$ be two sequences of parameters. In the expansion

$$s_{\mu;a} = \sum_{\nu} c_{\mu\nu}(a,b) \, s_{\nu;b}$$

the coefficients $c_{\mu\nu}(a,b)$ vanish unless $\nu \subseteq \mu$ and both diagrams have the same number of diagonal squares: $d(\nu) = d(\mu)$.

Assume $d(\nu) = d(\mu) = d$ and write both diagrams in the Frobenius notation,

$$\mu = (p_1, \dots, p_d | q_1, \dots, q_d), \quad \nu = (p'_1, \dots, p'_d | q'_1, \dots, q'_d).$$

Then

$$c_{\mu\nu}(a,b) = \det[c_{p_i,p_j'}(a,b)] \det[c_{q_i,q_j'}(\widehat{a},\widehat{b})],$$

where the determinants are of order d and the coefficients of the form $c_{pp'}(a, b)$ are defined above.

In particular, taking b = 0 we get a nice expression of the multiparameter Schur functions $s_{\mu;a}$ through the conventional Schur functions. From this expression we get

Corollary 8. Let $p = \mu_1 - 1$ and $q = \mu'_1 - 1$. The function $s_{\mu;a}$ depends only of p + q parameters (a_1, \ldots, a_p) and $(\widehat{a}_1, \ldots, \widehat{a}_q) = (-a_0, \ldots, -a_{q-1})$.

Let us specially state the particular case of Theorem 7 corresponding to $a_i = i - \frac{1}{2}$ and $b \equiv 0$: **Theorem 9.** We have

$$FS_{(p_1,\dots,p_d|q_1,\dots,q_d)} = \sum_{\substack{p_1' \le p_1,\dots,p_d' \le p_d \\ q_1' \le q_1,\dots,q_d' \le q_d}} \det[c_{p_i,p_j'}] \det[c_{q_i,q_j'}] s_{(p_1',\dots,p_d'|q_1',\dots,q_d')},$$

where

$$c_{pp'} = (-1)^{p-p'} e_{p-p'}(\underbrace{\frac{1}{2}, \dots, \frac{2p-1}{2}}_{p}).$$

Vanishing, interpolation, characterization. For a nonempty Young diagram written in the Frobenius notation

$$\lambda = (p_1(\lambda), \dots, p_d(\lambda) \mid q_1(\lambda), \dots, q_d(\lambda)), \qquad d = d(\lambda),$$

we set

$$(x(\lambda); y(\lambda)) = (a_{p_1(\lambda)+1}, \dots, a_{p_d(\lambda)+1}; \widehat{a}_{q_1(\lambda)+1}, \dots, \widehat{a}_{q_d(\lambda)+1}) \in \mathbb{C}^d \times \mathbb{C}^d \subset \mathbb{C}_0^\infty \times \mathbb{C}_0^\infty.$$

Note that for the special parameter sequence $a_i = i - \frac{1}{2}$, we have

$$(x(\lambda); y(\lambda)) = (p_1(\lambda) + \frac{1}{2}, \dots, p_d(\lambda) + \frac{1}{2}; q_1(\lambda) + \frac{1}{2}, \dots, q_d(\lambda) + \frac{1}{2}).$$

We also agree that $(x(\emptyset); y(\emptyset)) = (0; 0)$.

Theorem 10. If $\mu \not\subseteq \lambda$ then $s_{\mu;a}(x(\lambda); y(\lambda)) = 0$.

Theorem 11. We have

$$s_{\mu;a}(x(\mu);y(\mu)) = \prod_{(i,j)\in\mu} (a_{\mu_i-i+1} - a_{j-\mu'_j}) = \prod_{(i,j)\in\mu} (a_{\mu_i-i+1} + \widehat{a}_{\mu'_j-j+1}).$$

If the numbers a_i are pairwise distinct then $s_{\mu;a}(x(\mu); y(\mu)) \neq 0$ for all μ .

For the Frobenius–Schur functions this reduces to

$$FS_{\mu}(p_1(\mu) + \frac{1}{2}, \dots, p_d(\mu) + \frac{1}{2}; q_1(\mu) + \frac{1}{2}, \dots, q_d(\mu) + \frac{1}{2}) = \prod_{(i,j)\in\mu} (\mu_i - j + \mu'_j - i + 1).$$

By the hook formula, the last expression equals $\dim \mu/|\mu|!$, which agrees with Theorem 2.

As a corollary of Theorem 10 and Theorem 11 we get

Theorem 12. Assume that the parameters a_i are pairwise distinct. Given an arbitrary element $f \in \Lambda$ of degree $\leq n$, consider its expansion in the multiparameter Schur functions of degree $\leq n$: $f = \sum_{\lambda: |\lambda| \leq n} c(\lambda) s_{\lambda;a}$. Then the coefficients $c(\lambda)$ can be found by recurrence on $|\lambda|$, as follows:

 $c(\varnothing) = f(0;0), \qquad c(\lambda) = \frac{f(x(\lambda); y(\lambda)) - \sum_{\mu: \mu \subset \lambda, \mu \neq \lambda} c(\mu) s_{\mu;a}(x(\lambda); y(\lambda))}{s_{\lambda;a}(x(\lambda); y(\lambda))}.$

In other words, the multiparameter Schur functions naturally arise in the following Newton-type *interpolation process*: we consider the space of supersymmetric functions on $\mathbb{C}_0^{\infty} \times \mathbb{C}_0^{\infty}$ with its natural filtration by degree and take the points of interpolation of the form (0;0) and $(a_{i_1},\ldots,a_{i_d};\hat{a}_{j_1},\ldots,\hat{a}_{j_d})$. Here $d = 1, 2, \ldots$; $i_1 > \cdots > i_d \ge 1$ and $j_1 > \cdots > j_d \ge 1$ are d-tuples of natural numbers; (a_1, a_2, \ldots) and $(\hat{a}_1, \hat{a}_2, \ldots)$ are two fixed sequences of complex parameters such that $a_i \ne a_j$ and $\hat{a}_i \ne \hat{a}_j$ for $i \ne j$, and $a_i \ne -\hat{a}_j$ for all i, j (the condition $a_i \ne -\hat{a}_j$ is natural because of the cancellation property).

Note that the Frobenius–Schur functions correspond to the interpolation process with the points of interpolation of the form (0;0) and $(x_1 > \cdots > x_d; y_1 > \cdots > y_d)$, where x_i, y_j are taken from the set $\{\frac{1}{2}, \frac{3}{2}, \frac{5}{2}, \ldots\}$.

Theorem 13. Assume that the parameters a_i are pairwise distinct. Then the function $F = s_{\mu;a} \in \Lambda$ can be characterized by the following properties (i) and (ii) as well as by the properties (i') and (ii'):

(i) F vanishes at $(x(\lambda); y(\lambda))$ for all diagrams $\lambda \neq \mu$ such that $|\lambda| \leq |\mu|$;

(ii) F is of degree $\leq |\mu|$ and its value at $(x(\mu); y(\mu))$ is the same as in Theorem 11.

(i') F vanishes at $(x(\lambda); y(\lambda))$ for all diagrams λ with $|\lambda| < |\mu|$;

(ii') the top degree homogeneous component of F coincides with the Schur function s_{μ} .

In the particular case of the Frobenius–Schur functions, the results of this section are equivalent to similar claims for the shifted Schur functions contained in [Ok1], [OO, §3]. Interpolation of arbitrary polynomials in terms of Schubert polynomials is discussed in [L2]. A very general scheme of Newton interpolation for symmetric polynomials is developed in [Ok2]. In the latter paper one can also find the references to earlier works by Knop, Sahi, and Okounkov.

The combinatorial formula. Let $\mathbb{Z}' = \{\ldots, -\frac{3}{2}, -\frac{1}{2}, \frac{1}{2}, \frac{3}{2}, \ldots\}$ stand for the set of proper half-integers. For any $\varepsilon \in \mathbb{Z}'$ set $a'_{\varepsilon} = a_{\varepsilon+1/2}$. Note that $(\widehat{a})'_{\varepsilon} = a'_{-\varepsilon}$. For the special parameter sequence, $a'_{\varepsilon} = \varepsilon$.

Let ν be a skew hook (in another terminology, a ribbon). That is, ν is a connected skew Young diagram containing no 2×2 block of squares. We attach to ν a polynomial $f_{\nu;a}(u, v)$ in two variables u, v, of degree $|\nu|$, as follows.

Consider the interior sides s of ν : each interior side is adjacent to two squares of ν ; the total number of the interior sides is equal to $|\nu| - 1$. To each interior side s we attach the coordinates (ε, δ) of its midpoint,¹ and we write $s = (\varepsilon, \delta)$. Note that one of the coordinates is always in \mathbb{Z}' while another coordinate is integral. Specifically, if s is a vertical side then $\varepsilon \in \mathbb{Z}'$, $\delta \in \mathbb{Z}$, and the ends of s are the points $(\varepsilon - 1/2, \delta)$ and $(\varepsilon + 1/2, \delta)$; if s is a horizontal side then $\varepsilon \in \mathbb{Z}$, $\delta \in \mathbb{Z}'$, and the ends of s are the points $(\varepsilon, \delta - 1/2)$, $(\varepsilon, \delta + 1/2)$.

For both vertical and horizontal sides, $\delta - \varepsilon \in \mathbb{Z}'$, so that $a'_{\delta - \varepsilon} = a_{\delta - \varepsilon + 1/2}$ is well defined. We set

$$f_{\nu;a}(u,v) = (u+v) \prod_{\substack{\text{vertical interior}\\\text{sides } s = (\varepsilon,\delta) \text{ of } \nu}} (u-a'_{\delta-\varepsilon}) \prod_{\substack{\text{horizontal interior}\\\text{sides } s = (\varepsilon,\delta) \text{ of } \nu}} (v+a'_{\delta-\varepsilon}) dv_{\delta-\varepsilon}$$

¹As in [Ma1], we assume that the first coordinate axis is directed downwards and the second coordinate axis is directed to the right.

For instance, if $\nu = (4, 2, 2)/(1, 1)$ then there are 6 squares and 5 interior sides with midpoints

$$(\frac{5}{2},1), (2,\frac{3}{2}), (1,\frac{3}{2}), (\frac{1}{2},2), (\frac{1}{2},3),$$

and we have

$$f_{\nu;a}(u,v) = (u+v)(u-a'_{-3/2})(u-a'_{3/2})(u-a'_{5/2})(v+a'_{-1/2})(v+a'_{1/2})$$

= $(u+v)(u-a_{-1})(u-a_{2})(u-a_{3})(v+a_{0})(v+a_{1})$

Note that $f_{\nu;a}(v,u) = f_{\nu';\hat{a}}(u,v)$.

More generally, if ν is an arbitrary skew diagram with no 2×2 block of squares, then ν is a disjoint union of skew hooks and we define $f_{\nu;a}(u, v)$ as the product of the polynomials attached to the connected components of ν .

We also agree that $f_{\emptyset}(u, v) \equiv 1$. Note that if $\nu \neq \emptyset$ then $f_{\nu;a}(0; 0) = 0$, because of the factor u + v.

Given a Young diagram μ , a diagonal-strict tableau of shape μ is a function $\mathcal{T}(i,j)$ from the squares (i,j) of μ to $\{1,2,\ldots\}$ such that the numbers $\mathcal{T}(i,j)$ weakly increase both along the rows i = const (from left to right) and down the columns j = const, and strictly increase along the diagonals j - i = const. Each subset of form $\mathcal{T}^{-1}(k)$ is a skew diagram with no 2×2 block of squares, so that the polynomial $f_{\mathcal{T}^{-1}(k)}(u, v)$ makes sense.

Theorem 14. The following combinatorial formula holds

$$s_{\mu;a}(x;y) = \sum_{\mathcal{T}} \prod_{k \ge 1} f_{\mathcal{T}^{-1}(k);a}(x_k, y_k),$$

summed over all diagonal-strict tableaux of shape μ .

If n is so large that $x_{n+1} = x_{n+2} = \cdots = y_{n+1} = y_{n+2} = \cdots = 0$ then only tableaux \mathcal{T} with values in $\{1, \ldots, n\}$ really contribute. Further, each product in the above formula is actually finite, because $\mathcal{T}^{-1}(k) = \emptyset$ for sufficiently large k (and then $f_{\mathcal{T}^{-1}(k)} \equiv 1$). Thus, the right-hand side of the combinatorial formula makes sense.

The formula is compatible with the involution ω , because the definition of a diagonal-strict tableau is symmetric with respect to transposition and $f_{\nu;a}(v,u) = f_{\nu';\hat{a}}(u,v)$.

The formula can be restated in terms of tableaux of a different kind, see [ORV]. The proof of Theorem 14 is due to Vladimir Ivanov.

Note that for the first time, an "inhomogeneous" combinatorial formula probably appeared in [BL1], [BL2], see also [CL]. Other examples can be found in [GG], [Ma2], [Mo], [Ok1], [OO], [Ok2]. See also further references in [Ok2] to works by Knop, Okounkov, and Sahi about combinatorial formulas for interpolation Jack and Macdonald polynomials.

The Sergeev–Pragacz formula. In this section we assume that (x; y) ranges over $\mathbb{C}^n \times \mathbb{C}^n \subset \mathbb{C}_0^\infty \times \mathbb{C}_0^\infty$. Thus, we deal with supersymmetric polynomials in n + n variables. Then $s_{\mu;a}(x; y)$ vanishes unless $d(\mu) \leq n$.

In the space of all polynomials in n + n variables there is a natural action of $\mathfrak{S}_n \times \mathfrak{S}_n$, the product of two copies of the symmetric group \mathfrak{S}_n . For $w = (w_1, w_2) \in$

 $\mathfrak{S}_n \times \mathfrak{S}_n$ we write $\varepsilon(w) = \operatorname{sgn}(w_1) \operatorname{sgn}(w_2)$. Set

$$V(x_1, \dots, x_m) = \prod_{1 \le i < k \le m} (x_i - x_k), \qquad V(y_1, \dots, y_n) = \prod_{1 \le j < l \le n} (y_j - y_l),$$
$$(x \mid a)^m = \begin{cases} (x - a_1) \dots (x - a_m), & m \ge 1, \\ 1, & m = 0, \end{cases}$$

and $(k)_{+} = \max(k, 0)$.

Theorem 15. Fix n = 1, 2, ... and let μ be an arbitrary Young diagram such that $d = d(\mu) \leq n$. We have

$$s_{\mu;a}(x_1,\ldots,x_n;y_1,\ldots,y_n) = \frac{\sum\limits_{w\in\mathfrak{S}_n\times\mathfrak{S}_n}\varepsilon(w)\,w[g_{\mu;a}(x_1,\ldots,x_n;y_1,\ldots,y_n)]}{V(x_1,\ldots,x_n)V(y_1,\ldots,y_n)}\,,$$

where

$$g_{\mu;a}(x_1, \dots, x_n; y_1, \dots, y_n) = \prod_{i=1}^d (x_i \mid a)^{\mu_i - i} x_i^{(n-\mu_i)_+} (y_i \mid \widehat{a})^{\mu'_i - i} y_i^{(n-\mu'_i)_+}$$
$$\times \prod_{i=d+1}^n x_i^{n-i} y_i^{n-i} \cdot \prod_{\substack{i,j \le n \\ (i,j) \in \mu}} (x_i + y_j).$$

When $a \equiv 0$, this turns into the conventional Sergeev–Pragacz formula, see [PT], [Ma3], [Ma1, Ex. I.3.23]. Our proof is an adaptation of the argument of [PT].

Taking n = d we get an analog of the factorization property [BR]:

Corollary 16. Let $\mu = (p_1, \ldots, p_d | q_1, \ldots, q_d)$ be a Young diagram of depth d. Then

$$s_{\mu;a}(x_1,\ldots,x_d;y_1,\ldots,y_d) = \frac{\det[(x_i \mid a)^{p_j}]_{i,j=1}^d}{V(x_1,\ldots,x_d)} \cdot \frac{\det[(y_i \mid \widehat{a})^{q_j}]_{i,j=1}^d}{V(y_1,\ldots,y_d)} \cdot \prod_{i,j=1}^d (x_i + y_j) \,.$$

Schur's Q-functions. As is well known, there exists a deep analogy between the conventional Schur functions (also called Schur's S-functions) and Schur's Qfunctions. The latter are related to projective representations of symmetric groups just in the same way as the former are related to ordinary representations. Schur's Q-functions span a subalgebra in Λ , whose elements can be characterized by a supersymmetry property of another kind, see, e.g., [Ma1, §III.8], [P]. The results of the present paper have counteparts for Schur's Q-functions: these are due to Vladimir Ivanov ([I1], [I2], and a paper in preparation).

References

- [BR] A. Berele and A. Regev, Hook Young diagrams with applications to combinatorics and to representations of Lie superalgebras, Adv. Math. **64** (1987), 118–175.
- [BL1] L. C. Biedenharn and J. D. Louck, A new class of symmetric polynomials defined in terms of tableaux, Advances in Appl. Math. 10 (1989), 396–438.

- [BL2] _____, Inhomogeneous basis set of symmetric polynomials defined by tableaux, Proc. Nat. Acad. Sci. U.S.A. 87 (1990), 1441–1445.
- [BO] A. Borodin and G. Olshanski, Harmonic functions on multiplicative graphs and interpolation polynomials, Preprint, 1999, available via http://arXiv.org/abs/math/9912124.
- [CL] W. Y. C. Chen and J. D. Louck, The factorial Schur function, J. Math. Phys. 34 (1993), 4144–4160.
- [GG] I. Goulden and C. Greene, A new tableau representation for supersymmetric Schur functions, J. Algebra 170 (1994), 687–703.
- [I1] V. N. Ivanov, Dimension of skew shifted Young diagrams and projective representations of the infinite symmetric group, Representation theory, dynamical systems, combinatorial and algorithmical methods II (A. M. Vershik, ed.), Zapiski Nauchnykh Seminarov POMI 240, Nauka, St. Petersburg, 1997, pp. 115–135 (Russian); English transl. to appear in J. Math. Sci.
- [I2] V. N. Ivanov, Combinatorial formula for factorial Schur Q-functions, in: Representation theory, dynamical systems, combinatorial and algorithmical methods III (A. M. Vershik, ed.), Zapiski Nauchnykh Seminarov POMI 256, Nauka, St. Petersburg, 1999, pp. 73–94 (Russian); English transl. to appear.
- [K] S. Kerov, Gaussian limit for the Plancherel measure of the symmetric group, Comtes Rendus Acad. Sci. Paris Sér. I 316 (1993), 303–308.
- [KO] S. Kerov and G. Olshanski, Polynomial functions on the set of Young diagrams, Comptes Rendus Acad. Sci. Paris Sér. I 319 (1994), 121–126.
- [KOO] S. Kerov, A. Okounkov, G. Olshanski, The boundary of Young graph with Jack edge multiplicities, Intern. Math. Res. Notices (1998), no. 4, 173–199.
- [KOr] S. V. Kerov and O. A. Orevkova, Random processes with common cotransition probabilities, in: Studies in Mathematical Statistics IX, Zapiski Nauchnyh Seminarov LOMI 184 (1990), pp. 169–181 (Russian).
- [KV1] S. Kerov, A. Vershik, The characters of the infinite symmetric group and probability properties of the Robinson-Schensted-Knuth algorithm, SIAM J. Alg. Discr. Meth. 7 (1986), 116-124.
- [KV2] _____, The Grothendieck group of the infinite symmetric group and symmetric functions with the elements of the K₀-functor theory of AF-algebras, Representation of Lie groups and related topics (A. M. Vershik and D. P. Zhelobenko, eds.), Adv. Stud. Contemp. Math. 7, Gordon and Breach, 1990, pp. 36–114.
- [L1] A. Lascoux, Puissances extérieurs, déterminants et cycles de Schubert, Bull. Soc. Math. France 102 (1974), 161–179.
- [L2] _____, Notes on interpolation in one and several variables, Preprint, available via http://schubert.univ-mlv.fr/~al/MAIN/publications.html.
- [Ma1] I. G. Macdonald, Symmetric functions and Hall polynomials, 2nd edition, Oxford University Press, 1995.
- [Ma2] _____, Schur functions: theme and variations, Publ. I.R.M.A. Strasbourg, 498/S–27, Actes 28-e Séminaire Lotharingien (1992), 5–39.
- [Ma3] _____, Notes on Schubert polynomials, Publ. LACIM, Université du Quebec, Montréal, (1991).
- [Mo] A. Molev, Factorial supersymmetric Schur functions and super Capelli identities, Kirillov's Seminar on Representation Theory (G. Olshanski, ed.), American Mathematical Society Translations (2), Vol. 181, Amer. Math. Soc., Providence, R.I., 1997, pp. 109–137.
- [Ok1] A. Okounkov, Quantum immanants and higher Capelli identities, Transformation Groups 1 (1996), 99–126.
- [Ok2] _____, On Newton interpolation of symmetric functions: a characterization of interpolation Macdonald polynomials, Adv. Appl. Math. **20** (1998), 395–428.
- [OO] A. Okounkov and G. Olshanski, *Shifted Schur functions*, Algebra i Analiz 9 (1997), no. 2, 73–146 (Russian); English translation: St. Petersburg Math. J. 9 (1998), 239–300.
- [Ol] G. Olshanskii, Unitary representations of (G, K)-pairs connected with the infinite symmetric group $S(\infty)$, Algebra i Analiz **1** (1989), no. 4, 178–209 (Russian); English translation in Leningrad [St. Petersburg] Math. J. **1** (1990), no. 4, 983-1014.
- [ORV] G. Olshanski, A. Regev, and A. Vershik (with an appendix by V. Ivanov), *Frobenius–Schur* functions, to appear in the arXiv.

- [P] P. Pragacz, Algebro-geometric applications of Schur S- and Q-polynomials, Topics in Invariant Theory, Seminaire d'Algèbre Paul Dubriel et Marie-Paule Malliavin, Lecture Notes in Math., vol. 1478, Springer-Verlag, New-York/Berlin, 1991, pp. 130–191.
- [PT] P. Pragacz and A. Thorup, On a Jacobi-Trudi identity for supersymmetric polynomials, Adv. Math. (1992), 8–17.
- [S] J. R. Stembridge, A characterization of supersymmetric polynomials, J. Algebra 95 (1985), 439–444.
- [T] E. Thoma, Die unzerlegbaren, positive-definiten Klassenfunktionen der abzählbar unendlichen, symmetrischen Gruppe, Math. Zeitschr. 85 (1964), 40-61.
- [VK] A. M. Vershik and S. V. Kerov, Asymptotic theory of characters of the symmetric group, Funct. Anal. Appl. 15 (1981), no. 4, 246–255.
- [W] A. J. Wassermann, Automorphic actions of compact groups on operator algebras, Thesis, University of Pennsylvania (1981).

G. OLSHANSKI: Dobrushin Mathematics Laboratory, Institute for Information Transmission Problems, Bolshoy Karetny 19, Moscow 101447, GSP-4, Russia.

E-mail address: olsh@iitp.ru, olsh@glasnet.ru

A. REGEV: Department of Theoretical Mathematics, Weizmann Institute of Science, Rehovot 76100, Israel.

E-mail address: regev@wisdom.weizmann.ac.il

A. VERSHIK: Steklov Mathematical Institute (POMI), Fontanka 27, St. Petersburg 191011, Russia.

E-mail address: vershik@pdmi.ras.ru