

AN ARAKELOV THEORETIC PROOF OF THE EQUALITY OF CONDUCTOR AND DISCRIMINANT

SINAN ÜNVER

1. INTRODUCTION

Let K be a number field, \mathcal{O}_K be the ring of integers of K , and S be $\text{Spec}(\mathcal{O}_K)$. Let $f : X \rightarrow S$ be an arithmetic surface. By this we mean a regular scheme, proper and flat over S , of relative dimension one. We also assume that the generic fiber of X has genus ≥ 1 , and that X/S has geometrically connected fibers.

Let ω_X be the dualizing sheaf of X/S . The Mumford isomorphism ([Mumf], Theorem 5.10)

$$\det Rf_*(\omega_X^{\otimes 2}) \otimes K \rightarrow (\det Rf_*\omega_X)^{\otimes 13} \otimes K,$$

which is unique up to sign, gives a rational section Δ of

$$(\det Rf_*\omega_X)^{\otimes 13} \otimes (\det Rf_*(\omega_X^{\otimes 2}))^{\otimes -1}.$$

The discriminant $\Delta(X)$ of X/S is defined as the divisor of this rational section ([Saito]). If \mathfrak{p} is a closed point of S , we denote the coefficient of \mathfrak{p} in $\Delta(X)$ by $\delta_{\mathfrak{p}}$.

On the other hand X/S has an Artin conductor $\text{Art}(X)$ (cf. [Bloch]), which is similarly a divisor on S . We denote the coefficient of \mathfrak{p} in $\text{Art}(X)$ by $\text{Art}_{\mathfrak{p}}$. Let S' be the strict henselization of complete local ring at \mathfrak{p} , with field of fractions K' . Let s be its special point, η be its generic point, and $\bar{\eta}$ be a geometric generic point corresponding to an algebraic closure \bar{K}' of K' . Let ℓ be a prime different from the residue characteristic at \mathfrak{p} . Then

$$\begin{aligned} \text{Art}_{\mathfrak{p}}(X) &= \sum_{i \geq 0} (-1)^i \dim_{\mathbb{Q}_\ell} H_{\acute{e}t}^i(X_{\bar{\eta}}, \mathbb{Q}_\ell) - \sum_{i \geq 0} (-1)^i \dim_{\mathbb{Q}_\ell} H_{\acute{e}t}^i(X_s, \mathbb{Q}_\ell) \\ &\quad + \sum_{i \geq 0} (-1)^i \text{Sw}_{\bar{K}'/K'}(H_{\acute{e}t}^i(X_{\bar{\eta}}, \mathbb{Q}_\ell)), \end{aligned}$$

where $\text{Sw}_{\bar{K}'/K'}$ denotes the Swan conductor of the Galois representation of \bar{K}'/K' . Both of these divisors are supported on the primes of bad reduction of X . We give another proof of Saito's theorem ([Saito], Theorem 1) in the number field case.

Theorem 1. *For any closed point $\mathfrak{p} \in S$, we have $\delta_{\mathfrak{p}} = -\text{Art}_{\mathfrak{p}}$.*

Fix a Kähler metric on X , this gives metrics on $\Omega_{X_\nu}^1$'s, for each $\nu \in S(\mathbb{C})$. For a hermitian coherent sheaf \mathcal{E} , we endow $\det Rf_*(\mathcal{E})$ with its Quillen metric. The proof of the theorem has the following corollaries.

Date: May 30, 2000.

Proposition 1. *We have*

$$\deg \det Rf_*\omega_X = \frac{1}{12} [\deg f_*(\widehat{c}_1(\omega_X)^2) + \log \text{Norm}(-\text{Art}(X))] \\ [K : \mathbb{Q}](g-1)[2\zeta'(-1) + \zeta(-1)],$$

with ζ the Riemann zeta function.

Proposition 1 is an arithmetic analogue of Noether's formula in which $\det Rf_*\omega_X$ is endowed with the Quillen metric. Faltings [Falt] and Moret-Bailly [M-B] proved a similar formula for the Faltings metrics.

Proposition 2. *We have*

$$\frac{1}{[K : \mathbb{Q}]} \sum_{\nu \in S(\mathbb{C})} \log \|\Delta_\nu\| = 12(1-g)[2\zeta'(-1) + \zeta(-1)].$$

In particular, the norm of the Mumford isomorphism does not depend on the metric.

2. PROOF

First we prove Proposition 1. By duality ([Deligne], Lemme 1.3), $\deg \det Rf_*\omega_X = \deg \det Rf_*\mathcal{O}_X$. By the arithmetic Riemann-Roch theorem of Gillet and Soulé ([G-S], Theorem 7), we get

$$\deg \det Rf_*\mathcal{O}_X = \deg f_*(\widehat{Td}(\Omega_X^1)^{(2)}) - \frac{1}{2} \sum_{\nu \in S(\mathbb{C})} \int_{X_\nu} Td(T_{X_\nu})R(T_{X_\nu})$$

Here Td and R are the Todd and Gillet-Soulé genera respectively, and the superscript (2) denotes the degree 2 component. Applying the definitions of these characteristic classes we obtain

$$\deg \det Rf_*\mathcal{O}_X = \frac{1}{12} \deg f_*(\widehat{c}_1(\Omega_X^1)^2 + \widehat{c}_2(\Omega_X^1)) + [K : \mathbb{Q}](g-1)[2\zeta'(-1) + \zeta(-1)]$$

Let Z denote the union of singular fibers of f , and let $c_{2,X}^Z(\Omega_X^1)$ be the localized Chern class of Ω_X^1 with support in Z (cf. [Bloch], [Fulton]). Chinburg, Pappas, and Taylor ([CPT], Proposition 3.1) prove the formula

$$\deg f_*(\widehat{c}_2(\Omega_X^1)) = \log \text{Norm}(c_{2,X}^Z(\Omega_X^1)).$$

Combining this with the fundamental formula of Bloch ([Bloch], Theorem 1)

$$-\text{Art}_p(X) = \deg_p c_{2,X}^Z(\Omega_X^1),$$

we obtain the desired formula. Note that, since $\det \Omega_X^1 = \omega_X$, $\widehat{c}_1(\Omega_X^1) = \widehat{c}_1(\omega_X)$. \square
Taking degrees in the Mumford isomorphism gives

$$13 \deg \det Rf_*\omega_X = \deg \det Rf_*(\omega_X^{\otimes 2}) + \log \text{Norm}(\Delta(X)) - \sum_{\nu \in S(\mathbb{C})} \log \|\Delta_\nu\|.$$

The arithmetic Riemann-Roch theorem gives

$$\deg \det Rf_*(\omega_X^{\otimes 2}) = \deg \det Rf_*\omega_X + \deg f_*(\widehat{c}_1(\omega_X)^2).$$

Therefore we get

$$(1) \quad \deg \det Rf_* \omega_X = \frac{1}{12} [\deg f_* (\widehat{c}_1(\omega_X)^2) + \log \text{Norm}(\Delta(X)) - \sum_{\nu \in S(\mathbb{C})} \log \|\Delta_\nu\|].$$

Subtracting (1) from the expression in the statement of Proposition 1, we obtain

$$(2) \quad \log\left(\frac{\text{Norm}(\Delta(X/S))}{\text{Norm}(-\text{Art}(X/S))}\right) = \sum_{\nu \in S(\mathbb{C})} \log \|\Delta_\nu\| + 12[K : \mathbb{Q}](g-1)[2\zeta'(-1) + \zeta(-1)].$$

Now X_K has semistable reduction after a finite base change K'/K . For semistable X'/S' , both $-\text{Art}_{\mathfrak{p}'}(X')$ ([Bloch]), and $\delta_{\mathfrak{p}'}(X')$ ([Falt], Theorem 6) are equal to the number of singular points in the geometric fiber over \mathfrak{p}' . Therefore $-\text{Art}_{\mathfrak{p}'} = \delta_{\mathfrak{p}'}$, and hence

$$(3) \quad \text{Norm}(\Delta(X'/S')) = \text{Norm}(-\text{Art}(X'/S')).$$

Applying this to a semistable model X' of $X \otimes_K K'$, and noting that the base change multiplies the right hand side of (2) by $[K' : K]$, we see that the right hand side of (2) is equal to zero, and hence that the equality

$$(4) \quad \text{Norm}(\Delta(X/S)) = \text{Norm}(-\text{Art}(X/S))$$

holds for X .

To prove the equality $\delta_{\mathfrak{p}} = -\text{Art}_{\mathfrak{p}}$ for an arbitrary closed point $\mathfrak{p} \in S$, we will use the following lemma.

Lemma 1. *Fix distinct closed points $\beta_1, \dots, \beta_s \in S$. And fix finite extensions L_i of the completions K_i of K at β_i 's, for each $1 \leq i \leq s$, such that $[L_i : K_i] = n$ for some n . Then there exists an extension L/K such that, for each $1 \leq i \leq s$, there is only one prime γ_i of L lying over β_i , and the completion of L at γ_i is isomorphic (over K_i) to L_i .*

Proof. The proof is an easy application of Krasner's lemma, and the approximation lemma. Details are omitted. \square

Take $\mathfrak{p} = \beta_1$, a prime of bad reduction. Denote the remaining primes of bad reduction by β_i , $2 \leq i \leq s$. Choose extensions L_i of the local fields K_i , for all $1 \leq i \leq s$, such that L_1 is unramified over K_1 , X has semistable reduction over L_i , for $2 \leq i \leq s$, and $[L_i : K_i] = n$, for some n . Applying the lemma to this data we obtain an extension L of K . Let $T = \text{Spec}(\mathcal{O}_L)$. The curve $X \otimes_K L$ has a proper, regular model Y over T such that

- (i) $Y \otimes_T T_{\gamma_1} \simeq X \otimes_S T_{\gamma_1}$, and
- (ii) Y is semistable at γ_i , for $2 \leq i \leq s$.

Applying (4) to Y gives the equality

$$\sum_{1 \leq i \leq s} \delta_{\gamma_i} \log \text{Norm}(\gamma_i) = \sum_{1 \leq i \leq s} -\text{Art}_{\gamma_i} \log \text{Norm}(\gamma_i).$$

On the other hand because of semistability, we have $\delta_{\gamma_i} = -\text{Art}_{\gamma_i}$, for $2 \leq i \leq s$. Hence we get $\delta_{\gamma_1} = -\text{Art}_{\gamma_1}$. Since T/S is étale at γ_1 , (i) implies

$$\delta_{\mathfrak{p}} = \delta_{\gamma_1} = -\text{Art}_{\gamma_1} = -\text{Art}_{\mathfrak{p}}.$$

Acknowledgements. I would like to thank A.Abbes for his many mathematical suggestions, and for his constant encouragement. I would like to thank my adviser P.Vojta for supporting me this academic year.

REFERENCES

- [Bloch] Bloch, S.: Cycles on arithmetic schemes and Euler characteristics of curves. Proc. of Sympos. Pure Math. 46 (1987) AMS, 421-450.
- [CPT] Chinburg, T., Pappas, G., Taylor, M.J.: ϵ -constants and Arakelov Euler characteristics. Preprint, (1999).
- [Deligne] Deligne, P.: Le déterminant de la cohomologie. Contemp. Math. 67 (1987), 93-177.
- [Falt] Faltings, G.: Calculus on arithmetic surfaces. Ann. Math. 119 (1984), 387-424.
- [Fulton] Fulton, W.: Intersection theory. Springer-Verlag, Berlin, 1984.
- [G-S] Gillet, H., Soulé, C.: An arithmetic Riemann-Roch theorem. Invent. Math. 110 (1992), 473-543.
- [M-B] Moret-Bailly, L.: La formule de Noether pour les surfaces arithmétiques. Invent. Math. 98 (1989), 499-509.
- [Mumf] Mumford, D.: Stability of projective varieties. Enseign. Math. 23 (1977), 39-100.
- [Saito] Saito, T.: Conductor, discriminant, and the Noether formula of arithmetic surfaces. Duke Math. J. 57 (1988), 151-173.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA-BERKELEY, BERKELEY, CA 94720

E-mail address: `sinan@math.berkeley.edu`