

# A Free Probability Analogue of the Wasserstein Metric on the Trace-State Space

PHILIPPE BIANE AND DAN VOICULESCU

*Incomplete preliminary version, June 1, 2000*

**Abstract.** We define a free probability analogue of the Wasserstein metric, which extends the classical one. In dimension one, we prove that the square of the Wasserstein distance to the semi-circle distribution is majorized by a modified free entropy quantity.

## 0 Introduction

The Wasserstein distance between two probability distributions  $\mu, \nu$  on  $\mathbb{R}^n$  is given by

$$W(\mu, \nu) = \inf_{\pi \in \Pi(\mu, \nu)} \left( \int |x - y|^2 d\pi(x, y) \right)^{\frac{1}{2}}$$

where  $\Pi(\mu, \nu)$  denotes the probability measures on  $\mathbb{R}^n \times \mathbb{R}^n$  with marginals  $\mu$  and  $\nu$ .

Following the usual free probability recipe we shall replace the set of probability measures by the trace-state space of a  $C^*$ -algebra and take marginals with respect to a free product. In this note we begin the study of the ensuing free Wasserstein metric.

An inequality of M. Talagrand ([7],[10]) relates the Wasserstein distance from a Gaussian distribution and relative entropy. In the one-variable case we prove a related free probability result to this inequality, where the semicircle law replaces the Gauss law and the logarithmic energy plays the role of entropy. Note that in the case of  $n$ -tuples of commuting selfadjoint variables the classical and the free Wasserstein distances are equal.

In the context of non-commutative geometry, there is a different noncommutative extension, due to A. Connes [5], of the related Monge-Kantorowitz metric. The Monge-Kantorowitz metric is a  $p = 1$ ,  $p$ -Wasserstein metric, but the definition which is extended is the dual definition based on Lipschitz functions, and the extension involves Fredholm-modules or derivations (recent work is surveyed in [9]).

# 1 The free Wasserstein metric

## 1.1 The distance on $n$ -tuples of variables

We will work in the framework of tracial  $C^*$ -probability spaces  $(M, \tau)$ , where  $M$  is a unital  $C^*$ -algebra and  $\tau$  is a trace state. The simplest is to define the metric at the level of noncommutative random variables. If  $(X_1, \dots, X_n)$  and  $(Y_1, \dots, Y_n)$  are two  $n$ -tuples of noncommutative random variables in tracial  $C^*$ -probability spaces  $(M_1, \tau_1)$  and  $(M_2, \tau_2)$ , we define

$$W_p((X_1, \dots, X_n), (Y_1, \dots, Y_n))$$

as the infimum of

$$\|(|X'_j - Y'_j|_p)_{1 \leq j \leq n}\|_p$$

over  $2n$ -tuples  $(X'_1, \dots, X'_n, Y'_1, \dots, Y'_n)$  of noncommutative random variables in some tracial  $C^*$ -probability space  $(M_3, \tau_3)$  such that the  $n$ -tuples  $(X'_1, \dots, X'_n)$ ,  $(X_1, \dots, X_n)$  and respectively  $(Y'_1, \dots, Y'_n)$ ,  $(Y_1, \dots, Y_n)$  have the same  $*$ -distributions. Here  $|\cdot|_p$  is the  $p$ -norm in a tracial  $C^*$ -probability space, while  $\|\cdot\|_p$  is the  $p$ -norm on  $\mathbb{R}^n$ . Like in the classical case, if  $p = 2$  we call  $W_p$  the free Wasserstein metric and we will also use the notation  $W$  for  $W_2$ . We shall refer to  $W_p$  as the free  $p$ -Wasserstein metric. Note also that if

$$X_j = D_j + iE_j, \quad Y_j = F_j + iG_j$$

where  $D_j, E_j, F_j, G_j$  are self-adjoint, then

$$W((X_1, \dots, X_n), (Y_1, \dots, Y_n)) = W((D_1, \dots, D_n, E_1, \dots, E_n), (F_1, \dots, F_n, G_1, \dots, G_n))$$

Note also that  $W_p((X_1, \dots, X_n), (Y_1, \dots, Y_n))$  depends only on the  $*$ -distributions of  $(X_1, \dots, X_n)$  and  $(Y_1, \dots, Y_n)$ . If we consider  $n$ -tuples with the same  $*$ -distribution as equivalent; then  $W_p$  will be a distance between equivalence classes of  $n$ -tuples.

## 1.2 The distance on trace states

We pass now to trace-state spaces  $TS(A)$ , where  $A$  is a unital  $C^*$ -algebra. We will assume  $A$  is finitely generated and we will assume such a generator  $(a_1, \dots, a_n)$  has been specified.

The  $p$ -Wasserstein metric on  $TS(A)$  is given by

$$W_p(\tau', \tau'') = W_p((a'_1, \dots, a'_n), (a''_1, \dots, a''_n))$$

where  $\tau', \tau'' \in TS(A)$  and  $(a'_1, \dots, a'_n)$  and  $(a''_1, \dots, a''_n)$  denote the variables defined by  $(a_1, \dots, a_n)$  in  $(A, \tau')$  and respectively  $(A, \tau'')$ .

This definition can be rephrased using free products. If  $A_1, A_2$  are unital  $C^*$ -algebras, we denote by  $\sigma_j : A_j \rightarrow A_1 * A_2$  the canonical injection of  $A_j$  into the full free product  $C^*$ -algebra (this presumes amalgamation over  $\mathbb{C}1$ ). If  $\tau_j \in TS(A_j)$ , ( $1 \leq j \leq 2$ ) we define

$$TS(A_1 * A_2; \tau_1, \tau_2) = \{\tau \in TS(A_1 * A_2) \mid \tau \circ \sigma_j = \tau_j, j=1, 2\} .$$

Remark that  $\tau_1 * \tau_2 \in TS(A_1 * A_2; \tau_1, \tau_2)$ .

It is easy to see that

$$W_p(\tau', \tau'') = \inf\{\|(|\sigma_1(a_j) - \sigma_2(a_j)|_{p,\tau})_{1 \leq j \leq n}\|_p \mid \tau \in TS(A * A; \tau', \tau'')\}$$

where  $|\cdot|_{p,\tau}$  denotes the  $p$ -norm in  $L^p(A; \tau)$ .

Remark also that the distance on  $n$ -tuples of variables can be obtained from the definition for trace-states. Assume for simplicity  $X_j = X_j^*$ ,  $Y_j = Y_j^*$  and  $R \geq \|X_j\|$ ,  $R \geq \|Y_j\|$ ,  $1 \leq j \leq n$ . Let then  $A = (C[-R, R])^{*n}$  (the free product of  $n$  copies) and  $\sigma_k(a) = a_k$ , where  $a$  is the identical function in  $C[-R, R]$ . Let  $\rho_j : A \rightarrow M_j$ ,  $j=1, 2$  be the  $*$ -homomorphisms such that  $\rho_1(a_k) = X_k$ ,  $\rho_2(a_k) = Y_k$  where the  $X_k$ 's are in  $(M_1, \tau_1)$  and the  $Y_k$ 's in  $(M_2, \tau_2)$ . Then

$$W_p(\tau', \tau'') = W_p((X_1, \dots, X_n), (Y_1, \dots, Y_n))$$

where  $\tau' = \tau_1 \circ \rho_1$ ,  $\tau'' = \tau_2 \circ \rho_2$ .

**1.3 Theorem.**  $W_p$  is a metric.

**Proof.** To check that  $W_p$  is a metric on the set of equivalence classes of  $n$ -tuples of variables or equivalently on a trace-state space  $ST(A)$  like in 1.2, the nontrivial assertion is the triangle inequality. Indeed that  $W_p((X_1, \dots, X_n), (Y_1, \dots, Y_n)) = 0 \Leftrightarrow (X_1, \dots, X_n), (Y_1, \dots, Y_n)$  have the same  $*$ -distribution or

$$W_p(\tau', \tau'') = 0 \Leftrightarrow \tau' = \tau''$$

are easy to see. For the triangle inequality it will suffice to prove it in the context of 1.1.

Let  $(X'_1, \dots, X'_n, Y'_1, \dots, Y'_n)$  in  $(M_{12}, \tau_{12})$  and  $(Y''_1, \dots, Y''_n, Z''_1, \dots, Z''_n)$  in  $(M_{23}, \tau_{23})$  be  $2n$ -tuples in tracial  $W^*$ -probability spaces such that  $(X'_1, \dots, X'_n) \sim (X_1, \dots, X_n)$ ,  $(Y'_1, \dots, Y'_n) \sim (Y''_1, \dots, Y''_n) \sim (Y_1, \dots, Y_n)$ ,  $(Z''_1, \dots, Z''_n) \sim (Z_1, \dots, Z_n)$ , where  $\sim$  means the  $n$ -tuples have equal  $*$ -distribution. There is a trace-preserving automorphism of  $W^*(Y'_1, \dots, Y'_n)$  and  $W^*(Y''_1, \dots, Y''_n)$  which identifies  $Y'_j$  and  $Y''_j$ . Abusing notations we shall denote by  $M_2$  the von Neumann subalgebras of  $M_{12}$  and  $M_{23}$  generated by  $(Y'_1, \dots, Y'_n)$  and respectively  $(Y''_1, \dots, Y''_n)$  identified as above. Let  $E'$  and  $E''$  be the conditional expectations of  $M_{12}$  and respectively  $M_{23}$  onto  $M_2$ .

Let  $(M_{123}, E) = (M_{12}, E') *_{M_2} (M_{23}, E'')$  and  $\tau_{123} = \tau_2 \circ E$  where  $\tau_2 = \tau_{12}|_{M_2} = \tau_{23}|_{M_2}$  (see 3.8 in [14]). Further, with  $\rho_{12} : M_{12} \rightarrow M_{123}$ ,  $\rho_{23} : M_{23} \rightarrow M_{123}$  denoting the canonical embeddings, let  $X'''_j = \rho_{12}(X'_j)$ ,  $Z'''_j = Z''_j$ . Then  $\rho_{12}(Y'_j) = \rho_{23}(Y''_j)$  implies

$$\|X'''_j - Z'''_j\|_{p, \tau_{123}} \leq \|X'''_j - \rho_{12}(Y'_j)\|_{p, \tau_{123}} + \|\rho_{23}(Y''_j) - Z'''_j\|_{p, \tau_{123}} = \|X'_j - Y'_j\|_{p, \tau_{12}} + \|Y''_j - Z''_j\|_{p, \tau_{23}}$$

which is precisely what we need to establish the triangle inequality

$$W_p((X_1, \dots, X_n), (Y_1, \dots, Y_n)) + W_p((Y_1, \dots, Y_n), (Z_1, \dots, Z_n)) \geq W_p((X_1, \dots, X_n), (Z_1, \dots, Z_n)) .$$

□

Let us also record as a proposition some easy consequences of the compacity of the trace-state space. The proof is left to the reader.

**1.4 Proposition.** (a) *The infimum in the definition of  $W_p$  is attained (both in the 1.1 and 1.2 contexts).*

(b) *Let  $\tau_1^{(k)}, \tau_1, \tau_2^{(k)}, \tau_2 \in TS(A)$  and assume  $\tau_j^{(k)}$  converges weakly to  $\tau_j$  as  $k \rightarrow \infty$  ( $j = 1, 2$ ). Then*

$$\liminf_{k \rightarrow \infty} W_p(\tau_1^{(k)}, \tau_2^{(k)}) \geq W_p(\tau_1, \tau_2) .$$

(c) *Let  $(X_1^{(k)}, \dots, X_n^{(k)}), (X_1, \dots, X_n)$ ,  $(Y_1^{(k)}, \dots, Y_n^{(k)}), (Y_1, \dots, Y_n)$  be  $n$ -tuples of variables in tracial  $C^*$ -probability spaces and assume that  $\|X_j^{(k)}\| \leq R$ ,  $\|X_j\| \leq R$ ,  $\|Y_j^{(k)}\| \leq R$ ,  $\|Y_j\| \leq R$ , and that  $(X_1^{(k)}, \dots, X_n^{(k)}), (Y_1^{(k)}, \dots, Y_n^{(k)})$  converge in  $*$ -distribution to  $(X_1, \dots, X_n)$  and respectively  $(Y_1, \dots, Y_n)$ . Then*

$$\liminf_{k \rightarrow \infty} W_p((X_1^{(k)}, \dots, X_n^{(k)}), (Y_1^{(k)}, \dots, Y_n^{(k)})) \geq W_p((X_1, \dots, X_n), (Y_1, \dots, Y_n)) .$$

If  $(X_1, \dots, X_n)$  are commuting self-adjoint variables in a tracial  $C^*$ -probability space, then their distribution  $\mu_{X_1, \dots, X_n}$  is a compactly supported probability measure on  $\mathbb{R}^n$ .

**1.5 Theorem.** *Let  $(X_1, \dots, X_n)$  and  $(Y_1, \dots, Y_n)$  be two  $n$ -tuples of commuting self-adjoint variables in tracial  $C^*$ -probability spaces. Then the free and classical Wasserstein distances are equal:*

$$W((X_1, \dots, X_n), (Y_1, \dots, Y_n)) = W(\mu_{X_1, \dots, X_n}, \mu_{Y_1, \dots, Y_n}) .$$

**Proof.** The left-hand side is  $\leq$  the right-hand side, since the classical Wasserstein distance can be defined the same way as the free one, with the only difference that the  $2n$ -tuples  $(X'_1, \dots, X'_n, Y'_1, \dots, Y'_n)$  in the infimum are required to live in commutative tracial  $C^*$ -probability spaces. We therefore only need to prove  $\geq$ .

Let  $(X'_1, \dots, X'_n, Y'_1, \dots, Y'_n)$  be a  $2n$ -tuple in the infimum defining the free distance. Passing to the von Neumann algebra completion, we may assume  $(M_3, \tau_3)$ , where  $X'_j, Y'_j$  live, is a  $W^*$ -probability space with a normal faithful trace state. Let  $A = W^*(X'_1, \dots, X'_n) \subset M_3$ ,  $B = W^*(Y'_1, \dots, Y'_n) \subset M_3$  and let  $E_A$  be the canonical conditional expectation onto  $A$ . Then the unital trace-preserving completely positive map  $\varphi = E_A|_B : B \rightarrow A$  gives rise to a state  $\nu : A \otimes B \rightarrow \mathbb{C}$ , on a commutative algebra, defined by

$$\nu(a \otimes b) = \tau_3(a\varphi(b)) .$$

The positivity of  $\nu$ ,

$$\tau_3\left(\sum_{i,j} a_i a_j^* \varphi(b_i b_j^*)\right) \geq 0 ,$$

is easily inferred from the positivity of the matrix  $(\varphi(b_i b_j^*))_{i,j}$ . Alternatively, probabilistically,  $\nu$  is the probability measure on  $\mathbb{R}^{2n}$  obtained by integrating w.r.t.  $\mu_{X_1, \dots, X_n}$  the kernel of probability measures describing

$$\varphi : L^\infty(\mathbb{R}^n, \mu_{Y_1 \dots Y_n}) \rightarrow L^\infty(\mathbb{R}^n, \mu_{X_1 \dots X_n}) .$$

Then

$$\begin{aligned}
\sum_{1 \leq j \leq n} \nu((X'_j - Y'_j)^2) &= \sum_{1 \leq j \leq n} \tau_3(X_j'^2 + \varphi(Y_j'^2) - X'_j \cdot \varphi(Y'_j) - Y'_j \varphi(X'_j)) \\
&= \sum_{1 \leq j \leq n} \tau_3(X'_j + Y_j'^2 - 2E_A(X'_j Y'_j)) \\
&= \sum_{1 \leq j \leq n} \tau_3((X'_j - Y'_j)^2) .
\end{aligned}$$

Since  $A \otimes B$  is commutative this proves the theorem.  $\square$

## 2 Cost of transportation to the semicircle distribution

### 2.1 The complex quasilinear differential equation

Let  $X, S$  in  $(M, \tau)$  be self-adjoint and freely independent and assume  $S$  is  $(0,1)$  semicircular. The purpose of section 2 is to estimate  $W(X, S)$ . We begin by studying variables  $X(t) = e^{-t/2}X + (1 - e^{-t})^{\frac{1}{2}}S$  which have the same distribution as the variables in the free Ornstein-Uhlenbeck process. For technical reasons, and without extra work, the complex PDE will be derived under the more general assumption that  $X$  is unbounded self-adjoint affiliated with  $M$  (see [1]).

If  $Y$  is self-adjoint affiliated with  $M$ , we denote by  $\mu_X$  its distribution and by  $G_{\mu_Y}(z)$  or  $G_Y(z)$  the Cauchy transform of  $\mu_Y$ , which equals  $\tau((zI - X)^{-1})$ .

If  $Y(r) = X + r^{\frac{1}{2}}S$ , let  $\tilde{G}(r, z) = G_{Y(r)}(z)$  and  $G(t, z) = G_{X(t)}(z)$ ,  $\text{Im } z > 0$ ,  $r \geq 0$ ,  $t \geq 0$ . Then  $\tilde{G}$  satisfies the complex Burgers equation (see [3],[12])

$$\frac{\partial \tilde{G}}{\partial r} + \tilde{G} \frac{\partial \tilde{G}}{\partial z} = 0 .$$

Like  $\tilde{G}(t, z)$  also  $G(t, z)$  is  $C^1$  on  $[0, \infty) \times \{z \in \mathbb{C} \mid \text{Im } z > 0\}$  and holomorphic in  $z$  for fixed  $t$ . Note that  $X(t) = e^{-t/2}Y(e^t)$  and that  $G_{\alpha Y}(z) = \alpha^{-1}G(\alpha^{-1}z)$ . It follows that  $G(t, z) = e^{t/2}\tilde{G}(e^t, e^{t/2}z)$ . The complex Burgers equation then gives

$$\frac{\partial G}{\partial t} + \left(G - \frac{z}{2}\right) \frac{\partial G}{\partial z} - \frac{1}{2}G = 0 \tag{1}$$

with initial data  $G(0, z) = G_X(z)$ .

## 2.2 The transport equation

Here we shall assume that the distribution of  $X$  is of the form  $P_\lambda * \mu$  where  $P_\lambda$  is the Cauchy distribution with density  $\pi^{-1}\lambda(\lambda^2 + x^2)^{-1}$  ( $\lambda > 0$ ) and  $\mu$  has compact support. Since  $P_\lambda * \mu = P_\lambda \boxplus \mu$  ([1]) this is equivalent to replacing  $X$  with  $X + \lambda C$  where  $X$  is bounded,  $X$  and  $C$  are free and  $C$  has a Cauchy distribution  $P_1$ . Note that  $\mu_{X+\lambda C+r\frac{1}{2}S} = \mu_{X+r\frac{1}{2}S} * P_\lambda$ ,  $G_{X+\lambda C+r\frac{1}{2}S}(z) = G_{X+r\frac{1}{2}S}(z + i\lambda)$ , etc. Thus, if the distribution of  $X$  is of the form  $P_\lambda * \mu$  then the equation (1) is satisfied on an extended domain

$$\{(t, z) \in [0, \infty) \times \mathbb{C} \mid \text{Im } z > -e^{t/2}\lambda\} .$$

Let  $-\pi^{-1}G(x, t) = q(x, t) + ip(x, t)$  where  $x \in \mathbb{R}$ . Then  $p(\cdot, t)$  is the density of  $\mu_{X(t)}$  and is analytic. For fixed  $t$  and  $k \geq 0$  we have

$$\left| \frac{\partial^k}{\partial x^k} p(x, t) \right| = O((1 + |x|)^{-2-k}) \quad \text{and} \quad \left| \frac{\partial^k}{\partial x^k} q(x, t) \right| = O((1 + |x|)^{-1-k}) .$$

Moreover these bounds are uniform for  $t$  in a compact set.

Equation (1) gives

$$\begin{aligned} q_t &= \pi(qq_x - pp_x) + 2^{-1}(xq_x + q) \\ p_t &= \pi(pq_x + qp_x) + 2^{-1}(xp_x + p) \\ q &= -Hp \end{aligned} \tag{2}$$

where  $H$  denotes the Hilbert transform.

Since  $p(x, t) > 0$  we infer that

$$f(a, t) = \int_{-\infty}^a p(x, t) dx$$

is a  $C^\infty$ -diffeomorphism  $f(\cdot, t) : \mathbb{R} \rightarrow (0, 1)$  which transports  $\mu_{X(t)}$  to Lebesgue measure. Hence  $\varphi_{s,t}(\cdot) = f^{-1}(f(\cdot, s), t)$  ( $0 < s < t$ ) will be a  $C^\infty$ -diffeomorphism  $\mathbb{R} \rightarrow \mathbb{R}$ , which transports  $\mu_{X(s)}$  to  $\mu_{X(t)}$ . This is the same as saying that  $X(t)$  and  $\varphi_{s,t}(X(s))$  have the same distribution.

It is easily seen that

$$\frac{\partial}{\partial t} f^{-1}(y, t) = \frac{-\left(\frac{\partial}{\partial t} f\right)(f^{-1}(y, t), t)}{p(f^{-1}(y, t), t)} .$$

Using (2) to compute  $\frac{\partial}{\partial t} f$  we find

$$\frac{\partial}{\partial t} f(a, t) = \int_{-\infty}^a (\pi(pq)_x + 2^{-1}(xp)_x) dx = \pi(pq)(a, t) + 2^{-1}ap(a, t) .$$

Hence

$$\frac{\partial}{\partial t} f^{-1}(y, t) = -\pi q(f^{-1}(y, t), t) - 2^{-1}f^{-1}(y, t) .$$

For  $y = f(x, s)$  we get the transport equation

$$\frac{\partial}{\partial t} \varphi_{s,t}(x) = \pi(Hp(\cdot, t))(\varphi_{s,t}(x)) - 2^{-1}\varphi_{s,t}(x) \quad (3)$$

with initial condition  $\varphi_{s,s}(x) = x$ .

By the  $L^m$ -continuity ( $1 < m < \infty$ ) results for the density (see Corollary 2 in [ ]) applied to  $\mu \boxplus \mu_{r, \frac{1}{2}S}$  as a function of  $r$ , we infer after convolutions with Cauchy distributions the continuity of

$$(0, \infty) \ni t \longrightarrow Hp(\cdot, t) \in L^m(\mathbb{R})$$

(the  $L^m$ -space w.r.t. Lebesgue measure). The reader should keep these facts in mind in computations where we shall use (3).

**Lemma 2.3** *Assume  $X$  has distribution  $\mu * P_\lambda$ , where  $\mu$  has compact support and let  $X(t) = e^{-t/2}X + (1 + e^{-t})^{\frac{1}{2}}S$  with  $S$   $(0, 1)$ -semicircular and free from  $X$ . Let  $g \in C^\infty(\mathbb{R})$  be such that  $\|g\|_\infty < \infty$ ,  $\|g'\|_\infty \leq 1$  and assume  $g'$  has compact support. Then*

$$(t - s)^2 W(g(X(s)), g(X(t)))^2 \leq \sup_{s \leq h \leq t} \int_{\text{supp } g'} (\pi Hp(\cdot, h)(x) - 2^{-1}x)^2 p(x, h) dx .$$



**Proof.** We have

$$\begin{aligned}
& W(g(X(s)), g(X(t)))^2 \\
& \leq \int_{\mathbb{R}} |g(x) - g(\varphi_{s,t}(x))|^2 p(x, s) dx \\
& \leq \int_{\mathbb{R}} \left( \int_s^t g'(\varphi_{s,h}(x)) (\pi H p(\cdot, h)(\varphi_{sh}(x)) - 2^{-1} \varphi_{s,h}(x)) dh \right)^2 p(x, s) dx \\
& \leq (t-s) \int_{\mathbb{R}} \int_s^t (g'(\varphi_{s,h}(x)))^2 (\pi H p(\cdot, h)(\varphi_{sh}(x)) - 2^{-1} \varphi_{s,h}(x))^2 dh p(x, s) dx \\
& = (t-s) \int_s^t \left( \int_{\mathbb{R}} (g'(\varphi_{s,h}(x)))^2 (\pi H p(\cdot, h)(\varphi_{s,h}(x)) - 2^{-1} \varphi_{s,h}(x))^2 dh p(x, s) dx \right) dh \\
& = (t-s) \int_s^t \int_{\mathbb{R}} (g'(x))^2 (\pi H p(\cdot, h)(x) - 2^{-1} x)^2 p(x, h) dx dh \\
& \leq (t-s)^2 \sup_{s \leq h \leq t} \int_{\text{supp } g'} (\pi H p(\cdot, h)(x) - 2^{-1} x)^2 p(x, h) dx .
\end{aligned}$$

□

**2.4.** Assume  $X$  is bounded and the semicircular variable  $S$  is free w.r.t.  $X$ . Then the distribution  $\mu_{X(t)}$  of  $X(t) = e^{-t/2}X + (1 - e^t)^{\frac{1}{2}}S$  has  $L^\infty$ -density  $p(\cdot, t)$  w.r.t. Lebesgue measure (see any of the papers [1],[2],[3],[11],[12]).

**Lemma.** Assume  $X$  is bounded,  $S$  is  $(0, 1)$  semicircular,  $X$  and  $S$  are free and let  $p(\cdot, t)$  be the density of  $\mu_{X(t)}$ , where  $X(t) = e^{-t/2}X + (1 - e^t)^{\frac{1}{2}}S$ . Then

$$(t-s)^{-2} W(X(s), X(t))^2 \leq \sup_{s \leq h \leq t} \int (\pi H p(\cdot, h)(x) - 2^{-1} x)^2 p(x, h) dx .$$

**Proof.** Let  $C$  be a variable with Cauchy distribution and free w.r.t  $\{X, S\}$ . Let  $g \in C^\infty(\mathbb{R})$  be such that  $\|g'\|_\infty \leq 1$ ,  $g(x) = x$  if  $|x| \leq \|X\| + 1$  and  $g'(x) = 0$  if  $|x| \geq \|X\| + 2$ . We shall apply Lemma 2.3 to  $X + \lambda C$  in place of  $X$ . Let

$$Z(t, \lambda) = e^{-t/2}(X + \lambda C) + (1 - e^{-t})^{\frac{1}{2}}S = X(t) + e^{-t/2}\lambda C .$$

Then  $g(Z(t, \lambda))$  is an operator of norm  $\leq \|X\| + 2$  and converges in distribution to  $X(t)$ . Moreover the distribution of  $Z(t, \lambda)$  is given by the density  $P_{e^{-t/2}\lambda} * p(\cdot, t)$  and will be

denoted by  $p(\cdot, t, \lambda)$ . In view of the  $L^m$ -continuity of  $p(\cdot, t)$  ( $1 < m < \infty$ ) ([12]) it is easy to see that

$$\begin{aligned} & \limsup_{\lambda \downarrow 0} \left( \sup_{s \leq h \leq t} \int_{\text{supp } g'} (\pi H p(\cdot, h, \lambda)(x) - 2^{-1}x)^2 p(x, h, \lambda) dx \right) \\ & \leq \sup_{s \leq h \leq t} \int (\pi H p(\cdot, h)(x) - 2^{-1}x)^2 p(x, h) dx . \end{aligned}$$

□

**2.5.** From now on we return to the context of bounded variables  $X$ . If the distribution of  $X$  is Lebesgue absolutely continuous and has density  $p$  which is  $L^3$ , then  $\frac{1}{2}\mathcal{J}(X) = \pi H p(X)$  where  $\mathcal{J}(X)$  is the conjugate variable (a.k.a. free Brownian gradient, a.k.a. noncommutative Hilbert transform) (see [13]) and

$$\Phi(X) = \tau(\mathcal{J}(X)^2) = 4\pi^2 \int (H p(x))^2 p(x) dx = \frac{4}{3}\pi^2 \int p^3(x) dx$$

is the free Fisher information (see [11],[13] up to different normalizations). The quantity occurring in Lemma 2.4,

$$I(X) = 4 \int (\pi H p(x) - 2^{-1}x)^2 p(x) dx = \tau((\mathcal{J}(X) - X)^2) = \Phi(X) - 2 + \tau(X^2) ,$$

is a generalization of the free Fisher information for Ornstein-Uhlenbeck processes (see [4]). The inequality in Lemma 2.4 can also be written

$$4(t-s)^{-2} W(X(s), X(t))^2 \leq \sup_{s \leq h \leq t} I(X(h)) . \quad (4)$$

## 2.6 The free entropy

The free entropy of  $X$  with distribution  $\mu = \mu_X$  is

$$\chi(X) = \iint \log |s - t| d\mu(s) d\mu(t) + \frac{3}{4} + \frac{1}{2} \log(2\pi)$$

(see [11],[13] up to different constants) and we have

$$\chi(\alpha X) = \chi(X) + \log |\alpha| \quad \text{and} \quad \lim_{\varepsilon \downarrow 0} \varepsilon^{-1} (\chi(X + \varepsilon^{\frac{1}{2}} S) - \chi(X)) = 2^{-1} \Phi(X) .$$

The quantity we shall use in estimating the distance to the semicircle distribution is a modified free entropy adapted to the free Ornstein-Uhlenbeck process ([4]):

$$\begin{aligned}\tilde{\Sigma}(X) &= -\chi(X) + \chi(S) + \frac{1}{2}\tau(X^2) - \frac{1}{2} \\ &= \frac{1}{2}\tau(X^2) - \iint d\mu(s)d\mu(t) \log |s - t| - \frac{3}{4}.\end{aligned}$$

We have  $\lim_{t \rightarrow \infty} \tilde{\Sigma}(X(t)) = 0$  and

$$\begin{aligned}\frac{d}{dt}\tilde{\Sigma}(X(t)) &= \frac{d}{dt} \left( \frac{t}{2} - \chi(X + (e^t - 1)^{\frac{1}{2}}S) + \frac{1}{2}e^{-t}\tau(X^2) + \frac{1}{2}(1 - e^{-t}) \right) \\ &= 2^{-1} \left( 1 - e^t\Phi(X + (e^t - 1)^{\frac{1}{2}}S) - e^{-t}\tau(X^2) + e^{-t} \right) \\ &= 2^{-1}(1 - \Phi(X(t)) - \tau(X(t)^2) + 1) = -2^{-1}I(X).\end{aligned}$$

Note also that in [4] using the logarithmic Sobolev inequality for  $\chi$  (Prop. 7.9 in [13]), it is shown that

$$\tilde{\Sigma}(X(t)) \leq 2^{-1}I(X(t)) \tag{5}$$

which is a logarithmic Sobolev inequality for the Ornstein-Uhlenbeck process.

**Lemma 2.7** *Assume  $X, Y$  are bounded and self-adjoint, then if  $t > 0$  we have*

$$\limsup_{\varepsilon \rightarrow 0} |\varepsilon|^{-1} |W(Y, X(t + \varepsilon)) - W(Y, X(t))| \leq 2^{-1}(I(X(t)))^{\frac{1}{2}}$$

**Proof.** By the triangle inequality for  $W$ , we have

$$|W(Y, X(t + \varepsilon)) - W(Y, X(t))| \leq W(X(t), X(t + \varepsilon)).$$

The lemma then follows from (4) and the continuity of  $I(X(h))$  ( $h > 0$ ), which is a consequence of the continuity of  $\Phi(X(h))$  (Corollary 2 in [12]).  $\square$

We now have all ingredients to get an estimate for  $W(X, S)$  which is similar in the free context to an inequality of Talagrand in the classical setting ([7],[10]).

**Theorem 2.8**  $W(X, S)^2 \leq \tilde{\Sigma}(X)$ .

**Proof.** Because of the semicircular maximum for  $\chi$  we have  $\chi(X) \leq \chi(S) + 2^{-1} \log(\tau(X^2))$  so that  $\tilde{\Sigma}(X) \geq 2^{-1}(\tau(X^2) - (1 + \log \tau(X^2))) \geq 0$ . Thus it will suffice to prove that  $W(X, S) - (\tilde{\Sigma}(X))^{\frac{1}{2}} \leq 0$ .

By Lemma 2.7, the inequality (5) and the formula for the derivative of  $\tilde{\Sigma}(X(t))$ , we have for  $t > 0$ ,

$$\begin{aligned} & \liminf_{\varepsilon \rightarrow 0} \varepsilon^{-1} (W(X(t+\varepsilon), S) - (\tilde{\Sigma}(X(t+\varepsilon)))^{\frac{1}{2}} - W(X(t), S) + (\tilde{\Sigma}(X(t)))^{\frac{1}{2}}) \\ & \geq -2^{-1}(I(X(t)))^{\frac{1}{2}} + 2^{-2}I(X(t))(\tilde{\Sigma}(X(t)))^{-\frac{1}{2}} \\ & \geq 2^{-1}(I(X(t)))^{\frac{1}{2}} + 2^{-2+1}I(X(t))(I(X(t)))^{-\frac{1}{2}} = 0. \end{aligned}$$

Hence  $W(X(t), S) - (\tilde{\Sigma}(X(t)))^{\frac{1}{2}}$  is an increasing function and we have

$$\lim_{t \rightarrow \infty} (W(X(t), S) - (\tilde{\Sigma}(X(t)))^{\frac{1}{2}}) = 0$$

because of the semicircular maximum and lower semicontinuity of  $\chi$ . It follows that

$$W(X(t), S) - (\tilde{\Sigma}(X(t)))^{\frac{1}{2}} \leq 0$$

if  $t > 0$ . To get the inequality for  $t = 0$ , remark that  $X(t)$  is norm-continuous so that  $W(X(t), S)$  tends to  $W(X, S)$  as  $t \rightarrow 0$ . On the other hand, by lower semicontinuity of  $\chi$ ,

$$\liminf_{t \downarrow 0} (-(\tilde{\Sigma}(X(t)))^{\frac{1}{2}}) \geq -(\tilde{\Sigma}(X))^{\frac{1}{2}}.$$

□

## 2.9 Remark.

Because of the coincidence of the free and classical Wasserstein distance for single self-adjoint variables, the preceding theorem can also be written in terms of probability measures for the classical distance. Let  $\mu$  be a compactly supported probability measure on  $\mathbb{R}$  and  $\sigma$  a (0,1)-semicircle distribution. Then we have

$$(W(\mu, \sigma))^2 \leq \frac{1}{2} \int x^2 d\mu(x) - \iint d\mu(s) d\mu(t) \log |s - t| - \frac{3}{4}.$$

**Acknowledgment.** This research was conducted by Dan Voiculescu for the Clay Mathematics Institute. He was also supported in part by National Science Foundation grant DMS95–00308.

## References

- [1] Bercovici, H., Voiculescu, D. Free convolution of measures with unbounded support. *Indiana Univ. Math. J.* **42** (1993), no. 3, 733–773.
- [2] Biane, P. Processes with free increments. *Math. Z.* **227** (1998), 143–174.
- [3] Biane, P. On the free convolution with a semicircular distribution. *Indiana Univ. Math. J.* **46** (1997), 705–717.
- [4] Biane, P., Speicher R. Free diffusions, free entropy and free Fisher information. Preprint DMA, ENS 99–33 (1999).
- [5] Connes, A. Compact metric spaces, Fredholm modules and hyperfiniteness. *Ergodic Theory and Dynamical Systems* **9** (1989), 207–220.
- [6] Evans, L.C. Partial differential equations and Monge-Kantorovich mass transfer. Preprint.
- [7] Otto, F., Villani, C. Generalization of an inequality by Talagrand, and links with the logarithmic Sobolev inequality. Preprint DMA, ENS 99–23 (1999).
- [8] Rachev, S.T. *Probability Metrics and the Stability of Stochastic Models*, Wiley series in probability and mathematical statistics (1991).
- [9] Rieffel, M.A. Metrics on state space. Preprint (1999).
- [10] Talagrand, M. Transportation cost for Gaussian and other product measures. *Geom. Funct. Anal.* **6** (1996), 587–600.
- [11] Voiculescu, D. The analogues of entropy and of Fisher’s information measure in free probability theory, I. *Commun. Math. Phys.* **155** (1993), 71–92.

- [12] Voiculescu, D. The derivative of order  $1/2$  of a free convolution by a semicircle distribution. *Indiana Univ. Math. J.* **46** (1997), 697–703.
- [13] Voiculescu, D. The analogues of entropy and of Fisher’s information measure in free probability theory, V: noncommutative Hilbert transforms. *Invent. Math.* **132** (1998), 189–227.
- [14] Voiculescu, D., Dykema, K., Nica, A. *Free Random Variables*. CRM Monograph Series no. 1, American Mathematical Society, Providence, RI (1992).

Philippe Biane  
CNRS, DMA, École Normale Supérieure  
45 Rue d’Ulm  
75005 Paris, France  
BIANE@DMI.ENS.FR

Dan Voiculescu  
Department of Mathematics  
University of California  
Berkeley, California 94720–3840 USA  
dvv@math.berkeley.edu