

SUBNORMAL EMBEDDING THEOREMS FOR GROUPS

VAHAGN H. MIKAELIAN

To the memory of my brother Sassoun Mikaelian

ABSTRACT. In this paper we establish some subnormal embeddings of groups into groups with additional properties; in particular embeddings of countable groups into 2-generated groups with some extra properties. The results obtained are generalizations of theorems of P. Hall, R. Dark, B. Neumann, Hanna Neumann, G. Higman on embeddings of that type. Considering subnormal embeddings of finite groups into finite groups with additional properties we bring some illustration to a result of H. Heineken and J. Lennox.

1. INTRODUCTION AND THE MAIN RESULTS

The aim of this paper is to introduce a method of subnormal “verbal” embeddings of groups into larger groups with additional properties. First we prove the following main result:

THEOREM 1. *Let H be an arbitrary infinite group and $V \subseteq F_\infty$ be an arbitrary non-trivial word set. Then*

A. *there exists a group G of the same cardinality as H such that G has a subnormal subgroup $\tilde{H} \cong H$ and $\tilde{H} \subseteq V(G)$;*

B. *if the group H is countable the (countable) group G can be chosen two-generated;*

C. *if the group H is solvable the group G of the statement **A** or the two-generated group of the statement **B** can be chosen as well solvable.*

This theorem is also true for finite groups, as a special case of countable groups. However not each finite group H can be subnormally embedded into some finite group G such that the isomorphic image of H lies in $V(G)$. We will return to the case of finite groups later.

To avoid repetition we will sometimes talk of the group H and of its image \tilde{H} in G as of the same group.

In fact the statement **A** establishes a subnormal embedding of defect two. Thus it is natural to ask whether this defect can be reduced to one, i. e. whether the “verbal” embedding of H in G can be *normal*. The answer is contained in Theorem 2 in Section 3.

The statement **B** is a generalization of a theorem of G. Higman, B. Neumann and Hanna Neumann on embeddability of an arbitrary countable group in a 2-generated

Date: 30 May 1998.

2000 Mathematics Subject Classification. 20E07, 20E10.

JOURNAL OF THE LONDON MATHEMATICAL SOCIETY, 62.2 (2000), 398–406.

group (1947, [8]). The theorem, called in the book of Robinson [16] “probably the most famous of all embedding theorems”, was a stimulus for further research on embeddings into 2-generated groups and related problems. See the papers of Ol’shanskii [14] [15], Galvin [2], B. Neumann [11], Wilson and Zalesskii [17], Levin and Rosenberger [9], and literature cited there.

The idea that this embedding can be subnormal belongs to P. Hall and R. Dark [1]. In fact the Hall–Dark construction can be slightly modified to get a subnormal embedding into the *derived group* of some two-generated group.

The statement **C** generalizes the result of B. Neumann and Hanna Neumann [12] on embeddability of *solvable* countable groups in the second derived groups of *solvable* two-generated groups. Considering in some sense more “economical” embeddings B. Neumann and Hanna Neumann proved that the solvable group H of length n can be embedded into a two-generated solvable group G of length $n + 2$ but not in general of length $n + 1$, this is Corollary 5.2 of Theorem 5.1 [12] saying that each group of variety \mathfrak{U} can be embedded into some two-generated group of variety $\mathfrak{U}\mathfrak{A}^2$, where \mathfrak{A} is the variety of all abelian groups. This result cannot be strengthened by the requirement to embed into a verbal subgroup (see below). Nevertheless a simplified version of our construction can be used in order to make both embeddings of the mentioned Theorem 5.1 and Corollary 5.2 [12] *subnormal* (respectively Theorems 3 and 4 in Section 4).

The importance of the case of solvable groups follows, in particular, from the fact that we could *not* obtain the analogs of the mentioned result for *abelian* or *nilpotent* groups because each subgroup of a two-generated abelian group should be as well two-generated, and because each subgroup of a finitely generated nilpotent group should be as well finitely generated [16]. The countable abelian group, say, \mathbb{Q} of rational numbers is infinitely generated and cannot be contained in any finitely generated nilpotent group.

The statement **A** of Theorem 1 is in contrast with the *negative* results regarding subnormal embeddings of *finite* groups into *finite* groups. P. Hall and R. Dark have shown that for the symmetric group S_3 there is no finite G such that $S_3 \triangleleft\triangleleft G$ and $S_3 \subseteq G'$ (see in [1]). This result is strongly generalized by Heineken and Lennox [5] who proved that for *no* finite solvable complete group H there exists a finite G such that $H \triangleleft\triangleleft G$ and $H \subseteq G'$ (see also the extension of the latter result by Heineken and Soules [7]).

We make this contrast more visible in Section 5 and show that the result of Heineken and Lennox finds almost the “smallest” possible finite groups of that type. Namely we prove first that for each finite *nilpotent* group H and each non-trivial set of words V there exists a finite (and nilpotent) group G such that $H \triangleleft\triangleleft G$ and $H \subseteq V(G)$ (Proposition 1).

Thus it is very natural to ask whether there exist non-nilpotent (non-solvable) finite groups H for which it is possible (impossible) to find a finite group G such that $H \triangleleft\triangleleft G$ and $H \subseteq V(G)$. Proposition 2 says something more general; for *each* variety \mathfrak{U} different from the variety \mathfrak{D} of all groups there exist finite groups *not* from \mathfrak{U} that have (or do not have) embeddings of the mentioned type. And it is possible to find such groups in some very small classes of groups (Proposition 3).

This shows that the question of existence of subnormal “verbal” embeddings of finite groups into finite groups is far from being a trivial task. We dare formulate it as a *problem* at the end of the current paper.

Acknowledgments

It is a great pleasure for the author to express his gratitude to Professor Hermann Heineken for his attention to my research project and for useful considerations while I enjoyed the very warm hospitality of the Mathematics Institute of the Würzburg University, Germany. The author is thankful to the German Academic Exchange Service for financial support (Grant Nr. A/97/13683).

2. THE MAIN CONSTRUCTION

The following steps prove Theorem 1.

a. Let L be a nilpotent countable group such that its verbal subgroup $V(L)$ contains a subgroup isomorphic to \mathbb{Z} . To construct L we note that the set of all nilpotent groups generates the variety \mathfrak{D} of all groups (free groups are residually finite p -groups). The variety

$$\mathfrak{D} = \text{var}(F_\infty/V(F_\infty))$$

corresponding to the word set V is different from \mathfrak{D} . Thus there exists a nilpotent group $L = F_n(\mathfrak{N}_c)$ of some class c and some rank n such that $L \notin \mathfrak{D}$. An element a of infinite order in non-trivial and torsion-free subgroup $V(L)$ generates an isomorphic copy of $\mathbb{Z} = \langle 1_{\mathbb{Z}} \rangle \cong \langle a \rangle$. $1_{\mathbb{Z}}$ is not the trivial element of \mathbb{Z} (unlike in the notions that will come later) but the integer 1.

b. In order to define a special linear order \prec on the countable group L we establish the one-to-one correspondence α between the additive group of integers $\mathbb{Z} = \langle 1_{\mathbb{Z}} \rangle$ and L :

$$\alpha(i) = \alpha(i \cdot 1_{\mathbb{Z}}) \text{ is equal to } \begin{cases} a^i & \text{if and only if } i \in \{\dots, -2, -1, 0\}, \\ \text{an arbitrary } x \in L \setminus \{a^j\}_{j \leq 0} & \text{if } i \in \{1, 2, \dots\} \end{cases}$$

in such a manner that α is, in particular, a bijection between the sets

$$\mathbb{Z} \setminus \{\dots, -2, -1, 0\} \quad \text{and} \quad L \setminus \{\dots, a^{-2}, a^{-1}, 1_L\}.$$

Now define

$$\forall l_1, l_2 \in L, \quad l_1 \prec l_2 \quad \text{if and only if} \quad \alpha^{-1}(l_1) \leq \alpha^{-1}(l_2).$$

c. Consider the (cartesian) wreath product $H \text{Wr} L$ and assume that the copies of H in the base group $H^{(L)}$ are also ordered according to \prec . To be brief we write simply h_i instead of $h_{\alpha(i)}$ for element contained in $\alpha(i)$ 'th copy of H . Now we define the sets D_k ($k \in \mathbb{N} \cup \{0\}$) of elements of the base group as follows: $\bar{h} = (\dots, h_{-1}, h_o, h_1, \dots) \in D_k$ if and only if there exist integers n_1, n_2, \dots, n_k such that $n_i < n_{i+1}$, $i = 1, \dots, k-1$ and

$$h_i = \begin{cases} h_{n_1} & \text{if } i \leq n_1, \\ h_{n_2} & \text{if } n_1 < i \leq n_2, \\ \vdots & \vdots \\ h_{n_k} & \text{if } n_{k-1} < i \leq n_k, \\ h_{n_{k+1}} & \text{if } n_k < i. \end{cases}$$

Clearly $\text{card}(D_k) = \text{card}(H)$, because for each $\bar{h} = (\dots, h_{-1}, h_o, h_1, \dots) \in D_k$ we have only finitely many "jump" point indices n_1, n_2, \dots, n_k ; for each such an

index one can find only countably many “places” and because for each index n_i ($i = 1, \dots, k$) and for index $n_k + 1$ the values of h_i have only $\text{card}(H)$ possibilities.

The group L is countable. Therefore the subgroup

$$G_1 = \langle L, \bigcup_{i=1}^{\infty} D_k \rangle$$

of $H \text{ Wr } L$ is of cardinality $\text{card}(H)$ for it is generated by an infinite set of cardinality $\text{card}(H)$.

d. The group H is isomorphic to its first copy in the base group $H^{(L)}$. This copy lies in G_1 because

$$\bar{h}_0 = (\dots, 1_H, h, 1_H, \dots) \in D_2$$

(the element h has the number 0). Let us further take for an arbitrary $h \in H$ an element $\theta(h) \in H^{(L)}$:

$$\theta(h) = (\dots, h^{-1}, h^{-1}, h^{-1}, 1_H, 1_H, 1_H, \dots) \in D_1 \subseteq G_1$$

(the first element 1_H stands in the place number 0). Since

$$a \in V(L) \subseteq V(G_1)$$

and since verbal subgroups of groups are characteristic, we assert that the element

$$a^{\theta(h)} = a \cdot (\dots, h, h, 1_H, 1_H, \dots)^a \cdot (\dots, h^{-1}, h^{-1}, 1_H, 1_H, \dots)$$

belongs to $V(G_1)$. The order relation \prec is defined over L in such a manner that the elements standing in the places number $\dots, -2, -1, 0$ go under operation of a (right regular representation) one step “higher” and the elements standing in the places number $1, 2, 3, \dots$ go to some new places with numbers again from the set $\{1, 2, 3, \dots\}$. Thus

$$(\dots, h, h, 1_H, 1_H, \dots)^a = (\dots, h, h, h, 1_H, \dots),$$

where in the second string the first element 1_H stands now in the place number 1. We get

$$a^{\theta(h)} = a \cdot (\dots, 1_H, h, 1_H, \dots) = a \cdot \bar{h}_0 \in V(G_1).$$

Thus $\bar{h}_0 \in V(G_1)$ for each $h \in H$. This proves the statement **A** of Theorem 1 because the first copy of H in the base group is clearly subnormal in wreath product $H \text{ Wr } L$ and in $G = G_1$. And it is clear that if the group H is solvable, the group $G \subseteq H \text{ Wr } L$ and its subgroup G are solvable, too.

e. Next we need a group M such that $\text{card}(M) = \text{card}(G_1)$ and $G_1 \triangleleft\triangleleft M'$. We can simply use the statement **A** we just proved. But for this case of the *commutator* word V we could build a much smaller construction. That will be done in the proof of Theorem 3.

f. Finally we use an idea of P. Hall from [3]. Let us H be a countable group subnormally embedded into the derived subgroup of the group M and assume that the elements of the countable group M are linearly ordered:

$$M = \{m(0), m(1), m(2), \dots\}.$$

We define:

$$m_i = \begin{cases} m(s) & \text{if } i = 2^s, \text{ for some } s \in \{0, 1, 2, \dots\}, \\ 1_M & \text{if } i \neq 2^s, \text{ for arbitrary } s \in \{0, 1, 2, \dots\} \end{cases}$$

and

$$\bar{m} = (\dots, m_{-1}, m_0, m_1, \dots) \in M^{(\mathbb{Z})} \subseteq M \text{Wr } \mathbb{Z}.$$

Then for the intersection $T = \langle -1_{\mathbb{Z}}, \bar{m} \rangle \cap M^{(\mathbb{Z})}$ we have according [3]:

$$T' = \overline{M'}^{(\mathbb{Z})} = K,$$

where the latter means *direct* product of copies of M' .

K is normal in $G_2 = \langle -1_{\mathbb{Z}}, \bar{m} \rangle$ for it is normal even in $M \text{Wr } \mathbb{Z}$. Therefore the subnormal embedding we are looking for can be given as:

$$H \triangleleft\triangleleft G_1 \triangleleft\triangleleft M' \triangleleft K \triangleleft G_2.$$

(As we mentioned above, in order to avoid huge formulas, we talk of a group and its isomorphic image in the larger group as of the same group.) This proves the statement **B** of Theorem 1. And the part of the statement **C** concerning embeddability into two-generated solvable groups follows from the fact that all groups evolved in points **e** and **f** above are solvable, too.

Theorem 1 is proved. \square

3. THE DEFECTS OF EMBEDDINGS, THE CASE OF NORMAL EMBEDDABILITY

The embedding of $H \triangleleft\triangleleft G_1$ constructed in points **a**, ..., **d** of Section 2 is of defect 2. The first copy of H is subnormal of defect 2 in $H \text{Wr } L$ and G_1 is a subgroup of $H \text{Wr } L$ containing this first copy.

A natural question which arises is: whether this defect *can* (or *cannot*) be made smaller (namely 1), i. e.

Do there exist groups H that can (or cannot) be *normally* embedded into appropriate groups G such that the isomorphic image \tilde{H} of H lies in $V(G)$ for the given non-trivial word set V ?

The general answer to this question (not only for the case of infinite groups) is given in our co-work with Heineken [6], we can assert:

THEOREM 2. *Let us H be an arbitrary infinite group and $V \subseteq F_\infty$ be an arbitrary non-trivial word set. Then there exists a group G of the same cardinality as H such that G has a subnormal subgroup $\tilde{H} \cong H$ of defect at most 2 and $\tilde{H} \subseteq V(G)$. This embedding is of defect 1 (i. e. normal) if and only if $V(\text{Aut}(H)) \supseteq \text{Inn}(H)$.*

Proof. This follows from Theorem 1, from the remark made above, from Theorem 1 of [6] and from the fact that if the infinite group H is normally embeddable into some group G and lies in $V(G)$, then without loss of generality G is of the same cardinality as H . Let us show it.

The infinite group H has a system of generators of the same cardinality as H . So the image $\tilde{H} \subseteq V(G)$ is generated by the appropriate images, say, \tilde{h}_i , $i \in I$. For each element $\tilde{h}_i \in V(G)$ there is a representation

$$\tilde{h}_i = \left(w_1^{(i)}(g_{11}^{(i)}, \dots, g_{1q_1}^{(i)}) \right)^{\delta_1^{(i)}} \cdots \left(w_u^{(i)}(g_{u1}^{(i)}, \dots, g_{uq_u}^{(i)}) \right)^{\delta_u^{(i)}},$$

where $w_1^{(i)}, \dots, w_u^{(i)}$ are some words from V , elements $\delta_u^{(i)}$ take values 1 or -1 , and where $g_{jk}^{(i)} \in G$. Let us G_1 be the subgroup of G generated by this set Q of elements $g_{jk}^{(i)}$. Of course $\tilde{H} \subseteq V(G_1)$ and $\tilde{H} \triangleleft G_1$. Furthermore Q is of the same cardinality as H , and so is G_1 . \square

For concrete examples of groups that do not have normal embeddings of mentioned type see [6].

The obtained embedding of countable groups into two-generator groups has a larger defect, namely 6 (because the defect of H in G_1 is 2 and because the defect of the latter in M' is also 2).

Computation of an other nature can reduce this defect to 4. But we will not include that here since that would demand an apparatus strongly different from our main construction.

4. THE “ECONOMICAL” EMBEDDINGS OF SOLVABLE GROUPS INTO SOLVABLE GROUPS

The initial step here are the results of B. Neumann and Hanna Neumann, mentioned in the introduction, on embedding of a countable group of a variety \mathfrak{U} into some two-generated group of the variety $\mathfrak{U}\mathfrak{A}^2$ and on embedding of a countable solvable group of length l into a two-generated solvable group of length at most $l + 2$. (Theorem 5.1 and Corollary 5.2 in [12]) In fact these both embeddings are embeddings not only into the appropriate two-generated group G but into the *second derived group* of G .

Mentioned results are in some sense the best possible ones, since some solvable groups of length l *cannot* be embedded into two-generated solvable groups of length $l + 1$. As an example see Lemma 5.3 in [12].

As we mentioned above we cannot add the condition with verbal subgroups to these results. For if we take as V , say, the identity $\delta_l(x_1, \dots, x_{2^l})$ of solvability of length l : $\delta_0 = x$ and

$$\delta_{l+1}(x_1, \dots, x_{2^{l+1}}) = [\delta_l(x_1, \dots, x_{2^l}), \delta_l(x_{2^l+1}, \dots, x_{2^{l+1}})]$$

with $l \geq 3$, we get that for all $n \in \mathbb{N}$ the verbal subgroup $\delta_l(G)$ of each solvable group G of length $n + 2$ is of length $n + 2 - l < n$.

But we are able to use an easier version of our construction to make the embeddings of the mentioned results *subnormal*. Namely we build the construction mentioned in point **e** of Section 1 and prove:

THEOREM 3. *For an arbitrary countable group H of the variety \mathfrak{U} there exists in the variety $\mathfrak{U}\mathfrak{A}^2$ a two-generated group G with a subnormal subgroup \tilde{H} such that $\tilde{H} \cong H$.*

The following is an immediate consequence of Theorem 3 for the value $\mathfrak{U} = \mathfrak{A}^l$:

THEOREM 4. *For an arbitrary countable solvable group H of length l there exists a two-generated group G of length at most $l + 2$ (but not, in general, of length $l + 1$) with a subnormal subgroup \tilde{H} such that $\tilde{H} \cong H$.*

Proof of Theorem 3. We consider the wreath product $H \text{ Wr } \mathbb{Z}$. For each $h \in H$ we take a

$$\beta(h) = (\dots, 1_H, 1_H, h^{-1}, h^{-1} \dots) \in H^{(\mathbb{Z})}$$

(the first element h^{-1} is situated in the place number 0). Then it is easy to compute that

$$[1_{\mathbb{Z}}, \beta(h)] = (\dots, 1_H, h, 1_H, \dots)$$

(h is in the place number 0). We should now just note that the set of all strings on the right for all $h \in H$ forms the first copy of H in $H \text{ Wr } \mathbb{Z}$.

Let

$$M_1 = \langle \mathbb{Z} \cup \{\beta(h) \mid h \in H\} \rangle.$$

M_1 is a countable group and contains H as a subnormal subgroup. Besides an isomorphic copy of H lies in M'_1 . Therefore we can use the point \mathbf{f} from Section 2 in order to build a special element \bar{m}_1 in the wreath product $M_1 \text{ Wr } \mathbb{Z}$ such that the intersection

$$\langle -1_{\mathbb{Z}}, \bar{m}_1 \rangle \cap M_1^{(\mathbb{Z})} = \overline{M'_1}^{(\mathbb{Z})}$$

contains an isomorphic copy of H . Thus Theorem 3 is proved for the group $G = \langle -1_{\mathbb{Z}}, \bar{m}_1 \rangle \in \mathfrak{U}\mathfrak{Q}^2$. \square

We note that in these theorems, too, the group H is embedded not only into some appropriate two-generated group G but into the second derived group of the latter.

Our construction strongly depends on the fact that the active groups of our wreath products contain a subgroup isomorphic to \mathbb{Z} . Thus this construction cannot be used in embeddings of, say, finite groups into finite groups. B. Neumann and Hanna Neumann prove in [12] that arbitrary *finite* group H of the variety \mathfrak{U} is embeddable into (the second derived group) of some *finite* two-generated group $H \in \mathfrak{U}\mathfrak{Q}^2$. But the result of Heineken and Lennox [5] we mentioned shows that the “verbal” embeddings of finite groups into finite groups are in general *not* subnormal.

Finally we observe that the results we proved for solvable group have no analogs for the cases of abelian or nilpotent groups (see the arguments in the introduction).

5. ON “VERBAL” EMBEDDINGS OF FINITE GROUPS INTO FINITE GROUPS

In the sequence we consider embeddings of finite groups only. Since the mentioned results of P. Hall and R. Dark [1] and of Heineken and Lennox [5] concern finite *solvable* groups, the following Proposition 1 shows that those results describe in some sense the “smallest” possible finite groups without finite subnormal “verbal” embeddings.

PROPOSITION 1. *Let H be an arbitrary finite nilpotent group and V be an arbitrary non-trivial word set. Then there exists a finite nilpotent group G with a subnormal subgroup \tilde{H} such that*

$$\tilde{H} \cong H \quad \text{and} \quad \tilde{H} \subseteq V(G).$$

Moreover if H is a Π -group for the set Π of primes, the group G can be chosen a Π -group, too.

Proof. It is clear that if for each group of the family $\{H_i\}_{i \in I}$ there is an embedding of type stated in Proposition 1 (into a group G_i , $i \in I$), then for the (direct or cartesian) product $\prod_{i \in I} H_i$ there is an embedding of mentioned type as well, namely into the (direct or cartesian) product $\prod_{i \in I} G_i$.

And since each finite nilpotent group is a direct product of its p -subgroup, it is sufficient to prove Proposition 1 for finite p -groups.

Modifying Lemma 1 from [4] for the case of an arbitrary V we obtain for each power $m = p^k$ such a finite p -group P that its verbal subgroup $V(P)$ is of exponent

m and is from the center of P . Then, as it is shown in the proof of Theorem 3 in [4], the diagonal

$$D = \text{Diag}(H \text{ Wr } P) = \left\{ \prod_{t \in P} t^{-1}ht \mid h \in H \right\}$$

of wreath product $W = H \text{ Wr } P$ is contained in the verbal subgroup $V(W)$ of W . If H is a p -group W is a p -group, too. And there is an obvious embedding of H into W (onto D):

$$\nu : h \mapsto \prod_{t \in P} t^{-1}ht.$$

The base subgroup $H^{(P)}$ of W is normal in W and as a *nilpotent* group $H^{(P)}$ itself has $D \cong H$ as a subnormal subgroup. \square

Are there “many” non-nilpotent finite groups H for which there is a finite G such that $H \cong \tilde{H} \subseteq V(G)$? Are there “many” non-solvable finite groups H for which one cannot find a finite G with this property? The following statement shows something more.

PROPOSITION 2. 1. For each variety $\mathfrak{W} \neq \mathfrak{D}$ and an arbitrary non-trivial word set V there exists an infinite set of finite groups H such that $H \notin \mathfrak{W}$ and there are finite groups G and $\tilde{H} \triangleleft\triangleleft G$ such that $H \cong \tilde{H} \subseteq V(G)$;
2. For each variety $\mathfrak{W} \neq \mathfrak{D}$ there exists an infinite set of finite groups H such that $H \notin \mathfrak{W}$ and there is no finite group G with a subgroup $\tilde{H} \cong H$ such that $\tilde{H} \triangleleft\triangleleft G'$.

Proof. 1. We have $\mathfrak{V} = \text{var}(F_\infty/V(F_\infty)) \neq \mathfrak{D}$. Thus $\mathfrak{W} \cdot \mathfrak{V} \neq \mathfrak{D}$ (see in [13]). Since the set of all finite groups generates the variety \mathfrak{D} of all groups, there is a finite group $G \notin \mathfrak{W} \cdot \mathfrak{V}$. Thus $V(G) \notin \mathfrak{W}$ because G is an extension of $V(G)$ by $G/V(G)$. The finite group $H = V(G)$ satisfies the requirement of the first point. We can now repeat this step for the variety

$$\mathfrak{W}_1 = \text{var}(\mathfrak{W}, H) \neq \mathfrak{D}$$

and obtain a new finite group $H_1 \notin \mathfrak{W}_1$. The next finite groups H_2, H_3, \dots are found by induction.

2. Let N be a finite complete group that cannot be subnormally embedded into the derived group of some finite group. We can find such a group according to the result of Heineken and Lennox [5]. And let R be a finite group with $R \notin \mathfrak{W}$. We can find such a group, too, because $\mathfrak{D} \neq \mathfrak{W}$ and the set of finite groups generates \mathfrak{D} . The group $H = N \times R$ does not belong to \mathfrak{W} and does not have the mentioned subnormal embedding for N does not have the latter. Taking other values for R we will obtain infinitely many examples of group H . \square

Let us observe that since the set of all varieties (i. e. the set of all word sets) is of continuum cardinality [13] and since the set of all finite groups is countable, there exist finite groups that satisfy the statement 1 or 2 of previous proposition for \aleph word sets simultaneously.

We can repeat the argument of Proposition 2 to obtain finite groups with needed embedding “between” two varieties that can be very “near” one to other.

PROPOSITION 3. Let \mathfrak{U} be an arbitrary non-trivial locally finite variety and V be a non-trivial word set such that $\mathfrak{V} = \text{var}(F_\infty/V(F_\infty))$ is a locally finite variety, too.

Then for each subvariety \mathfrak{W} of \mathfrak{U} there is a finite group in the set $\mathfrak{U} \setminus \mathfrak{W}$ such that there exists a finite G with a subnormal subgroup $\tilde{H} \cong H$, $\tilde{H} \subseteq V(G)$.

Proof. Since $\mathfrak{U}, \mathfrak{W} \neq \mathfrak{D}$ and since multiplication of varieties is a monotonic operation we have

$$\mathfrak{W} \cdot \mathfrak{V} < \mathfrak{U} \cdot \mathfrak{V}$$

(see 21.21 in [13]). The product of two locally finite varieties is locally finite (O. Yu. Schmidt [16]). Thus the sets $\mathcal{F}_{\mathfrak{W} \cdot \mathfrak{V}}$ and $\mathcal{F}_{\mathfrak{U} \cdot \mathfrak{V}}$ of finite groups of products $\mathfrak{W} \cdot \mathfrak{V}$ and $\mathfrak{U} \cdot \mathfrak{V}$ are not equal too. Let $1 \neq Q \in \mathcal{F}_{\mathfrak{U}} \setminus \mathcal{F}_{\mathfrak{W}}$. Since $Q/V(Q) \in \mathfrak{V}$, necessarily $H = V(Q) \notin \mathfrak{W}$. \square

We can get similar facts for even finer layers between classes of groups [10], but it is already clear that the finite groups we are looking for are “many” and that the classes of such group are “unrestricted”. Let us set the question we are interested in as a *problem*:

Let H be a finite group and $V \subseteq F_\infty$ be a non-trivial word set. Find a criterion under which there exists a finite group G with a subnormal subgroup \tilde{H} such that the latter is isomorphic to H and lies in the verbal subgroup $V(G)$.

REFERENCES

- [1] R. Dark, *On subnormal embedding theorems of groups*, J. London Math Soc. 43 (1968), 387–390.
- [2] F. Galvin, *Embedding countable groups in 2-generator groups*. Amer. Math. Monthly 100 (1993), no. 6, 578–580.
- [3] P. Hall, *The Frattini subgroups of finitely generated groups*, Proc. London Math. Soc., (3) 11 (1961), 327–352.
- [4] H. Heineken, *Normal embeddings of p -groups into p -groups*, Proc. Edinburgh Math. Soc. 35 (1992), 309–314.
- [5] H. Heineken, J. C. Lennox, *The subnormal embedding of complete groups*, J. Algebra, 90 (1984), 435–445.
- [6] H. Heineken, V. H. Mikaelian, *On normal verbal embeddings of groups*, (to appear in the volume dedicated to 70th Birthday of A. I. Kostrikin, Plenum Pubs.).
- [7] H. Heineken, P. Soules, *The subnormal embedding of complete groups*, J. Austral. Math. Soc. (Series A) 45 (1988), 389–400.
- [8] G. Higman, B. Neumann, Hanna Neumann, *Embedding theorems for groups*, J. London Math. Soc. (3), 24 (1947), 247–254.
- [9] F. Levin, G. Rosenberger, *A class of SQ-universal groups*, Group theory (Singapore, 1987), 409–415, de Gruyter, Berlin-New York, 1989.
- [10] V. H. Mikaelian, *On identities of extensions of groups and extensions of finite groups*, Dokladi National’noii Akademii Nauk Armenii, 95 (1995), 72–75. (Russian)
- [11] B. H. Neumann, *Embedding theorems for groups*. Nieuw Arch. Wisk. (3) 16 (1968), 73–78.
- [12] B. Neumann, Hanna Neumann, *Embedding theorems for groups*, J. London Math. Soc., 34 (1959), 465–479.
- [13] Hanna Neumann, *Varieties of Groups*, Springer Verlag, Berlin (1968)

- [14] A. Yu. Ol'shanskii, Efficient embeddings of countable groups. Vestnik Moskov. Univ. Ser. I Mat. Mekh., 105, no. 2 (1989), 28–34. (Russian)
- [15] A. Yu. Ol'shanskii, Embedding of countable periodic groups in simple 2-generator periodic groups, Ukrain. Mat. Zh. 43 (1991), no. 7-8, 980–986 (Russian); translation in Ukrainian Math. J. 43 (1991), no. 7-8, 914–919 (1992).
- [16] D. J. S. Robinson, A Course in the Theory of Groups, second edition, Springer-Verlag, New York, Berlin, Heidelberg (1996).
- [17] J. S. Wilson, P. A. Zalesskii, An embedding theorem for certain residually finite groups, Arch. Math. (Basel) 67 (1996), no. 3, 177–182.

Mathematisches Institut
Universität Würzburg
Am Hubland
D-97074 Würzburg
Germany

e-mail: mikaelian@e-math.ams.org
vahagn.mikaelian@mail.uni-wuerzburg.de
<http://www.crosswinds.net/~mikaelian/home.html>