

**COMBINATORICS OF GEOMETRICALLY DISTRIBUTED  
RANDOM VARIABLES:  
VALUE AND POSITION OF LARGE LEFT-TO-RIGHT MAXIMA**

HELMUT PRODINGER

ABSTRACT. For words of length  $n$ , generated by independent geometric random variables, we consider the average value and the average position of the  $r$ th left-to-right maximum *counted from the right*, for fixed  $r$  and  $n \rightarrow \infty$ . This complements previous research [5] where the analogous questions were considered for the  $r$ th left-to-right maximum *counted from the left*.

1. INTRODUCTION

In this paper we study sequences with respect to the *left-to-right maxima* of the elements of the sequence.

This paper is a companion paper to [5], and these two papers should be read together.

In [6] the number of left-to-right maxima was investigated in the model of *words* (strings)  $a_1 \dots a_n$ , where the letters  $a_i \in \mathbb{N}$  are independently generated according to the geometric distribution with

$$\mathbb{P}\{X = k\} = pq^{k-1}, \quad \text{where } p + q = 1. \quad (1)$$

(An element is a left-to-right maximum if it is larger than all the elements to the left. The version where ‘larger’ is replaced by ‘larger or equal’ would be also of interest but is not considered here.)

In [5] we considered the two parameters ‘value’ and ‘position’ of the  $r$ th left-to-right maximum for geometric random variables and obtained the asymptotic formulæ  $\frac{r}{p}$  and  $\frac{1}{(r-1)!} \left(\frac{r}{q} \log_Q n\right)^{r-1}$ .

These results cover the instances of small left-to-right maxima; therefore we investigate here the instance of *large* left-to-right maxima. We say that the last left-to-right maximum is the instance  $r = 1$ , the previous left-to-right maximum is the instance  $r = 2$ , and so on.

As can be seen in the forthcoming book [4], in the limit  $q \rightarrow 1$  the model becomes the model of *random permutations*. The concept of ‘value’ does not carry over, but the concept of ‘position’ does.

The largest element in a random permutation of  $n$  elements (which is  $n$ ) is expected to be in the middle. The second largest is then expected to be in the middle of the first half, and so on. In general, the average position should be about  $\frac{n}{2^r}$ .

We find it useful to use the additional notations  $Q := q^{-1}$ ,  $L := \log Q$ , and  $\llbracket i \rrbracket := 1 - (1 - q^i)z$ .

In [6], compare also [8], the average value of the *height* appeared; it is asymptotic to  $\log_Q n$ , which is also the instance  $r = 1$  of the average value of left-to-right maxima, counted from the right. Intuitively, for fixed  $r$ , the same asymptotic formula is to be expected, the dependency of  $r$  only appearing in the lower order terms.

It should be noted that not all random strings of length  $n$  have  $r$  left-to-right maxima. However, as explained already in [5], this happens with low probability and can be ignored.

## 2. THE AVERAGE VALUE OF THE $r$ TH LEFT-TO-RIGHT MAXIMUM FROM THE RIGHT

The following expression for the average value of the  $r$ th left-to-right maximum from the right is obtained as in [5]; just observe that we assume that it is  $h$ , and that  $r - 1$  more left-to-right maxima follow to the right, whereas to the left everything is smaller than  $h$ :

$$\begin{aligned}
E_n^{(r)} &= [z^n] \sum_{1 \leq h < i_1 < \dots < i_{r-1}} h \frac{1}{\llbracket h-1 \rrbracket} \frac{zpq^{h-1}}{\llbracket h \rrbracket} \frac{zpq^{i_1-1}}{\llbracket i_1 \rrbracket} \cdots \frac{zpq^{i_{r-1}-1}}{\llbracket i_{r-1} \rrbracket} \\
&= \left(\frac{p}{q}\right)^r [z^n] \sum_{1 \leq h < i_1 < \dots < i_{r-1}} h \frac{1}{\llbracket h-1 \rrbracket} \frac{zq^h q^{i_1} z}{\llbracket h \rrbracket \llbracket i_1 \rrbracket} \cdots \frac{q^{i_{r-1}} z}{\llbracket i_{r-1} \rrbracket} \\
&= (-1)^r \left(\frac{p}{q}\right)^r (-1)^n [w^n] (1-w)^n \times \\
&\quad \times \sum_{1 \leq h < i_1 < \dots < i_{r-1}} \frac{1}{1-wq^{h-1}} \frac{wq^h h}{1-q^h w} \frac{q^{i_1} w \dots q^{i_{r-1}} w}{(1-q^{i_1} w) \dots (1-q^{i_{r-1}} w)} \\
&= (-1)^r \left(\frac{p}{q}\right)^r \sum_{k=r}^n \binom{n}{k} (-1)^k \times \\
&\quad \times [w^k] \sum_{1 \leq h < i_1 < \dots < i_{r-1}} \frac{hwq^h}{(1-wq^{h-1})(1-wq^h)} \frac{q^{i_1} w \dots q^{i_{r-1}} w}{(1-q^{i_1} w) \dots (1-q^{i_{r-1}} w)}.
\end{aligned}$$

For the computation of the inner sum we need the following formula, which can be computed by elementary manipulations. For  $n \geq m$  we have

$$[w^n] \sum_{h \geq 1} \frac{hwq^h}{(1-wq^{h-1})(1-wq^h)} (wq^h)^m = \frac{q}{p} \frac{q^m - q^n}{(1-q^n)^2}.$$

Using this and the principle that

$$[w^n] \sum_i a_i f(b_i w) = \sum_i a_i b_i^n \cdot [w^n] f(w)$$

we find that

$$\begin{aligned} [w^k] & \sum_{1 \leq h < i_1 < \dots < i_{r-1}} \frac{hwq^h}{(1-wq^{h-1})(1-wq^h)} \frac{q^{i_1}w \dots q^{i_{r-1}}w}{(1-q^{i_1}w) \dots (1-q^{i_{r-1}}w)} \\ & = \frac{q}{p} \frac{1}{(1-q^k)^2} \sum_{0 < l_1 < \dots < l_{r-1} < k} \frac{1}{Q^{l_1} - 1} \dots \frac{1}{Q^{l_{r-1}} - 1} (q^{l_{r-1}} - q^k). \end{aligned}$$

It is worthwhile to note the instance  $r = 1$ , which is the *maximum*:

$$E_n^{(1)} = \sum_{k=1}^n \binom{n}{k} (-1)^{k-1} \frac{1}{1-q^k}.$$

This checks with previous results, compare [6, 8], but it seems to be a folklore result.

Now define the function

$$\phi_r(k) := \sum_{0 < l_1 < \dots < l_{r-1} < k} \frac{1}{Q^{l_1} - 1} \dots \frac{1}{Q^{l_{r-1}} - 1} (q^{l_{r-1}} - q^k).$$

With this notation, we have

$$E_n^{(r)} = (-1)^r \left(\frac{p}{q}\right)^{r-1} \sum_{k=r}^n \binom{n}{k} (-1)^k \frac{\phi_r(k)}{(1-q^k)^2}.$$

Since our quantities come out as alternating sums, the appropriate treatment of them is *Rice's method* which is surveyed in [3]; the key point is the following Lemma.

**Lemma 1.** *Let  $\mathcal{C}$  be a curve surrounding the points  $1, 2, \dots, n$  in the complex plane and let  $f(z)$  be analytic inside  $\mathcal{C}$ . Then*

$$\sum_{k=1}^n \binom{n}{k} (-1)^k f(k) = -\frac{1}{2\pi i} \int_{\mathcal{C}} [n; z] f(z) dz,$$

where

$$[n; z] = \frac{(-1)^{n-1} n!}{z(z-1) \dots (z-n)} = \frac{\Gamma(n+1)\Gamma(-z)}{\Gamma(n+1-z)}.$$

Extending the contour of integration it turns out that under suitable growth conditions on  $f(z)$  (compare [3]) the asymptotic expansion of the alternating sum is given by

$$\sum \text{Res}([n; z]f(z)) + \text{smaller order terms}$$

where the sum is taken over all poles  $z_0$  different from  $1, \dots, n$ . Poles that lie more to the left lead to smaller terms in the asymptotic expansion.

The range  $1, \dots, n$  for the summation is not sacred; if we sum, for example, over  $k = 2, \dots, n$ , the contour must encircle  $2, \dots, n$ , etc.  $\square$

Therefore a key issue is to extend sequences like  $\phi_r(k)$  to the complex plane, in other words, give a meaningful definition of  $\phi_r(z)$ , for  $z \in \mathbb{C}$ .

Note that

$$\begin{aligned} \sum_{0 < i_1 < \dots < i_s < j} \frac{1}{Q^{i_1} - 1} \cdots \frac{1}{Q^{i_s} - 1} &= [x^s] \prod_{i=1}^{j-1} \left( 1 + \frac{x}{Q^i - 1} \right) \\ &= [x^s] \prod_{i=1}^{j-1} \frac{Q^i - 1 + x}{Q^i - 1} \\ &= [x^s] \prod_{i=1}^{j-1} \frac{1 - (1-x)q^i}{1 - q^i} \\ &= [x^s] \frac{((1-x)q; q)_{j-1}}{(q; q)_{j-1}}. \end{aligned}$$

In this computation we use the standard notation for  $q$ -factorials,

$$(a; q)_n := (1-a)(1-aq) \dots (1-aq^{n-1}), \quad (a; q)_\infty := (1-a)(1-aq) \dots,$$

see e. g. [1]. In the following we need Cauchy's theorem ( $q$ -binomial theorem), viz.

$$\sum_{n \geq 0} \frac{(a; q)_n}{(q; q)_n} t^n = \frac{(at; q)_\infty}{(t; q)_\infty}.$$

Then

$$\begin{aligned} \phi_r(k) &= [x^{r-2}] \sum_{1 \leq l \leq k} \frac{((1-x)q; q)_{l-1} q^l - q^k}{(q; q)_{l-1} Q^l - 1} \\ &= [x^{r-2}] \sum_{1 \leq l \leq k} \frac{((1-x)q; q)_{l-1} (q^{2l} - q^{k+l})}{(q; q)_l} \\ &= [x^{r-2}] [t^k] \frac{1}{1-t} \sum_{l \geq 1} \frac{((1-x)q; q)_{l-1} (q^{2l} - q^{k+l}) t^l}{(q; q)_l} \\ &= [x^{r-2}] [t^k] \frac{1}{1-t} \frac{1}{x} \left( - (1 - q^k) + \sum_{l \geq 0} \frac{(1-x; q)_l}{(q; q)_l} (q^{2l} - q^{k+l}) t^l \right) \\ &= [x^{r-1}] [t^k] \frac{1}{1-t} \left( - (1 - q^k) + \frac{((1-x)q^2t; q)_\infty}{(q^2t; q)_\infty} - q^k \frac{((1-x)qt; q)_\infty}{(qt; q)_\infty} \right) \\ &= -[x^{r-1}] [t^k] \frac{1}{1-t} \left( (1 - q^k) + (q^k - 1) \frac{((1-x)qt; q)_\infty}{(qt; q)_\infty} + \frac{((1-x)q^2t; q)_\infty}{(qt; q)_\infty} xqt \right) \\ &= -\delta_{r,1} (1 - q^k) - [x^{r-1}] [t^k] \left( (q^k - 1) \frac{((1-x)qt; q)_\infty}{(t; q)_\infty} + \frac{((1-x)q^2t; q)_\infty}{(t; q)_\infty} xqt \right) \\ &= -\delta_{r,1} (1 - q^k) - [x^{r-1}] \left( (q^k - 1) \frac{((1-x)q; q)_k}{(q; q)_k} + xq \frac{((1-x)q^2; q)_{k-1}}{(q; q)_{k-1}} \right) \end{aligned}$$

$$= -\delta_{r1}(1 - q^k) - [x^{r-1}] \left( (q^k - 1) \frac{((1-x)q; q)_k}{(q; q)_k} + \frac{xq(1 - q^k)}{1 - (1-x)q} \frac{((1-x)q^2; q)_k}{(q; q)_k} \right).$$

Hence

$$\frac{\phi_r(k)}{1 - q^k} = -\delta_{r1} + [x^{r-1}] \left( \frac{((1-x)q; q)_k}{(q; q)_k} - \frac{xq}{1 - (1-x)q} \frac{((1-x)q^2; q)_k}{(q; q)_k} \right)$$

and

$$\begin{aligned} \left. \frac{\phi_r(k)}{1 - q^k} \right|_{k=0} &= -\delta_{r1} + [x^{r-1}] \left( 1 - \frac{xq}{1 - (1-x)q} \right) \\ &= -[x^{r-1}] \frac{xq}{1 - (1-x)q} \\ &= -\frac{q}{p} [x^{r-2}] \frac{1}{1 + \frac{xq}{p}} \\ &= (-1)^{r-1} \left( \frac{q}{p} \right)^{r-1}. \end{aligned}$$

To get lower order terms, we first note that

$$\frac{((1-x)q; q)_z}{(q; q)_z} \sim 1 + L \sum_{j \geq 1} \frac{xq^j}{(1 - q^j)(1 - (1-x)q^j)} \cdot z \quad (z \rightarrow 0)$$

and (for  $r \geq 2$ )

$$[x^{r-1}] L \sum_{j \geq 1} \frac{xq^j}{(1 - q^j)(1 - (1-x)q^j)} \cdot z = L(-1)^r \sum_{j \geq 1} \frac{q^{j(r-1)}}{(1 - q^j)^r} \cdot z.$$

Also note that

$$\begin{aligned} \frac{xq}{1 - (1-x)q} \frac{((1-x)q^2; q)_z}{(q; q)_z} &\sim \frac{xq}{1 - (1-x)q} + \frac{xq}{1 - (1-x)q} L \sum_{j \geq 1} \frac{xq^{j+1}}{(1 - q^j)(1 - (1-x)q^{j+1})} \cdot z \\ &= \frac{xq}{1 - (1-x)q} + xqL \sum_{j \geq 1} \frac{q^j}{(1 - q^j)^2} \left[ -\frac{1}{1 + \frac{qx}{p}} + \frac{1}{1 + \frac{q^{j+1}x}{1 - q^{j+1}}} \right] \cdot z \end{aligned}$$

and (for  $r \geq 2$ )

$$\begin{aligned} [x^{r-1}] xqL \sum_{j \geq 1} \frac{q^j}{(1 - q^j)^2} \left[ -\frac{1}{1 + \frac{qx}{p}} + \frac{1}{1 + \frac{q^{j+1}x}{1 - q^{j+1}}} \right] \cdot z \\ = qL(-1)^{r-1} \sum_{j \geq 1} \frac{q^j}{(1 - q^j)^2} \left[ \left( \frac{q}{p} \right)^{r-2} - \left( \frac{q^{j+1}}{1 - q^{j+1}} \right)^{r-2} \right] \cdot z. \end{aligned}$$

Hence

$$\frac{\phi_r(z)}{1-q^z} \sim (-1)^{r-1} \left(\frac{q}{p}\right)^{r-1} + \beta_r \cdot z \quad (z \rightarrow 0),$$

with

$$\beta_r = L(-1)^r \sum_{j \geq 1} \frac{q^{j(r-1)}}{(1-q^j)^r} + qL(-1)^r \sum_{j \geq 1} \frac{q^j}{(1-q^j)^2} \left[ \left(\frac{q}{p}\right)^{r-2} - \left(\frac{q^{j+1}}{1-q^{j+1}}\right)^{r-2} \right].$$

Computing the residue of

$$\frac{1}{z} \frac{1}{1-q^z} \frac{\phi_r(z)}{1-q^z}$$

at  $z = 0$  eventually leads to

**Theorem 1.** *The average value of the  $r$ th left-to-right maximum from the right is given by*

$$E_n^{(r)} = \log_Q n + \sigma_r + \delta_r(\log_Q n) + o(1),$$

where  $\delta_r(x)$  is a small periodic function of  $\log_Q n$  (originating from the poles at  $2k\pi i / \log Q$ ,  $k \in \mathbb{Z} \setminus \{0\}$ ). The constant  $\sigma_r$  is given by

$$\sigma_r = \frac{\gamma}{L} + \frac{1}{2} - p^{r-1} \sum_{j \geq 1} \frac{q^{(j-1)(r-1)}}{(1-q^j)^r} - p \sum_{j \geq 1} \frac{q^j}{(1-q^j)^2} + p^{r-1} \sum_{j \geq 1} \frac{q^{j(r-1)}}{(1-q^j)^2(1-q^{j+1})^{r-2}}.$$

This holds for  $r = 1$  as well, when the 3 sums combined evaluate to 0.

For interest, let us note the following limit:

$$\lim_{q \rightarrow 1} (1-q)\sigma_r = \sum_{k=3}^{r-3} (r-1-k)\zeta(k) + (r-1)\zeta(2) + \zeta(r-2) - \zeta(r) - \frac{r(r-3)}{2}.$$

### 3. THE AVERAGE POSITION OF THE $r$ TH LEFT-TO-RIGHT MAXIMUM FROM THE RIGHT

We consider the generating function

$$\sum_{h \geq 1} \frac{zpq^{h-1}}{(1-zv(1-q^{h-1}))[[h]]} \sum_{h < i_1 < \dots < i_{r-1}} \frac{zpq^{i_1-1}}{[[i_1]]} \dots \frac{zpq^{i_{r-1}-1}}{[[i_{r-1}]]},$$

where  $v$  marks the elements to the left of the  $r$ th left-to-right maximum from the right. The actual position is then one more than the number of  $v$ 's, but that is overshadowed by the error term anyway. To get the average, we must differentiate this with respect to  $v$ , and then replace  $v$  by 1. The result is

$$\begin{aligned}
 & \sum_{h \geq 1} \frac{zpq^{h-1}z(1-q^{h-1})}{[[h-1]]^2[[h]]} \sum_{h < i_1 < \dots < i_{r-1}} \frac{zpq^{i_1-1}}{[[i_1]]} \cdots \frac{zpq^{i_{r-1}-1}}{[[i_{r-1}]]} \\
 &= \left(\frac{p}{q}\right)^r \sum_{h \geq 1} \frac{zq^h z(1-q^{h-1})}{[[h-1]]^2[[h]]} \sum_{h < i_1 < \dots < i_{r-1}} \frac{zq^{i_1}}{[[i_1]]} \cdots \frac{zq^{i_{r-1}}}{[[i_{r-1}]]} \\
 &= (-1)^{r-1} \left(\frac{p}{q}\right)^r (1-w) \sum_{h \geq 1} \frac{wq^h w(1-q^{h-1})}{(1-wq^{h-1})^2(1-wq^h)} \sum_{h < i_1 < \dots < i_{r-1}} \frac{wq^{i_1}}{1-wq^{i_1}} \cdots \frac{wq^{i_{r-1}}}{1-wq^{i_{r-1}}}.
 \end{aligned}$$

Therefore

$$\begin{aligned}
 E_n^{(r)} &= (-1)^n [w^n] (-1)^{r-1} \left(\frac{p}{q}\right)^r (1-w)^n \sum_{h \geq 1} \frac{wq^h w(1-q^{h-1})}{(1-wq^{h-1})^2(1-wq^h)} \times \\
 &\quad \times \sum_{h < i_1 < \dots < i_{r-1}} \frac{wq^{i_1}}{1-wq^{i_1}} \cdots \frac{wq^{i_{r-1}}}{1-wq^{i_{r-1}}} \\
 &= (-1)^{r-1} \left(\frac{p}{q}\right)^r \sum_{k=r+1}^n \binom{n}{k} (-1)^k \times \\
 &\quad \times [w^k] \sum_{1 \leq h < i_1 < \dots < i_{r-1}} \frac{wq^h w(1-q^{h-1})}{(1-wq^{h-1})^2(1-wq^h)} \frac{wq^{i_1}}{1-wq^{i_1}} \cdots \frac{wq^{i_{r-1}}}{1-wq^{i_{r-1}}}.
 \end{aligned}$$

For the computation of the inner sum we need

$$\begin{aligned}
 [w^k] \sum_{h \geq 1} \frac{(wq^h)^M}{(1-wq^{h-1})^2(1-wq^h)} \\
 = -\frac{q^{M+1}}{(1-q)^2} \frac{1}{1-q^k} + \frac{q^M}{1-q} \frac{1}{1-q^k} (k+1-M) + \frac{q^2}{(1-q)^2} \frac{1}{Q^k-1}
 \end{aligned}$$

and also (as a consequence)

$$\begin{aligned}
 [w^k] \sum_{h \geq 1} \frac{(wq^h)^M w(1-q^{h-1})}{(1-wq^{h-1})^2(1-wq^h)} \\
 = -\frac{q^{m+2}}{(1-q)^2} \frac{1}{1-q^{k-1}} + \frac{q^{m+1}}{1-q} \frac{1}{1-q^{k-1}} (k-m-1) + \frac{q^2}{(1-q)^2} \frac{1}{Q^{k-1}-1} \\
 + \frac{q^{m+2}}{(1-q)^2} \frac{1}{1-q^k} - \frac{q^{m+1}}{1-q} \frac{1}{1-q^k} (k-m-1) - \frac{q^2}{(1-q)^2} \frac{1}{Q^k-1}.
 \end{aligned}$$

The righthand side will be denoted by  $\tau(k, m)$  and can alternatively be written as

$$\begin{aligned}
 \tau(k, m) &= -\frac{q^{m+2}}{p^2} \frac{1}{Q^{k-1}-1} + \frac{q^{m+1}}{p} \frac{1}{Q^{k-1}-1} (k-m-1) + \frac{q^2}{p^2} \frac{1}{Q^{k-1}-1} \\
 &\quad + \frac{q^{m+2}}{p^2} \frac{1}{Q^k-1} - \frac{q^{m+1}}{p} \frac{1}{Q^k-1} (k-m-1) - \frac{q}{p^2} \frac{1}{Q^k-1}.
 \end{aligned}$$

Consequently

$$\begin{aligned} [w^k] \sum_{h \geq 1} \frac{wq^h w(1 - q^{h-1})}{(1 - wq^{h-1})^2 (1 - wq^h)} &= \sum_{h < i_1 < \dots < i_{r-1}} \frac{wq^{i_1}}{1 - wq^{i_1}} \cdots \frac{wq^{i_{r-1}}}{1 - wq^{i_{r-1}}} \\ &= \sum_{0 < l_1 < \dots < l_{r-1} < k-1} \frac{1}{Q^{l_1} - 1} \cdots \frac{1}{Q^{l_{r-1}} - 1} \tau(k, l_{r-1}). \end{aligned}$$

It is instructive to start with the case  $r = 1$ :

$$\begin{aligned} E_n^{(1)} &= \frac{p}{q} \sum_{k=2}^n \binom{n}{k} (-1)^k [w^k] \sum_{h \geq 1} \frac{wq^h w(1 - q^{h-1})}{(1 - wq^{h-1})^2 (1 - wq^h)} \\ &= \frac{p}{q} \sum_{k=2}^n \binom{n}{k} (-1)^k \left[ \frac{q}{p} \frac{k-1}{Q^{k-1} - 1} - \frac{q}{p} \frac{k}{Q^k - 1} \right] \\ &= \sum_{k=2}^n \binom{n}{k} (-1)^k \left[ \frac{k-1}{Q^{k-1} - 1} - \frac{k}{Q^k - 1} \right]. \end{aligned}$$

When applying Rice's formula, it turns out that the dominant singularity is at  $z = 1$ . This matches the intuition, since the position should be roughly at  $\approx \text{const} \cdot n$ .

Indeed, apart from fluctuations (coming from complex poles at  $z = 1 + 2k\pi i / \log Q$ , with  $k \in \mathbb{Z} \setminus \{0\}$ ) we get  $E_n^{(1)} \sim n(\frac{1}{L} - \frac{q}{p})$ .

Formally, for  $q \rightarrow 1$ , we get  $\frac{n}{2}$ , as it should.

Now, for the general case, define the function

$$\psi_r(k) = \sum_{0 < l_1 < \dots < l_{r-1} < k-1} \frac{1}{Q^{l_1} - 1} \cdots \frac{1}{Q^{l_{r-1}} - 1} \tau(k, l_{r-1}).$$

We need the value  $\psi_r(1)$ , because of the result (apart from fluctuations)

$$E_n^{(r)} \sim (-1)^{r-1} \left(\frac{p}{q}\right)^r \psi_r(1) n. \quad (2)$$

We want to study the constants in more detail. Assume  $r = 2$ . Since  $\tau(z, z-1) = 0$ , the adequate extension is

$$\psi_2(z) = \sum_{l \geq 1} \frac{1}{Q^l - 1} \tau(z, l) - \sum_{l \geq 1} \frac{1}{Q^{l+z-1} - 1} \tau(z, l+z-1).$$



An elementary computation yields

$$\begin{aligned}\tau(z, l) &\sim \frac{-(l-1)q^{l+2} + lq^{l+1} - q^2}{Lp^2} \frac{1}{z-1} \\ &\quad + \frac{q^{l+1}}{pL} + \frac{1}{2p^3} [-3q^2 + q^3 + lq^{l+1} + q^{l+2} - (l-1)q^{l+3}] + \dots, \\ \tau(z, z+l-1) &\sim \frac{-(l-1)q^{l+2} + lq^{l+1} - q^2}{Lp^2} \frac{1}{z-1} \\ &\quad + \frac{1}{2p^3} [-3q^2 + q^3 + 3lq^{l+1} - (4l-3)q^{l+2} + lq^{l+3}], \\ \frac{1}{Q^{l+z-1} - 1} &\sim \frac{1}{Q^l - 1} + \frac{LQ^l}{(Q^l - 1)^2} (z-1) + \dots.\end{aligned}$$

Hence

$$\begin{aligned}\psi_2(1) &= \frac{1}{pL} \sum_{l \geq 1} \frac{q^{l+1}}{Q^l - 1} + \frac{1}{p^3} \sum_{l \geq 1} \frac{1}{Q^l - 1} [-lq^{l+1} + (2l-1)q^{l+2} - (l-1)q^{l+3}] \\ &\quad + \frac{1}{p^2} \sum_{l \geq 1} \frac{Q^l}{(Q^l - 1)^2} [(l-1)q^{l+2} - lq^{l+1} + q^2] \\ &= \frac{q}{pL} \sum_{l \geq 1} \frac{1}{Q^l - 1} - \frac{q^2}{p^2L} + \frac{q^2(q+1)}{p^3} - \frac{q}{p} \sum_{l \geq 1} \frac{lQ^l}{(Q^l - 1)^2}\end{aligned}$$

and thus

$$-\frac{p^2}{q^2} \psi_2(1) = -\frac{p}{qL} \sum_{l \geq 1} \frac{1}{Q^l - 1} + \frac{p}{q} \sum_{l \geq 1} \frac{lQ^l}{(Q^l - 1)^2} + \frac{1}{L} - \frac{1}{p} - \frac{q}{p}.$$

It is our aim to show now that for  $q \rightarrow 1$  this quantity approaches  $\frac{1}{4}$ .

For this, we write  $q = e^{-t}$  and study the behaviour at  $t \rightarrow 0$ . The *Mellin transform* allows us to write

$$\begin{aligned}\sum_{l \geq 1} \frac{1}{Q^l - 1} &= \frac{1}{2\pi i} \int_{3-i\infty}^{3+i\infty} \zeta^2(s) \Gamma(s) t^{-s} ds, \\ \sum_{l \geq 1} \frac{lQ^l}{(Q^l - 1)^2} &= \frac{1}{2\pi i} \int_{3-i\infty}^{3+i\infty} \zeta^2(s-1) \Gamma(s) t^{-s} ds.\end{aligned}$$

Shifting the line of integration and collecting residues gives us the following asymptotic expansions (see [2] for a friendly explanation of the subject):

$$\sum_{l \geq 1} \frac{1}{Q^l - 1} = -\frac{\log t}{t} + \frac{\gamma}{t} + \frac{1}{4} + \mathcal{O}(t),$$

$$\sum_{l \geq 1} \frac{lQ^l}{(Q^l - 1)^2} = -\frac{\log t}{t^2} + \frac{1 + \gamma}{t^2} + \mathcal{O}(1).$$

Using that, we find

$$-\frac{p^2}{q^2} \psi_2(1) = \frac{1}{4} - \frac{t}{9} + \mathcal{O}(t^2) = \frac{1}{4} + \frac{1}{9} \log q + \mathcal{O}(\log^2 q).$$

In the general case, the suitable extension is

$$\psi_r(z) = \sum_{l \geq 1} g_{r-2}(l) \frac{\tau(z, l)}{Q^l - 1} - \sum_{l \geq 1} g_{r-2}(l + z - 1) \frac{\tau(z, l + z - 1)}{Q^{l+z-1} - 1},$$

with

$$g_s(l) = \sum_{1 \leq i_1 < \dots < i_s < l} \frac{1}{Q^{i_1} - 1} \cdots \frac{1}{Q^{i_s} - 1},$$

and the suitable extension of it, which is, at least theoretically, no problem, since we can, as already explained in [5], use powersum symmetric functions (see [7]) or proceed as in the last section.

For instance,

$$g_1(z) = \sum_{i \geq 1} \frac{1}{Q^i - 1} - \sum_{i \geq 1} \frac{1}{Q^{i+z-1} - 1}$$

and

$$g_1(l + z - 1) = g_1(l) + L \sum_{i \geq l} \frac{Q^i}{(Q^i - 1)^2} (z - 1) + \dots,$$

as  $z \rightarrow 1$ .

Using this, one could write an explicit formula for  $\psi_3(1)$  which is already extremely messy.

Since already the quantity  $\psi_2(1)$  cannot be simplified, there is no hope for closed forms here.

It is possible to make the translation to permutations (the limit  $q \rightarrow 1$ ) at an earlier stage.

Note that

$$\lim_{q \rightarrow 1} (1 - q) \tau(k, m) = \frac{1}{2} - \frac{(m+1)m}{2k} + \frac{m(m-1)}{2(k-1)}.$$

Hence the constant in (2) translates into

$$\lambda_r := (-1)^{r-1} \sum_{0 < l_1 < \dots < l_{r-1} \leq k-1} \frac{1}{l_1 \dots l_{r-1}} \left[ \frac{1}{2} - \frac{(l_{r-1} + 1)l_{r-1}}{2k} + \frac{l_{r-1}(l_{r-1} - 1)}{2(k-1)} \right] \Big|_{k=1}.$$

Observe that

$$\sum_{0 < l_1 < \dots < l_{r-2} < l} \frac{1}{l_1 \dots l_{r-2}} = [x^{r-2}] \prod_{i=1}^{l-1} \left( 1 + \frac{x}{i} \right) = [x^{r-2}] \frac{\Gamma(l+x)}{\Gamma(1+x)\Gamma(l)}.$$

Hence

$$\begin{aligned} \lambda_r &= -\frac{1}{2} [x^{r-2}] \sum_{1 \leq l \leq k-1} \frac{\Gamma(l-x)}{\Gamma(1-x)\Gamma(l)} \left[ \frac{1}{l} - \frac{l+1}{k} + \frac{l-1}{k-1} \right] \Big|_{k=1} \\ &= [x^{r-2}] \frac{1}{2x} \left[ \frac{2\Gamma(k+1-x)}{\Gamma(k+1)\Gamma(3-x)} - 1 \right] \Big|_{k=1} \\ &= [x^{r-2}] \frac{1}{4} \frac{1}{1 - \frac{x}{2}} = 2^{-r}, \end{aligned}$$

as it should be.

In this permutation case we can even do better and compute the next term (of order 1), because of

$$E_n^{(r)} \sim (-1)^{r-1} \left( \frac{p}{q} \right)^r \left( \psi_r(1)n - \psi_r(0) \right).$$

All we have to do is to compute the following quantity:

$$\begin{aligned} \mu_r &= (-1)^{r-1} \sum_{0 < l_1 < \dots < l_{r-1} \leq k-1} \frac{1}{l_1 \dots l_{r-1}} \left[ \frac{1}{2} - \frac{(l_{r-1} + 1)l_{r-1}}{2k} + \frac{l_{r-1}(l_{r-1} - 1)}{2(k-1)} \right] \Big|_{k=0} \\ &= [x^{r-2}] \left( \frac{1}{1-x} - \frac{1}{4} \frac{1}{1 - \frac{x}{2}} \right) = 1 - 2^{-r}. \end{aligned}$$

Since there are no more poles to the left and a technical growth condition (see [3]) is satisfied, there are no more terms, and we have the exact formula for the average position of the  $r$ th left-to-right maximum, counted from the right (remember that we counted one less than the position!):

$$E_n^{(r)} + 1 = \frac{n+1}{2^r}.$$

**Theorem 2.** *The average position of the  $r$ -th left-to-right maximum from the right is given by*

$$E_n^{(r)} = (-1)^r \left( \frac{p}{q} \right)^{r-1} \psi_r(1)n + \varpi_r(\log_Q n) + o(n),$$

where the constants are defined in the text, and  $\varpi_r(x)$  is a small periodic function, originating from the complex poles  $1 + 2k\pi i / \log Q$ .

In the limit  $q \rightarrow 1$  (permutations), this simplifies to

$$E_n^{(r)} = \frac{n}{2^r} + o(n).$$

#### REFERENCES

- [1] G. Andrews. *The Theory of Partitions*, volume 2 of *Encyclopedia of Mathematics and its Applications*. Addison–Wesley, 1976.
- [2] P. Flajolet, X. Gourdon, and P. Dumas. Mellin transforms and asymptotics: Harmonic sums. *Theoretical Computer Science*, 144:3–58, 1995.
- [3] P. Flajolet and R. Sedgewick. Mellin transforms and asymptotics: Finite differences and Rice’s integrals. *Theoretical Computer Science*, 144:101–124, 1995.
- [4] P. Flajolet and R. Sedgewick. *Analytic Combinatorics*. Cambridge University Press, Cambridge, 2005.
- [5] A. Knopfmacher and H. Prodinger. Combinatorics of geometrically distributed random variables: Value and position of the  $r$ th left-to-right maximum. *Discrete Mathematics*, 2000.
- [6] H. Prodinger. Combinatorics of geometrically distributed random variables: Left-to-right maxima. *Discrete Mathematics*, 153:253–270, 1996.
- [7] R. Stanley. *Enumerative combinatorics. Vol. 2*. Cambridge University Press, Cambridge, 1999.
- [8] W. Szpankowski and V. Rego. Yet another application of a binomial recurrence. Order statistics. *Computing*, 43(4):401–410, 1990.

HELMUT PRODINGER, THE JOHN KNOPFMACHER CENTRE FOR APPLICABLE ANALYSIS AND NUMBER THEORY, DEPARTMENT OF MATHEMATICS, UNIVERSITY OF THE WITWATERSRAND, P. O. WITS, 2050 JOHANNESBURG, SOUTH AFRICA, EMAIL: [helmut@gauss.cam.wits.ac.za](mailto:helmut@gauss.cam.wits.ac.za),  
 HOMEPAGE: <http://www.wits.ac.za/helmut/index.htm>.