Harmonic maps between generalized Lagrange spaces

Mircea Neagu

Abstract

In Section 1 the author defines the notion of harmonic map between generalized Lagrange spaces. Section 2 analyses the particular case when the generalized Lagrange spaces are Lagrange spaces of electrodynamics. In Section 3 it is proved that for certain systems of differential or partial differential equations, the solutions are harmonic maps between certain generalized Lagrange spaces, in the sense of Section 1. Section 4 describes the main properties of the generalized Lagrange spaces constructed in Section 3.

Mathematics Subject Classification: 53C60, 49N45, 35R30

Key words: generalized Lagrange spaces, harmonic maps, geodesics, differential equations, partial differential equations.

1 Introduction

Let $(M^m, g_{\alpha\beta})$ and (N^n, h_{ij}) be two generalized Lagrange spaces, where m, respectively n, is the dimension of M, respectively N. The manifold M, respectively N, is coordinated by (a^{α}) , respectively (x^i) . On $M \times N$, the first m coordinates are indexed by $\alpha, \beta, \gamma, \ldots$ and the last n coordinates are indexed by i, j, k, \ldots . The fundamental metric tensors are expressed locally by

i) $g_{\alpha\beta} = g_{\alpha\beta}(a,b), \forall \alpha, \beta = \overline{1,m}$, where $(a,b) = (a^{\mu}, b^{\mu})$ are adapted coordinates on TM.

ii) $h_{ij} = h_{ij}(x, y), \ \forall \ i, j = \overline{1, n}$, where $(x, y) = (x^k, y^k)$ are adapted coordinates on TN.

On $M \times N$, we consider an arbitrary tensor of type (1,2), denoted by P, with all components null except $P_{\alpha i}^{\beta}(a,x)$ and $P_{\alpha i}^{j}(a,x)$, where $\alpha, \beta = \overline{1,m}, i, j = \overline{1,n},$ which will be called *tensor of connection*. The connection tensor P allows one to build the directions b and y of the metric tensors $g_{\alpha\beta}(a,b)$ and $h_{ij}(x,y)$ used in the construction of the energy functional E whose extremals will be the harmonic maps between the generalized Lagrange spaces M and N. In conclusion, this tensor makes the connection between the metric structures of the spaces M, respectively N, and the harmonic maps that we will define.

We assume that the manifold M is connected, compact, orientable and endowed with a Riemannian metric $\varphi_{\alpha\beta}$. These conditions assure the existence of a volume element and, implicitly, of a theory of integration on M. Using the generalized Lagrange metrics $g_{\alpha\beta}$ and h_{ij} and the tensor of connection P, we can define the following $\left(egin{array}{cc} P & \ g & arphi & h \end{array}
ight)$ -energy functional,

$$E = E_{g\varphi h}^{i} : C^{\infty}(M, N) \to R,$$

$$E_{g\varphi h}^{P}(f) = \frac{1}{2} \int_{M} g^{\alpha\beta}(a, b(a, f^{k}, f^{k}_{\gamma}))h_{ij}(f(a), y(a, f^{k}, f^{k}_{\gamma}))f^{i}_{\alpha}f^{j}_{\beta}\sqrt{\varphi} \, da,$$
where
$$\begin{cases}
f^{i} = x^{i}(f), \ f^{i}_{\alpha} = \frac{\partial f^{i}}{\partial a^{\alpha}}, \ \varphi = det(\varphi_{\alpha\beta}), \\
b(a, f^{k}, f^{k}_{\gamma}) = b^{\gamma}(a) \ \frac{\partial}{\partial a^{\gamma}}\Big|_{a} \stackrel{\text{def}}{=} \varphi^{\alpha\beta}(a)f^{i}_{\alpha}(a)P^{\gamma}_{\beta i}(a, f(a)) \ \frac{\partial}{\partial a^{\gamma}}\Big|_{a}, \\
y(a, f^{k}, f^{k}_{\gamma}) = y^{k}(a) \ \frac{\partial}{\partial x^{k}}\Big|_{f(a)} \stackrel{\text{def}}{=} \varphi^{\alpha\beta}(a)f^{i}_{\alpha}(a)P^{k}_{\beta i}(a, f(a)) \ \frac{\partial}{\partial x^{k}}\Big|_{f(a)}.$$

Definition. A map $f \in C^{\infty}(M, N)$ is $\begin{pmatrix} P \\ g & \varphi & h \end{pmatrix}$ -harmonic iff f is a critical point for the functional E^{P}_{ach} .

point for the functional $E_{g\varphi h}^{P}$. **Particular cases.** i) If $g_{\alpha\beta}(a,b) = \varphi_{\alpha\beta}(a)$ and $h_{ij}(x,y) = h_{ij}(x)$ are Riemannian metrics and the connection tensor is an arbitrary one, it recovers the classical definition of a harmonic map between two Riemannian manifolds [1]. We remark that, in this case, the definition of harmonic maps is independent of the connection tensor field P.

ii) If we take N = R, $h_{11} = 1$ and the tensor of connection is of the form $P = (\delta^{\alpha}_{\beta}, P^{1}_{\beta 1})$, we obtain $C^{\infty}(M, N) = \mathcal{F}(M)$ and the energy functional becomes

$$E_{g\varphi 1}^{P}(f) = \frac{1}{2} \int_{M} g^{\alpha\beta}(a, grad_{\varphi}f) f_{\alpha} f_{\beta} \sqrt{\varphi} \, da \, , \, \forall f \in \mathcal{F}(M).$$

iii) If we consider $M = [a, b] \subset R$, $\varphi_{11} = g_{11} = 1$ and the tensor of connection is $P = (P_{1i}^1, \delta_i^k)$, we obtain $C^{\infty}(M, N) = \{x : [a, b] \to N \mid x - C^{\infty} \text{ differentiable}\}$. Denoting $C^{\infty}(M, N) = \Omega_{a,b}(N)$, the energy functional should be

$$E_{11h}^P(x) = \frac{1}{2} \int_a^b h_{ij}(x(t), \dot{x}(t)) \frac{dx^i}{dt} \frac{dx^j}{dt} dt, \ \forall x \in \Omega_{a,b}(N).$$

In conclusion, the $\begin{pmatrix} P \\ 1 & 1 \end{pmatrix}$ -harmonic curves are exactly the geodesics of the generalized Lagrange space $(N, h_{ij}(x, y))$ [2].

2 Harmonic maps between Lagrange spaces of electrodynamics

Let (M^m, L_M) and (N^n, L_N) be Lagrange spaces with the Lagrangians

$$L_M(a,b) = g_{\alpha\beta}(a)b^{\alpha}b^{\beta} + g_{\alpha\beta}(a)U^{\alpha}(a)b^{\beta} + F(a),$$

$$L_N(x,y) = h_{ij}(x)y^iy^j + h_{ij}(x)V^i(x)y^j + G(x),$$

where

- $g_{\alpha\beta}$ (resp. h_{ij}) is Riemannian metric on M (resp. N) representing the gravitational potentials on M (resp. N).
- U^{α} (resp. V^{i}) is a vector field on M (resp. N) representing the electromagnetic potentials.
- F (resp. G) is a smooth function on M (resp. N) representing the potential function.

The fundamental metric tensors of these Lagrangians are

$$g_{\alpha\beta}(a) = \frac{\partial^2 L_M}{\partial b^{\alpha} \partial b^{\beta}}, \ h_{ij}(x) = \frac{\partial^2 L_N}{\partial y^i \partial y^j}.$$

Taking an arbitrary tensor of connection P, the energy functional becomes

$$E_{g\varphi h}(f) = \frac{1}{2} \int_{M} g^{\alpha\beta}(a) h_{ij}(f(a)) f^{i}_{\alpha} f^{j}_{\beta} \sqrt{\varphi} \, da.$$

We remark that, in this case, the energy functional is independent of the connection tensor P.

The Euler-Lagrange equations will be, obviously, the equations of harmonic maps, namely

$$g^{\alpha\beta}\left\{f_{\alpha\beta}^{k} - \left[G_{\alpha\beta}^{\gamma} + \frac{1}{2}\frac{\partial}{\partial a^{\alpha}}\left(\ln\frac{g}{\varphi}\right)\delta_{\beta}^{\gamma}\right]f_{\gamma}^{k} + H_{ij}^{k}f_{\alpha}^{i}f_{\beta}^{j}\right\} = 0 , \ \forall \ k = \overline{1, n},$$

where

•
$$f^k_{\alpha\beta} = \frac{\partial^2 f^k}{\partial a^{\alpha} \partial a^{\beta}}$$
, $g = det(g_{\alpha\beta})$, $\varphi = det(\varphi_{\alpha\beta})$.

- $G_{\alpha\beta}^{\gamma}$ are the Christoffel symbols of the metric $g_{\alpha\beta}$.
- H_{ij}^k are the Christoffel symbols of the metric h_{ij} .

Remarks. i) If $g_{\alpha\beta} = \varphi_{\alpha\beta}$, we recover the classical equations of harmonic maps

Remarks. If $\ln g_{\alpha\beta} = r_{\alpha\beta}$, between two Riemannian manifolds. ii) Denoting $\Delta^{\gamma}_{\alpha\beta} = G^{\gamma}_{\alpha\beta} + \frac{1}{2} \frac{\partial}{\partial a^{\alpha}} \left(\ln \frac{g}{\varphi} \right) \delta^{\gamma}_{\beta}$, we remark that $\Delta^{\gamma}_{\alpha\beta}$ represent the components of a linear connection induced by the metrics $g_{\alpha\beta}$ and $\varphi_{\alpha\beta}$.

By the last remark, we can give the following

Definition. Let q, φ be Riemannian metrics on M. The curve $c : I \to M$, expressed locally by $c(t) = (a^{\alpha}(t))$, is a (g, φ) -geodesic iff c is an autoparallel curve of the connection $\Delta_{\alpha\beta}^{\gamma}$ induced by the metrics g and φ , namely

$$\frac{d^2 a^{\gamma}}{dt^2} = -\Delta^{\gamma}_{\alpha\beta} \frac{da^{\alpha}}{dt} \frac{da^{\beta}}{dt}.$$

Remarks. i) If $q = \varphi$, then we recover the classical definition of a geodesic on the Riemannian manifold $(M, g = \varphi)$.

ii) It is obviously that a (q, φ) -geodesic is a reparametrized geodesic of the metric q.

Theorem. Let $f : (M, L_M) \to (N, L_N)$ be a smooth map which carries (g, φ) -geodesics into h-geodesics. Then f is harmonic map.

Proof. Let $c: I \subset R \to M$, $c(t) = (a^{\alpha}(t))$ be a (g, φ) -geodesic. Then we have $\frac{d^2 a^{\gamma}}{dt^2} = -\Delta_{\alpha\beta}^{\gamma} \frac{da^{\alpha}}{dt} \frac{da^{\beta}}{dt}$. Because $\overline{c}(t) = f(c(t))$ is *h*-geodesic, it follows that $\frac{d^2 \overline{c}^k}{dt^2} + H_{ij}^k \frac{d\overline{c}^i}{dt} \frac{d\overline{c}^j}{dt} = 0$. But $\frac{d\overline{c}^k}{dt} = f_{\alpha}^k(c(t)) \frac{da^{\alpha}}{dt} \Rightarrow \frac{d^2 \overline{c}^k}{dt^2} = f_{\alpha\beta}^k \frac{da^{\alpha}}{dt} \frac{da^{\beta}}{dt} + f_{\alpha}^k \frac{d^2 a^{\alpha}}{dt^2}$. In conclusion we obtain

$$\frac{d^2 a^{\alpha}}{dt^2} f^k_{\alpha} + f^k_{\alpha\beta} \frac{da^{\alpha}}{dt} \frac{da^{\beta}}{dt} + H^k_{ij} f^i_{\alpha} f^j_{\beta} \frac{da^{\alpha}}{dt} \frac{da^{\beta}}{dt} = 0 \Rightarrow$$
$$\left(f^k_{\alpha\beta} - \Delta^{\gamma}_{\alpha\beta} f^k_{\gamma} + H^k_{ij} f^i_{\alpha} f^j_{\beta}\right) \frac{da^{\alpha}}{dt} \frac{da^{\beta}}{dt} = 0 , \ \forall k = \overline{1, n} \Box$$

3 Geometrical interpretation of solutions of certain PDEs of order one

The problem of finding a geometrical structure of Riemannian type on a manifold M such that the orbits of an arbitrary vector field X should be geodesics, was intensively studied by Sasaki. The results were not satisfactory, but, in his study, Sasaki discovered the well known almost contact structures on a manifold of odd dimension [6]. After the introduction of generalized Lagrange spaces by Miron [2], the same problem is resumed by Udrişte [8, 9]. This succeded to discover a Lagrange structure on M, depending of the vector field X and an associated (1,1)-tensor field, such that the orbits of C^2 class should be geodesics. Moreover, he formulated a more general problem [9], namely

1) Are there structures of Lagrange type such that the solutions of certain PDEs of order one should be *harmonic maps*?

2) What is a *harmonic map* between two generalized Lagrange spaces?

A partial answer of these questions is offered by author in his paper [4], using the notion of harmonic map on a direction between a Riemannian manifold and a generalized Lagrange manifold. The notion of *harmonic map* between two generalized Lagrange spaces introduced in [7] allows one to extend the results of previous papers [4], [8], [9] and to obtain a beautiful geometrical interpretation of the solutions of the certain PDEs of order one.

For every smooth map $f \in C^{\infty}(M, N)$, we use the following notation

$$\delta f = f^i_{\alpha} da^{\alpha} |_a \otimes \frac{\partial}{\partial y^i} \Big|_{f(a)} \in \Gamma(T^*M \otimes f^{-1}(TN)).$$

On $M \times N$, let T be one tensor of type (1, 1) with all components equal to zero except $(T^i_{\alpha})_{i=\overline{1,m} \atop \alpha=\overline{1,m}}$. Let the system of partial differential equations

(E)
$$\delta f = T$$
 expressed locally by $\frac{\partial f^i}{\partial a^{\alpha}} = T^i_{\alpha}(a, f(a)).$

If $(M, \varphi_{\alpha\beta})$ and (N, ψ_{ij}) are Riemannian manifolds, we can build a scalar product on $\Gamma(T^*M \otimes f^{-1}(TN))$ by $\langle T, S \rangle = \varphi^{\alpha\beta}(a)\psi_{ij}(f(a))T^i_{\alpha}S^j_{\beta}$, where $T = T^i_{\alpha}da^{\alpha} \otimes \frac{\partial}{\partial y^i}$ and $S = S^j_{\beta}da^{\beta} \otimes \frac{\partial}{\partial y^j}$.

Under these conditions, we can prove the following

Theorem. If $(M, \varphi), (N, \psi)$ are Riemannian manifolds and $f \in C^{\infty}(M, N)$ is a solution of the system (E), then f is a solution of the variational problem associated to the functional $\mathcal{L}_T : C^{\infty}(M, N) \setminus \{f \mid \exists a \in M \text{ such that } < \delta f, T > (a) = 0\} \rightarrow R_+,$

$$\mathcal{L}_T(f) = \frac{1}{2} \int_M \frac{\|\delta f\|^2 \|T\|^2}{\langle \delta f, T \rangle^2} \sqrt{\varphi} \, da = \frac{1}{2} \int_M \frac{\|T\|^2}{\langle \delta f, T \rangle^2} \varphi^{\alpha\beta} \psi_{ij} f^i_{\alpha} f^j_{\beta} \sqrt{\varphi} \, da.$$

Proof. In the space $\Gamma(T^*M \times f^{-1}(TN))$, the Cauchy inequality for the scalar product \langle , \rangle holds. It follows that the following inequality is true, $\langle T, S \rangle^2 \leq ||T||^2 ||S||^2$, $\forall T, S \in \Gamma(T^*M \times f^{-1}(TN))$, with equality if and only if there exists $\mathcal{K} \in \mathcal{F}(M)$ such that $T = \mathcal{K}S$. Consequently, for every $f \in C^{\infty}(M, N)$ we have

$$\mathcal{L}(f) = \frac{1}{2} \int_M \frac{\|\delta f\|^2 \|T\|^2}{\langle \delta f, T \rangle^2} \sqrt{\varphi} \, da \ge \frac{1}{2} \int_M \sqrt{\varphi} \, da = \frac{1}{2} Vol_{\varphi}(M)$$

Obviously, if f is a solution of the system (E), we obtain $\mathcal{L}_T(f) = \frac{1}{2} Vol_{\varphi}(M)$, that is, f is a global minimum point for $\mathcal{L}_T \square$.

Remarks. i) In certain particular cases of the system (*E*), the functional \mathcal{L}_T becomes exactly a functional of type $\begin{pmatrix} P \\ g & \varphi & h \end{pmatrix}$ -energy.

ii) The global minimum points of the functional \mathcal{L}_T are solutions of the system $\delta f = \mathcal{K}T$, where $\mathcal{K} \in \mathcal{F}(M)$, not necessarily with $\mathcal{K} = 1$.

iii) Replacing the Riemannian metric ψ_{ij} by a pseudo-Riemannian metric, the preceding theorem survives because the form of the Euler-Lagrange equations remains unchanged. The difference is that the solutions of the system (E), in the pseudo-Riemannian case, are not the global minimum points for the functional \mathcal{L}_T . Moreover, the statement (ii), of above, does not hold.

Fundamental examples.

1. Orbits

For M = ([a, b], 1) and $T = \xi \in \Gamma(x^{-1}(TN))$, the system (E) becomes

(E₁)
$$\frac{dx^i}{dt} = \xi^i(x(t)), \ x: [a, b] \to N,$$

that is the system of orbits for ξ , and the functional \mathcal{L}_{ξ} is

$$\mathcal{L}_{\xi}(x) = \frac{1}{2} \int_{a}^{b} \frac{\|\xi\|_{\psi}^{2}}{[\xi^{b}(\dot{x})]^{2}} \psi_{ij} \frac{dx^{i}}{dt} \frac{dx^{j}}{dt} dt,$$

where $\xi^b = \xi_i dx^i = \psi_{ij} \xi^j dx^i$. Hence the functional \mathcal{L}_{ξ} is a $\begin{pmatrix} P \\ 1 & 1 & h \end{pmatrix}$ -energy (see (iii) of first particular cases of this paper), where

$$h_{ij}: TN \setminus \{y \mid \xi^b(y) = 0 \text{ for some } y\} \to R$$

is defined by

$$h_{ij}(x,y) = \frac{\|\xi\|_{\psi}^2}{[\xi^b(y)]^2} \psi_{ij}(x) = \psi_{ij}(x) \exp\left[2\ln\frac{\|\xi\|_{\psi}}{|\xi^b(y)|}\right]$$

This case is studied in other way by Udrişte in [8]-[9].

2. Pfaffian systems

For N = (R, 1) and $T = A \in \Lambda^1(T^*M)$, the system (E) becomes

$$(E_2) df = A, \ f \in \mathcal{F}(M)$$

that is a Pfaffian system, and the functional \mathcal{L}_T reduces to

$$\mathcal{L}_A(f) = \frac{1}{2} \int_M \frac{\|A\|_{\varphi}^2}{[A(grad_{\varphi}f)]^2} \varphi^{\alpha\beta} f_{\alpha} f_{\beta} \sqrt{\varphi} da.$$

Hence, the functional \mathcal{L}_A is a $\begin{pmatrix} P \\ g & \varphi & 1 \end{pmatrix}$ -energy (see (ii) of first particular cases of this paper), where $g_{\alpha\beta}: TM \setminus \{b \mid A(b) = 0 \text{ for some } b\} \to R$ is defined by

$$g_{\alpha\beta}(a,b) = \frac{[A(b)]^2}{\|A\|_{\varphi}^2} \varphi_{\alpha\beta}(a) = \varphi_{\alpha\beta}(a) \exp\left[2\ln\frac{|A(b)|}{\|A\|_{\varphi}}\right].$$

3. Pseudolinear functions

We suppose that $T_{\beta}^{k}(a, x) = \xi^{k}(x)A_{\beta}(a)$, where ξ^{k} is vector field on N and A_{β} is 1-form on M. In this case the system (E) is

(E₃)
$$\frac{\partial f^k}{\partial a^\beta} = \xi^k(f) A_\beta(a)$$

and the functional \mathcal{L}_T is expressed by

$$\mathcal{L}_{T}(f) = \frac{1}{2} \int_{M} \frac{\|\xi\|_{\psi}^{2} \|A\|_{\varphi}^{2}}{[A(b)]^{2}} \varphi^{\alpha\beta} \psi_{ij} f_{\alpha}^{i} f_{\beta}^{j} \sqrt{\varphi} da =$$
$$= \frac{1}{2} \int_{M} g^{\alpha\beta}(a, b) h_{ij}(f(a)) f_{\alpha}^{i} f_{\beta}^{j} \sqrt{\varphi} da,$$

where $h_{ij}(x) = \|\xi\|^2_{\psi}\psi_{ij}(x)$, the tensor of connection is $P^{\gamma}_{i\beta}(x) = \delta^{\gamma}_{\beta}\xi_i(x)$, $b^{\gamma} = \varphi^{\alpha\beta}f^i_{\alpha}P^{\gamma}_{i\beta}$ and the Lagrange metric tensor

$$g_{\alpha\beta}: TM \setminus \{b \mid A(b) = 0 \text{ for some } b\} \to R$$

is defined by

$$g_{\alpha\beta}(a,b) = \frac{[A(b)]^2}{\|A\|_{\varphi}^2} \varphi_{\alpha\beta}(a) = \varphi_{\alpha\beta}(a) \exp\left[2\ln\frac{|A(b)|}{\|A\|_{\varphi}}\right]$$

It follows that the functional \mathcal{L}_T becomes a $\begin{pmatrix} P \\ g & \varphi & h \end{pmatrix}$ -energy.

Remark. Take M to be an open subset in $(R^n, \varphi = \delta)$ and $N = (R, \psi = 1)$, the system from the third example is

(PL)
$$\frac{\partial f}{\partial a^{\alpha}} = \xi(a) A_{\alpha}(f(a)), \ \forall \ \alpha = \overline{1, m}.$$

Supposing that $(grad f)(a) \neq 0, \forall a \in M$, the solutions of this system are the well known *pseudolinear functions* [5]. These functions have the following property,

-for every fixed point $x_0 \in M$, the hypersurface of constant level

$$M_{f(x_0)} = \{ x \in M \mid f(x) = f(x_0) \}$$

is totally geodesic [5] (i. e. the second fundamental form vanishes identically).

In conclusion, the pseudolinear functions are examples of harmonic maps between the generalized Lagrange spaces $\left(M, g_{\alpha\beta}(a, b) = \frac{[A(b)]^2}{\|A\|^2} \delta_{\alpha\beta}\right)$ and $(R, h(x) = \xi^2(x))$. For example, we have the following pseudolinear functions [5]:

3. 1. $f(a) = e^{\langle v, a \rangle + w}$, where $v \in M$, $w \in R$, is solution for the system (*PL*) with $\xi(a) = 1$ and A(f(a)) = f(a)v.

3. 2.
$$f(a) = \frac{\langle v, a \rangle + w}{\langle v', a' \rangle + w'}$$
, where $v, v' \in M$, $w, w' \in R$, is solution for (PL) th $\xi(a) = \frac{1}{1}$ and $A(f(a)) = v - f(a)w$.

with $\xi(a) = \frac{1}{\langle v', a \rangle + w'}$ and A(f(a)) = v - f(a)w. **Bemark** The preceding cases appear also in [7] and $f(a) = \frac{1}{\langle v', a \rangle + w'}$

Remark. The preceding cases appear also in [7] and, from another point of view, in [4]. The following case is the main novelty of this paper.

4. The general case

If we have $T^i_{\alpha}(a,x) = \sum_{r=1}^t \xi^i_r(x) A^r_{\alpha}(a)$, where $\{\xi_r\}_{r=\overline{1,t}} \subset \mathcal{X}(N)$ is a family of vector fields on N and $\{A^r\}_{r=\overline{1,t}} \subset \Lambda^1(T^*M)$ is a family of 1-forms on M, the system of equations (E) reduces to

(E₄)
$$\frac{\partial f^i}{\partial a^{\alpha}} = \sum_{r=1}^t \xi^i_r(f) A^r_{\alpha}(a).$$

Without loss of generality, we can suppose that $\{\xi_r\}_{r=\overline{1,t}} \subset \mathcal{X}(N)$ (resp. $\{A^r\}_{r=\overline{1,t}} \subset \Lambda^1(T^*M)$) are linearly independent. In these conditions, we shall have $t \leq \min\{m, n\}$, where $m = \dim M$ and $n = \dim N$.

4. 1. Assume that $\{\xi_r\}_{r=\overline{1,t}} \subset \mathcal{X}(N)$ is an orthonormal system of vector fields with respect to the Riemannian metric ψ_{ij} on N. Let $B \in \Lambda^1(T^*M)$ be an arbitrary unit 1-form on M. With our assumptions, by a simple calculation, we obtain

$$||T||^{2} = \varphi^{\alpha\beta}\psi_{ij}\xi_{r}^{i}A_{\alpha}^{r}\xi_{s}^{j}A_{\beta}^{s} = \sum_{r,s=1}^{t} \langle \xi_{r}, \xi_{s} \rangle_{\psi} \langle A^{r}, A^{s} \rangle_{\varphi} = \sum_{r=1}^{t} ||A^{r}||_{\varphi}^{2}$$
$$\langle \delta f, T \rangle = \varphi^{\alpha\beta}\psi_{ij}f_{\alpha}^{i}\xi_{r}^{j}A_{\beta}^{r}B^{\mu}B_{\mu}.$$

Defining the tensor of connection by $P_{i\beta}^{\gamma}(a,x) = \psi_{ij}(x)\xi_r^j(x)A_{\beta}^r(a)B^{\gamma}(a)$ and $b^{\gamma} = \varphi^{\alpha\beta}f_{\alpha}^i P_{i\beta}^{\gamma}$, the functional \mathcal{L}_T takes the form

$$\mathcal{L}_T(f) = \frac{1}{2} \int_M \frac{\sum_{r=1}^t \|A^r\|_{\varphi}^2}{[B(b)]^2} \varphi^{\alpha\beta} \psi_{ij} f^i_{\alpha} f^j_{\beta} \sqrt{\varphi} \, da = \frac{1}{2} \int_M g^{\alpha\beta}(a,b) \psi_{ij}(f(a)) f^i_{\alpha} f^j_{\beta} \sqrt{\varphi} \, da,$$

where the Lagrange metric tensor $g_{\alpha\beta}$: $TM \setminus \{b \mid B(b) = 0 \text{ for some } b\} \to R$ is expressed by

$$g_{\alpha\beta}(a,b) = \frac{[B(b)]^2}{\sum_{r=1}^t \|A^r\|_{\varphi}^2} \varphi_{\alpha\beta}(a) = \varphi_{\alpha\beta}(a) \exp\left[2\ln\frac{|B(b)|}{\sqrt{\sum_{r=1}^t \|A^r\|_{\varphi}^2}}\right]$$

Consequently, the functional \mathcal{L}_T is a $\begin{pmatrix} P \\ g & \varphi & \psi \end{pmatrix}$ -energy.

4. 2. As above, we assume that the system $\{A^r\}_{r=\overline{1,t}}$ of 1-forms is orthonormal with respect to the metric $\varphi^{\alpha\beta}$ and we choose an arbitrary unit vector field $X \in \mathcal{X}(N)$. By analogy to **4. 1** we shall have

$$||T||^2 = \sum_{r=1}^t ||\xi_r||_{\psi}^2 \text{ and } < \delta f, T > = \varphi^{\alpha\beta} \psi_{ij} f_{\alpha}^i \xi_r^j A_{\beta}^r X^k X_k.$$

Using the notations $P_{i\beta}^k(a,x) = \psi_{ij}(x)\xi_r^j(x)A_\beta^r(a)X^k(x)$ and $y^k = \varphi^{\alpha\beta}f_{\alpha}^i P_{i\beta}^k$ we obtain the following expression of the functional \mathcal{L}_T ,

$$\mathcal{L}_T(f) = \frac{1}{2} \int_M \frac{\sum_{r=1}^t \|\xi_r\|_{\psi}^2}{[X^b(y)]^2} \varphi^{\alpha\beta} \psi_{ij} f_{\alpha}^i f_{\beta}^j \sqrt{\varphi} \, da = \frac{1}{2} \int_M \varphi^{\alpha\beta}(a) h_{ij}(f(a), y) f_{\alpha}^i f_{\beta}^j \sqrt{\varphi} \, da,$$

where the Lagrange metric tensor $h_{ij}: TN \setminus \{y \mid X^b(y) = 0 \text{ for some } y\} \to R$ is

$$h_{ij}(x,y) = \frac{\sum_{r=1}^{t} \|\xi_r\|_{\psi}^2}{[X^b(y)]^2} \psi_{ij}(x) = \psi_{ij}(x) \exp\left[2\ln\frac{\sqrt{\sum_{r=1}^{t} \|\xi_r\|_{\psi}^2}}{|X^b(y)|}\right]$$

Obviously, the functional \mathcal{L}_T is a $\begin{pmatrix} P \\ \varphi & \varphi & h \end{pmatrix}$ -energy.

Remarks. i) For the use of the above, we assume a priori a Riemannian metric φ or ψ on M or N such that the system of covectors $\{A^r\}_{r=\overline{1,t}}$ or of vectors $\{\xi_r\}_{r=\overline{1,t}}$ is orthonormal. This fact is always possible. In conclusion, our assumptions on the orthonormality of these systems do not restrict the generality of problem.

ii) In the particular case, n = r = 1, N = R, $\psi_{11} = 1$, $\xi = \frac{d}{dx}$, and $A \in \Lambda^1(T^*M)$ is an arbitrary 1-form on M, we recover the Pfaffian system df = A. In this situation, taking $B \in \Lambda^1(T^*M)$ to be an arbitrary unit 1-form, we obtain, for the functional \mathcal{L}_T , the expression

$$\mathcal{L}_T(f) = \frac{1}{2} \int_M \frac{\|A\|^2}{[B(b)]^2} \varphi^{\alpha\beta} f_\alpha f_\beta \sqrt{\varphi} \, da = \frac{1}{2} \int_M \frac{\|A\|^2}{[A(grad_\varphi f)]^2} \varphi^{\alpha\beta} f_\alpha f_\beta \sqrt{\varphi} \, da,$$

where the tensor of connection is defined by $P_{1\beta}^{\gamma} = A_{\beta}B^{\gamma}$ and the direction b is $b^{\gamma} = \varphi^{\alpha\beta}f_{\alpha}^{i}A_{\beta}B^{\gamma} = B^{\gamma}A(grad_{\varphi}f).$

Since the last integral is the functional \mathcal{L}_A from the example **1**, we remark that the solutions of the Pfaffian system df = A can be regarded in an infinity manner as $\begin{pmatrix} P \\ g & \varphi & 1 \end{pmatrix}$ -harmonic maps. This fact appears because the tensor of connection P

and the generalized Lagrange metric g are dependent of the arbitrary unit covector field B.

iii) Analog to ii), if we take m = r = 1, M = [a, b], $g_{11} = \varphi_{11} = 1$, A = dt and $\xi \in \mathcal{X}(N)$ is an arbitrary vector field on N, we find the system of orbits for ξ , that is,

$$\frac{dx^i}{dt} = \xi^i(x(t)), \ x: [a, b] \to N.$$

Starting with $X \in \mathcal{X}(N)$ an arbitrary unit vector field, the functional \mathcal{L}_T becomes

$$\mathcal{L}_T(x) = \frac{1}{2} \int_a^b \frac{\|\xi\|_{\psi}^2}{[X^b(y)]^2} \psi_{ij} \frac{dx^i}{dt} \frac{dx^j}{dt} dt = \frac{1}{2} \int_a^b \frac{\|\xi\|_{\psi}^2}{[\xi^b(x)]^2} \psi_{ij} \frac{dx^i}{dt} \frac{dx^j}{dt} dt,$$

where the tensor of connection is $P_{i1}^k = \xi_i^b X^k$ and $y^k = \xi^b(\dot{x}) X^k$. Because we can vary the tensor of connection P and the generalized Lagrange metric h by the arbitrary unit vector field X, we remark that the trajectories of the vector field ξ can be also regarded in an infinity manner as $\begin{pmatrix} P \\ 1 & 1 & h \end{pmatrix}$ -harmonic maps.

4 Lagrange geometry associated to PDEs of order one

We first remark that, in all above cases, the solutions of C^2 class of the system $\delta f = T$ becomes harmonic maps between generalized Lagrange spaces, in the sense defined in this paper. Moreover, the above generalized Lagrange structures are of type $(M^n, e^{2\sigma(x,y)}\gamma_{ij}(x))$, where $\sigma : TM \setminus \{\text{Hyperplane}\} \to R$ is a smooth function. Using the ideas exposed in [2], in these spaces, we can construct a Lagrange geometry and field theory. This geometrical Lagrange theory will be regarded as a natural one associated to the PDE system $\delta f = T$, in the sense of the first Udrişte's question.

Now, we assume that a generalized Lagrange space $(M^n, g_{ij}(x, y))$ satisfies the following axioms:

a. 1. The fundamental tensor field $g_{ij}(x, y)$ is of the form

$$g_{ij}(x,y) = e^{2\sigma(x,y)}\gamma_{ij}(x).$$

a. 2. The space is endowed with the non-linear connection

$$N_{i}^{i}(x,y) = \Gamma_{ik}^{i}(x)y^{k},$$

where $\Gamma^{i}_{ik}(x)$ are the Christoffel symbols for the Riemannian metric $\gamma_{ij}(x)$.

Under these assumptions, our space verifies a constructive-axiomatic formulation of General Relativity due to Ehlers, Pirani and Schild [2]. This space represents a convenient relativistic model, since it has the same conformal and projective properties as the Riemannian space (M, γ_{ij}) .

In the Lagrangian theory of electromagnetism, the electromagnetic tensors F_{ij} and f_{ij} are

$$F_{ij} = \left(g_{ip}\frac{\delta\sigma}{\delta x^j} - g_{jp}\frac{\delta\sigma}{\delta x^i}\right)y^p , \ f_{ij} = \left(g_{ip}\frac{\partial\sigma}{\partial y^j} - g_{jp}\frac{\partial\sigma}{\partial y^i}\right)y^p.$$

Developping the formalism presented in [2], [3] and denoting by r_{ikl}^i the curvature tensor field of the metric $\gamma_{ij}(x)$, the following Maxwell equations of the electromagnetic tensors hold

$$\begin{cases} F_{ij|k} + F_{jk|i} + F_{ki|j} = -\sum_{(ijk)} g_{ip} r_{qjk}^{h} \frac{\partial \sigma}{\partial y^{h}} y^{p} y^{q}, \\ F_{ij|k} + F_{jk|i} + F_{ki|j} = -(f_{ij|k} + f_{jk|i} + f_{ki|j}), \\ f_{ij|k} + f_{jk|i} + f_{ki|j} = 0, \end{cases}$$

where $|_i$ (resp. $|_a$) represents the h- (resp. v-) covariant derivative induced by the

non-linear connection N_j^i . Suppose $\sigma = \sigma(x)$. Then the v- electromagnetic tensor is $f_{ij} = 0$, the hcovariant operator " $_{i}$ " becomes the covariant derivative with respect to Levi-Civita connection of the metric $g_{ij}(x,y) = e^{2\sigma(x)}\varphi_{ij}(x)$, the *h*- electromagnetic tensor F_{ij} is the same with the classical electromagnetic tensor and the Maxwell's equations reduce to the classical ones.

In the construction of the gravitational field equations, we shall use the notations

$$\begin{cases} r_{ij} = r_{ijk}^k, \ r = \gamma^{ij} r_{ij}, \ \frac{\delta}{\delta x^i} = \frac{\partial}{\partial x^i} - N_i^j \frac{\partial}{\partial y^j}, \\ \sigma^{\scriptscriptstyle H} = \gamma^{kl} \frac{\delta\sigma}{\delta x^k} \frac{\delta\sigma}{\delta x^l}, \ \sigma^{\scriptscriptstyle V} = \gamma^{ab} \frac{\partial\sigma}{\partial y^a} \frac{\partial\sigma}{\partial y^b}, \ \overline{\sigma} = \gamma^{ij} \sigma_{ij}, \ \dot{\sigma} = \gamma^{ab} \dot{\sigma}_{ab} \end{cases}$$

where $\sigma_{ij} = \frac{\delta\sigma}{\delta x^i}|_j + \frac{\delta\sigma}{\delta x^i}\frac{\delta\sigma}{\delta x^j} - \frac{1}{2}\gamma_{ij}\sigma^{\scriptscriptstyle H}, \ \dot{\sigma}_{ab} = \left.\frac{\partial\sigma}{\partial y^a}\right|_b + \frac{\partial\sigma}{\partial y^a}\frac{\partial\sigma}{\partial y^b} - \frac{1}{2}\gamma_{ab}\sigma^{\scriptscriptstyle V}.$

In these conditions, the Einstein's equations of the space $(M, g_{ij}(x, y))$ take the form

$$\begin{cases} r_{ij} - \frac{1}{2}r\gamma_{ij} + t_{ij} = \mathcal{K}T^{H}_{ij} \\ (2-n)(\dot{\sigma}_{ab} - \dot{\sigma}\gamma_{ab}) = \mathcal{K}T^{V}_{ab}, \end{cases}$$

where $T_{ij}^{\scriptscriptstyle H}$ and $T_{ab}^{\scriptscriptstyle V}$ are the h- and the v- components of the energy momentum tensor field, \mathcal{K} is the gravific constant and

$$t_{ij} = (n-2)(\gamma_{ij}\overline{\sigma} - \sigma_{ij}) + \gamma_{ij}r_{st}y^s\gamma^{tp}\frac{\partial\sigma}{\partial y^p} + \frac{\partial\sigma}{\partial y^i}r^a_{tja}y^t - \gamma_{is}\gamma^{ap}\frac{\partial\sigma}{\partial y^p}r^s_{tja}y^t.$$

It is clear that the general metric $g_{ij}(x,y) = e^{2\sigma(x,y)}\varphi_{ij}(x)$ implies Einstein equations which differ from the classical ones by the additional tensor t_{ij} .

Finally, we remark that, in certain particular cases, it is possible to build a generalized Lagrange geometry and field theory naturally attached to a system of partial differential equations. In these geometrical structures, the solutions of C^2 class of the PDE system become harmonic maps. This idea was suggested by Udriste in private discussions and in [9], [10].

Open problem. Because the generalized Lagrange structure constructed in this paper is not unique, it arises a natural question:

-Is it possible to build a unique generalized Lagrange geometry naturally asociated to a given PDEs system?

An answer to this question will be offered by author in a subsequent paper, using a more general Lagrange geometry, naturally attached to a multidimensional Lagrangian defined on the jet fibration of order one.

Acknowledgements. I would like to express my gratitude to the reviewer of Southeast Asian Bulletin of Mathematics and Prof. Dr. C. Udrişte for their valuable comments and very useful suggestions.

References

- J. Eells, L. Lemaire, A Report on Harmonic Maps, Bull. London Math. Soc. 10 (1978), 1-68.
- [2] R. Miron, M. Anastasiei, The Geometry of Lagrange Spaces: Theory and Applications, Kluwer Academic Publishers, 1994.
- [3] R. Miron, R. K. Tavakol, V. Balan, I. Roxburgh, Geometry of Space-Time and Generalized Lagrange Gauge Theory, Publ. Math., Debrecen, Hungary, 42, 3-4 (1992), 215-224.
- [4] M. Neagu, Solutions of Inverse Problems for Variational Calculus, Workshop on Diff. Geom., Global Analysis, Lie Algebras, Aristotle University of Thessaloniki, Greece, June 24-28, 1998; BSG Proceedings 4, pp 180-187, Editor:Prof. Dr. Grigorios Tsagas, Geometry Balkan Press, Bucharest, 2000.
- [5] T. Rapcsák, Smooth Nonlinear Optimization in Rⁿ, Kluwer Academic Publishers, 1997.
- [6] S. Sasaki, Almost Contact Manifolds, I, II, III, Mathematical Institute Tohoku University, 1965, 1967, 1968.
- [7] C. Udrişte, M. Neagu, Geometrical Interpretation of Solutions of Certain PDEs, Balkan Journal of Geometry and Its Applications, 4, 1 (1999), 145-152.
- [8] C. Udrişte, A. Udrişte, *Electromagnetic Dynamical Systems*, Balkan Journal of Geometry and Its Applications, 2, 1 (1997), 129-140.
- [9] C. Udrişte, Geometric Dynamics, Second Conference of Balkan Society of Geometers, Aristotle University of Thessaloniki, Greece, June 23-26, 1998; Southeast Asian Bulletin of Mathematics, Springer-Verlag, 24 (2000), 1-11.
- [10] C. Udrişte, Nonclassical Dynamics, Proceedings of Conference on Mathematics in honour of Professor Radu Roşca at the occasion of his ninetieth birthday, Katholieke Universiteit Brussel, Katholieke Universiteit Leuven, Dec. 11-16, 1999, to appear.

University POLITEHNICA of Bucharest Department of Mathematics I Splaiul Independentei 313 77206 Bucharest, Romania e-mail:mircea@mathem.pub.ro