

Some remarks on $\mathcal{U}_q(\mathfrak{sl}(2, R))$ at root of unity

PAVEL ŠŤOVÍČEK

*Department of Mathematics
Faculty of Nuclear Science, Czech Technical University
Trojanova 13, 120 00 Prague, Czech Republic
stovicek@km1.fjfi.cvut.cz*

Abstract

We discuss a modification of $\mathcal{U}_q(\mathfrak{sl}(2, \mathbb{R}))$ and a class of its irreducible representations when q is a root of unity.

1 Introduction

Nowadays q -deformed universal enveloping algebras $\mathcal{U}_q(\mathfrak{g})$ are understood in depth in the case when \mathfrak{g} is a complex simple Lie algebra belonging to one of the four principal series. The same is true for compact forms of these Lie algebras (see, e.g., monographs [1], [2], [3]). On the other hand, attempts to introduce q -deformed enveloping algebras for non-compact real Lie algebras frequently lead to serious difficulties though several particular cases have been already studied (see, e.g., [4], [5], [6]). In this note we discuss one of the simplest examples with $\mathfrak{g} = \mathfrak{sl}(2, \mathbb{R})$ as a real form of $\mathfrak{sl}(2, \mathbb{C})$. The deformation parameter q is supposed to be a root of unity,

$$q = \exp(i\pi P/Q),$$

where $Q \in \mathbb{N}$ is odd, $P \in \{1, \dots, Q-1\}$, and P and Q are relatively prime integers. So $q^{2j} \neq 1$, $j = 1, \dots, Q-1$, and $q^{2Q} = 1$.

We use the standard definition of the Hopf algebra $\mathcal{U}_q(\mathfrak{sl}(2, \mathbb{C}))$ with the generators K, K^{-1}, E, F , the defining relations

$$K K^{-1} = K^{-1} K = 1, \quad K E = q E K, \quad K F = q^{-1} F K, \\ [E, F] = \frac{1}{q - q^{-1}}(K^2 - K^{-2}),$$

the comultiplication

$$\Delta K = K \otimes K, \quad \Delta E = K \otimes E + E \otimes K^{-1}, \quad \Delta F = K \otimes F + F \otimes K^{-1},$$

the antipode

$$S(K) = K^{-1}, \quad S(E) = -q^{-1}E, \quad S(F) = -qF,$$

and the counit

$$\varepsilon(K) = 1, \quad \varepsilon(E) = \varepsilon(F) = 0.$$

A real form is determined by a $*$ -involution; an element X of a complex Hopf algebra belongs to a real form if and only if $X^* = S(X)$. Particularly, $\mathcal{U}_q(\mathfrak{sl}(2, \mathbb{R}))$ is determined by the $*$ -involution

$$K^* = K, \quad E^* = -q^{-1}E, \quad F^* = -qF. \quad (1)$$

Necessarily, q is a complex unit, $\bar{q} = q^{-1}$.

Usually it is more convenient to deal with the complexification of a real form. In that case one regards the real form as the original complex Hopf algebra but endowed, in addition, with the $*$ -involution in question. We shall adopt this point of view and treat $\mathcal{U}_q(\mathfrak{sl}(2, \mathbb{R}))$ as the complex Hopf algebra $\mathcal{U}_q(\mathfrak{sl}(2, \mathbb{C}))$ with the $*$ -involution (1).

2 A modification of $\mathcal{U}_q(\mathfrak{sl}(2, \mathbb{R}))$

Let \mathcal{U} be a $*$ -Hopf subalgebra of $\mathcal{U}_q(\mathfrak{sl}(2, \mathbb{R}))$ generated by X, Y, Z, Z^{-1} , where

$$X = -i q^{-1} E K^{-1}, \quad Y = -i q F K^{-1}, \quad Z = K^{-2}.$$

Thus \mathcal{U} is defined by the relations

$$Z X = q^{-2} X Z, \quad Z Y = q^2 Y Z, \quad q^{-1} X Y - q Y X = -\frac{1}{q - q^{-1}}(1 - Z^2), \quad (2)$$

with the comultiplication

$$\Delta Z = Z \otimes Z, \Delta X = 1 \otimes X + X \otimes Z, \Delta Y = 1 \otimes Y + Y \otimes Z,$$

the antipode

$$S(Z) = Z^{-1}, S(X) = -X Z^{-1}, S(Y) = -Y Z^{-1},$$

and the counit

$$\varepsilon(Z) = 1, \varepsilon(X) = \varepsilon(Y) = 0.$$

Furthermore, all the generators are Hermitian,

$$Z^* = Z, X^* = X, Y^* = Y.$$

It is also straightforward to check that

$$C = X Y Z^{-1} - \frac{1}{(q - q^{-1})^2} (Z^{-1} + q^2 Z)$$

is an Casimir element in \mathcal{U} .

Unfortunately, there exists no non-trivial irreducible representations ρ of \mathcal{U} . Actually, Z^Q belongs to the center of \mathcal{U} and is Hermitian. Thus, by the Schur lemma, $\rho(Z)^Q = cI$ for some real $c \neq 0$. Consequently, the self-adjoint operator $\rho(Z)$ is a multiple of the identity as well. The commutation relations then imply that $\rho(X) = \rho(Y) = 0$, $\rho(Z) = \pm I$.

To improve this situation we propose a modification of \mathcal{U} that we call here \mathcal{U}^\natural . As a Hopf algebra, \mathcal{U} is extended to \mathcal{U}^\natural by adding another generator, T , which satisfies

$$T^2 = 1, \Delta T = T \otimes T, S(T) = T, \varepsilon(T) = 1.$$

A $*$ -involution on \mathcal{U}^\natural is defined as follows:

$$X^* = T X T, Y^* = T Y T, Z^* = T Z T, T^* = T.$$

So \mathcal{U} is a Hopf subalgebra of \mathcal{U}^\natural but not a $*$ -Hopf subalgebra. On the other hand, \mathcal{U} may be obtained from \mathcal{U}^\natural by specializing T to 1.

3 A class of representations of \mathcal{U}^\natural

Next we present a class of irreducible representations of the $*$ -algebra \mathcal{U}^\natural while the question of a complete classification of irreducible representations of \mathcal{U}^\natural is proposed as an open problem. Though it is not excluded that the definition of \mathcal{U}^\natural should be further modified in order to get a reasonable theory. In this section most steps are only outlined with some details omitted.

The representation ρ depends on an integer parameter $n \in \{1, 2, \dots, Q\}$ and its dimension d equals $Q + 1 - n$. The matrices $\rho(X)$, $\rho(Y)$, $\rho(Z)$ are tridiagonal with non-vanishing entries

$$\begin{aligned}\rho(Z)_{m-1,m} &= -(q^2 - q^{-2})q^{2m+n-1}a_m, \\ \rho(Z)_{m+1,m} &= (q^2 - q^{-2})q^{-2m-n-1}b_{m+1}, \\ \rho(Z)_{mm} &= (q + q^{-1})c_m,\end{aligned}\tag{3}$$

$$\begin{aligned}\rho(X)_{m-1,m} &= (q + q^{-1})a_m, \\ \rho(X)_{m+1,m} &= (q + q^{-1})b_{m+1}, \\ \rho(X)_{mm} &= d_m,\end{aligned}\tag{4}$$

$$\begin{aligned}\rho(Y)_{m-1,m} &= (q + q^{-1})q^{2(2m+n-1)}a_m, \\ \rho(Y)_{m+1,m} &= (q + q^{-1})q^{-2(2m+n+1)}b_{m+1}, \\ \rho(Y)_{mm} &= -d_m,\end{aligned}\tag{5}$$

$m = 0, 1, \dots, d - 1$. Here

$$\begin{aligned}a_m &= b_m \\ &= \frac{1}{q^{2m+n-1} + q^{-2m-n+1}} \sqrt{\frac{[m]_{q^2}[m+n-1]_{q^2}}{(q^{2m+n-2} + q^{-2m-n+2})(q^{2m+n} + q^{-2m-n})}}, \\ c_m &= \frac{q^{n-1} + q^{-n+1}}{(q^{2m+n-1} + q^{-2m-n+1})(q^{2m+n+1} + q^{-2m-n-1})}, \\ d_m &= \frac{q^{n-1} + q^{-n+1}}{(q^{2m+n-1} + q^{-2m-n+1})(q^{2m+n+1} + q^{-2m-n-1})} [2m+n]_q.\end{aligned}$$

The quantum numbers are defined as usual,

$$[x]_q = \frac{q^x - q^{-x}}{q - q^{-1}}.$$

The matrix $\rho(T)$ is diagonal,

$$\rho(Z) = \text{diag}(\tau_0, \tau_1, \dots, \tau_{d-1})$$

where $\tau_0 = 1$ and

$$\frac{\tau_m}{\tau_{m-1}} = \text{sgn} \left(\frac{[m]_{q^2} [m+n-1]_{q^2}}{(q^{2m+n-2} + q^{-2m-n+2})(q^{2m+n} + q^{-2m-n})} \right), \quad (6)$$

$m = 1, 2, \dots, d-1$.

Let us remark that a source of difficulties when working with real forms comes from the fact that the deformation parameter q is forced to be a complex unit. In that case the sign τ_m/τ_{m-1} in (6) may equal -1 for particular values of m . Concerning the representation ρ , it is worth mentioning that the matrix $\rho((qX - q^{-1}Y)Z^{-1})$ is diagonal and

$$\rho((qX - q^{-1}Y)Z^{-1})_{mm} = [2m+n]_q.$$

The verification of the commutation relations (2) is straightforward. This may be done even in the case when q is generic and the tridiagonal matrices (3), (4), (5) are infinite with $m = 0, 1, 2, \dots$. One then finds that relations (2) are satisfied if and only if the coefficients c_m obey a recursive equation,

$$(q^{2m+n+3} + q^{-2m-n-3})c_{m+1} - (q + q^{-1})(q^{2m+n} + q^{-2m-n})c_m + (q^{2m+n-3} + q^{-2m-n+3})c_{m-1} = 0,$$

and d_m and $a_m b_m$ are expressed in terms of c_m ,

$$\begin{aligned} d_m &= -\frac{(q^{2m+n-1} + q^{-2m-n+1})c_m - (q^{2m+n-3} + q^{-2m-n+3})c_{m-1}}{(q - q^{-1})^2}, \\ a_m b_m &= (q - q^{-1})^{-4} (q + q^{-1})^{-2} \\ &\quad \times (q^{2m+n} + q^{-2m-n})^{-1} (q^{2m+n-2} + q^{-2m-n+2})^{-1} \\ &\quad \times ((q^{2m+n+1} + q^{-2m-n-1})^2 c_m^2 + (q^{2m+n-3} + q^{-2m-n+3})^2 c_{m-1}^2 \\ &\quad - (q^2 + q^{-2})(q^{2m+n-3} + q^{-2m-n+3})(q^{2m+n+1} + q^{-2m-n-1})c_m c_{m-1} \\ &\quad + (q - q^{-1})^2). \end{aligned}$$

Equivalently,

$$d_m = -\frac{(q^{2m+n+3} + q^{-2m-n-3})c_{m+1} - (q^{2m+n+1} + q^{-2m-n-1})c_m}{(q - q^{-1})^2}.$$

To verify the irreducibility we shall show that even the restriction of ρ to the subalgebra \mathcal{U} is irreducible. This will become obvious as soon as we prove that ρ is equivalent to $\tilde{\rho}$ with

$$\tilde{\rho}(X) = \begin{pmatrix} 0 & 0 & \dots & 0 & 0 \\ x_1 & 0 & \dots & 0 & 0 \\ 0 & x_2 & \dots & 0 & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & \dots & x_{d-1} & 0 \end{pmatrix}, \quad \tilde{\rho}(Y) = \begin{pmatrix} 0 & y_1 & 0 & \dots & 0 \\ 0 & 0 & y_2 & \dots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \dots & y_{d-1} \\ 0 & 0 & 0 & \dots & 0 \end{pmatrix},$$

and

$$\tilde{\rho}(Z) = \begin{pmatrix} z_0 & 0 & \dots & 0 \\ 0 & z_1 & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & z_{d-1} \end{pmatrix},$$

where

$$x_j = q^{-n-2j+1}[n+j-1]_q, \quad y_j = [j]_q, \quad z_j = q^{-n-2j}.$$

Note that $x_j \neq 0$, $y_j \neq 0$, for $j = 1, \dots, d-1$.

The equivalence in turn follows from a more geometrical realization of the representation ρ which is closely related to the twisted adjoint action [7], [8]. The vector space \mathcal{M} of meromorphic functions in variable w on the complex plane becomes a left \mathcal{U} module with respect to the action

$$\begin{aligned} X \cdot f(w) &= -i \frac{q^{-1}w}{q - q^{-1}} (q^n f(w) - q^{-n} f(q^{-2}w)), \\ Y \cdot f(w) &= i \frac{q^{-n+1}}{(q - q^{-1})w} (f(w) - f(q^{-2}w)), \\ Z \cdot f(w) &= q^{-n} f(q^{-2}w). \end{aligned}$$

Set

$$\psi_m(w) = \frac{\prod_{j=0}^{m-1} (q^{2j+n}w - i)}{\prod_{j=0}^{n+m-1} (q^{2j-2m-n}w + i)}, \quad m = 0, 1, 2, \dots$$

Then the vector space $\text{span}\{\psi_0, \psi_1, \dots, \psi_{d-1}\}$, $d = Q+1-n$, may be checked to be \mathcal{U} invariant. After renormalization of the basis vectors, $\psi_m = \lambda_m \psi_m$, with the factors λ_m being determined by $\lambda_0 = 1$ and

$$\lambda_m = \sqrt{\frac{(q^{2m+n} + q^{-2m-n})[m+n-1]_{q^2}}{(q^{2m+n-2} + q^{-2m-n+2})[m]_{q^2}}} \lambda_{m-1},$$

we get the representation ρ .

Consider now a point set,

$$M = \{1, q^2, \dots, q^{2Q-2}\} \subset \mathbb{C}.$$

Note that for any function f , the values of the function $A \cdot f$ on the set M depend only on the restriction $f|_M$ where A is any of the generators X, Y or Z . Thus the vector space $\mathcal{F} \cong \mathbb{C}^Q$ of functions on M becomes a \mathcal{U} module and the restriction map $\mathcal{M} \rightarrow \mathcal{F} : f \mapsto f|_M$ is a surjective morphism of \mathcal{U} modules. The representation ρ corresponds to the submodule $\mathcal{R} = \text{span}\{\tilde{\psi}_0|_M, \tilde{\psi}_1|_M, \dots, \tilde{\psi}_{d-1}|_M\}$ with the distinguished basis. Omitting the details we claim that another basis in \mathcal{R} may be chosen as $\{\phi_0|_M, \phi_1|_M, \dots, \phi_{d-1}|_M\}$ where

$$\phi_j(w) = \left(\frac{1}{i}\right)^j q^{\frac{1}{2}j(j-1)+nj} w^{-Q+j}.$$

Expressing operators in the latter basis we get the representation $\tilde{\rho}$. This proves the equivalence of ρ and $\tilde{\rho}$ and consequently that the representation ρ is irreducible.

References

- [1] V. Chari, A. Pressley: *A Guide to Quantum Groups*. Cambridge University Press, Cambridge, 1994.
- [2] A. Klimyk, K. Schmüdgen: *Quantum Groups and Their Representations*. Springer-Verlag, Berlin Heidelberg, 1997.
- [3] A. Joseph: *Quantum Groups and Their Primitive Ideals*. Springer-Verlag, Berlin Heidelberg, 1995.
- [4] S.L. Woronowicz: *Lett. Math. Phys.* **23** (1991) 251.
- [5] V.K. Dobrev, P. Moylan: *Phys. Lett. B* **315** (1993) 292.
- [6] L.I. Korogodsky: *Commun. Math. Phys.* **163** (1994) 433.
- [7] P. Šťovíček: in *Proc. Internat. Sympos. Quantum Theory and Symmetries*, Goslar 1999 (Eds. H.-D. Doebner et al.), World Scientific, Singapore, 2000, p.470.

[8] P. Šťovíček: Lett. Math. Phys. **47** (1999) 125.