EXAMPLES IN CONCORDANCE

CHARLES LIVINGSTON

ABSTRACT. In this paper we present a series of examples of new phenomena in the classical knot concordance group. First we show that for (almost) every Seifert form there is an infinite family of knots, distinct in concordance, having that form. Next we demonstrate that a number of results that are known to hold in higher dimensional concordance fail in the classical case. These include: (1) examples of knots with Seifert forms that split as direct sums of Seifert forms but the knots are not concordant to corresponding connected sums, and (2) knots with Alexander polynomials that factor as products of Alexander polynomials (with resultant 1) but the knots are not concordant to corresponding connected sums. We also provide examples showing that: (3) for almost every metabolic Seifert form M and for every Seifert form V, there are knots with Seifert form $V \oplus M$ which are not concordant to knots with Seifert forms V and (4) there are pairs of irreducible algebraically concordant Seifert forms V and W such that there are knots with Seifert form V that are not concordant to any knot with Seifert form W.

In the course of proving these main results we also prove several new knot theoretic results. It has been known that the set of cyclic branched covers of a knot are all homology spheres if and only if the Alexander polynomial of the knot is trivial. Here we prove that the set of all *prime power* cyclic branched covers consists only of homology spheres if and only if every factor of the Alexander polynomial is a cyclotomic polynomial $\phi_n(t)$ with *n* divisible by three distinct primes. It has also been known that there are infinite sets of unit complex numbers for which the corresponding Levine–Tristram signature functions, $\sigma_{\omega}(K)$, are linearly independent. We show that, in fact, for any infinite set of unit complex numbers not containing a conjugate pair, the corresponding set of signature functions is linearly independent.

Contents

0

Inti	roduction.	- 2
1.	Knot Theory	7
2.	Concordance and Casson–Gordon Invariants	8
3.	Independence of the Signature Functions	10
4.	Iterated Satellite Knots	11
5.	Knots with all Prime Power Covers Homology Spheres	14
6.	Knots with the Same Seifert Forms	16
7.	Connected Sums	17
8.	Products of Alexander Polynomials	19
9.	Irreducible Knots with Reducible Seifert Forms	22
10.	Inequivalent Knots with Algebraically Equivalent, Irreducible Seifert	
	Forms	23

Date: November 16, 2018.

¹⁹⁹¹ Mathematics Subject Classification. 57M25, 57N70.

Key words and phrases. Concordance.

11. Conjectures and Addenda References

INTRODUCTION.

In 1969 Levine [Le1, Le2] presented an algebraic classification of the higher dimensional knot concordance group. A key element of his work was the proof that if the Seifert form of a knot is metabolic then the knot is slice. A few years later Casson and Gordon [CG1, CG2] demonstrated the distinction between classical knot concordance and higher dimensional concordance by constructing knots in S^3 that have metabolic Seifert forms but are not slice. However, Levine's results imply a number of other properties of higher dimensional concordance that have remained open in the classical setting. We will demonstrate the failure of several of these.

In Section 1 we will summarize the necessary background in knot theory. For now, recall that associated to a knot K in S^3 and a choice of Seifert surface for K, there is a Seifert form, V_K . In general, a Seifert form is an integer matrix Vsatisfying det $(V-V^t) = \pm 1$. An equivalence relation called *S*-equivalence is defined on the set of Seifert matrices [Tr2, Le3]; a basic result concerning S-equivalence states that if V_1 and V_2 are different Seifert matrices for a given knot, then $V_1 =_s V_2$ where $=_s$ denotes S-equivalence.

The Alexander polynomial of a Seifert form, $\Delta_V(t)$, is defined to be $\det(V - tV^t) \in \mathbf{Z}[t, t^{-1}]$ and the Alexander polynomial of a knot, $\Delta_K(t)$, is defined to be $\Delta_{V_K}(t)$; it is well defined up to multiplication by $\pm t^k$. We will normalize Alexander polynomials so that $\Delta_V(t) \in \mathbf{Z}[t, t^{-1}]$ and $\Delta_V(0) \neq 0$.

The resultant of two integral polynomials, Res(f, g), is an integer function of f and g; a precise definition is given in Section 5.

We will prove each of the following theorems for knots in dimension three. In each case the analogous higher dimensional statement is false. These theorems are discussed individually later in this introduction.

Theorem 0.1. Suppose that $\Delta_V(t)$ has a nontrivial irreducible factor that is not an n-cyclotomic polynomial with n divisible by three distinct primes. Then there is an infinite collection of knots $\{K_i\}$ with $V_{K_i} = V$ for all i, and no two K_i are concordant. In particular, if $\Delta_V(t)$ satisfies this property and V is metabolic, then there is an infinite collection of nonslice knots with Seifert form V, all of which are distinct up to concordance.

Theorem 0.2. There exists a knot K with $V_K = V_1 \oplus V_2$ but K is not concordant to a connected sum $K_1 \# K_2$ with $V_{K_i} =_s V_i$, i = 1, 2.

Theorem 0.3. There exists a knot K with $\Delta_K = \Delta_{V_1} \Delta_{V_2}$ and $Res(\Delta_{V_1}, \Delta_{V_2}) = 1$ but K is not concordant to a connected sum of knots, $K_1 \# K_2$, with $\Delta_{K_i} = \Delta_{V_i}$.

Theorem 0.4. There exists a knot K and a nontrivial Seifert form V that is algebraically concordant to V_K , but such that K is not concordant to any knot K_1 with $V_{K_1} =_s V$. More generally:

A: If M is a metabolic Seifert form and $\Delta_M(t)$ has a factor that is not a cyclotomic polynomial $\phi_n(t)$ with n divisible by three distinct primes, then for every Seifert form V there is a knot K with $V_K = V \oplus M$ but K is not concordant to a knot with Seifert form S-equivalent to V.

B: There exist algebraically concordant reduced Seifert forms V_K and V (eg. neither contains a metabolic summand) and a knot K with Seifert form V_K such that K is not concordant to a knot with Seifert form S-equivalent to V.

The technical condition on cyclotomic factors that appears in Theorem 0.1 is based on the following result which appears to be new and is of interest in its own right.

Theorem 0.5. A knot K has a prime power branched cover with nontrivial first homology if and only if $\Delta_K(t)$ has a nontrivial irreducible factor that is not a ncyclotomic polynomial with n divisible by three distinct primes.

In the course of proving these results it will be useful to know that the classical Tristram-Levine signature functions of knots are independent. The signature $\sigma_{\omega}(K)$ is defined to be the signature of the Hermitian matrix $(1-\omega)V_K + (1-\bar{\omega})V_K^t$ (with a slight modification if $\Delta_K(\omega) = 0$). Tristram and Levine both observed that there is an infinite set of ω for which the corresponding signature functions are linearly independent on the concordance group. We will show:

Theorem 0.6. For the set of unit complex numbers with positive imaginary part the corresponding set of signature functions is linearly independent.

0.1. Background of Results. We will be working in the topological locally flat category; all the results carry over to the smooth category. Basic results for knot theory can be found in [Ro, BZ]. For results on concordance references include [Le1, Le2, CG1, CG2]. An excellent survey of the knot theory we will be using appears in [Go]. In Sections 1 and 2 we will give a summary of the necessary results in classical knot theory, concordance, and Casson–Gordon invariants.

If F is a Seifert surface for a knot K then there is a Seifert form $V_K : H_1(F, \mathbf{Z}) \times H_1(F, \mathbf{Z}) \to \mathbf{Z}$ satisfying $\det(V - V_K) = \pm 1$. In general, an integral form V on a finitely generated free \mathbf{Z} -module H is called a Seifert form if $\det(V - V^t) = \pm 1$. We will usually not distinguish between a form and a matrix representation for that form.

There is an equivalence relation on the set of Seifert matrices called S-equivalence. It is generated by congruence of matrices and an algebraic enlargement of the matrix corresponding to the geometric construction of adding a hollow handle to the surface. It is essentially an observation of Murasugi [Mu] that any two Seifert matrices for a given knot are S-equivalent. Other references for S-equivalence include [Tr1, Tr2].

Definition 0.7. If Seifert forms V_1 and V_2 are S-equivalent we write $V_1 =_s V_2$.

The S-equivalence class of a Seifert form completely determines the homological properties of its cyclic branched and infinite cyclic covering spaces.

A Seifert form V is called *metabolic* if it vanishes on some half-dimensional summand $W \subset H$. Forms V_1 and V_2 are called *algebraically concordant* if $V_1 \oplus -V_2$ is metabolic. The set of algebraic concordance classes of Seifert forms is a group \mathcal{G} under direct sum called the *algebraic concordance group*, first defined by Levine in [Le1].

Knots K_1 and K_2 in are called *concordant* if there is a pair $(S^3 \times [0, 1], A)$ having boundary the disjoint union $(S^3, K_1) \coprod -(S^3, K_2)$, where A is homeomorphic to

 $S^1 \times [0,1]$. Equivalently, they are concordant if $K_1 \# - K_2$ is *slice*; that is, if it bounds an embedded disk in B^4 . The set of concordance classes of knots forms a group, C, with its operation induced by connected sum. This group was first defined by Fox in [F2]; see also [FM] for details.

Levine [Le1] proved that in the analogous higher dimensional setting of knotted 2n-1 spheres in S^{2n+1} the following result holds:

Theorem (Levine). The map $K \to V_K$ induces an injective homomorphism from C to $\mathcal{G}_{(-1)^n}$, n > 1.

(Levine defined groups $\mathcal{G}_{\pm 1}$ and we are abbreviating \mathcal{G}_{-1} simply by \mathcal{G} in this paper. He also proved that the map is surjective if n > 2 and is onto an index 2 subgroup if n = 2.)

Corollary. For n > 1, if the Seifert form of a knot $K \subset S^{2n+1}$ is metabolic then K is concordant to an unknot.

Casson and Gordon's proof [CG1, CG2] that Levine's theorem does not apply in the classical dimension consisted of finding examples to demonstrate that this corollary does not hold for n = 1:

Theorem (Casson and Gordon). There exists a knot $K \subset S^3$ with V_K metabolic but K is not concordant to the unknot.

Levine's theorem implies a number of other corollaries that we will see also do not apply in dimension three. Each of our theorems implies the Casson–Gordon theorem but is not a consequence of it. The distinguishing feature of our theorems is that to prove each one it is necessary to show that a given knot is not in any of an infinite family of concordance classes, all of which are algebraically identical. An added difficulty is that in most cases all that is known about that infinite family comes from its algebraic concordance class.

We will now restate each main result and discuss it briefly.

0.2. Distinct Knots with the Same Seifert Forms. Theorem 0.1 provides an indication of the complexity of the other main theorems. Since each theorem rules out knots being concordant to any knot with a particular Seifert form, it follows from Theorem 0.1 that we are having to deal with an infinite collection of concordance classes in each case.

Before discussing the general import of Theorem 0.1 we should describe the need for the technical condition on the Alexander polynomial. This condition arises from the result of Theorem 0.5.

Theorem 0.5 A knot K has a prime power branched cover with nontrivial first homology if and only if $\Delta_K(t)$ has a nontrivial irreducible factor that is not a ncyclotomic polynomial with n divisible by three distinct primes.

The principal tool for distinguishing knots that are algebraically concordant has been Casson–Gordon invariants. These can be applied only when some branched cover of prime power order has nontrivial homology. It is well known that for knots with trivial Alexander polynomial all covers have trivial homology, so this is clearly a class on which Casson–Gordon techniques do not apply. Of course, it has since been shown by Freedman [Fr, FQ], that if $\Delta_V(t) = 1$ then any knot with Seifert

4

form V is concordant to the unknot. Here we describe, in terms of Alexander polynomials, the exact class of knots for which the Casson–Gordon constraint on prime powers applies. Furthermore we show, via Theorem 0.1, that for any Alexander polynomial for which it is possible that Casson–Gordon methods apply we can in fact construct infinite families of knots for which they do apply.

Theorem 0.1. If $\Delta_V(t)$ has a nontrivial irreducible factor that is not an ncyclotomic polynomial with n divisible by three distinct primes, then there is an infinite collection of knots $\{K_i\}$ with $V_{K_i} = V$ for all i, and no two K_i are concordant. In particular, if $\Delta_V(t)$ satisfies this property and V is metabolic, then there is an infinite collection of nonslice knots with Seifert form V, all of which are distinct up to concordance.

Corollary. If V is metabolic and $\Delta_V(t)$ satisfies the given condition, then there is a nonslice knot K with $V_K = V$.

In general, in order to prove that an algebraically slice knot is not slice, it is necessary to analyze the full set of metabolizers for the Seifert form (or, as discussed below, metabolizers for linking forms on associated covering spaces). Hence, past examples have been drawn from situations in which the relevant metabolizers are easily listed. In contrast to this, we show that for every metabolic form an analysis is possible despite the possible complexity of its set of metabolizers.

The first examples of infinite families of algebraically slice knots that are distinct in concordance were presented in [J], though basic examples of this also follow quickly from the results of [CG1, CG2]. In these cases the relevant knots have distinct Seifert forms (in fact, the 2–fold branched covers have distinct homology groups) and this is key to distinguishing the knots up to concordance. In our setting similar methods cannot be applied.

0.3. Direct Sums of Seifert Forms. The result here is the following:

Theorem 0.2. There exists a knot K with $V_K = V_1 \oplus V_2$ but K is not concordant to a connected sum $K_1 \# K_2$ with $V_{K_i} =_s V_i$, i = 1, 2.

That such examples cannot exist in higher dimensions follows immediately from Levine's theorem. The only known invariants that can distinguish algebraically concordant knots up to concordance are Casson–Gordon style invariants, and it is initially not clear what properties of Casson–Gordon invariants are special for connected sums. We will see that it is an additivity property, discovered by Gilmer [G], that is essential here. We will again have to confront the fact that nothing is known concerning the knots K_i other than their Seifert forms.

It is interesting to note that according to [KLk, L] every knot is concordant to a prime knot.

0.4. Products of Alexander Polynomials. Here the result is:

Theorem 0.3. There exists a knot K with $\Delta_K = \Delta_{V_1} \Delta_{V_2}$ where the resultant satisfies, $\operatorname{Res}(\Delta_{V_1}, \Delta_{V_2}) = 1$ but K is not concordant to a connected sum of knots, $K_1 \# K_2$, with $\Delta_{K_i} = \Delta_{V_i}$.

Stoltzfus proved in [St, Corollary 6.5] that in higher dimensions Theorem 0.3 is false. The difficulty of Stolzfus' theorem is made clear by noting that the Alexander polynomial does not determine the algebraic concordance class of a knot and that

CHARLES LIVINGSTON

the factoring of the Alexander polynomial does not even imply the splitting of the Seifert form. Furthermore, the result fails without the condition on the resultant.

The difficulty here is much greater than in Theorem 0.2; there are many distinct Seifert matrices yielding the same Alexander polynomial, and it is not clear how the combination of being a connected sum and having a particular polynomial constrains the invariants used to show that knots are not concordant. By way of contrast, Theorem 0.2 will be proved using 2–fold branched covers and \mathbf{Z}_{3-} coefficients. The simplest example for Theorem 0.3 that we have found is based on 11–fold covers and \mathbf{Z}_{67-} coefficients.

0.5. **Reduction of Seifert Forms.** Our main result concerning the reduction of Seifert forms is the following:

Theorem 0.4. There exists a knot K and a nontrivial Seifert form V that is algebraically concordant to V_K , but such that K is not concordant to any knot K_1 with $V_{K_1} = V$. More generally:

- A: If M is a metabolic Seifert form and $\Delta_M(t)$ has a factor that is not a cyclotomic polynomial $\phi_n(t)$ with n divisible by three distinct primes, then for every Seifert form V there is a knot K with $V_K = V \oplus M$ but K is not concordant to a knot with Seifert form S-equivalent to V.
- **B:** There exist algebraically concordant reduced Seifert forms V_K and V (eg. neither contains a metabolic summand) and a knot K with Seifert form V_K such that K is not concordant to a knot with Seifert form S-equivalent V.

The only notation in the statement of Theorem 0.4 that may be unfamiliar is the definition of reduced. The precise definition as given in [Ke] is the following: a Seifert form is reduced if it is nonsingular and there is no proper submodule on which $V + V^t$ vanishes that is invariant under $V^{-1}V^t$. It is proved in [Ke] that every Seifert form is concordant to a reduced form. The same result appears in [Le2, Lemma 13] where it is shown that any Seifert form is equivalent to one that contains no proper totally isotropic subspace. Hence, in this language, every Seifert form is concordant to an anisotropic form. As a special case, it follows immediately that a Seifert form on a rank two **Z**-module is reduced if and only if it is not metabolic; that is, if and only if there is no nonzero element on which the form V vanishes. In general, forms with metabolic summands are reducible.

Notice that without the assumption of nontriviality in the statement of Theorem 0.4, Casson and Gordon's example satisfies condition **A**. Again, in that setting it was sufficient to show that some particular knot is not concordant to the unknot. As in the previous examples, here we have to show that a given knot does not lie in each of an infinite collection of concordance classes. We will see that the construction of Theorem 0.4 **A** is one of the simpler ones that we face; the presence of the metabolic summand greatly aids the construction.

For Theorem 0.4 **B** the proof is harder. Simply distinguishing reduced but algebraically concordant Seifert forms can be nontrivial. Here we will take advantage of the fact that the linking forms on covers associated with knots with given reduced Seifert forms can be different, though these too will be anisotropic and stably equivalent.

Notice that if W is a metabolic form then for any knot K, K is concordant to a knot K_1 with $V_{K_1} = V_K \oplus W$. Simply let $K_1 = K \# K'$ where K' is a slice knot with $V_{K'} = W$. Such a K' is easily constructed.

0.6. **Outline.** Section 1 presents the basic results from classical knot theory that we will be using, along with the relevant notation. We will review concordance and Casson–Gordon invariants in Section 2. In Section 3 we prove Theorem 0.6 showing that the signature functions are all linearly independent. A key result used for building knots with specified Casson–Gordon invariants is proved in Section 4. The construction depends on an iterated companionship argument.

Section 5 presents the proof of Theorem 0.5 stating that a knot has a prime power cover with nontrivial first homology if and only if some irreducible factor of the Alexander polynomial is not a cyclotomic polynomial $\phi_n(t)$ with *n* divisible by three distinct primes. This is used in Section 6 to prove Theorem 0.1 stating that most Seifert forms are realized by an infinite family knots that are distinct in concordance.

Section 7 gives the construction of examples to demonstrate Theorem 0.2. Alternative examples could be built using the knots constructed in Section 8 to prove Theorem 0.3. We include the examples of Section 7 since they are fairly explicit whereas those of 8 are not, and they also provide useful concrete examples needed to understand the abstract constructions of the later sections. As mentioned earlier, the added complexity of these later examples over those used in Section 7 to prove Theorem 0.3 is readily apparent: in Section 7 we can work with 2–fold covers and \mathbf{Z}_3 –valued characters; in Section 8 we must work with an 11–fold cover and \mathbf{Z}_{67} –valued characters. Sections 9 and 10 are devoted to proving Theorem 0.4.

Thanks are due to Pat Gilmer for helpful conversations regarding this paper. Jim Davis, Darrell Haile, and Michael Larsen assisted me with Theorem 0.5.

1. KNOT THEORY

In this section we will review basic results from classical knot theory. Our goal is to present the basic notation and background needed for the rest of our work. More detailed presentations can be found in the references [BZ, Go, Ro].

We will be working in the category of locally flat oriented manifolds and pairs throughout this paper; the results all apply in the smooth category.

1.1. Knots and Branched Covering Spaces. A knot is formally defined to be the oriented homeomorphism class of an oriented pair (S, K), where S is homeomorphic to S^3 and K is homeomorphic to S^1 . As is standard, we will abbreviate an equivalence class of a pair (S, K) by simply K and assume that $S = S^3$. This definition of knot is equivalent to the definition that states that a knot is the isotopy class of an oriented connected 1-submanifold of S^3 . Again, we are working in the locally flat category, but for classical knot theory this can be shown to be equivalent to the smooth category. The same is not true for concordance.

For a knot K we will denote its q-fold cyclic branched cover and its infinite cyclic cover by $M_{K,q}$ and $M_{K,\infty}$, respectively. The group of deck transformations is generated by an homeomorphism denoted $T_{K,q}$ or $T_{K,\infty}$.

is generated by an homeomorphism denoted $T_{K,q}$ or $T_{K,\infty}$. A Seifert surface for (S^3, K) is an oriented surface $F \subset S^3$ with $\partial F = K$. Associated to F and a choice of basis for $H_1(F, \mathbb{Z})$ there is a Seifert matrix V_K . It satisfies the basic property that $\det(V_K - V_K^t) = 1$. More generally, any matrix satisfying this condition on the determinant will be called a Seifert matrix.

The preimage of $S^3 - F$ in the infinite cyclic cover, $M_{K,\infty}$, has an infinite set of components, $\{X_i\}_{i=-\infty,\dots,\infty}$. A Mayer–Vietoris argument shows that $H_1(M_{K,\infty}, \mathbb{Z})$ is generated by the homology groups of the X_i . Furthermore, as a $\mathbb{Z}[t, t^{-1}]$ –module

 $H_1(M_{K,\infty}, \mathbf{Z})$ is generated by $H_1(X_0, \mathbf{Z})$. The Mayer–Vietoris argument shows that with respect to the appropriate basis of $H_1(X_0, \mathbf{Z})$ the homology is presented by the matrix $V_K - tV_K^t$.

The homology of the q-fold cyclic cover of $S^3 - K$ is a $\mathbf{Z}[\mathbf{Z}_q]$ module; since $1+T_{K,q}+T_{K,q}^2+\cdots+T_{K,q}^{q-1}$ acts trivially on $H_1(M_{K,q},\mathbf{Z})$ it follows that $H_1(M_{K,q},\mathbf{Z})$ is a $\mathbf{Z}[\zeta_q]$ module, where ζ_q is a primitive q-root of unity.

The Alexander polynomial of K is defined to be $\Delta_K(t) = \det(V_K - tV_K^t)$. More generally, for any Seifert matrix V, the Alexander polynomial is defined to be $\Delta_V(t) = \det(V - tV^t)$.

Two basic theorems we will be using are the following. The second may be less well known; it was first proved by Fox [F1] and a proof can be found in [BZ].

Theorem 1.1. Multiplication by $\Delta_K(t)$ annihilates $H_1(M_{K,\infty}, \mathbf{Z})$. Multiplication by $\Delta_K(\zeta_q)$ annihilates $H_1(M_{K,q}, \mathbf{Z})$.

Theorem 1.2. $|H_1(M_{K,q}, \mathbf{Z})| = \prod_{i=1,...,q-1} \Delta(\zeta_q^i).$

Corollary 1.3. If q is a prime power, then $M_{K,q}$ is a rational homology sphere: $|H_1(M_{K,q}, \mathbf{Q})| = 0.$

Proof. This is usually viewed as a consequence of the Milnor exact sequence [M]; such a proof is contained in [CG1]. The result also follows immediately from Theorem 1.2. To see this, suppose that $\Delta_K(\zeta_q^i) = 0$ for some prime power $q = p^r$. Then the q-cyclotomic polynomial, $\phi_q(t) = (t^{p^r} - 1)/(t^{p^{r-1}} - 1)$ would divide $\Delta(t)$. But $\phi_q(1) = p$ by L'Hopital's rule, while $\Delta(1) = 1$.

1.2. Linking Forms. Let M be a rational homology sphere. There is a linking form lk: $H_1(M, \mathbb{Z}) \times H_1(M, \mathbb{Z}) \to \mathbb{Q}/\mathbb{Z}$. This pairing is symmetric and is nonsingular, meaning that the induced map $H_1(M, \mathbb{Z}) \to \operatorname{Hom}(H_1(M, \mathbb{Z}), \mathbb{Q}/\mathbb{Z})$ is an isomorphism. Using this we will identify class of order p^n in $H_1(M, \mathbb{Z})$ with characters on $H_1(M, \mathbb{Z})$ taking value in \mathbb{Z}_{p^n} .

2. Concordance and Casson-Gordon Invariants

References for details regarding the general theory of concordance include [Le1, Le2]. References for the theory of Casson–Gordon invariants include the original papers [CG1, CG2] and also [G, GL]

2.1. **Concordance.** A knot (S, K) is called *slice* if it bounds a proper pair (B, D) where B is homeomorphic to B^4 and D is homeomorphic to B^2 . Knots (S_1, K_1) and (S_2, K_2) are called *concordant* if the connected sum $(S_1, K_1)\# - (S_1, K_2)$ is slice, where the connected sum is as usual for oriented pairs. The set of concordance classes forms an abelian group called the *concordance group*, C, under the operation induced by connected sum.

Alternatively, knots (S_1, K_1) and (S_2, K_2) knots are called concordant if there is a pair, $(S^3 \times [0, 1], A)$, where A is homeomorphic to $S^1 \times [0, 1]$ and $\partial(S^3 \times [0, 1], A) \cong (S_1, K_1) \prod -(S_2, K_2)$.

(In working in the smooth category there is a similarly defined concordance group, C_s . Since smooth submanifolds are locally flat, there is a homomorphism $C_s \to C$ that is surjective. It is not an isomorphism.)

2.2. Algebraically Slice Knots and Algebraic Concordance. The basic result concerning slice knots is the following.

Theorem 2.1. If a knot K is slice then for any Seifert surface F its Seifert form is metabolic; that is, there is a half-dimensional summand of $H_1(F, \mathbb{Z})$ on which the Seifert form vanishes.

A knot is called *algebraically slice* if its Seifert from is metabolic. A Seifert matrix is called *algebraically slice* if it is metabolic. Two Seifert matrices, V and W, are called *algebraically concordant* if $V \oplus -W$ is algebraically slice. This defines an equivalence relation on the set of Seifert forms and the operation of direct sum places a group structure on the set of algebraic concordance classes. This group is called Levine's *algebraic concordance group* and is denoted \mathcal{G} .

The following is a theorem of Levine [Le1].

Theorem 2.2. The map $K \to V_K$ induces a well defined surjective homomorphism from C to G.

2.3. Signature Functions. Let W(t) be a nonsingular matrix with coefficients in $\mathbf{C}(t)$, Hermitian with respect to complex conjugation and the involution $t \to t^{-1}$. There is a Witt group of such matrices, $W(\mathbf{C}(t))$. For a unit complex number ω one has the signature $\sigma(W(\omega))$. Although W(t) is assumed nonsingular, $W(\omega)$ may be singular, and hence, even if W(t) is metabolic, $\sigma(W(\omega))$ may be nonzero. For this reason, taking signatures at unit numbers ω is not a well defined function on the Witt group $W(\mathbf{C}(t))$. However, the limit $\lim_{s\to 0} (\sigma(W(\omega e^{si}) + \sigma(W(\omega e^{-si}))/2 \text{ does yield a well defined signature function, <math>\sigma_{\omega}$, on $W(\mathbf{C}(t))$.

Definition 2.3. The Tristram-Levine signature function of a knot K, $\sigma_{\omega}(K)$, is defined to be the signature function of the form $V_K(t) = (1-t)V_K + (1-t^{-1})V_K^t$.

We will usually be interesting in evaluating this function at roots of unity, $\omega = e^{2\pi i \frac{p}{q}}$. To simplify notation we have:

Definition 2.4. $\sigma_{p/q}(K) = \sigma_{\omega}(K)$ where $\omega = e^{2\pi i \frac{p}{q}}$.

It is worth noting at this point the following result.

Theorem 2.5. The Hermitian form $V_K(\omega)$ with ω a unit complex number is singular only if $\omega = 1$ or $\Delta_K(\omega) = 0$.

Proof. We can write

$$(1-\omega)V_K + (1-\omega^{-1})V_K^t = (1-\omega)(V_K + \frac{(1-\omega^{-1})}{(1-\omega)}V_K^t)$$

The coefficient on V_K^t equals $-\omega^{-1}$, The determinant is hence $(1 - \omega)^{2g} \Delta_K(\omega^{-1})$, where 2g is the dimension of V_K . Since the Alexander polynomial is symmetric, this determinant is 0 if and only if $\omega = 1$ or ω is a root of the Alexander polynomial.

2.4. **Casson–Gordon Invariants.** Let K be a knot in S^3 with q-fold branched cover $M_{K,q}$, q a prime power. Let $\chi: H_1(M_{K,q}, \mathbb{Z}) \to \mathbb{Z}_{p^r}$ be a homomorphism where p is a prime. Casson and Gordon [CG1] defined an invariant, denoted $\sigma_1 \tau(K, \chi) \in \mathbb{Q}$. To simplify notation we will define:

Definition 2.6. $\sigma(K, q, \chi) = \sigma_1 \tau(K, \chi).$

The precise definition of $\sigma(K, q, \chi)$ will not be needed here; we now summarize the essential properties that we will be using.

Recall that the generator of the group of deck transformations of $M_{K,q}$ is denoted $T_{K,q}$; $T_{K,q}$ is of order q. The linking form on $H_1(M_{K,q}, \mathbb{Z})$ is invariant under the group of deck transformations. The main result of [CG1] is the following.

Theorem 2.7. If K is slice there is a subgroup $H \subset H_1(M_{K,q}, \mathbb{Z})$ with $|H|^2 = |H_1(M_{K,q}, \mathbb{Z})|$ and such that for any character χ with values in \mathbb{Z}_{p^r} that vanishes on H, $\sigma(K, q, \chi) = 0$. Furthermore, it can be assumed that H is a metabolizer (self-annihilating) for the linking form on $M_{K,q}$ and that H is invariant under the action of $T_{K,q}$.

The observation that H is $T_{K,q}$ invariant is not made in [CG1]. However, in the proof given in [CG1], H is defined to be the kernel of the inclusion of the homology of $M_{K,q}$ into a 4-manifold over which the action of $T_{K,q}$ extends. Hence, the invariance of H follows.

A simple result concerning the Casson–Gordon invariant is that $\sigma(K, q, \tau) = \sigma(K, q, -\chi)$. This follows from the fact that signatures are unchanged under complex conjugation. A much deeper result is the additivity result proved by Gilmer [G], as we now describe. If $K = K_1 \# K_2$, then $H_1(M_{K,q}, \mathbf{Z}) = H_1(M_{K_1,q}, \mathbf{Z}) \oplus H_1(M_{K_2,q}, \mathbf{Z})$ and any character χ on $H_1(M_{K,q}, \mathbf{Z})$ can be written as $\chi = \chi_1 \oplus \chi_2$ with χ_i a character on $H_1(M_{K_i,q}, \mathbf{Z})$.

Theorem 2.8. $\sigma(K, q, \tau) = \sigma(K_1, q, \chi_1) + \sigma(K_2, q, \chi_2).$

Another basic result concerning the Casson–Gordon invariant concerns its value at the trivial character, $\sigma(K, q, 0)$. The result was first proved by Litherland [Lt2, Corollary B2].

Theorem 2.9. $\sigma(K, q, 0) = 0$.

3. INDEPENDENCE OF THE SIGNATURE FUNCTIONS

3.1. Independence of the Signature Functions. To say that an infinite set of elements in Hom(\mathcal{G}, \mathbf{Z}) is independent means that no nontrivial finite linear combination of these elements is trivial. The independence of the signature functions $\{\sigma_{\omega}\}$ for an infinite set of ω was observed independently by Tristram and Levine [Le1, T]; in a different guise this appears in [M]. It will be useful later to have a more general result.

Theorem 3.1. The set of functions $\{\sigma_{\omega}\}_{\omega \in U_+}$ with U_+ the set of all unit complex numbers with positive imaginary parts forms a linearly independent set in $Hom(\mathcal{G}, \mathbf{Z})$ (and hence in $Hom(\mathcal{C}, \mathbf{Z})$).

This theorem is clearly equivalent to the following corollary, which we prove.

Theorem 3.2. Let $\{\omega_i\}$ be a finite set of unit complex numbers with $Im(\omega_i) > 0$ for all *i* and $Re(\omega_i) \neq Re(\omega_j)$ for $i \neq j$. The set of functions $\{\sigma_{\omega_i}\}$ is linearly independent in $Hom(\mathcal{G}, \mathbf{Z})$ (and hence in $Hom(\mathcal{C}, \mathbf{Z})$).

Proof. By a result of Matumoto [Ma] (see also [Lt2]), the signature function has jumps only at roots of the Alexander polynomial. Furthermore, if the root is of multiplicity one the jump is ± 2 .

We will show that there is a set of unit complex numbers S which is dense on the unit circle and such that for all $\eta \in S$ there is an Alexander polynomial having

10

 η and $\bar{\eta}$ as its only complex roots, both with multiplicity one. For a knot K_{η} with such an Alexander polynomial, the signature function $\sigma_{\omega}(K_{\eta})$ is 0 for $\operatorname{Re}(\omega) > \operatorname{Re}(\eta)$ and is ± 2 for $\operatorname{Re}(\omega) < \operatorname{Re}(\eta)$; we can assume it is positive in this case by taking the mirror image if need be. For a given ω_i , choose η_1 and η_2 so that the interval on the unit circle with endpoints η_1 and η_2 contains ω_i , and no other ω_j . The knot $K = K_{\eta_2} \# - K_{\eta_1}$ thus satisfies $\sigma_{\omega_i}(K) = 2$ and $\sigma_{\omega_j}(K) = 0$ for all other $\omega_j \in S$. The desired linear independence of the signature functions σ_{ω_i} follows.

Because an Alexander polynomial $\Delta(t)$ is symmetric and real, if τ is a root then so are $\bar{\tau}, 1/\tau$, and $1/\bar{\tau}$. It follows that if Δ is of degree four and has two complex (nonreal) roots and two real roots, then the complex roots must be unit, of multiplicity 1.

Consider the polynomial

$$F_r(t) = (t-1)^2(t^2 - 2rt + 1) = t^4 + (-2 - 2r)t^3 + (4r+2)t^2 + (-2 - 2r)t + 1.$$

It is easily seen that $F_r(t)$ has a pair of complex roots with real part r in the case -1 < r < 1. For a given ϵ there is a δ such that a perturbation of the coefficients of F_r by less the δ moves the roots less than ϵ . Choose a rational approximation a/b, (b > 0) to r so that replacing r in the coefficients of F_r by a/b changes the coefficients by less than $\delta/2$. Furthermore, choose b large enough so that $1/b < \delta/2$. Then the roots of

$$G(t) = t^4 + (-2 - 2\frac{a}{b})t^3 + (4\frac{a}{b} + 2 - \frac{1}{b})t^2 + (-2 - 2\frac{a}{b})t + 1$$

are within ϵ of those of F_r . Multiplying through by b yields the polynomial

$$\Delta(t) = bt^4 + (-2b - 2a)t^3 + (4a + 2b - 1)t^2 + (-2b - 2a)t + b.$$

Since this is a symmetric integral polynomial and $\Delta(1) = -1$, it is an Alexander polynomial whose roots are within ϵ of those of $F_r(t)$. Furthermore, since $\Delta(1) = -1$, $\Delta(t)$ has at least two real roots. Hence, the remaining roots are conjugate unit complex numbers with real part within ϵ of r. The result follows.

We will need to use knots with large signature functions in later arguments. The following corollary provides the needed examples.

Corollary 3.3. For any prime p and constant C there is a knot J for which the signatures $\sigma_{k/p}(J) > C$ for all k, 0 < k < p.

Proof. The proof is immediate from Theorem 3.2.

4. Iterated Satellite Knots

4.1. **Basic Results.** Let $L = U \coprod L_1$ be a two component link in S^3 with U the unknot. Let K be a knot in S^3 . The satellite of K, K^* , with embellishment L_1 is constructed as follows: form the union $(S^3 - N^0(K)) \cup (S^3 - N^0(U))$ where N^0 denotes an open tubular neighborhood and the union identifies the meridian of U with the longitude of K and the longitude of U with the meridian of K. The image of L_1 in this union is denoted K^* . Furthermore, K is called the *companion* of K^* .

In [Lt2] Litherland described the relationship between the Casson–Gordon invariants of a satellite knot and those of its component parts. A slight reformulation of the construction of satellite knots will facilitate the use of Litherland's results under iteration. Further details are presented in [GL]. We will describe these results only in the special case that we will be needing. Suppose that L is an unknotted circle in S^3 in the complement of K and that L is null homologous in $S^3 - K$. (More precisely, we only need that L is trivial in $H_1(S^3 - K, \mathbb{Z}_q)$.) Let J be a knot in S^3 . Construct the union $(S^3 - N^0(L)) \cup (S^3 - N^0(J))$ where the union if formed via a homeomorphism that interchanges meridians and longitudes. The resulting manifold is easily seen to be homeomorphic to S^3 : it is the union of a knot complement with a solid torus. The image of K in the union represents a perhaps different knot, say K^* in S^3 . A fairly simple geometric argument shows that K^* is a satellite of J with embellishment K, and hence that J is a companion of K^* . Intuitively, the effect of this construction is that the portion of K that passes through L is tied into the knot J.

To understand the effect of this construction on covers and the associated invariants of the knots, let $M_{K,q}$ be the q-fold branched cover of K. The unknotted curve L lifts to a set of distinct curves, $\{\tilde{L_0}, T_{K,q}(\tilde{L_0}), T_{K,q}^2(\tilde{L_0}), \ldots, T_{K,q}^{q-1}(\tilde{L_0})\} =$ $\{\tilde{L_i}\}_{i=0...q-1}$, in $M_{K,q}$. Thus, $M_{K^*,q}$ is constructed from $M_{K,q}$ by removing neighborhoods of each $\tilde{L_i}$ and replacing them with copies of the complement of J. Since we are simply removing a homology circle and replacing it with another homology circle via maps that are homologically identical, the construction leaves the homology of the space unchanged and there is a natural correspondence between the homology, and cohomology, groups of $M_{K,q}$ and $M_{K^*,q}$. In particular we can identify a character χ on $H_1(M_{K^*,q}, \mathbb{Z})$ with a character also denoted χ on $H_1(M_{K,q}, \mathbb{Z})$. In this situation the Casson–Gordon invariants of K^* are related to the Tristram– Levine signatures of J, as described in [Lt2] and [GL].

Theorem 4.1. $\sigma(K^*, \chi) = \sigma(K, \chi) + \sum_{i=0}^{q-1} \sigma_{\chi(\tilde{L}_i)/p}(J).$

In the case of q = 2 one has that $T_{K,2}$ acts by multiplication by -1 on homology and hence we have:

Corollary 4.2.
$$\sigma(K^*, \chi) = \sigma(K, \chi) + 2\sigma_{\chi(\tilde{L_0})/p}(J)$$
.

4.2. Iterated Companionship. If the knot L in Theorem 4.1 is replaced with a link $\{L^j\}_{j=1...k}$ in $S^3 - K$ which forms the unlink in S^3 and for which each component is trivial in $H_1(S^3 - K, \mathbb{Z}_q)$ then via iteration we have, using similar notation and assuming that the neighborhood of L^j is replaced with the complement of J_j we have the following result.

Theorem 4.3. $\sigma(K^*, \chi) = \sigma(K, \chi) + \sum_{j=1}^k \sum_{i=0}^{q-1} \sigma_{\chi(\tilde{L}_i^j)/p}(J_j).$

Notice that if in the above construction each component of L is in the complement of a fixed Seifert surface F for K (and in particular is trivial in $H_1(S^3 - K)$), then F represents a Seifert surface for K^* , which we will denote F^* , and that for the Seifert forms constructed via these surfaces we have $V_K = V_{K^*}$.

In our constructions we will need to control the class represented by components of \tilde{L}_i^j in $H_1(M, \mathbb{Z})$. To do this we must start with the classes themselves and build L. The following theorem gives situations in which this can be done.

Theorem 4.4. (A) Let $\{\tilde{\gamma}^j\}$ be a collection of homology classes in $H_1(M_{K,q}, \mathbb{Z})$. Each γ^j can be represented by a simple closed curve which is one of the q components of the preimage of an unknotted curve γ^j in $S^3 - K$. The γ^j can be chosen to form an unlink in S^3 with each component trivial in $H_1(S^3 - K, \mathbb{Z})$.

(B) If F is a Seifert surface for K and each $\tilde{\gamma}^j$ can be represented by a curve in $M_{K,q} - p^{-1}(F)$, then it can furthermore be arranged that each component γ^j is in $S^3 - F$.

Proof. Represent each class by a curve, which we will denote $\tilde{\gamma}^j$ also. By transversality we can assume that the entire collection $\{T^i(\tilde{\gamma}^j)\}_{i=0...q-1, j=1...k}$ consists of disjoint embedded curves. If the $\tilde{\gamma}^j$ are in the complement of $p^{-1}(F)$, then the same will be true after the small homotopy is performed to achieve transversality.

With this, the collection $\{\gamma^j\}$ forms a link in $S^3 - K$, and since the preimage of each component consists of q components, each element is trivial in $H_1(M_{K,q}, \mathbf{Z}_q)$. Furthermore, if we were working in the complement of the preimage of the Seifert surface, then $\{\gamma^j\}$ is in the complement of the Seifert surface; in particular we will have in this case that the γ^j are trivial in $H_1(M, \mathbf{Z})$.

A series of crossing changes in the γ^j results in an unlink. These crossing changes can be performed (in S^3) by a homotopy, and this homotopy can be performed in the complement of the 1-skeleton of the Seifert surface. But since the Seifert surface itself isotopes into a small neighborhood of its 1-skeleton, the homotopy that makes $\{\gamma^j\}$ into an unlink can take place in the complement of F, and in particular in the complement of K. If follows that the classes in $H_1(S^3 - K, \mathbb{Z})$ are unchanged. The homotopy lifts to one of the $\tilde{\gamma}^j$ so the homology classes represented by these is unchanged.

This construction may appear clearer if presented in terms of diagrams. After an isotopy the surface F can be placed in a small neighborhood of its 1–skeleton. The crossings in the diagram of $\{\gamma^j\}$ all can then be assumed to occur at points that are far from F. Hence, a change in the crossings of that diagram can be performed in the complement of F as well.

It remains to assure that the γ^j are trivial in $H_1(S^3 - K, \mathbf{Z})$, not just in $H_1(S^3 - K, \mathbf{Z}_q)$. In the case that the curves are disjoint from the Seifert surface, Case **B**, this is automatic. If any γ^j is nontrivial in homology, the figure below indicates a move that can be made on a curve γ^j that changes the homology class represented by γ^j in $H_1(S^3 - K, \mathbf{Z})$ by q but doesn't change the homology classes of the preimages of γ^j in M. (In the figure, q = 3.) The effect of this move is to have the $\tilde{\gamma}^j$ pass through the branch set. This move can be used to assure that the γ^j are trivial in $H_1(M, \mathbf{Z})$.



Figure 1. Inserting twists

Theorem 4.5. Under the conditions of Theorem 4.4 we have that $\Delta_{K^*} = \Delta_K$. Under the conditions of part (**B**) of Theorem 4.4 we further have that $V_{K^*} = V_K$ for the Seifert surface F. *Proof.* The Alexander polynomial of a satellite knot is determined by that of its companion and its embellishment. This was first proved by Seifert [Se], and a modern proof based on infinite cyclic covering spaces follows from [LM]. In the case of winding number 0 the formula states simply that the Alexander polynomial of the satellite knot is equal to the polynomial of the embellishment.

In the case that the link $\{L^j\}$ is in the complement of F, the Seifert form is unaffected by the satellite construction.

4.3. Knots with Positive Casson–Gordon Invariants. By applying the results of the previous section, the following result is easily proved.

Theorem 4.6. For every Seifert form V, positive number C, and prime power q, there is a knot K with $V_K = V$ and $\sigma(K, q, \chi) > C$ for all nontrivial χ .

Proof. Pick a knot K with $V_K = V$. Observe that the finite set of values of $\sigma(K, q, \chi)$ is bounded: say $-C_0 < \sigma(K, q, \chi) < C_0$ for all χ . Pick a set of generators for $H_1(M_{K,q}, \mathbb{Z})$ that can be represented by curves in the complement of the preimage of the Seifert surface for K (having Seifert form V). Next, apply Theorem 4.4 to construct the unlink $\{\gamma^j\}$. Replace each component of this collection of curves with the complement of a knot J for which all relevant p-signatures are greater than $C + C_0$. Any nontrivial χ must evaluate nontrivially on one of the associated Casson–Gordon invariant has increased by at least $C + C_0$, making its value at least C.

5. KNOTS WITH ALL PRIME POWER COVERS HOMOLOGY SPHERES

Our goal in this section is to prove Theorem 0.5:

Theorem 0.5. A knot K has a prime power branched cover with nontrivial first homology if and only if $\Delta_K(t)$ has a nontrivial irreducible factor that is not a ncyclotomic polynomial with n divisible by three distinct primes.

The proof depends on properties of the resultant of polynomial. The resultant reappears in Section 8. For this reason we will present a general summary of properties of the resultant in the first section below. A good reference for the resultant is [La].

5.1. **Resultants of Polynomials.** The resultant $\operatorname{Res}(f, g)$ is defined for integral polynomials as follows. Let $f(x) = \sum_{i=0...n} a_i x^i$ and $g(x) = \sum_{j=0...m} b_j x^j$ with $a_n \neq 0 \neq b_m$. Let the complex roots of f be $\{s_i\}_{i=1...n}$ and let g have roots $\{t_i\}_{i=1...n}$, in both cases appearing with multiplicity. With this we have:

Definition 5.1. Res $(f,g) = a_n^m b_m^n \prod_{i,j} (s_i - t_j).$

Notice that $\operatorname{Res}(f,g) = \pm \operatorname{Res}(g,f)$. By factoring f and g over \mathbb{C} one quickly attains two useful formula, in which this symmetry is not longer apparent.

Lemma 5.2. $b_m^n \prod_j f(t_j) = \operatorname{Res}(f,g) = \pm a_n^m \prod_i g(s_i).$

One of the main results concerning resultants is the following theorem. Proofs are given in [La] as well as in most other texts in advanced algebra.

Theorem 5.3. For polynomials f and g as above, there exist integral polynomials $\phi(x)$ and $\psi(x)$ such that $\phi(x)f(x) + \psi(x)g(x) = \operatorname{Res}(f,g)$.

Corollary 5.4. Res(f,g) is an integer.

Proof. From its definition the resultant is a complex number. On the other hand, the sum $\phi(x)f(x) + \psi(x)g(x)$ is an integral polynomial.

Theorem 5.5. If g (or f) is monic then there exist integral polynomials $\phi(x)$ and $\psi(x)$ such that $\phi(x)f(x) + \psi(x)g(x) = 1$ if and only if $\operatorname{Res}(f,g) = 1$.

Proof. By Theorem 5.3 one direction follows immediately, without the assumption of f or g being monic. To prove the other direction, assume g is monic and suppose that $\phi(x)f(x) + \psi(x)g(x) = 1$. Then $\prod_j \phi(t_j)f(t_j) + \psi(t_j)g(t_j) = 1$. Since the t_j are roots of g, this gives that $\prod_j \phi(t_j)f(t_j) = 1$. Since $b_m = 1$, this gives that $\operatorname{Res}(\phi, g)\operatorname{Res}(\psi, g) = 1$. Since both are integers, the result follows.

5.2. Homology Groups of Prime Power Covers. In this section we will prove Theorem 0.5, explaining the constraint on the Alexander polynomial given in many of the later results.

Proof of Theorem 0.5. The argument consists of a series of observations that reduce the result to a purely number theoretic lemma that follows.

- 1. According to Riley [Ri] the order of the homology of the k-fold cyclic branched cover of a knot K grows exponentially if the Alexander polynomial has a root that is not a root of unity. Hence, we only need to consider the case that all irreducible factors of the Alexander polynomial are cyclotomic polynomials, $\phi_n(t)$.
- By Fox's theorem, Theorem 1.2, the order of the homology of the p^k-branched cover of a knot K is given by Π_{i=0...p^k-1} Δ_K(ωⁱ_{p^k}), where ω_{p^k} is a primitive p^k-root of unity. This reduces the general proof to the case in which Δ_K(t) is irreducible, and hence to the case Δ_K(t) = φ_n(t) for some n.
 Recall that for p prime, φ_{p^k}(t) = (t^{p^k} 1)/(t^{p^{k-1}} 1). By L'Hopital's rule it
- 3. Recall that for p prime, $\phi_{p^k}(t) = (t^{p^k} 1)/(t^{p^{k-1}} 1)$. By L'Hopital's rule it follows that $\phi_{p^k}(1) = p$. For a knot K, $\Delta_K(1) = \pm 1$, and hence we only need to consider the case of $\Delta_K(t) = \phi_n(t)$ with n divisible by at least two distinct primes.
- 4. We now have that the order of the homology of the p^k -fold branched cover of K is given by the product of $\phi_n(t)$ evaluated at all (not necessarily primitive) p^k -roots of unity; that is, over the set of roots of the polynomial $t^{p^k} 1$. This is equal to the resolvent, $\operatorname{Res}(\phi_n(t), t^{p^k} 1)$. The symmetry of the resolvent implies that the order of the homology is also given by the product of $t^{p^k} 1$ evaluated at all primitive *n*-roots of unity.
- 5. The p^k power of primitive n root of unity is a primitive $(n/\gcd(n, p^k))$ -root of unity. Hence, the product of $t^{p^k} - 1$ evaluated at all primitive n-roots of unity equals $(\prod(\omega - 1))^d$, where the product is over all primitive $n/(\gcd(n, p^k))$ roots of unity. The integer d is given by $p^l - p^{l-1}$ if k is greater than or equal to the exponent l of p in n. This is the only situation in which we need d explicitly. This value of d is obtained by noting that the homomorphism induced by raising numbers to the p^k powers maps the multiplicative group of primitive n-roots of unity, G_n , onto the group of primitive $n/\gcd(n, p^k)$ roots of unity. If $n = p_1^{k_1} p_2^{k_2} \dots p_r^{k_r}$ then the order of the group is $(p_1^{k_1} - p_1^{k_1-1})(p_2^{k_2} - p_1^{k_2-1}) \dots (p_r^{k_r} - p_r^{k_r-1})$. It follows readily that the kernel of the homomorphism has order d as described.

CHARLES LIVINGSTON

- 6. The product $(\prod(\omega-1))$ over all primitive *m*-roots of unity is given by $\phi_m(1)$. To see this, note that the product $(\prod(\omega-t))$ gives $\phi_m(t)$.
- 7. Combining the observations of the previous two items we have that the order of the p^k -fold branched cover of a knot with Alexander polynomial $\phi_n(t)$ is given by $(\phi_{(n/\gcd(n,p^k))}(1))^d$; if p has exponent l in n, then $d = p^l - p^{l-1}$ in the case that $k \geq l$.
- 8. Using the lemma below which gives $\phi_n(1)$, we now have the following calculations which imply the theorem. If a knot K has Alexander polynomial $\phi_{p^rm}(t)$ with p not dividing m, then the order of the homology of the p^k -fold branched cover of K is: (A) 1 if r = 0 (in which case m must be composite); (B) 1 if r > 0 and m is composite; (C) 1 if r > 0 and k < r; (D) $q^{p^r - p^{r-1}}$ if $k \ge r$ and m is a power of a prime q.

Lemma 5.6. $\phi_n(1) = p$ if n is a power of a prime number p. If n is composite then $\phi_n(1) = 1$.

Proof. As noted before, if $n = p^k$ then $\phi_n(t) = (t^n - 1)/(t^{n-1} - 1)$ and the result follows from L'Hopital's rule.

If n is a composite, we have that $\prod \phi_m(t) = (t^n - 1)/(t - 1)$ if the product is taken over all divisors m of n other than 1. By L'Hopital's rule the right side evaluated at 1 yields n. The factors ϕ_m on the left side with m a prime power each contribute a prime factor to the product, and these together multiply to give n. Hence the remaining terms on the left side of the equation must each equal 1.

6. KNOTS WITH THE SAME SEIFERT FORMS

To repeat the goal of this section, we have:

Theorem 0.1. If $\Delta_V(t)$ has a nontrivial factor that is not an n-cyclotomic polynomial with n divisible by three distinct primes, then there is an infinite collection of knots $\{K_i\}$ with $V_{K_i} = V$ for all i, and no two K_i are concordant. In particular, if $\Delta_V(t)$ satisfies this property and V is metabolic, then there is an infinite collection of nonslice knots with Seifert form V, all of which are distinct up to concordance.

Begin by picking a knot K with Seifert form V and a prime power q so that for the q-fold branched of K, $M_{K,q}$, the homology group $H_1(M_{K,q}, \mathbf{Z})$ is nontrivial. Pick a prime p dividing the order of $H_1(M_{K,q}, \mathbf{Z})$.

Let $X = M_{K,q} - p^{-1}(F)$, where $p^{-1}(F)$ is the preimage in $M_{K,q}$ of the Seifert surface F for K. Also, let X_0 be a component of X. Represent each class in $H_1(X_0, \mathbf{Z})$ by a curve \tilde{L}^j . Apply Theorem 4.4 to assure that the collection of projections, say $\{L^j\}$, is an unlink consisting of curves in the complement of F. Let K_i be the knot formed by replacing each L^j with a copy of the complement of a knot J_i where J_i is chosen as now described.

First, choose J_1 to build K_1 so that $\sigma(K_1, q, \chi) > 0$ for all nontrivial \mathbb{Z}_p -valued χ . The existence of such a J_i is assured by Theorem 4.6. Similarly, pick a sequence of J_i so that the resulting knots K_i satisfy $\sigma(K_{i+1}, q, \chi) > \sigma(K_i, q, \chi')$ for all nontrivial χ and for all χ' .

16

Theorem 6.1. No two knots in the set $\{K_i\}_{i>0}$ are concordant. All these knots have the same Seifert form.

Proof. By construction the knots have the same Seifert forms. Suppose that K_i and K_j , j > i, are concordant. Then $K_i \# - K_j$ is slice, and some Casson–Gordon invariant for some nontrivial \mathbb{Z}_p -valued character on the q-fold cover must vanish. Write such a χ as $\chi_1 \oplus \chi_2$ with respect to the natural direct sum decomposition of the cover. Using the additivity of Casson–Gordon invariants we have $\sigma(K, \chi_1) - \sigma(K, \chi_2) = 0$. But either χ_1 or χ_2 is nontrivial. If χ_j is nontrivial then by the fact that $\sigma(K_j, q, \chi) > \sigma(K_i, q, \chi')$ for all nontrivial χ and for all χ' this difference could not be trivial. On the other hand, if χ_j is trivial then χ_i is nontrivial and the given difference is nonzero.

7. Connected Sums

Our goal in this section is to prove Theorem 0.2. We do this by constructing an explicit example of a knot with Seifert form that splits as a direct sum of Seifert forms but for which the knot is not concordant to a connected sum of knots with corresponding Seifert forms. Further examples follow immediately from those of Section 8, but the construction there is much more abstract. The argument here is included because it provides intuition regarding the latter constructions and because of its simplicity.

Consider the knot K^* drawn in Figure 2. It is built from the knot K (the connected sum of two trefoils, T # T) by removing a neighborhood of an unknot L and replacing it with the complement of a knot, J. The Seifert form of K^* is

$$\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} \oplus \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$$

For shorthand we write this form $V \oplus V$. Our main result is:

Theorem 7.1. If $\sigma_{1/3}(J) \neq 0$, then K^* is not concordant to a connected sum of knots $K_1 \# K_2$ with $V_{K_i} =_s V$.



Figure 2. The Knot K^*

Proof. The proof of this theorem occupies the rest of this section.

Suppose that such a concordance existed. Then the knot $K\# - K_1\# - K_2$ would be slice. The first homology of the 2-fold branched cover of $K\# - K_1\# - K_2$ is easily computed using the Seifert form to be $(\mathbf{Z}_3 \oplus \mathbf{Z}_3) \oplus (\mathbf{Z}_3) \oplus (\mathbf{Z}_3)$, with the direct sum decomposition of the homology corresponding to the direct sum decomposition of the knot. The linking form of the cover is determined by the Seifert form. Since there are only two possible nonsingular linking forms on \mathbb{Z}_3 it follows that (up to sign) the linking form on $(\mathbb{Z}_3 \oplus \mathbb{Z}_3) \oplus (\mathbb{Z}_3) \oplus (\mathbb{Z}_3)$ is given in \mathbb{Z}_3 by

$$lk((a_1, a_2, a_3, a_4), (b_1, b_2, b_3, b_4)) = a_1b_1 + a_2b_2 - a_3b_3 - a_4b_4$$

An algebraic manipulation of generators for a metabolizer shows that the only possible metabolizers are of the following form:

 $<(1,0,\pm 1,0),(0,1,0,\pm 1)>$ or $<(1,0,0,\pm 1),(0,1,\pm 1,0)>.$

(Suppose that the metabolizer is of the form

 $\langle A, B \rangle = \langle (a_1, a_2, a_3, a_4), (b_1, b_2, b_3, b_4) \rangle$.

The sum of two nontrivial squares in \mathbb{Z}_3 is nontrivial. The fact that lk(A, A) = 0 quickly implies that either a_1 or a_2 , is nonzero, and hence can assumed to be 1. Suppose that $a_1 \neq 0$. By subtracting a multiple of A from B if necessary, it can be assumed that $b_1 = 0$ and hence that $b_2 = 1$. The self linking condition now implies that exactly one of the remaining entries, a_3 or a_4 is nonzero, and similarly for b_3 and b_4 . Finally, considering the linking of A with B gives the desired form.)

We will consider the case of the metabolizer $H = \langle (1, 0, 1, 0), (0, 1, 0, 1) \rangle$; the other cases follow similarly.

We next need to consider the set of \mathbb{Z}_3 valued characters on homology that vanish on H. These can be identified with H itself via the linking form. Similarly, we will identify characters on the homology of the covers of K, K_1 , and K_2 , as elements in the homology of the covers, again via the linking form. For clarity, we will write χ_x for the character determined by a homology class x, for whatever space being considered at the time.

Consider the two characters $\chi_{(1,1,1,1)}$ and $\chi_{(1,-1,1,-1)}$ in H. The next figure illustrates the 2-fold branched cover of K^* , $M_{K^*,q}$, drawn using the algorithm of Akbulut and Kirby [AK]. On $M_{K^*,q}$ the character $\chi_{(1,1)}$ takes value 1 on the lifts \tilde{L}_1 and \tilde{L}_2 of L. The character $\chi_{(1,-1)}$ takes value 0 on both lifts.



Figure 3. The 2–Fold Branched Cover, $M_{K^*,q}$

Applying the necessary vanishing of the Casson–Gordon invariant and its additivity yields the following equalities when applied to the character $\chi_{(1,1,1,1)}$:

$$0 = \sigma(K^* \# - K_1 \# - K_2, 2, \chi_{(1,1,1,1)}) = \sigma(K^*, 2, \chi_{(1,1)}) - \sigma(K_1, 2, \chi_{(1)}) - \sigma(K_2, 2, \chi_{(1)}).$$

Applying the surgery formula (Theorem 4.2) now yields:

$$0 = \sigma(K, 2, \chi_{(1,1)}) + 2\sigma_{1/3}(J) - \sigma(K_1, 2, \chi_{(1)}) - \sigma(K_2, 2, \chi_{(1)}).$$

Since the knot K is the connected sum of a knot (the trefoil, T) with itself, K = T # T, we have

$$0 = 2\sigma(T, 2, \chi_{(1)}) + 2\sigma_{1/3}(J) - \sigma(K_1, 2, \chi_{(1)}) - \sigma(K_2, 2, \chi_{(1)}).$$

Via a similar calculation, working with the character $\chi_{(1,-1,1,-1)}$ along with the fact the changing the sign of a character doesn't change the value of the Casson–Gordon invariant, yields the equation:

$$0 = 2\sigma(T, \chi_{(1)}) - \sigma(K_1, \chi_{(1)}) - \sigma(K_2, \chi_{(1)}).$$

Notice that the $\sigma_{1/3}(J)$ term does not appear since $\chi_{(1,-1)}$ vanishes on the \tilde{L}_i .

If $\sigma_{1/3}(J) \neq 0$ then the two formula are contradictory and the proof is complete.

8. PRODUCTS OF ALEXANDER POLYNOMIALS

We begin by restating Theorem 0.3. The goal of the section is to provide examples demonstrating this result.

Theorem 0.3 There exists a knot K with $\Delta_K = \Delta_{V_1} \Delta_{V_2}$ where $Res(\Delta_{V_1}, \Delta_{V_2}) = 1$ but K is not concordant to a connected sum of knots, $K_1 \# K_2$, with $\Delta_{K_i} = \Delta_{V_i}$.

The argument here will be far more subtle than that of the previous section, and in particular we will not be able to draw explicit diagrams of the knots in question. Instead we will rely on Theorem 4.4. The necessary knot is provided by constructing a knot as follows.

Theorem 8.1. There exists a knot K^* with $\Delta_K(t) = (2t^2 - 3t + 2)(3t^2 - 5t + 3)$ but K^* is not concordant to a connected sum $K_1 \# K_2$ with $\Delta_{K_1}(t) = 2t^2 - 3t + 2$ and $\Delta_{K_2}(t) = 3t^2 - 5t + 3$.

Notice that $\operatorname{Res}(2t^2 - 3t + 2, 3t^2 - 5t + 3) = 1$. This follows easily from a direct calculation using the definition.

8.1. Construction of K^* . To begin the construction, let K_1 be knot with Alexander polynomial $2t^2 - 3t + 2$ and let K_2 be a knot with Alexander polynomial $3t^2 - 5t + 3$. We will abbreviate our standard notation by writing M_i for $M_{K_i,11}$, the 11-fold branched cover of S^3 branched over K_i . We will also write T for the deck transformation $T_{K,11}$ and will abbreviate the Casson–Gordon invariant $\sigma(K, 11, \chi)$ by $\sigma(K, \chi)$.

Theorem 8.2. $H_1(M_i, \mathbb{Z}) \cong (\mathbb{Z}_{67})^2$. As a \mathbb{Z}_{67} vector space, $H_1(M_i, \mathbb{Z})$ splits as the direct sum of eigenspaces of the transformation T. For J_1 the eigenvectors, v_1 and v_2 , have eigenvalues 40 and 62. For J_2 the eigenvectors, w_1 and w_2 , have eigenvalues 9 and 15.

Proof. Using Theorem 1.2, the order of homology of the q-fold branched cover of a knot with Alexander polynomial Δ is given by $\prod_{i=1,\ldots,q-1} \Delta(\zeta^i)$, where ω is a primitive q root of unity. This value can be computed explicitly for both knots, using for instance Maple. In both cases one find the homology to be of order 67².

Since the homology of an odd-fold branched cover of a knot is always of the form $G \oplus G$, (see, eg. [Ro]), it follows that $H_1(M_i, \mathbb{Z}) \cong (\mathbb{Z}_{67})^2$, for i = 1, 2.

From Theorem 1.1, $2T^2 - 3T + 2$ annihilates the homology of M_1 . Over \mathbf{Z}_{67} the polynomial $2T^2 - 3T + 2$ factors as 2(T - 40)(T - 62), and hence the homology does split as the sum of a 40 and a 62 eigenspace.

We next need to show that both these eigenspaces are nontrivial. To see this we first observe that the linking form of M_1 vanishes on each eigenspace of T, as follows. For eigenvectors of eigenvalue λ , x and y, lk(x, y) = lk(Tx, Ty) = $lk(\lambda x, \lambda y) = \lambda^2 lk(x, y)$. Since $\lambda^2 \neq 1$ we have the vanishing of the linking form on the eigenspace. But since the linking form on a 3-manifold is nonsingular, the eigenspace must be of dimension at most one. Since this holds for both eigenspaces, both eigenspaces must be exactly one dimensional, as desired.

A similar argument applies for K_2 where we have $3T^2 - 5T + 3 = 3(T - 9)(T - 15) \mod 67$.

Let $K = K_1 \# K_2$. In this case, the 11-fold branched cover, $M = M_1 \# M_2$, and, from Theorem 8.2 we have the homology of M is generated by T-eigenvectors, $\{v_1, v_2, w_1, w_2\}$ as described in that theorem.

Let each of these eigenvectors be represented by curves $\{\tilde{L}^1, \tilde{L}^2, \tilde{L}^3, \tilde{L}^4\}$ and let $v_2 + w_1$ be represented by \tilde{L}^5 . By Theorem 4.4 we can assume that the \tilde{L}_i arise as the lifts of components of an unlink $\{L_1, L_2, L_3, L_4, L_5\}$ in S^3 , each component of which is trivial in $H_1(S^3 - K, \mathbb{Z})$.

A knot J will be chosen based on the needed properties of its 67-signatures, as developed later in this section. Given this, we will let K^* be the knot formed by replacing neighborhoods of the L_i in S^3 with the complement of the knot $J \subset S^3$.

8.2. Characters and Casson–Gordon Invariants of K^* . Let $\{\chi_1, \chi_2, \chi_3, \chi_4\}$ be the \mathbb{Z}_{67} -valued characters on $H_1(M, \mathbb{Z})$ that are dual to the basis $\{v_1, v_2, w_1, w_2\}$. We will use the same names for the corresponding characters on $H_1(M^*, \mathbb{Z})$. We have the following values of the Casson–Gordon invariants; these follow immediately from the definitions along with Theorem 4.3.

F1:
$$\sigma(K^*, \chi_1) = \sigma(K, \chi_1) + \sum_{i=0}^{10} \sigma_{40^i/67}(J).$$

F2: $\sigma(K^*, \chi_2) = \sigma(K, \chi_2) + 2\sum_{i=0}^{10} \sigma_{62^i/67}(J).$
F3: $\sigma(K^*, \chi_3) = \sigma(K, \chi_3) + 2\sum_{i=0}^{10} \sigma_{9^i/67}(J).$
F4: $\sigma(K^*, \chi_4) = \sigma(K, \chi_4) + \sum_{i=0}^{10} \sigma_{15^i/67}(J).$

Notice that each of the summations that occurs above is in fact the same sum, being taken over the full set of 11-roots of unity in \mathbb{Z}_{67} . The factors of 2 appear in **F2** and **F3** because χ_2 and χ_3 evaluate nontrivially on L_5 .

We will also need the following two values, which again follow readily from the construction and Theorem 4.3.

$$\mathbf{F5:} \ \sigma(K^*, \chi_2 + \chi_3) = \sigma(K, \chi_2 + \chi_3) + \sum_{i=0}^{10} \sigma_{62^i/67}(J) + \sum_{i=0}^{10} \sigma_{9^i/67}(J) + \sum_{i=0}^{10} \sigma_{62^i+9^i/67}(J).$$

$$\mathbf{F6:} \ \sigma(K^*, \chi_2 - \chi_3) = \sigma(K, \chi_2 - \chi_3) + \sum_{i=0}^{10} \sigma_{62^i/67}(J) + \sum_{i=0}^{10} \sigma_{9^i/67}(J) + \sum_{i=0}^{10} \sigma_{9^i/67}(J).$$

The difference between the two formulas occurs only in the last terms, since $\chi_2 \pm \chi_3$ evaluates to be $62^i \pm 9^i$ on $T^i(L_5)$.

These calculations lead to the following result.

Theorem 8.3.

$$\sigma(K^*, \chi_2 + \chi_3) - \sigma(K^*, \chi_2 - \chi_3) = \sum_{i=0}^{10} \sigma_{(62^i + 9^i)/67}(J) - \sum_{i=0}^{10} \sigma_{(62^i - 9^i)/67}(J).$$

Proof. We have by the additivity of Casson–Gordon invariants that $\sigma(K, \chi_2 + \chi_3) = \sigma(K_1, \chi_3) + \sigma(K_2, \chi_3)$ and $\sigma(K, \chi_2 - \chi_3) = \sigma(K_1, \chi_2) + \sigma(K_2, -\chi_3)$. Since $\sigma(K_2, \chi_3) = \sigma(K_2, -\chi_3)$, the result follows.

8.3. Identifying Metabolizers of $K^* - \#K'_1 \# - K'_2$. Suppose that K^* is concordant to a the connected sum of knots $K'_1 \# K'_2$ where K'_1 has Alexander polynomial $2t^2 - 3t + 2$ and K'_2 has Alexander polynomial $3t^2 - 5t + 3$. The covers of the K'_i have the identical homological properties as described by Theorem 8.2 for the M_i , since the proof depended only on Alexander polynomials. We use the exact same notation for these knots (including the characters), only with the addition of the primes throughout.

We are now assuming that $K^* \# - K'_1 \# - K'_2$ is slice; denote $K^* \# - K'_1 \# - K'_2$ by \bar{K} . Our next goal is to identify possible *T*-invariant metabolizers for the linking form of the 11-fold branched cover over \bar{K} . Since the homology of the cover (which is isomorphic to $(\mathbf{Z}_{67})^8$) is annihilated by (T - 40)(T - 62)(T - 9)(T - 15), a metabolizer, $H \cong (\mathbf{Z}_{67})^4$, is also annihilated by the same polynomial. It follows that *H* splits into a direct sum of eigenspaces.

Via the linking form we can identify characters with homology classes. The linking form on homology induces a linking form on the group of characters. The only change is that a homology class of eigenvalue λ corresponds to a character of eigenvalue λ^{-1} . Since it will be simpler to work on the level of characters instead of homology classes, move to that setting now. All the arguments can be translated back to the level of homology if desired.

Case 1: Two dimensional eigenspaces in the metabolizer. The two dimensional eigenspaces are spanned by a pair of characters $\{\chi_j, \chi'_j\}$ for some j. Hence, in this case some character χ_j appears in the metabolizer. As a result we would have:

R1: $\sigma(\bar{K}, \chi_j) = 0$ for some $j \in \{1, 2, 3, 4\}$.

Case 2: All eigenspace in the metabolizer are one dimensional. Each eigenvector can be expressed as $a\chi_j + b\chi'_j$ for some $a, b \in \mathbb{Z}_{67}$ and $j \in \{1, 2, 3, 4\}$. As an example, suppose that $a\chi_1 + b\chi'_1 \in H$. We would also have that $a'\chi_2 + b'\chi'_2 \in H$ for some $a', b' \in \mathbb{Z}_{67}$. The linking number of these two characters is aa' - bb', and this should be trivial since we are in a metabolizer.

Notice that if a = 0 then $b \neq 0$ so b' = 0 and hence $a' \neq 0$. Hence the character χ_2 will be in H. It follows that if any character of the form χ'_j is in the metabolizer, then some character of the form χ_j is necessarily in the metabolizer, and we have the equation **R1** above must hold.

The more interesting and the only other possibility is that the metabolizer is spanned by characters $\{\chi_1 + a_1\chi'_1, \chi_2 + a_2\chi'_2, \chi_3 + a_3\chi'_3, \chi_4 + a_4\chi'_4\}$ with all $a_i \neq 0$. In particular we have that H contains the characters: $\chi_2 + a_2\chi'_2 + \chi_3 + a_3\chi'_3$ and $\chi_2 + a_2\chi'_2 - \chi_3 - a_3\chi'_3$. Hence for some nonzero a_2 and a_3 we have

R2: $\sigma(\bar{K}, \chi_2 + a_2\chi'_2 + \chi_3 + a_3\chi'_3) = 0.$ **R3:** $\sigma(\bar{K}, \chi_2 + a_2\chi'_2 - \chi_3 - a_3\chi'_3) = 0.$ 8.4. Computing the obstructions. In the first case, in which formula **R1** holds we can use the additivity of the Casson–Gordon invariant along with the formula **Fi** above to achieve one of the following equations, depending on the value of j.

G1: $\sigma(K_1, \chi_1) + \sigma(K_2, 0) - \sigma(K'_1, 0) - \sigma(K'_2, 0) + \sum_{i=0}^{10} \sigma_{40^i/67}(J) = 0$. **G2:** $\sigma(K_1, \chi_2) + \sigma(K_2, 0) - \sigma(K'_1, 0) - \sigma(K'_2, 0) + 2\sum_{i=0}^{10} \sigma_{62^i/67}(J) = 0$. **G3:** $\sigma(K_1, 0) + \sigma(K_2, \chi_3) - \sigma(K'_1, 0) - \sigma(K'_2, 0) + 2\sum_{i=0}^{10} \sigma_{9^i/67}(J) = 0$. **G4:** $\sigma(K_1, 0) + \sigma(K_2, \chi_4) - \sigma(K'_1, 0) - \sigma(K'_2, 0) + \sum_{i=0}^{10} \sigma_{15^i/67}(J) = 0$.

(Of course these simplify further since, by Theorem 2.9 the terms $\sigma(\cdot, 0)$ all equal 0.)

The more interesting case is when equations **R2** and **R3** hold. These lead to the following identities.

$$\mathbf{G5:} \ \sigma(K_1,\chi_2) + \sigma(K_2,\chi_3) - \sigma(K'_1,a_2\chi'_2) - \sigma(K'_2,a_3\chi'_3) + \sum_{i=0}^{10} \sigma_{62^i/67}(J) + \sum_{i=0}^{10} \sigma_{9^i/67}(J) + \sum_{i=0}^{10} \sigma_{(62^i+9^i)/67}(J) = 0.$$

$$\mathbf{G6:} \ \sigma(K_1,\chi_2) + \sigma(K_2,\chi_3) - \sigma(K'_1,a_2\chi'_2) - \sigma(K'_2,a_3\chi'_3) + \sum_{i=0}^{10} \sigma_{62^i/67}(J) + \sum_{i=0}^{10} \sigma_{15^i/67}(J) + \sum_{i=0}^{10} \sigma_{(62^i-9^i)/67}(J) = 0.$$

Summarizing, we have the following result.

Theorem 8.4. If $K^* - \#K'_1 \# - K'_2$ is slice, either one of the four equations, G1, G2, G3, or G4, holds, or else $\sum_{i=0}^{10} \sigma_{(62^i+9^i)/67}(J) - \sum_{i=0}^{10} \sigma_{(62^i-9^i)/67}(J) = 0.$

Proof. We have already seen that either one of these equations (G1, G2, G3, or G4) must hold, or else G5 and G6 are true. If the differences of the two equations in G5 and G6 is computed then most terms cancel and the only remaining terms yield the difference of summations as desired. \Box

8.5. Finding the appropriate J and completion of the proof. We begin by expanding the summation of signatures that occur in formulas G1 through G4, say $P_1(J)$, and in the difference, $P_2(J) = \sum_{i=0}^{10} \sigma_{(62^i+9^i)/67}(J) - \sum_{i=0}^{10} \sigma_{(62^i-9^i)/67}(J)$.

- $P_1(J) = \sum_{i=0}^{10} \sigma_{40^i/67}(J) = \sigma_{1/67}(J) + \sigma_{3/67}(J) + \sigma_{5/67}(J) + \sigma_{8/67}(J) + \sigma_{9/67}(J) + \sigma_{14/67}(J) + \sigma_{15/67}(J) + \sigma_{22/67}(J) + \sigma_{24/67}(J) + \sigma_{25/67}(J) + \sigma_{27/67}(J).$
- $P_2(J) = \sum_{i=0}^{10} \sigma_{(62^i+9^i)/67}(J) \sum_{i=0}^{10} \sigma_{(62^i-9^i)/67}(J) = (\sigma_{1/67}(J) + \sigma_{2/67}(J) + \sigma_{4/67}(J) + \sigma_{11/67}(J) + \sigma_{12/67}(J) + \sigma_{16/67}(J) + \sigma_{17/67}(J) + \sigma_{21/67}(J) + \sigma_{27/67}(J) + \sigma_{28/67}(J) + \sigma_{30/67}(J)) (\sigma_{0/67}(J) + \sigma_{26/7}(J) + \sigma_{10/67}(J) + \sigma_{11/67}(J) + \sigma_{11/67}(J) + \sigma_{17/67}(J) + \sigma_{17/67}(J) + \sigma_{24/67}(J) + \sigma_{25/67}(J) + \sigma_{27/67}(J) + \sigma_{32/67}(J)).$

That a knot J can be chosen so that none of the conditions given in Theorem 8.4 is satisfied now follows immediately from the independence of the signature function, 3.2.

9. IRREDUCIBLE KNOTS WITH REDUCIBLE SEIFERT FORMS

Our goal in this section is to prove the following theorem, Part \mathbf{A} or Theorem 0.4.

Theorem 9.1. If M is a metabolic Seifert form and $\Delta_M(t)$ has a factor that is not a cyclotomic polynomial $\phi_n(t)$ with n divisible by three distinct primes, then for every Seifert form V there is a knot K with $V_K = V \oplus M$ with the property that Kis not concordant to a knot with Seifert form S-equivalent to V. *Proof.* Begin with an arbitrary knot K_1 having Seifert form M. For some prime power q the branched cover $M_{K_1,q}$ has nontrivial homology, with the order of the homology divisible by p for some prime p. Let K_2 be a knot with Seifert form Vwith q-fold branched cover $M_{K_2,q}$. Let K^* be the knot formed by the iterated companionship construction on $K_1 \# K_2$ to assure that the Seifert form is unchanged but that all nontrivial Casson–Gordon invariants on $M_{K^*,q}$ are positive, as described in Theorem 4.6. We show that K^* has the desired properties.

Let J be a knot with Seifert form $V_J =_s V$ and assume that J is concordant to K^* . Then $K^* \# - J$ is slice and has Seifert form S-equivalent to $(M \oplus V) \oplus -V$. The homology and linking form of the q-fold cover of $K^* \# - J$ thus has a corresponding decomposition as $(H_1 \oplus H_2) \oplus -H_2$ where H_1 is nontrivial.

Let W be a metabolizer for the linking form on $(H_1 \oplus H_2) \oplus -H_2$. We will show that for some character that vanishes on that metabolizer, the Casson–Gordon invariant is nontrivial, thus contradicting Theorem 2.7.

For each $h \in -H_2$ let W_h denote the set of elements in $g \in H_1 \oplus H_2$ such that $(g,h) \in W$. Clearly, if W_h is nontrivial it is a coset of W_0 . It is also easily seen that W is the union of the set of all elements $\{(g,h)|g \in W_h\}$. It follows that W has order at most $|W_0||H_2|$. On the other hand, the order of W is given by the square root of the order of $(H_1 \oplus H_2) \oplus -H_2$, which is greater than the order of H_2 , since we have arranged that H_1 is nontrivial. It follows that W_0 is nontrivial, and in particular there is a homology class in W of the form (g, 0) with $g \in H_1 \oplus H_2$ and $0 \in -H_2$.

Let χ be given by linking with the class (g, 0). Because it is in W, which is a metabolizer for the linking form, linking χ certainly vanishes on W. If χ is not of prime power order, some multiple of it is, so we can assume it is of prime power order. By the additivity of Casson–Gordon invariants we have $\sigma(K^* \# J, q, \chi) =$ $\sigma(K^*, q, \chi') + \sigma(J, q, 0)$, where χ' is the restriction of χ to $H_1(M_{K^*,q}, \mathbb{Z})$. But Casson–Gordon invariants vanish for the trivial character (Theorem 2.9) and we have arranged that $\sigma(K^*, q, \chi') > 0$. Hence the Casson–Gordon invariant is nontrivial, and K^* is not concordant to J as desired.

10. INEQUIVALENT KNOTS WITH ALGEBRAICALLY EQUIVALENT, IRREDUCIBLE SEIFERT FORMS

Part **B** of Theorem 0.4 has the following restatement. The proof of this result occupies the entire section.

Theorem 10.1. There exists a knot K with Seifert form V_K such that V_K is reduced and algebraically concordant to a second reduced form, V, but K is not concordant to any knot K_1 with $V_{K_1} =_s V$.

We will consider the two Seifert matrices

$$V_1 = \begin{pmatrix} 3 & 2 \\ 1 & 3 \end{pmatrix}$$
 and $V_2 = \begin{pmatrix} 1 & 2 \\ 1 & 9 \end{pmatrix}$.

We will build a knot K with Seifert form V_1 and show that K_1 is not concordant to any knot K_2 with $V_{K_2} =_s V_2$, despite the fact that, as we will soon see, V_1 and V_2 are algebraically concordant and reduced. 10.1. **Irreducibility of Seifert Forms.** We will begin by giving the precise needed definitions. Further information can be found in the basic references [Ke, Le2].

Definition 10.2. A Seifert form V is called reducible if it is nonsingular (det $\neq 0$) and the (rational) transformation $S = -V^{-1}V^t$ has a nontrivial invariant subspace on which the form $Q = V + V^t$ vanishes. Otherwise it is called irreducible.

This following result, though not needed here, is proved in [Le2, Section 8].

Theorem 10.3. Every Seifert form is algebraically concordant to a nonsingular form: one with $det(V) \neq 0$.

We can now state formally the irreducibility of our forms.

Theorem 10.4. The Seifert forms V_1 and V_2 as defined above are irreducible.

Proof. In both cases the characteristic polynomial of $-V_i^{-1}V_i^t$ is given by $t^2 - 13t + 49$. The roots are complex, so there is no nontrivial invariant rational subspace.

10.2. Properties of the Seifert Forms and Related Covering Spaces.

Theorem 10.5. The Seifert matrices V_1 and V_2 are algebraically concordant.

Proof. We want to see that $V_1 \oplus -V_2$ is metabolic. According to Levine, [Le2, Proposition 14], a Seifert form V with Alexander polynomial of the form $\lambda(t)^e$ where $\lambda(t)$ is quadratic is null concordant if and only if the rational pairing given by $V + V^t$ is metabolic. In our case the Alexander polynomial is $7t^2 - 13t + 7$. The symmetrized forms are

$$W_1 = \begin{pmatrix} 6 & 3 \\ 3 & 6 \end{pmatrix}$$
 and $W_2 = \begin{pmatrix} 2 & 3 \\ 3 & 18 \end{pmatrix}$.

Using simultaneous row and column operations to clear the off–diagonal entries yields

$$W'_1 = \begin{pmatrix} 6 & 0 \\ 0 & 9/2 \end{pmatrix}$$
 and $W'_2 = \begin{pmatrix} 2 & 0 \\ 0 & 27/2 \end{pmatrix}$.

Working over the rationals, any diagonal entry can multiplied by a rational square, r^2 , corresponding to multiplying a basis element by r. For both matrices we can multiply the second entry by 4/9 to get the matrices

$$W_1' = \begin{pmatrix} 6 & 0 \\ 0 & 2 \end{pmatrix}$$
 and $W_2' = \begin{pmatrix} 2 & 0 \\ 0 & 6 \end{pmatrix}$.

These are obviously equivalent, so the difference is metabolic, as desired.

(The reader might prefer a direct proof of $V_1 \oplus -V_2$ being metabolic. One such proof is just to observe that a metabolizer is given by < (0, -3, 6, -1,), (2, -1, 0, 1) >. Checking that this is a metabolizer is easy; finding it is a lengthy exercise that need not be repeated here.)

Theorem 10.6. Let K_i be a knot with Seifert form S-equivalent to V_i as above, i = 1, 2.

A.: The 2-fold branched cover of S^3 over K_1 , M_1 , has $H_1(M_1, \mathbb{Z}) \cong \mathbb{Z}_9 \oplus \mathbb{Z}_3$ and the homology is generated by elements x and y of order 9 and 3 respectively with lk(x, x) = 5/9, lk(y, y) = 2/3, and lk(x, y) = 0.

B.: The 2-fold branched cover of S^3 over K_2 , M_2 , has $H_1(M_2, \mathbf{Z}) \cong \mathbf{Z}_{27}$ and the homology is generated by an element z with lk(z, z) = 2/27.

Proof. **A.** The homology of the 2-fold branched cover of the knot K_1 is given by W_1 , [Ro]. Note that the determinant is 27 so the order of $H_1(M_1, \mathbb{Z})$ is 27. The matrix W_1 is given with respect to a generating set $\{a, b\}$. A second generating set is given by $\{x, y\} = \{2b - a, b - a\}$. These generate, since x - y = b. Subtracting twice the first row of W_1 from the second shows that 9a = 0. Adding the two rows shows that 9a + 9b = 0, so 9b = 0. Subtracting the rows shows that 3a - 3b = 0. It then follows that 9x = 0 and 3y = 0, as needed.

The linking form with respect to the original generating set $\{a, b\}$ is given by

$$W_1^{-1} = \begin{pmatrix} 6/27 & -3/27 \\ -3/27 & 6/27 \end{pmatrix}.$$

(Such a description follows most readily from the Akbulut–Kirby [AK] surgery description of the 2–fold branched cover.) One now computes explicitly that the values of the linking form are as given in statement **A** of Theorem 10.6.

Proof. **B.** This proof is somewhat easier. The homology of M_2 is presented by W_2 . In this case row operations yield the matrix

$$\begin{pmatrix} 0 & 27 \\ 1 & 15 \end{pmatrix}$$

Hence the homology is \mathbf{Z}_{27} as desired and the second generator, the one corresponding to the second column of the presentation matrix W_2 , generates the homology; call this element z. Taking the inverse,

$$W_2^{-1} = \begin{pmatrix} 18/27 & -3/27 \\ -3/27 & 2/27 \end{pmatrix},$$

yields the desired self-linking, lk(z, z) = 2/27.

10.3. Metabolizers. If K_i has linking form V_i as above, then the connected sum, $K_1 \# - K_2$, has 2-fold branched cover $M = M_1 \# - M_2$ with homology $\mathbf{Z}_9 \oplus \mathbf{Z}_3 \oplus \mathbf{Z}_{27}$, generated by elements x, y, z as given in Theorem 10.6. As shorthand, we write the element ax + by + cz as (a, b, c). The linking form is diagonal, given by

$$lk((a, b, c), (a', b', c')) = \frac{(15aa' + 18bb' - 2cc')}{27} \in \mathbf{Q}/\mathbf{Z}$$

In what follows we will work with integer representatives of a, b, and c, taken from the intervals $0 \le a < 9$, $0 \le b < 3$, $0 \le c < 27$, with the linking form taking values in \mathbb{Z}_{27} .

Theorem 10.7. The linking form on $H_1(M, \mathbb{Z})$ has exactly two metabolizers. One metabolizer is generated by $\{(0, 1, 3), (3, 0, 0)\}$ while the second metabolizer is generated by $\{(0, -1, 3), (3, 0, 0)\}$. Both are isomorphic to $\mathbb{Z}_9 \oplus \mathbb{Z}_3$.

Proof. A metabolizer must be of order 27 since the order of $H_1(M, \mathbb{Z})$ is 27². We first observe that the metabolizer cannot be generated by a single element of order 27. Any such element is of the form (a, b, c) with c not divisible by 3. Then clearly $15a^2 + 18b^2 - 2c^2$ is not divisible by 3, and the self-linking is nontrivial.

_	
_	
_	
_	
_	
_	
_	
_	
_	

The metabolizer also cannot be isomorphic to $\mathbf{Z}_3 \oplus \mathbf{Z}_3 \oplus \mathbf{Z}_3$. There is a unique subgroup of this form, generated by $\{(3,0,0), (0,1,0), (0,0,9)\}$. The second of these generators does not have trivial self-linking.

The remaining case consists of a subgroup isomorphic to $\mathbf{Z}_9 \oplus \mathbf{Z}_3$ generated by two elements, say

$$\{(\alpha, \beta, \gamma), (\alpha', \beta', \gamma')\}$$

Via a change of basis we can assume that $\gamma' = 0$, so now our generating set is given by

$$\{(\alpha,\beta,\gamma),(\alpha',\beta',0)\}.$$

The self-linking of this second element is $(15{\alpha'}^2 + 18{\beta'}^2) \in \mathbb{Z}_{27}$. If α' is not divisible by 3, then $15{\alpha'}^2$ is not divisible by 9, where $18{\beta'}^2$ is divisible by 9, so the sum could not be divisible by 27. Hence, α' is divisible by 3, in which case $15{\alpha'}^2$ is divisible by 27, and β' must be 0 if the sum is to be divisible by 27. Hence, we have that the metabolizer is generated by elements

$$\{(\alpha,\beta,\gamma),(3\alpha'',0,0)\}.$$

Since $3\alpha'' \neq 0$, by multiplying the second generator by 2 if need be, we can assume that the second generator is (3, 0, 0). If the linking of these two generators is to be 0 then α must be divisible by 3, so after subtracting a multiple of the second generator from the first we have that the metabolizer is generated by

$$\{(0, \beta, \gamma), (3, 0, 0)\}$$

The first generator must be of order 9, so we can assume that $\gamma = 3$. Finally, for the self-linking of this first generator to be 0, we have that $\beta = \pm 1$ and the proof is complete.

10.4. Evaluating Casson–Gordon Invariants and building K. We again identify characters on the first homology of the relevant 3–manifolds with the first homology via the linking form, and write χ_x for the character given by linking with x in the first homology.

Theorem 10.8. Suppose that K_1 has Seifert form S-equivalent to V_1 , K_2 has Seifert form S-equivalent to V_2 , and K_1 and K_2 are concordant. Then $\sigma(K_1, \chi_{(0,1)}) = \sigma(K_1, \chi_{(3,1)})$.

Proof. By the computation of the metabolizers in Theorem 10.7, the metabolizer for $K_1 \# - K_2$ (based on 2–fold covers) contains the elements $(0, 1, \pm 3)$ and $(3, 1, \pm 3)$. Hence, computing Casson–Gordon invariants and using additivity, we have:

$$\sigma(K_1, \chi_{(0,1)}) = \sigma(K_2, \chi_{(3)})$$
 and $\sigma(K_1, \chi_{(3,1)}) = \sigma(K_2, \chi_{(3)}).$

In particular, $\sigma(K_1, \chi_{(0,1)}) = \sigma(K_1, \chi_{(3,1)})$, as desired.

We now conclude the proof of Theorem 10.1 by describing a knot K with Seifert form V_1 for which the identity given in Theorem 10.8 fails. Begin with an arbitrary knot K_1 having Seifert form V_1 . As we have seen, with M_1 the 2-fold branched cover of S^3 over K_1 the homology group $H_1(M_1, \mathbb{Z}) = \mathbb{Z}_9 \oplus \mathbb{Z}_3$. Represent the class $(1,0) \in H_1(M_1, \mathbb{Z})$ by a curve \tilde{L} . Apply Theorem 4.4 to assure that \tilde{L} is the lift of a null homologous unknot, L, in $S^3 - K_1$, and then replace a neighborhood of Lwith the complement of a knot J, the properties of which will soon be presented.

26

Using the linking form for M_1 we have that $\chi_{(1,0)}(1,0) = 15/27 \in \mathbf{Q}/\mathbf{Z}$ and $\chi_{(0,1)}(1,0) = 0 \in \mathbf{Q}/\mathbf{Z}$. Hence, with values taken in \mathbf{Q}/\mathbf{Z} :

- $\chi_{(0,1)}(L) = 0$
- $\chi_{(3,1)}(L) = 45/27 = 2/3$

By Theorem 4.2 and the fact that $\sigma_{1/3}(J) = \sigma_{2/3}(J)$ for all knots J, we have:

- $\sigma(K, \chi_{(0,1)}) = \sigma(K_1, \chi_{(0,1)})$
- $\sigma(K, \chi_{(3,1)}) = \sigma(K_1, \chi_{(3,1)}) + 2\sigma_{1/3}(J)$

These two will be equal if

$$2\sigma_{1/3}(J) = \sigma(K_1, \chi_{(0,1)}) - \sigma(K_1, \chi_{(3,1)}).$$

It follows that to construct the desired example of the knot K with $\sigma(K, \chi_{(0,1)}) \neq \sigma(K, \chi_{(3,1)})$, one need simply pick J so that this equality does not hold. This can be done since the expression on the right is determined by the fixed knot K_1 , and knots with 1/3 signatures arbitrarily large are easily constructed.

11. Conjectures and Addenda

Recent work of Cochran, Orr, and Teichner have generalized Casson–Gordon invariants. With this, the constraint on working with prime power covers is not always necessary. Thus, it may be possible to remove the constraint that occurs in this paper that the Alexander polynomial has a factor that is not a cyclotomic polynomial $\phi_n(t)$ with n divisible by at least three distinct primes. The first such polynomial that occurs is $\phi_{30}(t) = t^8 + t^7 - t^5 - t^4 - t^3 + t + 1$. The order of the homology of the 6–fold cover of knot having this polynomial is 25. We would conjecture that for a Seifert form, say V, with this polynomial there would be an infinite number of nonconcordant knots having Seifert form V. Nothing of this sort has been proved yet; the strongest result in this direction has been proved by Taehee Kim; he has constructed algebraically slice knots with Alexander polynomial $\phi_{30}(t)^2$ that are not slice.

In a different direction, Theorem 0.1 provides an infinite family of knots, $\{K_i\}_{i>0}$. It should be the case that this set, or another set that satisfies the conditions of the theorem, is affinely independent. That is, that the set of knots $\{K_i - K_1\}_{i>1}$ is independent in the concordance group.

Much deeper questions concern the relationship of the Seifert form of a knot and its concordance class. In particular, is it possible that the Seifert form of a knot contains information about the concordance class of the knot that is not contained in its algebraic concordance class? The following conjecture makes this concrete. Based on the examples of Section 10, consider the Seifert forms

$$V_1 = \begin{pmatrix} 3 & 2 \\ 1 & 3 \end{pmatrix}$$
 and $V_2 = \begin{pmatrix} 1 & 2 \\ 1 & 9 \end{pmatrix}$.

Conjecture 11.1. If K_i has Seifert form V_i for i = 1, 2, then K_1 and K_2 are not concordant.

The conjecture could be further strengthened by requiring only that the Seifert forms of the knots be S-equivalent to these forms. The example of Section 10 is the strongest known result indicating the possibility that a Seifert form contains more information about the concordance class of a knot than does the algebraic concordance class of the Seifert form.

CHARLES LIVINGSTON

References

- [AK] S. Akbulut and R. Kirby. Branched covers of surfaces in 4-manifolds. Math. Ann. 252 (1979/80), 111–131.
- [BZ] G. Burde and H. Zieschang. Knots. de Gruyter Studies in Mathematics, 5. Walter de Gruyter & Co., Berlin-New York, 1985.
- [CG1] A. Casson and C. Gordon. Cobordism of classical knots. A la recherche de la Topologie perdue, ed. by Guillou and Marin, Progress in Mathematics, Volume 62, 1986. (Originally published as Orsay Preprint, 1975.)
- [CG2] A. Casson and C. Gordon. On slice knots dimension three. Algebraic and geometric topology (Proc. Sympos. Pure Math., Stanford Univ., Stanford, Calif., 1976), Part 2, pp. 39–53 Proc. Sympos. Pure Math., XXXII, Amer. Math. Soc., Providence, R.I., 1978.
- [COT] T. Cochran, K. Orr, and P. Teichner. Knot concordance, Whitney towers and L² signatures preprint 1999. (front.math.ucdavis.edu/math.GT/9908117.)
- [CF] P. E. Conner and E. E. Floyd. *Differentiable periodic maps.* Ergebnisse der Mathematik und ihrer Grenzgebiete, N. F., Band 33 Academic Press Inc., Publishers, New York; Springer-Verlag, Berlin-Gttingen-Heidelberg 1964.
- [F1] R. Fox Free differential calculus. III. Subgroups. Ann. of Math. (2) 64 1956 407-419.
- [F2] R. Fox, A quick trip through knot theory. *Topology of 3-Manifolds* ed. by M. K. Fort, Prentice Hall (1962), 120–167.
- [FM] R. Fox and J. Milnor, Singularities of 2-spheres in 4-space and cobordism of knots. Osaka J. Math. 3 (1966) 257–267.
- [Fr] M. Freedman. The topology of four-dimensional manifolds. J. Differential Geom. 17 (1982), 357–453.
- [FQ] M. Freedman and F. Quinn. Topology of 4-manifolds. Princeton Mathematical Series, 39 Princeton University Press, Princeton, NJ, 1990.
- [G] P. Gilmer. Slice knots in S³. Quart. J. Math. Oxford Ser. (2) 34 (1983), no. 135, 305–322.
- [GL] P. Gilmer and C. Livingston. The Casson–Gordon invariant and link concordance. *Topology* 31 (1992), no. 3, 475–492.
- [Go] C. McA. Gordon. Some aspects of classical knot theory. Knot theory (Proc. Sem., Planssur-Bex, 1977), pp. 1–60 Lecture Notes in Math., 685, Springer, Berlin, 1978.
- [J] B. Jiang. A simple proof that the concordance group of algebraically slice knots is infinitely generated. Proc. Amer. Math. Soc. 83 (1981), 189–192.
- [Ka] L. Kauffman. Branched coverings, open books and knot periodicity. *Topology* 13 (1974), 143–160.
- [Ke] M. Kervaire. Knot cobordism in codimension two. Manifolds-Amsterdam 1970 (Proc. Nuffic Summer School) pp. 83–105 Lecture Notes in Mathematics, Vol. 197 Springer, Berlin (1971).
- [KLk] R. Kirby and W. B. R. Lickorish. Prime knots and concordance. Math. Proc. Cambridge Philos. Soc. 86 (1979), no. 3, 437–441.
- [KLv] P. Kirk and C. Livingston. Concordance and mutation. preprint 2000 online at http://front.math.ucdavis.edu/math.GT/9912174
- [La] S. Lang. Algebra, Addison-Wesley Publishing Co., Reading, Mass., 1984.
- [Le3] J. Levine. An algebraic classification of some knots of codimension two. Comment. Math. Helv. 45 (1970) 185–198.
- [Le1] J. Levine. Knot cobordism groups in codimension two. Comment. Math. Helv. 44 (1969), 229–244.
- [Le2] J. Levine. Invariants of knot cobordism. Invent. Math. 8 (1969), 98-110.
- [Le3] J. Levine. Metabolic and hyperbolic forms from knot theory. J. Pure Appl. Algebra 58 (1989), no. 3, 251–260.
- [Lt1] R. Litherland. A formula for the Casson–Gordon invariant of a knot. Preprint.
- [Lt2] R. Litherland. Cobordism of satellite knots. *Four–Manifold Theory*, Contemporary Mathematics, eds. C. Gordon and R. Kirby, American Mathematical Society, Providence RI 1984, 327–362.
- [L] C. Livingston. Homology cobordisms of 3-manifolds, knot concordances, and prime knots. Pacific J. Math. 94 (1981), no. 1, 193–206.
- [LM] C. Livingston and P. Melvin. Abelian invariants of satellite knots. Geometry and topology (College Park, Md., 1983/84) pp. 217–227. Lecture Notes in Math., 1167, Springer, Berlin-New York, 1985.

- [Ma] T. Matumoto. On the signature invariants of a non-singular complex sesquilinear form. J. Math. Soc. Japan 29 (1977), no. 1 67–71.
- [M] J. Milnor. Infinite Cyclic Coverings. Topology of Manifolds. Complementary Series in Mathematics vol. 13, ed. J. G. Hocking, Prindle, Weber & Schmidt. Boston, 1968.
- [Mu] K. Murasugi. On a certain numerical invariant of link types. Trans. Amer. Math. Soc. 117 (1965), 387–422.
- [Ri] R. Riley. Growth of order of homology of cyclic branched covers of knots. Bull. London Math. Soc. 22 (1990), no. 3, 287–297.
- [Ro] D. Rolfsen. Knots and Links. Publish or Perish, Berkeley CA (1976).
- [Se] H. Seifert. On the homology invariants of knots. Quart. J. Math., Oxford Ser. (2) 1, (1950), 23–32.
- [St] N. Stoltzfus Unraveling the integral knot concordance group. Memoirs of the AMS (1977), no. 192.
- [T] A. Tristram. Some cobordism invariants for links. Proc. Camb. Phil. Soc. 66 (1969), 251–264.
- [Tr1] H. F. Trotter. Homology of group systems with applications to knot theory. Ann. of Math. (2) 76 (1962), 464–498.
- [Tr2] H. F. Trotter. On S-equivalence of Seifert matrices. Invent. Math. 20 (1973), 173-207.

Department of Mathematics, Indiana University, Bloomington, IN 47405 $E\text{-}mail\ address:\ \texttt{livingst@indiana.edu}$