

In order to state the results in the strongest form, throughout this note, we shall consider a non-trivial smooth 1-dimensional family of K3 surfaces

$$\varphi : \mathcal{X} \rightarrow \Delta := \{t \in \mathbf{C} \mid |t| < 1\}.$$

One may consider the family φ both as a small part of the Kuranishi family of a given K3 surface and as a small part of a given global family of K3 surfaces, such as a universal family of polarized K3 surfaces of a given degree. We regard Δ as a topological space by the Euclidean topology unless stated otherwise. We denote the fiber $\varphi^{-1}(t)$ by \mathcal{X}_t .

For the statement, we choose a marking

$$\tau : R^2\varphi_*\mathbf{Z}_{\mathcal{X}} \rightarrow \Lambda \times \Delta.$$

Here Λ is the K3 lattice, i.e. the lattice isomorphic to $U^{\oplus 3} \oplus E_8(-1)^{\oplus 2}$. Set $\Lambda_t := \tau_t(\text{NS}(\mathcal{X}_t))$. By [Og], one can find a primitive sublattice $\Lambda^0 \subset \Lambda$ and an uncountable dense subset $\mathcal{G} \subset \Delta$ such that

- (1) $\mathcal{S} := \Delta - \mathcal{G}$ is countable and dense;
- (2) $\Lambda^0 \subset \Lambda_t$ for all $t \in \Delta$;
- (3) $\Lambda^0 = \Lambda_t$ for all $t \in \mathcal{G}$;
- (4) $\Lambda^0 \neq \Lambda_t$ for all $t \in \mathcal{S}$.

We call a point $t \in \mathcal{G}$ *generic* and a point $s \in \mathcal{S}$ *special*. Special points are nothing but the points s at which the Picard numbers $\rho(\mathcal{X}_s)$ jump above.

Under this notation, we can state our main result as follows:

Main Theorem. *Assume that $\varphi : \mathcal{X} \rightarrow \Delta$ is projective, i.e. there exists a φ -ample invertible sheaf \mathcal{L} . Then,*

- (1) *There exist a (possibly empty) finite subset $\mathcal{F} \subset \mathcal{S}$, a group G^0 , and a natural number N depending only on φ such that*

$$G^0 < \text{Aut}(\mathcal{X}_t) \text{ for all } t \in \Delta - \mathcal{F}$$

and

$$[G^0 : \text{Aut}(\mathcal{X}_t)] < N \text{ for all } t \in \mathcal{G}.$$

In particular, the map

$$\text{Aut} : \Delta \rightarrow \{\text{groups}\} / \cong; t \mapsto [\text{Aut}(\mathcal{X}_t)]$$

is "upper-semicontinuous" with respect to the co-finite topology of $\Delta - \mathcal{F}$.

- (2) *There exists a subset $\mathcal{D} \subset \mathcal{S}$ such that $\overline{\mathcal{D}} = \Delta$ and that*

$$|\text{Aut}(\mathcal{X}_t)| = \infty \text{ for all } t \in \mathcal{D}.$$

- (3) *There exists a projective family of K3 surfaces $\varphi : \mathcal{X} \rightarrow \Delta$ such that $\mathcal{F} \neq \emptyset$.*

As a direct consequence of the main Theorem, we obtain the following

Corollary. *Let $f : \mathcal{X} \rightarrow \Delta$ be a (not necessarily projective) non-trivial family of K3 surfaces. Then, there is a dense subset $\mathcal{D} \subset \Delta$ such that $|\text{Aut}(\mathcal{X}_t)| = \infty$ for all $t \in \mathcal{D}$. In particular, the nef cone $\overline{A}(\mathcal{X}_t)$ is not finite rational polyhedral if $t \in \mathcal{D}$.*

We shall prove the main Theorem and Corollary in Section 1.

The assertion (2) and Corollary are somewhat surprising. For instance, let us take the component \mathcal{H} of the Hilbertscheme consisting of quartic K3 surfaces and consider the universal family $u : \mathcal{U} \rightarrow \mathcal{H}$. Then for any sufficiently general $\Delta \rightarrow \mathcal{H}$, the induced family $\varphi : \mathcal{X} \rightarrow \Delta$ satisfies that $\text{Pic}(\mathcal{X}_t) = \mathbf{Z}\mathcal{L}_t$ for $t \in \mathcal{G}$, where \mathcal{L}_t is the plane section class of $\mathcal{X}_t \subset \mathbf{P}^3$. So $|\text{Aut}(\mathcal{X}_t)| < \infty$ if $t \in \mathcal{G}$. A bit more strongly, there also exists M with $|\text{Aut}(\mathcal{X}_t)| < M$, for all $t \in \mathcal{G}$ by the boundedness of the finite groups of automorphism groups of K3 surfaces ([Ni1], [Mu] see also [Kn2]). The statement (2) claims, however, that there exists a subset $\mathcal{D} \subset \mathcal{S}$ such that $\overline{\mathcal{S}} = \Delta$ and $|\text{Aut}(\mathcal{X}_t)| = \infty$ for all $t \in \mathcal{D}$ even in this family. This also provides an explicit example of a family in which the automorphism groups actually jump above. On the other hand, one has by Kollár ([Bo]): *If $f : \mathcal{Y} \rightarrow \Delta$ is a family of Calabi-Yau manifolds in $|-K_V|$ of a Fano manifold V , then $\overline{A}(\mathcal{X}_t)$ are finite rational polyhedral and $|\text{Aut}(\mathcal{Y}_t)| < \infty$ for all $t \in \Delta$ provided that $\dim V \geq 4$.* Therefore, the statement similar to Theorem (2) and Corollary do not hold for Calabi-Yau manifolds of higher dimension.

The assertion (1) mostly justifies the phrase quoted at the beginning, while the assertion (3) denies the phrase in the most strict sense. By the statement (1) and Corollary, one might expect something more geometric such as a covariant relation among jumping of automorphism groups, jumping of Picard numbers, those of smooth rational curves and those of elliptic pencils in a family of K3 surfaces. However, there are no such relations in general. Indeed, the second aim of this note is to point out the following:

Proposition. *There exist families of projective K3 surfaces $\varphi^i : \mathcal{X}^i \rightarrow \Delta$ ($1 \leq i \leq 5$) such that the central fiber \mathcal{X}_0^i satisfies that $\rho(\mathcal{X}_0^i) = 19$, $|\text{Aut}(\mathcal{X}_0^i)| < \infty$, contains finitely many smooth rational curves and admits finitely many elliptic pencils, but that generic fiber \mathcal{X}_t^i satisfies $\rho(\mathcal{X}_t^i) = 3$ and that:*

- (1) $|\text{Aut}(\mathcal{X}_t^1)| = \infty$ and \mathcal{X}_t^1 contains infinitely many smooth rational curves but admits no elliptic pencil.
- (2) $|\text{Aut}(\mathcal{X}_t^2)| = \infty$ and \mathcal{X}_t^2 contains no smooth rational curve but admits infinitely many elliptic pencils.
- (3) $|\text{Aut}(\mathcal{X}_t^3)| = \infty$ and \mathcal{X}_t^3 has infinitely many smooth rational curves and infinitely many elliptic pencils.
- (4) $|\text{Aut}(\mathcal{X}_t^4)| = \infty$ and \mathcal{X}_t^4 has neither smooth rational curve nor elliptic pencil.
- (5) $|\text{Aut}(\mathcal{X}_t^5)| < \infty$ and \mathcal{X}_t^5 has both (necessarily finitely many) smooth rational curves and elliptic pencils.

We shall construct these families in Section 2 by deforming a very remarkable K3 surface found by Nikulin [Ni2] (See also [Kn1]). In our construction, the morphisms φ^i are not projective.

ACKNOWLEDGEMENT

Main part of work has been done during the author's stay at Bayreuth in March 2001, under financial support by Germany. Idea of work has been brought to the

author by Professor Thomas Peternell through their discussion on another problem (in progress). The author would like to express his best thanks to him for his invitation to Bayreuth as well as valuable discussion.

PROOF OF THE MAIN THEOREM

Proof of Theorem (1). Set

$$l := \tau_t(\mathcal{L}_{\mathcal{X}_t}) \in \Lambda_t (\subset \Lambda).$$

Since l is independent of t , we have $l \in \Lambda^0$. We may assume that the class l is primitive. Let $A(\mathcal{X}_t)$ be the ample cone of \mathcal{X}_t and set

$$A_t := \tau_t(A(\mathcal{X}_t)) \subset \Lambda_t \otimes \mathbf{R},$$

$$A'_t := A_t \cap (\Lambda^0 \otimes \mathbf{R}),$$

$$G_t := \{g \in O(\Lambda^0) \mid g|(\Lambda^{0*}/\Lambda^0) = id, g(l) \subset A'_t\},$$

for each $t \in \Delta$. Here Λ^{0*} is the dual lattice of Λ^0 and we regard Λ^{0*} as an overlattice of Λ^0 via the cup product $(*, **)$, which is non-degenerate on Λ^0 by the Hodge index Theorem. Note that $l \in A'_t$ for each t . For simplicity of description, we often identify Λ_t with $\text{NS}(\mathcal{X}_t)$ by τ_t . For instance, we say that an element $a \in A_t$ is ample on \mathcal{X}_t . The next Claim completes the proof of Theorem (1).

Claim 1. *Choose $p \in \mathcal{G}$. Then, there exists a (possibly empty) finite subset $\mathcal{F} \subset \mathcal{S}$ such that*

- (1) $A'_t \subset A_p$ for all $t \in \Delta$, and $A_t = A_p$ for all $t \in \mathcal{G}$.
- (2) $G_p \subset G_t$ for all $t \in \Delta - \mathcal{F}$, and $G_t = G_p$ for all $t \in \mathcal{G}$.
- (3) *There is a natural embedding $\iota_t : G_t \hookrightarrow \text{Aut}(\mathcal{X}_t)$ for all $t \in \Delta$. Moreover, there exists a natural number N such that $[\iota_t(G_t) : \text{Aut}(\mathcal{X}_t)] \leq N$ for all $t \in \mathcal{G}$.*

Proof. The first assertion is clear if $\text{rank } \Lambda^0 = 1$. We assume $\text{rank } \Lambda^0 \geq 2$. Let $h \in A'_t$. Then $(h^2) > 0$ and $(h, l) > 0$. If h is not ample on \mathcal{X}_p , then there is an irreducible curve C on \mathcal{X}_p such that $([C], h) \leq 0$. This is due to the real version of the Nakai-Moishezon criterion of ampleness by Campana and Peternell [CP]. This C must be a smooth rational curve by the Hodge index Theorem and the adjunction formula. Moreover, since C is a curve on \mathcal{X}_p , we have $[C] \in \Lambda^0$. Thus $[C] \in \Lambda_t$ as well. Since $(C^2) = -2$, either $-[C]$ or $[C]$ represents an effective curve on \mathcal{X}_t by the Riemann-Roch Theorem. Since $([C], l) > 0$ by the ampleness of l plus effectiveness of C on \mathcal{X}_p and since l is also ample on \mathcal{X}_t , it is $[C]$ that is represented by the effective class on \mathcal{X}_t . (Here the assumption φ being projective is essential.) However, since $h \in A_t$, one would then have $([C], h) > 0$, a contradiction to the previous inequality. Here we used the fact that the values $(*, **)$ are independent on which K3 surfaces \mathcal{X}_t we are arguing. Therefore we have $A'_t \subset A_p$ for all $t \in \Delta$. (Note, however, that the other inclusion is false in general as we will see in the proof of (3).) Since $A'_t = A_t$ if $t \in \mathcal{G}$, replacing p by t , one has $A_t = A_p$ for all $t \in \mathcal{G}$. Now, by the definition of G_* , one has $G_t = G_p$ for $t \in \mathcal{G}$ as well.

We shall show the first assertion of (2). From now, we write A_p by A . Set

$$\mathcal{T} := \{t \in \mathcal{S} \mid C_t \not\subset C\}$$

We need to show that \mathcal{F} is finite. Recall that the group G_p is finitely generated, because G_p is a quotient of an arithmetic subgroup of $O(\Lambda^0)$ [BC]. Let $\{g_i | 1 \leq i \leq n\}$ be a set of generators of G_p . Set for each i :

$$\mathcal{F}_i := \{t \in \mathcal{S} | g_i \notin G_t\}.$$

It suffices to show that $|\mathcal{F}_i| < \infty$. Choose i and set $g := g_i$. Assume that $t \in \mathcal{F}_i$. Then, by the definition, $g(l) \notin A_t$. (Note that $g(l) \in \Lambda^0$ by $g \in G_p$.) Since $(g(l)^2) = (l^2) > 0$ and $(g(l), l) > 0$ by $l, g(l) \in A$, we see that $g(l)$ is in the positive cone and that $(g(l), l) > 0$ by the Hodge index Theorem. Then, applying the Riemann-Roch Theorem, there exists an effective divisor D on \mathcal{X}_t such that $g(l) = [D]$. Since $g(l)$ is not ample on a K3 surface \mathcal{X}_t , there exists a smooth rational curve C_t such that $(g(l), C_t) \leq 0$.

Let \tilde{g} be the element of $O(\Lambda)$ such that $\tilde{g}|_{\Lambda^0} = g$ and $\tilde{g}|_{\Lambda^{0\perp}} = id$, where $\Lambda^{0\perp}$ is the orthogonal complement of Λ^0 in Λ . Such an extension exists, because $g|_{\Lambda^{0*}/\Lambda^0} = id$. Since $T_{\mathcal{X}_t} \subset \Lambda^{0\perp}$, where $T_{\mathcal{X}_t}$ is the transcendental lattice of \mathcal{X}_t , this \tilde{g} is also an Hodge isometry of \mathcal{X}_t . Thus, $E := \tilde{g}^{-1}([C_t])$ is an integral $(1, 1)$ -class on \mathcal{X}_t and satisfies $([E]^2) = -2$. In particular, $[E]$ or $-[E]$ is represented by an effective curve on \mathcal{X}_t by the Riemann-Roch Theorem. Therefore, $(g(l), C_t) = (l, E) \neq 0$ by the ampleness of l , and we have $(g(l), C_t) < 0$. Then $C_t \subset \text{Supp } D$ and one has $(l, C_t) \leq (l, g(l))$. Note that $(l, g(l))$ is independent on t . So, if there are infinitely many t such that $t \in \mathcal{F}_i$, then there exists a component \mathcal{H} of the relative Hilbertscheme $\text{Hilb}_{\mathcal{X}/\Delta}^l$ to which infinitely many of C_t belong. This \mathcal{H} must dominates Δ and one has then an effective curve C on \mathcal{X}_p such that $([C], g(l)) = ([C_t], g(l)) < 0$ (for C and C_t belonging to \mathcal{H}). However, this contradicts the ampleness of $g(l)$ on \mathcal{X}_p . Hence $|\mathcal{F}_i| < \infty$ for each i and we are done for (2).

We shall show the assertion (3). As before, each element $g_t \in G_t$ extends an effective Hodge isometry \tilde{g}_t of \mathcal{X}_t such that $\tilde{g}_t|_{\Lambda^0} = g_t$ and $\tilde{g}_t|_{\Lambda^{0\perp}} = id$. There then exists a unique $f_t \in \text{Aut}(\mathcal{X}_t)$ such that $f_t^*|_{\Lambda^0} = g_t$ and $f_t^*|_{\Lambda^{0\perp}} = id$ by the global Torelli Theorem for K3 surfaces. The map ι_t is given by $g_t \mapsto f_t$. Now the assertion follows from the next facts:

- (1) $\tau_t(\text{NS}(\mathcal{X}_t)) = \Lambda^0$ for $t \in \mathcal{G}$;
- (2) Λ^{0*}/Λ^0 is finite (and is independant of t);
- (3) The subgroup $\text{Ker}(\text{Aut}(\mathcal{X}_t) \rightarrow O(\text{NS}(\mathcal{X}_t)^\perp))$ of $\text{Aut}(\mathcal{X}_t)$ is of finite index less than or equal to $66 \times |\text{Aut}(\Lambda^{0*}/\Lambda^0)|$ if $t \in \mathcal{G}$ ([Ni1], see also [St]). \square

Now we are done for Theorem (1). \square

Proof of Theorem (2). We may consider the case $|\text{Aut}(\mathcal{X}_t)| < \infty$ for all $t \in \mathcal{G}$. (Indeed, if otherwise, one has $|\text{Aut}(\mathcal{X}_t)| = \infty$ for all $t \in \Delta - \mathcal{F}$ by Theorem (1) and may put $\mathcal{D} = \mathcal{S} - \mathcal{F}$.) Since \mathcal{G} is dense, the statement will follow from:

Claim 2. *For any given $p \in \mathcal{G}$ and any open neighbourhood $p \in U$, there exists $t \in \mathcal{S} \cap U$ such that $|\text{Aut}(\mathcal{X}_t)| = \infty$.*

Proof of Claim 2. For abuse of notation, we shall write again $U = \Delta$ and $\varphi|_U = \varphi$. Assuming $|\text{Aut}(\mathcal{X}_t)| < \infty$ for all $t \in \Delta$, we shall derive a contradiction. Set $\mathcal{S} = \text{graph } \Lambda$. By shrinking, we may assume that $\mathcal{T} = \emptyset$. We shall argue dividing

into the following two cases (1) and (2) (by shrinking again, these two cover all the possible cases):

- (1) There exists a sequence $\{t_n\}_{n \geq 1} (\subset \mathcal{S})$ such that $\text{rank } \Lambda_{t_n} \geq 3$ for all n and that $\lim_{n \rightarrow \infty} t_n = p$;
- (2) $r_0 = 1$ and $\text{rank } \Lambda_t = 2$ for all $t \in \mathcal{S}$.

Case 1. Since $\text{rank } \Lambda_{t_n} \geq 3$ and $|\text{Aut}(\mathcal{X}_{t_n})| < \infty$, by [PSS], \mathcal{X}_{t_n} contains only finitely many smooth rational curves, say,

$$C_{n1}, C_{n2}, \dots, C_{nk(n)}.$$

By [Kv], the classes $[C_{ni}]$ generate the Kleiman-Mori cone $N_n := \overline{NE}(\mathcal{X}_{t_n})$ of \mathcal{X}_{t_n} . Recall also that the isomorphism classes of the Picard lattices and the dual graph of smooth rational curves of K3 surfaces with $|\text{Aut}(X)| < \infty$ and $\rho(X) \geq 3$ belong to finitely many of them. This deep result is due to Nikulin [Ni 2, 3, 4]. (*It is this finiteness that we need the case assumption essentially; remark that this finiteness is false if $\rho \leq 2$.*) Therefore we may assume that $\rho := \text{rank } \Lambda_{t_n}$ and $k := k(t_n)$ are constant and the intersection matrices (C_{ni}, C_{nj}) are also constant. (*However, we do not know apriori whether the classes $[C_{ni}]$ form constant subsystem of $\Lambda \times \Delta$.*)

Consider first the case of $r_0 \geq 3$. Then \mathcal{X}_p has finitely many smooth rational curves as well by $|\text{Aut}(\mathcal{X}_p)| < \infty$. Denote all of them by D_j ($1 \leq j \leq k'$). Then one has $l = \sum_{j=1}^{k'} a_j D_j$, where a_j are non-negative rational numbers. Since $[D_j] \in \Lambda^0 \subset \Lambda_t$, each D_j extends \mathbf{P}^1 -bundles $\mathcal{D}_j \rightarrow \Delta$ near p . (See for instance [Og].) Thus, for large n , one has $\mathcal{D}_{jt_n} \in \{C_{ni}\}_{i=1}^k$ and $l = \sum_{j=1}^{k'} a_j \mathcal{D}_{jt_n}$. Then the set $\{(C_{ni}, l)\}_{n \geq 1}$ is bounded for all i by the finiteness of dual graphs. Thus, using the existence of relative Hilbertscheme and arguing as before, one finds effective divisors C_{0i} ($i = 1, 2, \dots, k$) on \mathcal{X}_p whose dual graph is same as C_{ni} . However, we would then have $\text{rank } \Lambda^0 \geq \text{rank } \Lambda_t > r_0$, a contradiction.

Next consider the case of $r_0 = 2$. By $|\text{Aut}(\mathcal{X}_p)| < \infty$, the Kleiman-Mori cone N_p of \mathcal{X}_p is generated by either (1) two classes of smooth rational curves, (2) two classes of elliptic pencils, (3) the class of a smooth rational curve and the class of elliptic pencil, or (4) the class of a smooth rational curve and an irrational ray $\mathbf{R}_{\geq 0}e$ with $(e^2) = 0$ (See for instance [Kv]). In the first three cases, the same argument as above immediately gives a contradiction. Indeed, it suffices to observe the following facts instead:

- (1) Each elliptic pencil on \mathcal{X}_p also extends to a family of elliptic pencils near p . This is because $\Lambda^0 \subset \Lambda_t$. (See for instance [Og]).
- (2) Each elliptic pencil on \mathcal{X}_{t_n} has at least one fiber consisting of smooth rational curves. This follows from Shioda's formula [Sh] of the Mordell-Weil rank of the Jacobian and the assumption $\text{rank } \Lambda_{t_n} \geq 3$. Indeed, the Mordell-Weil rank must be 0 by $|\text{Aut}(\mathcal{X}_{t_n})| < \infty$.

So, we may consider the last case. Since the smooth rational curve C on \mathcal{X}_p extends sideways (by $[C] \in \Lambda^0$), we may assume that $[C] = [C_{n1}]$ for large n . This $[C]$ generates one of two boundary rays of N_p . Since A_{t_n} is finite rational polyhedral, so is $A'_n := A_{t_n} \cap (\Lambda^0 \otimes \mathbf{R})$. Therefore, by the case assumption, one has $A'_n \neq A_p$ (but one has $A'_n \subset A_p$ by Claim 1 (1).) There then exist a nef and big divisor H_n on \mathcal{X}_{t_n} and a smooth rational curve C_n on \mathcal{X}_{t_n} such that $(H_n, C_n) = 0$ and that

$[H_n] \in A_p$. (Note that $([H_n], [C_{n1}]) = ([H_n], [C]) > 0$ because $[H_n] \in A_p$.) If the set $\{\|C_{n2}\|\}_{n \geq 1}$ is bounded, then, since there are only finitely many lattice points with bounded norm, the set of intersection numbers $\{(C_{n2}, l)\}_{n \geq 1}$ is bounded as well. Then, by the existence of the relative Hilbertscheme, \mathcal{X}_p contains effective divisor D_{02} such that $[D_{02}] = [C_{n2}]$. However, the class $[D_{02}]$ gives then another rational boundary of N_p , a contradiction to the case assumption. Therefore, $\{\|C_{n2}\|\}_{n \geq 1}$ is unbounded. Then, passing to a subsequence, we have $\lim_{n \rightarrow \infty} \|C_{n2}\| = \infty$. Set $x_{n2} := C_{n2}/\|C_{n2}\|$. Since $\|x_{n2}\| = 1$, by passing to some subsequence, we find the limit $x_{02} := \lim_{n \rightarrow \infty} x_{n2}$ in $\Lambda \otimes \mathbf{R}$. We calculate that $\|x_{02}\| = 1$ and that

$$(x_{02}^2) = \lim_{n \rightarrow \infty} -2/\|C_n\| = 0.$$

Moreover, we calculate that

$$(x_{02} \cdot [\omega_{\mathcal{X}_p}]) = \lim_{n \rightarrow \infty} (x_{n2} \cdot [\omega_{\mathcal{X}_{t_n}}]) = 0.$$

Here ω_* is the 2-form on \mathcal{X}_* given by the base change of a local section ω of $\varphi_* \Omega_{\mathcal{X}/\Delta}^2$. Thus, x_{02} is also a real $(1, 1)$ -class on \mathcal{X}_p . Set $h_n := H_n/\|H_n\|$. Since $h_n \in A_p$, passing to a subsequence, we have the limit

$$h_0 := \lim_{n \rightarrow \infty} h_n \in \overline{A}_p.$$

Since $(h_n, x_{n2}) = 0$, one has $(h_0, x_{02}) = 0$ as well. Then, by the Hodge index Theorem, one has that $x_{02} = \alpha h_0$ and $x_{02} \in \overline{A}_p$. In particular, $\mathbf{R}_{>0} x_{02}$ gives another boundary of N_p . On the other hand, since (C_{n1}, C_{n2}) is constant, one calculates

$$([C], x_{02}) = \lim_{n \rightarrow \infty} ([C_{n1}], [C_{n2}])/\|C_{n2}\| = 0.$$

Since Λ^0 is of rank 2, this implies $\mathbf{R} x_{02} = [C]^\perp$. However, $\mathbf{R}_{>0} x_{02}$ would be then a rational boundary, a contradiction to the case assumption.

Finally consider the case where $r_0 = 1$. If $\{\|C_{ni}\|\}_{n \geq 1}$ is bounded for some i , then, as before, $\{(C_{ni}, l)\}_{n \geq 1}$ is bounded and we have an effective curve D such that $[D] = [C_{ni}]$ for large n . This class $[D]$ satisfies $[D] \in \text{NS}(\mathcal{X}_p)$ and $([D]^2) = -2$. However, this is impossible, because \mathcal{X}_p is projective and $\text{rank NS}(\mathcal{X}_p) = r_0 = 1$. Therefore the sets $\{\|C_{ni}\|\}_{n \geq 1}$ are unbounded for all i . We set $x_{ni} := C_{ni}/\|C_{ni}\|$. Passing to a subsequence, we have $\lim_{n \rightarrow \infty} \|C_{ni}\| = \infty$ and get the limit points $x_{0i} := \lim_{n \rightarrow \infty} x_{ni}$. Since (C_{ni}, C_{nj}) are all bounded, we have that $(x_{0i}^2) = (x_{0i}, x_{0j}) = 0$ for all i and j . Moreover, by the same argument as in the previous case, we see that x_{0i} are all real $(1, 1)$ -classes on \mathcal{X}_p . Then, by the Hodge index Theorem, there exists e with $(e^2) = 0$ such that $x_{0i} \in \mathbf{R}e$ for all i . Then, the Kleiman-Mori cones N_n , which are finite polyhedral cones generated by x_{ni} , accumulate to the ray $\mathbf{R}e$ under $n \rightarrow \infty$. However, since $l \in N_n$, we would then have $l \in \mathbf{R}e$, a contradiction to $(l^2) > 0$. Now we are done in Case 1. \square

Case 2. Recall that if $\text{rank } \Lambda_t = 2$, then $|\text{Aut}(\mathcal{X}_t)| = \infty$ if and only if the quadratic form of Λ_t represents neither 0 nor -2 . Let us consider the period map

For each $a \in \Lambda$, we set

$$H_a := \{[\omega] \in \mathcal{P} \mid (a, \omega) = 0\} \subset \mathcal{P}.$$

Then $\pi(\Delta) \subset H_l$. Moreover, by the case assumption, we have that $t \in \mathcal{S}$ if and only if there exists $a \in \Lambda - \mathbf{Z}l$ such that $\pi(t) \in H_a$ and that $\Lambda_t = \mathbf{Z}\langle l, a \rangle$. (Recall that l is assumed to be primitive.)

Claim 3. *There exist an element $h \in \Lambda$ and an arbitrarily large natural number n such that $\mathbf{Z}\langle l, na + h \rangle$ is primitive in Λ and represents neither 0 nor -2 .*

Proof. Set $(l, l) = 2A$, $(l, a) = B$ and $(a, a) = 2C$. Since the primitive embedding $\Lambda_t \hookrightarrow \Lambda = U^{\oplus 3} \oplus E_8(-1)^{\oplus 2}$ is unique up to isomorphism, we may assume that $l = e_{11} + Ae_{12}$, $a = Be_{12} + e_{21} + Ce_{22}$, where e_{i1} and e_{i2} ($1 \leq i \leq 3$) are the standard basis of $U^{\oplus 3}$. Set $h := e_{31} - me_{32}$. Then $(h^2) = -2m$ and $(h, l) = (a, l) = 0$. Put:

$$L_{m,n} := \mathbf{Z}\langle l, na + h \rangle.$$

This lattice is primitive in Λ and its quadratic form is

$$Q(x, y) := 2\{Ax^2 + nBxy + (n^2C - m)y^2\}.$$

If $A \geq 2$, we take $n = AN$ and $m = AM$. Then $Q(x, y)$ is divisible by $2A$. Thus $L_{m,n}$ does not represent -2 . Moreover, the discriminant of Q is

$$4A^2\{N^2(B^2 - 4AC) + 4M\}.$$

This is not square if M and N are general. Thus $L_{m,n}$ does not represent 0, either. If $A = 1$, we set $n = 4N$ and $m = 8M$. Then $Q(x, y)$ and its discriminant are

$$Q(x, y) = 2x^2 + 8NBxy + 8(4N^2C - M)y^2;$$

$$D := 64\{N^2(B^2 - 4C) + M\}.$$

Then, $Q(x, y) \equiv 2$ or 0 modulo 8. Thus $L_{m,n}$ does not represent -2 . Moreover, D is not square if M and N are general. Thus $L_{m,n}$ does not represent 0, either. \square

Let us return back to the original situation. If the lattice $\mathbf{Z}\langle l, a \rangle$ represents neither 0 nor -2 then we are done. If otherwise, using Claim 3, we find an element $h \in \Lambda$ and an element $n \in \mathbf{Z}$ such that $\mathbf{Z}\langle l, na + h \rangle$ is primitive in Λ and represents neither 0 nor -2 . Since $H_{na+h} = H_{a+(h/n)}$, the hyperplane H_{na+h} meets $\pi(\Delta)$ at t' where t' being close to t if n is sufficiently large. For this t' , one has $\Lambda_{t'} = \mathbf{Z}\langle l, na + h \rangle$, whence $|\text{Aut}(\mathcal{X}_{t'})| = \infty$. \square

Now we are done for Theorem (2). \square

Proof of Theorem (3). Let S be a K3 surface such that $\text{NS}(S) = U \oplus A_1(-1)$ and $|\text{Aut}(S)| < \infty$. Such a K3 surface exists by [Ni2]. Let us consider the Kuranishi family $u : \mathcal{U} \rightarrow \mathcal{K}$ of S . Then choosing a marking

$$\tau : R^2u_*\mathbf{Z}_{\mathcal{U}} \rightarrow \Lambda \times \mathcal{K},$$

one can define the period map

Since π is a local isomorphism by the local Torelli Theorem for K3 surfaces, we may identify \mathcal{K} with a small open set of the period domain \mathcal{P} (denoted again by \mathcal{P}) through π . Let Π be a general 2-plane in $U \oplus A_1(-1)$. Then Π represents neither 0 nor -2 . Let us take the subset $\mathcal{K}^0 \subset \mathcal{K} = \mathcal{P}$ defined by the equations

$$(a, \omega) = (b, \omega) = 0,$$

where a and b form an integral basis of Π . Consider the induced family $u^0 : \mathcal{U}^0 \rightarrow \mathcal{K}^0$. Here \mathcal{K}^0 is of dimension 18. By construction, we have $0 \in \mathcal{K}^0$ and $\text{NS}(\mathcal{U}_t^0) \simeq \Pi$, whence $|\text{Aut}(\mathcal{U}_t^0)| = \infty$ for generic t . Then, the induced family $\varphi : \mathcal{X} \rightarrow \Delta$ for general $0 \in \Delta \subset \mathcal{K}$ satisfies $0 \in \mathcal{F}$. In this example, we have $A'_0 \neq A_t$ for t being generic. Indeed, $\partial A'_0 = \partial(A_0 \cap \Pi)$ is rational, while ∂A_t is irrational. \square

Now we are done. Q.E.D. of the main Theorem. \square

PROOF OF COROLLARY

Since an ample divisor on \mathcal{X}_p extends sideways if $p \in \mathcal{G}$, f is locally projective around p . The result now follows from the Theorem together with the density of \mathcal{G} . \square

PROOF OF PROPOSITION

We shall construct explicit examples with required properties by deforming a remarkable K3 surface found by Nikulin and Kondo ([Ni2], [Ko1]). This is a K3 surface S which satisfies the following properties:

- (1) $\rho(S) = 19$ and $\text{NS}(S) = U \oplus E_8(-1) \oplus E_8(-1) \oplus A_1(-1)$ as a lattice,
- (2) S contains exactly 24 smooth rational curves and finitely many (> 0) elliptic pencils;
- (3) $\text{Aut}(S)$ is a finite group and is isomorphic to $S_3 \times \mu_2$.

As before, let us take the Kuranishi family $u : \mathcal{U} \rightarrow \mathcal{K}$ of S and choose a marking

$$\tau : R^2 u_* \mathbf{Z}_{\mathcal{U}} \rightarrow \Lambda \times \mathcal{K}.$$

Then $S = \mathcal{U}_0$ and one has the period map

$$\pi : \mathcal{K} \rightarrow \mathcal{P}.$$

As before, we identify \mathcal{K} with a small open set of the period domain \mathcal{P} (denoted again by \mathcal{P}) through π . Let

$$e, f, v_{ij} (i = 1, 2, 1 \leq j \leq 8) v$$

be the elements of Λ corresponding to the standard integral basis of $\text{Pic}(S) = U \oplus E_8(-1) \oplus E_8(-1) \oplus A_1(-1)$.

Proof of (1). Let us take the subset $0 \in \mathcal{K}^1 \subset \mathcal{K} = \mathcal{P}$ defined by the equations

where n is an integer non-divisible by 3. Consider the induced family $u^1 : \mathcal{U}^1 \rightarrow \mathcal{K}^1$. Here \mathcal{K}^1 is of dimension 17. By construction, $\text{NS}(\mathcal{U}_t^1) \simeq \langle e + 3nf, v_{11}, v_{21} \rangle$ and the intersection matrix is:

$$\begin{pmatrix} 6n & 0 & 0 \\ 0 & -2 & 0 \\ 0 & 0 & -2 \end{pmatrix},$$

if $t \in \mathcal{U}^1$ is generic. The discriminant of this matrix is $24n$. Since the quadratic form

$$Q_1(x, y) := 6nx^2 - 2y^2 - 2z^2$$

does not represent 0, \mathcal{U}_t^1 has no elliptic pencil. Since there are only finitely many isomorphism classes of lattices of rank 3 which are isomorphic to the Picard lattices of K3 surfaces X with $|\text{Aut}(X)| < \infty$ [Ni3], one has $|\text{Aut}(\mathcal{U}_t^1)| = \infty$ for n being sufficiently large. Moreover, \mathcal{U}_t^1 admits at least one smooth rational curve. This follows from the fact that $Q_1(x, y)$ represents -2 plus the Riemann-Roch Theorem. Then \mathcal{U}_t^1 contains infinitely many smooth rational curves by $|\text{Aut}(\mathcal{U}_t^1)| = \infty$ and by [PSS]. (See also [St] and [Kv].) Now the induced family over a generic linear section $0 \in \Delta \subset \mathcal{K}^1$ satisfies the required property of (1). \square

Proof of (2). Let us take the subset $0 \in \mathcal{K}^2 \subset \mathcal{K} = \mathcal{P}$ defined by the equations

$$(e + 2f, \omega) = (v_{11} + v_{13}, \omega) = (v_{21} + v_{23}, \omega) = 0.$$

Consider the induced family $u^2 : \mathcal{U}^2 \rightarrow \mathcal{K}^2$. Here \mathcal{K}^2 is of dimension 17. By construction, $\text{NS}(\mathcal{U}_t^2) \simeq \langle e + 2f, v_{11} + v_{13}, v_{21} + v_{23} \rangle$ and the intersection matrix is:

$$\begin{pmatrix} 4 & 0 & 0 \\ 0 & -4 & 0 \\ 0 & 0 & -4 \end{pmatrix},$$

if $t \in \mathcal{U}^2$ is generic. Since the quadratic form

$$Q_2(x, y) := 4x^2 - 4y^2 - 4z^2$$

does not represent -2 , \mathcal{U}_t^2 contains no smooth rational curve. In particular, the ample cone coincides with the positive cone. This already implies $|\text{Aut}(\mathcal{U}_t^2)| = \infty$, because the automorphism group is then isomorphic to the orthogonal group of the Picard lattice up to finite groups [SPP]. Moreover, since $Q_2(x, y)$ admits infinitely many primitive integral zero's, \mathcal{U}_t^2 admits infinitely many elliptic pencils. Now the induced family over a generic linear section $0 \in \Delta \subset \mathcal{K}^2$ satisfies the required property of (2). \square

Proof of (3). Let us take the subset $\mathcal{K}_3 \subset \mathcal{K} = \mathcal{P}$ defined by the equations

$$(e, \omega) = (f, \omega) = (v_{11} + v_{13} + v_{15} + v_{17}, \omega) = 0.$$

Consider the induced family $u^3 : \mathcal{U}^3 \rightarrow \mathcal{K}^3$. Here \mathcal{K}^3 is of dimension 17. By construction, one has $\text{NS}(\mathcal{U}_t^3) \simeq U \oplus \langle -8 \rangle$ if $t \in \mathcal{U}^3$ is generic. Then X admits a Jacobian pencil $f : X \rightarrow \mathbf{P}^1$, i.e. elliptic pencil with section, given by U . Here we set $X := \mathcal{U}_t^3$. Since f has no reducible fibers, the Mordell-Weil lattice $M(f)$ of f is isomorphic to the lattice $\langle 8 \rangle$. (See the excellent survey of Shioda [Sh] for details.)

for basic properties of the Mordell-Weil lattice.) Let g be the generator of $M(f)$ corresponding to -8 . Then g defines an automorphism of X of infinite order. In particular, $\text{Aut}(X)$ is infinite. Let C_0 be the zero-section of f and let C_1 be the section corresponding to g . Then $C_1 = g(C_0)$ and $g^n(C_0)$ are mutually distinct smooth rational curves on X . On the other hand, by the definition of the height pairing $\langle *, * \rangle$, we calculate

$$8 = \langle C_1, C_1 \rangle = 4 + 2\langle C_0, C_1 \rangle.$$

This gives $\langle C_0, C_1 \rangle = 2$ and $\langle C_n, C_{n+1} \rangle = 2$. Thus, $C_n + C_{n+1}$ is a divisor of Kodaira type I_2 or III and defines an elliptic pencil

$$f_n := \Phi_{|C_n + C_{n+1}|} : X \rightarrow \mathbf{P}^1.$$

Since an elliptic pencil on K3 surface admits at most 24 singular fibers, one has $|\{f_n\}_{n \geq 1}| = \infty$. Combining all these together, one finds that \mathcal{U}_t^3 has $\rho(\mathcal{U}_t^3) = 3$, infinitely many smooth rational curves, infinitely many elliptic pencils, and satisfies $|\text{Aut}(\mathcal{U}_t^3)| = \infty$ if $t \in \mathcal{K}^3$ is generic. Now the induced family over a generic linear section $0 \in \Delta \subset \mathcal{K}^3$ satisfies the required property of (3). \square

Proof of (4). Let us take the subset $\mathcal{K}^4 \subset \mathcal{K} = \mathcal{P}$ defined by the equations

$$(e + 6f, \omega) = (v_{11} + v_{13}, \omega) = (v_{21} + v_{23}, \omega) = 0.$$

Consider the induced family $u^4 : \mathcal{U}^4 \rightarrow \mathcal{K}^4$. Here \mathcal{K}^4 is of dimension 17. By construction, $\text{NS}(\mathcal{U}_t^4) \simeq \langle e + 6f, v_{11} + v_{13}, v_{21} + v_{23} \rangle$ and the intersection matrix is:

$$\begin{pmatrix} 12 & 0 & 0 \\ 0 & -4 & 0 \\ 0 & 0 & -4 \end{pmatrix},$$

if $t \in \mathcal{K}^4$ is generic. Since the quadratic form

$$Q_4(x, y) := 12x^2 - 4y^2 - 4z^2$$

represents neither 0 nor -2 , \mathcal{U}_t^4 has neither smooth rational curve nor elliptic pencil. Then as before, the ample cone coincides with the positive cone and one has $|\text{Aut}(\mathcal{U}_t^4)| = \infty$ by [PSS]. Now the induced family over a generic linear section $0 \in \Delta \subset \mathcal{K}^4$ satisfies the required property of (4). \square

Proof of (5). Let us take the subset $\mathcal{K}^5 \subset \mathcal{K} = \mathcal{P}$ defined by the equations

$$(e, \omega) = (f, \omega) = (v, \omega) = 0.$$

Consider the induced family $u^5 : \mathcal{U}^5 \rightarrow \mathcal{K}^5$. Here \mathcal{K}^5 is of dimension 17 and one has $\text{NS}(\mathcal{U}_{1,t}) = U \oplus A_1(-1)$ for generic $t \in \mathcal{U}_5$ and \mathcal{U}_t^5 enjoys all the properties required in (5) by [Ni2]. Now the induced family over a generic linear section $0 \in \Delta \subset \mathcal{K}_2$ satisfies the required property of (5). Note that no element $h \in U \oplus A_1(-1)$ is ample on $S = \mathcal{U}_0^5$, because such h is orthogonal to $E_8(-1) \oplus E_8(-1)$. Therefore, the family is not projective. \square

Now we are done. Q.E.D. for Proposition \square

REFERENCES

- [Bo] C. Borcea, *Homogeneous vector bundles and families of Calabi-Yau threefolds, II*, Proc. Sym. Pure Math. **52** (1991), 83–91.
- [BC] A. Borel and H. Chandra, *Arithmetic subgroups of algebraic groups*, Ann. Math **75** (1962), 485–535.
- [BPV] W. Barth, C. Peters, A. Van de Ven, *Compact complex surfaces*, Springer-Verlag (1984).
- [CP] F. Campana and T. Peternell, *Algebraicity of the ample cone of projective varieties*, J. reine angew. Math. **407** (1990), 160–166.
- [Gr] A. Grothendieck, *Fondements de la Géométrie Algébrique*, Sec. Math. Paris (1962).
- [Ke] J.H. Keum, *Automorphisms of Jacobian Kummer surfaces*, Compositio Math. **107** (1997), 269–288.
- [KK] J.H. Keum and S. Kondo, *The automorphism groups of Kummer surfaces associated with the product of two elliptic curves*, Trans. Amer. Math. Soc. **353** (2001), 1469–1487.
- [Kl] J. Kollár, *Rational curves on algebraic varieties. A series of Modern Surveys in Mathematics*, Springer-Verlag **32** (1996).
- [Kn1] S. Kondo, *Algebraic K3 surfaces with finite automorphism groups*, Nagoya Math. J. **116** (1989), 1–15.
- [Kn2] S. Kondo, *Niemeyer Lattices, Mathieu groups, and finite groups of symplectic automorphisms of K3 surfaces*, Duke Math. J. **92** (1998), 593–598.
- [Kn3] S. Kondo, *The automorphism groups of a generic Kummer surface*, J. Alg. Geom. **7** (1998), 589 – 609.
- [Kv] S. Kovacs, *The cone of curves of a K3 surfaces*, Math. Ann. **300** (1994), 681–691.
- [Mu] S. Mukai, *Finite groups of automorphisms of K3 surfaces and the Mathieu group*, Invent. Math. **94** (1988), 183–221.
- [Ni1] V. V. Nikulin, *Finite automorphism groups of Kähler K3 surfaces*, Trans. Moscow Math. Soc. **38** (1980), 71–135.
- [Ni2] V. V. Nikulin, *On the quotient groups of the automorphism groups of hyperbolic forms by the subgroups generated by the 2-reflections*, J. Soviet Math. **22** (1983), 1401–1476.
- [Ni3] V. V. Nikulin, *Surfaces of type K3 with finite automorphism groups and a Picard group of rank three*, Proc. Steklov Institute Math. **3** (1985), 131–155.
- [Ni4] V. V. Nikulin, *Discrete reflection groups in Lobachevsky spaces and algebraic surfaces, Proceedings of the International Congress of Mathematics, Berkley 1986*, Amer. Math. Soc. (1987), 654–671.
- [Ni5] V. V. Nikulin, *A remark on algebraic surfaces with polyhedral Mori cone*, Nagoya Math. J. **157** (2000), 73–92.
- [Og] K. Oguiso, *Picard numbers in a family of hyperkähler manifolds - a supplement to the article of R. Borcherds, L. Katzarkov, T. Pantev, N. I. Shephers-Barron, preprint (2000; AG/0011259)..*
- [PSS] I. Piatetski-Shapiro and I. R. Shafarevich, *A Torelli theorem for algebraic surfaces of type K3*, Math. USSR Izv. **5** (1971), 547–587.
- [Sh] T. Shioda, *Theory of Mordell-Weil lattices: in Proceedings of the International Congress of Mathematicians (Kyoto 1990)*, Math. Soc. Japan (1991), 473 – 489.
- [St] H. Sterk, *Finiteness results for algebraic K3 surfaces*, Math. Z. **189** (1985), 507–513.
- [Vi] E. B. Vinberg, *The two most algebraic K3 surfaces*, Math. Ann. **265** (1983), 1–21.

DEPARTMENT OF MATHEMATICAL SCIENCES UNIVERSITY OF TOKYO 153-8914 KOMABA ME-GURO TOKYO JAPAN

E-mail address: oguiso@ms.u-tokyo.ac.jp