

The dynamical zeta function and transfer operators for the Kac-Baker model

J. Hilgert* and D. Mayer†

October 30, 2001

1 Introduction

Dynamical zeta functions seem to play a rather special role in the general theory of zeta functions. They share many properties with the zeta functions of number theory and algebraic geometry and have lead to new approaches to some of the outstanding open problems in the theory of these functions. A well known such problem in this theory is the general Riemann hypothesis for these functions and a possible spectral interpretation of their zeros and poles in terms of some dynamical operator attached to the zeta function. There are several examples like the Selberg or the Artin-Mazur zeta functions where such an operator exists and which furthermore is indeed related to some dynamical system. In Selberg's case the underlying dynamical system is the geodesic flow on a surface of constant negative curvature and the operator in question is the transfer operator of this flow (see [Ma91]). For the Artin-Weil function the dynamical system is the Frobenius map on an algebraic variety and the operator is again the transfer operator for this map (see [Ro86], [Ru92]). One should also mention the recent approaches to Riemann's zeta function and more general L -functions by A. Connes ([Co96]) and C. Deninger ([De99]) where also a dynamical interpretation of these zeros is looked for.

In special cases there is even a simple physical interpretation of the zeros of such a zeta function in terms of energy eigenvalues of a hamiltonian system, like in the Selberg case, so that it is not surprising that also ideas from the theory of quantum chaos are being applied now to the Riemann zeros, which should be connected to the spectrum of some hamiltonian whose classical limit should be chaotic (cf. [Be86]). One could therefore speculate if not all the known zeta functions finally turn out to be dynamical zeta functions.

In the same spirit one can interpret also the work of M. Gutzwiller in [Gu82] on the semiclassical quantization of a particle moving in the anisotropic Kepler potential. There he related the energy spectrum of this particle to the zeros of the Ruelle zeta function of an abstract dynamical system known in ergodic theory as a subshift of finite type over two symbols. In statistical mechanics language this is a lattice spin system of Ising type with a 2-body interaction of the spins decaying exponentially fast with distance on the lattice. The system is known in the physical literature as the Kac-Baker model (cf. [Ka59] and [Ba61]). These authors introduced this model for a better understanding of phase transitions in weak long-range systems like the van der Waals gas.

It turns out that the Ruelle zeta function of this system, which is just a generating function for the finite lattice partition functions of the model, can be expressed in terms of Fredholm

*Institut für Mathematik, Technische Universität Clausthal, 38678 Clausthal-Zellerfeld, Germany. E-mail: hilgert@math.tu-clausthal.de

†MPI für Mathematik, D-53111 Bonn, on sabbatical from Institut für Theoretische Physik, Technische Universität Clausthal, 38678 Clausthal-Zellerfeld, Germany. E-mail: mayer @mpim-bonn.mpg.de or dieter.mayer@tu-clausthal.de

determinants of a transfer operator (see [Ka66], [Ma80]) and hence its zeros and poles have a nice spectral interpretation. There exist indeed two completely different such operators for this system: one is the Ruelle operator introduced for general lattice spin systems in dimension one with a rather straightforward physical interpretation (see [Ru68]), the other one was found by M. Kac (see [Ka66]), using the very special nature of the interaction of this model and its relation to the Ornstein-Uhlenbeck process. Indeed Kac considered his operator himself a nice “trick devoid of any physical significance”.

Whereas Ruelle’s operator is acting in a Banach space of observables closely related to the lattice spin system, the Kac operator is an abstract integral operator in the Hilbert space $L^2(\mathbb{R}, dx)$ not directly related to the spin system. It has a rather complicated kernel \mathcal{K}_β which in the form used also by Gutzwiller in [Gu82] is given by

$$\mathcal{K}_\beta(\xi, \eta) = \left(\frac{\cosh(\sqrt{\beta J} \xi) \cosh(\sqrt{\beta J} \eta)}{\pi \sinh \gamma} \right)^{\frac{1}{2}} \exp \left(-\frac{1}{4} \left(\tanh \frac{\gamma}{2} \right) (\xi^2 + \eta^2) - \frac{(\xi - \eta)^2}{4 \sinh \gamma} \right), \quad (1)$$

where β denotes “inverse temperature” and $J > 0$ and $0 < \gamma < 1$ are physical parameters characterizing the strength of the interaction respectively its decay rate as a function of distance. We should mention that the kernel used by Kac in [Ka66] is slightly different from the one given above since he worked with so called fixed boundary conditions whereas we will use periodic boundary conditions as Gutzwiller did in [Gu82].

The Ruelle operator $\mathcal{L}_\beta : B(D) \rightarrow B(D)$ on the other hand is acting in a Banach space of smooth observables holomorphic in some disc D in the complex plane and has the following rather simple looking form

$$\mathcal{L}_\beta g(z) = e^{\beta J z} g(\lambda + \lambda z) + e^{-\beta J z} g(-\lambda + \lambda z),$$

where $\lambda = \exp(-\gamma)$. Various aspects of this operator have been studied in [Ma80], [ViMa77] and [PTMT94]. For instance, it defines a family of nuclear operators holomorphic in the entire complex β -plane.

The Ruelle zeta function $\zeta_R(z, \beta)$ of the subshift of finite type (Ω_+, τ) , where $\Omega_+ = \{1, -1\}^{\mathbb{Z}_+}$ denotes the configuration space over the two symbols $\{1, -1\}$ and τ the shift on the lattice \mathbb{Z}_+ , is then defined as

$$\zeta_R(z, \beta) = \exp \sum_{n=1}^{\infty} \frac{z^n}{n} Z_n(\beta), \quad (2)$$

where the $Z_n(\beta)$ denote the so called finite lattice partition functions of the Kac-Baker model. In his diploma thesis [Mo89] B. Moritz showed that for real β this zeta function can be expressed through Fredholm determinants of both these two operators as

$$\zeta_R(z, \beta) = \frac{\det(1 - z\lambda\mathcal{G}_\beta)}{\det(1 - z\mathcal{G}_\beta)} = \frac{\det(1 - z\lambda\mathcal{L}_\beta)}{\det(1 - z\mathcal{L}_\beta)}, \quad (3)$$

where the operator \mathcal{G}_β is related to the Kac operator \mathcal{K}_β simply by $\mathcal{G}_\beta = \frac{1}{\sqrt{\lambda \exp \beta}} \mathcal{K}_\beta$. From this he concluded that the two operators have indeed the same spectra. However, he could not relate the two operators \mathcal{G}_β and \mathcal{L}_β in an explicit way so that the eigenfunctions of the two operators could be determined from each other directly.

In the present paper we solve this problem and give explicit formulas relating the two operators and their eigenfunctions to each other. Hence the two operators are closely connected to each other and it is not by accident that M. Kac could find his operator. Since the Fredholm determinant of the operator \mathcal{L}_β is an entire function in the complex variable β surprisingly also the Fredholm determinant of the operator \mathcal{G}_β can be extended to an entire function in β . This is certainly not true for the operator \mathcal{G}_β itself.

Our result could throw some light also on a more general problem arising in the transfer operator method in ergodic theory of dynamical systems: there are several cases known where the Ruelle operator when acting in some Banach space of smooth observables turns out, at least numerically, to have real spectrum for real “temperature” β . Unfortunately, there are no general criteria known in the theory of operators in Banach spaces which would guarantee such a behaviour. There are very special approaches to this problem for special systems (see [Rug94]), in several cases however reality of the spectrum of such transfer operators can be proved by relating the Ruelle operator to some selfadjoint operator in a Hilbert space. A typical example for this is the transfer operator for the Gauss map and its generalizations, which when restricted to a certain Hardy space of holomorphic functions can be conjugated to an integral operator with symmetric kernel given by Bessel functions (see [Ma90], [Is01]). One could therefore speculate if there does exist always such a relation in cases where the Ruelle operator has real spectrum for real β . A complete understanding of this rather mysterious behaviour of transfer operators however is certainly still missing.

In the present example of the Kac-Baker model the Ruelle transfer operator \mathcal{L}_β acting on a Banach space of holomorphic functions in a disc can be restricted to a Hilbert space of entire functions in the complex plane square integrable with respect to the weight function $\mu_t(z) = t \exp(-t |z|^2)$, the so called Fock space $\mathcal{HL}^2(\mathbb{C}, \mu_t)$ for an appropriate value of the parameter t depending obviously on β . This Hilbert space is isomorphic to the space $L^2(\mathbb{R}, d\xi)$ via the Segal-Bargmann transform $B_t : L^2(\mathbb{R}, d\xi) \rightarrow \mathcal{HL}^2(\mathbb{C}, \mu_t)$. It is basically this transformation for $t = 1$ which relates the Kac-Gutzwiller operator \mathcal{G}_β and the Ruelle operator \mathcal{L}_β for real β in a unitary way. For this special parameter value $t = 1$ the Segal-Bargmann transformation will be denoted simply by B and it is given by

$$Bf(z) = 2^{\frac{1}{4}} \int_{\mathbb{R}} f(\xi) \exp(2\pi\xi z - \pi\xi^2 - \frac{\pi}{2}z^2) d\xi.$$

This transformation allows us to relate also the eigenfunctions of the two operators in an explicit way.

By making use of both these two operators we can show that the Ruelle zeta function $\zeta_R(1, \beta)$ has infinitely many zeros and poles on the real axis unless some unexpected cancellations will take place, whereas there are also infinitely many trivial zeros at least for the special value $\lambda = \frac{1}{2}$ on the line $\text{Re}(\beta) = \ln 2$, again if there are not analogous cancellations. Indeed, there are no signs for such cancellations numerically. We expect a similar behaviour also for general $0 < \lambda < 1$. Obviously it would be interesting to know if there exists other zeros of this zeta function in the complex β -plane besides the ones mentioned above. If this is not the case, and indeed we have some numerical hints also for this, this dynamical zeta function would satisfy some kind of Riemann hypothesis well known for many zeta and L -functions of number theory. This would be a further sign for the basic role dynamical zeta functions play in the general theory of zeta functions.

2 The Kac-Gutzwiller and the Ruelle transfer operator

If $F = \{1, -1\}$ denotes the set of possible values of a classical spin variable σ the configuration space Ω_+ of all allowed spin configurations $\underline{\xi}$ on the half lattice $\mathbb{Z}_+ = \{0, 1, 2, \dots\}$ is defined as $\Omega_+ = F^{\mathbb{Z}_+} = \left\{ \underline{\xi} = (\xi_i)_{i \in \mathbb{Z}_+} \mid \xi_i \in F \forall i \in \mathbb{Z}_+ \right\}$. The shift $\tau : \Omega_+ \rightarrow \Omega_+$ is defined as $(\tau \underline{\xi})_i = \xi_{i+1}$ for all $i \in \mathbb{Z}_+$ if $\underline{\xi} = (\xi_i)_{i \in \mathbb{Z}_+} \in \Omega_+$. Consider next the two body interaction $\Phi(\xi_i, \xi_j) = -J \xi_i \xi_j \lambda^{|i-j|}$ between spins on lattice sites i and j , where $J > 0$ determines the strength of the interaction and $0 < \lambda < 1$ is its decay rate. Obviously this interaction decays exponentially fast with distance $|i - j|$. The energy $U_n(\xi)$ of a configuration $\xi \in \Omega_+$ when restricted to a finite sublattice $[0, n - 1]$

of length n is defined as

$$U_n(\underline{\xi}) = \sum_{i=0}^{n-1} \sum_{j=1}^{\infty} \Phi(\xi_i, \xi_{i+j}).$$

A configuration $\underline{\xi} = (\xi_i)_{i \in \mathbb{Z}_+} \in \Omega_+$ is called periodic with period n iff $\xi_{i+n} = \xi_i$ for all $i \in \mathbb{Z}_+$. Denote by Per_n the set of periodic configurations of period n , i.e.

$$\text{Per}_n = \{ \underline{\xi} \in \Omega_+ \mid \tau^n \underline{\xi} = \underline{\xi} \}.$$

The partition functions $Z_n(\beta)$ for the lattice spin system with interaction Φ for the finite sublattices $[0, n-1]$ with periodic boundary conditions are defined as

$$Z_n(\beta) = \sum_{\underline{\xi} \in \text{Per}_n} \exp(-\beta U_n(\underline{\xi})).$$

Inserting the explicit expression for the interaction Φ for the Kac-Baker model we get

$$Z_n(\beta) = \sum_{\underline{\xi} \in \text{Per}_n} \exp \left(\beta J \sum_{k=0}^{n-1} \sum_{j=1}^{\infty} \xi_k \xi_{k+j} \lambda^j \right).$$

The physical properties of the spin system are completely determined by these partition functions. For instance the free energy $f = f(\beta)$ is given by $f(\beta) = -\beta \lim_{n \rightarrow \infty} \frac{1}{n} \ln Z_n(\beta)$. The main problem of statistical mechanics is then to determine the analytic properties of this function in the temperature variable $\beta = \frac{1}{kT}$, where T denotes the absolute temperature and k Boltzmann's constant. A standard procedure for doing this is the so called transfer matrix method. There one looks for a matrix whose leading eigenvalue is closely related to this free energy. Indeed M. Kac found for his model an integral operator acting in some Hilbert space of square integrable functions whose traces can be related to the partition functions $Z_n(\beta)$ and hence determine also the free energy $f(\beta)$. As was mentioned before, Kac in [Ka66] did not work with periodic boundary conditions so that the kernel of his integral operator is slightly different from the operator $\mathcal{K}_\beta(\xi, \eta)$ for periodic boundary conditions given by (1), which we use in this paper and which was used also by M. Gutzwiller in [Gu82]. The kernel $\mathcal{K}_\beta(\xi, \eta)$ defines a trace class operator in the Hilbert space $L^2(\mathbb{R}, d\xi)$. Obviously $\mathcal{K}_\beta(\xi, \eta) = \mathcal{K}_\beta(\eta, \xi)$ and hence the operator \mathcal{K}_β is symmetric for real β . Its spectrum is therefore real for such β -values. In the following we will simply write β for the expression $J\beta$ since J is assumed to be fixed. For $\underline{\xi} \in \text{Per}_n$ denote by $\underline{\xi}^p$ the configuration $\underline{\xi}^p \in \Omega = F^{\mathbb{Z}}$ on the lattice \mathbb{Z} with $\xi_i^p = \xi_i$ for $i \in \mathbb{Z}_+$ and $\xi_{i+n}^p = \xi_i$ for all $i \in \mathbb{Z}$. Then one has

$$\frac{\beta}{2} \sum_{i=0}^{n-1} \sum_{j=-\infty}^{+\infty} \xi_i^p \xi_j^p \exp(-\gamma|i-j|) = \frac{1}{2} \underline{\xi}_n \cdot \mathbb{A} \underline{\xi}_n,$$

where \mathbb{A} is the $(n \times n)$ -matrix with matrix elements

$$\mathbb{A}_{i,j} = \beta \sum_{k=-\infty}^{+\infty} \exp(|i-j+nk|), \quad 0 \leq i, j \leq n-1$$

and $\underline{\xi}_n = (\xi_0, \dots, \xi_{n-1})$. On the other hand the following identity due to H. Cramér (see [Cr46], p. 118), used also by M. Kac and M. Gutzwiller, holds :

$$\exp\left(\frac{1}{2} \underline{\xi}_n \cdot \mathbb{A} \underline{\xi}_n\right) = (2\pi)^{-\frac{n}{2}} (\det \mathbb{A}^{-1})^{\frac{1}{2}} \int_{-\infty}^{+\infty} dz_0 \dots \int_{-\infty}^{+\infty} dz_{n-1} \exp\left(-\frac{1}{2} \underline{z} \cdot \mathbb{A}^{-1} \underline{z}\right) \exp(\underline{\xi}_n \cdot \underline{z})$$

with $\underline{\xi}_n = (\xi_0, \dots, \xi_{n-1})$ and $\underline{z} = (z_0, \dots, z_{n-1})$. From this one easily derives the identity

$$Z_n(\beta) = 2 \sinh\left(\frac{n\gamma}{2}\right) \exp\left(-\frac{n\beta}{2}\right) \text{trace}(\mathcal{K}_\beta^n),$$

where \mathcal{K}_β is the integral operator with kernel $\mathcal{K}_\beta(\xi, \eta)$ as given in the introduction, and where we used the identity $\frac{1}{2}(\underline{\xi}_n \cdot \mathbb{A}\underline{\xi}_n) = -\beta U_n(\xi) + \frac{n\beta}{2}$. Defining next the operator

$$\mathcal{G}_\beta : L^2(\mathbb{R}, d\xi) \rightarrow L^2(\mathbb{R}, d\xi)$$

with kernel $\mathcal{G}_\beta(\xi, \eta) = \frac{1}{\sqrt{\lambda \exp \beta}} \mathcal{K}_\beta(\xi, \eta)$ one finds finally for real β

$$Z_n(\beta) = (1 - \lambda)^n \text{trace } \mathcal{G}_\beta^n.$$

The operator \mathcal{G}_β hence serves as a transfer operator for the Kac-Baker model at least for real β . We call it the Kac-Gutzwiller operator.

Let us next briefly recall the Ruelle operator for a 1-dimensional lattice spin system. For this we denote by $C(\Omega_+)$ the space of continuous observables for the spin system. On this space the following family of linear operators $\mathcal{L}_\beta : C(\Omega_+) \rightarrow C(\Omega_+)$ can be defined

$$\mathcal{L}_\beta f(\underline{\xi}) = \sum_{\underline{\eta} \in \tau^{-1}(\underline{\xi})} \exp(-\beta U_1(\underline{\eta})) f(\underline{\eta}),$$

where $U_1(\underline{\eta})$ denotes the interaction energy of the configuration $\underline{\eta}$ on the sublattice consisting of the lattice site 0 only. Inserting the explicit form of $U_1(\underline{\eta})$ we find

$$\mathcal{L}_\beta f(\underline{\xi}) = \sum_{\sigma=\pm 1, -1} \exp\left(\beta\sigma \sum_{i=1}^{\infty} \xi_{i-1} \lambda^i\right) f((\sigma, \underline{\xi})),$$

where we have again replaced $J\beta$ by β . Obviously the operators \mathcal{L}_β leave invariant the subspace of functions f depending on the configuration $\underline{\xi}$ only through the variable $z = \sum_{i=1}^{\infty} \xi_{i-1} \lambda^i$. Indeed, they leave invariant also the space of such functions depending holomorphically on this variable for instance in the disc D_r with $r > \frac{\lambda}{1-\lambda}$. It was shown in [Ma80] that the operators \mathcal{L}_β are nuclear for all complex values of β in the sense of Grothendieck in the Banach space $B(D_r)$ of functions holomorphic in D_r and continuous on $\overline{D_r}$ with the supremum norm. Furthermore the analytic version of the Atiyah-Bott fixed point formula (see [AtBo67]) shows that

$$Z_n(\beta) = (1 - \lambda^n) \text{trace } \mathcal{L}_\beta^n.$$

Hence also this Ruelle operator can be used as a transfer operator for the Kac-Baker model. The Ruelle zeta function $\zeta_R(z, \beta)$ for the Kac-Baker model defined in (2) can indeed be expressed through Fredholm determinants of both these two transfer operators as described in (3). But the Fredholm determinant $\det(1 - z\mathcal{L}_\beta)$ of the nuclear operator \mathcal{L}_β is an entire function both in the variables z and β hence the Ruelle zeta function $\zeta(z, \beta)$ is a meromorphic function in the entire z - and β -plane. From this it follows immediately (see [Mo89]) that the operators \mathcal{G}_β and \mathcal{L}_β have the same spectrum for real β . To relate in this case the two operators and also their eigenfunctions in a more explicit way we have to investigate the Kac operator in more detail. We should mention that some of the following arguments have been used already by M. Gutzwiller in [Gu82].

3 Hermite Functions and Mehler's Formula

Consider the operators $Z = m_x + \frac{1}{2\pi} \frac{\partial}{\partial x}$ and $Z^* = m_x - \frac{1}{2\pi} \frac{\partial}{\partial x}$ on $L^2(\mathbb{R})$, where m_g is the multiplication operator $(m_g f)(x) = g(x)f(x)$. Then the Hermite functions $h_k \in L^2(\mathbb{R})$ with $k \in \mathbb{N}_0$ are given by (see [Fo89], p.51)

$$\begin{aligned} h_0(x) &= 2^{\frac{1}{4}} e^{-\pi x^2} \\ h_k(x) &= \sqrt{\frac{\pi^k}{k!}} (Z^*)^k h_0(x) = \frac{2^{\frac{1}{4}}}{\sqrt{k!}} \left(\frac{-1}{2\sqrt{\pi}} \right)^k e^{-\pi x^2} \left(\frac{\partial}{\partial x} \right)^k e^{-2\pi x^2}. \end{aligned}$$

One has

$$Z^* h_k = \sqrt{\frac{k+1}{\pi}} h_{k+1} \quad \text{and} \quad Z h_k = \sqrt{\frac{k}{\pi}} h_{k-1}.$$

The Hermite functions form a Hilbert basis for $L^2(\mathbb{R})$ (see [Fo89], p.53) and the Mehler formula (see [Fo89], p.55) is the identity

$$\sum_{k=0}^{\infty} \lambda^k h_k(x) h_k(y) = \left(\frac{2}{1-\lambda^2} \right)^{\frac{1}{2}} \exp \left(\frac{-\pi(1+\lambda^2)(x^2+y^2) + 4\pi\lambda xy}{1-\lambda^2} \right) \quad (4)$$

for $\lambda \in D_1$. Here $u = \frac{2}{1-\lambda^2}$ lies in the right half plane, and the square root is the branch which is positive for $u > 0$.

Proposition 3.1 For $\gamma \in \{z \in \mathbb{C} \mid \operatorname{Re}(z) > 0\}$ and $\lambda = e^{-\gamma}$ we have

$$\frac{-\pi(1+\lambda^2)(x^2+y^2) + 4\pi\lambda xy}{1-\lambda^2} = \frac{2\pi}{\sinh \gamma} \left(-\frac{1}{2}(\cosh \gamma)(x^2+y^2) + xy \right).$$

Proof. Set $a := \frac{-\pi(1+\lambda^2)}{1-\lambda^2}$ and $b := \frac{4\pi\lambda}{1-\lambda^2}$. Then $\frac{b}{a} = \frac{-4e^{-\gamma}}{1+e^{-2\gamma}} = \frac{-4}{e^{\gamma}+e^{-\gamma}} = \frac{2}{\cosh \gamma}$ so that

$$\begin{aligned} \frac{-\pi(1+\lambda^2)(x^2+y^2) + 4\pi\lambda xy}{1-\lambda^2} &= \frac{4\pi\lambda}{1-\lambda^2} \left(-\frac{1}{2}(\cosh \gamma)(x^2+y^2) + xy \right) \\ &= \frac{4\pi e^{-\gamma}}{1-e^{-2\gamma}} \left(-\frac{1}{2}(\cosh \gamma)(x^2+y^2) + xy \right) \\ &= \frac{2\pi}{\sinh \gamma} \left(-\frac{1}{2}(\cosh \gamma)(x^2+y^2) + xy \right) \end{aligned}$$

□

Proposition 3.2

$$-\frac{1}{4} \left(\tanh \frac{\gamma}{2} \right) (\xi^2 + \eta^2) - \frac{(\xi - \eta)^2}{4 \sinh \gamma} = \frac{1}{2 \sinh \gamma} \left(-\frac{1}{2}(\cosh \gamma)(\xi^2 + \eta^2) + \xi\eta \right).$$

Proof. Using $(\xi - \eta)^2 = \xi^2 + \eta^2 - 2\xi\eta$ we calculate

$$\begin{aligned} -\frac{1}{4} \left(\tanh \frac{\gamma}{2} \right) (\xi^2 + \eta^2) - \frac{(\xi - \eta)^2}{4 \sinh \gamma} &= -\frac{1}{4} \left(\tanh \frac{\gamma}{2} \right) (\xi^2 + \eta^2) - \frac{\xi^2 + \eta^2}{4 \sinh \gamma} + \frac{1}{2} \frac{\xi\eta}{\sinh \gamma} \\ &= -\frac{1}{4} \left(\tanh \frac{\gamma}{2} + \frac{1}{\sinh \gamma} \right) (\xi^2 + \eta^2) + \frac{1}{2} \frac{\xi\eta}{\sinh \gamma} \\ &= -\frac{1}{4} \left(\frac{e^{\gamma} - 1}{e^{\gamma} + 1} + \frac{2}{e^{\gamma} - e^{-\gamma}} \right) (\xi^2 + \eta^2) + \frac{\xi\eta}{e^{\gamma} - e^{-\gamma}}. \end{aligned}$$

Set $a := -\frac{1}{4} \left(\frac{e^\gamma - 1}{e^\gamma + 1} + \frac{2}{e^\gamma - e^{-\gamma}} \right)$ and $b := \frac{1}{e^\gamma - e^{-\gamma}}$. Then

$$\begin{aligned}
\frac{b}{a} &= -4 \left(\frac{e^\gamma - 1}{e^\gamma + 1} + \frac{2}{e^\gamma - e^{-\gamma}} \right)^{-1} \frac{1}{e^\gamma - e^{-\gamma}} \\
&= -4 \left(\frac{e^{2\gamma} - e^\gamma - 1 + e^{-\gamma}}{e^\gamma + 1} + 2 \right)^{-1} \\
&= -4 \left(\frac{e^{2\gamma} - e^\gamma - 1 + e^{-\gamma} + 2e^\gamma + 2}{e^\gamma + 1} \right)^{-1} \\
&= -4 (e^\gamma + e^{-\gamma})^{-1} \\
&= -\frac{2}{\cosh \gamma}
\end{aligned}$$

and therefore

$$-\frac{1}{4} \left(\tanh \frac{\gamma}{2} \right) (\xi^2 + \eta^2) - \frac{(\xi - \eta)^2}{4 \sinh \gamma} = \frac{1}{2 \sinh \gamma} \left(-\frac{1}{2} (\cosh \gamma) (\xi^2 + \eta^2) + \xi \eta \right).$$

□

Proposition 3.3 Set $\lambda = e^{-\gamma}$, $x = \xi \frac{1}{2\sqrt{\pi}}$ and $y = \eta \frac{1}{2\sqrt{\pi}}$. Then

$$\sum_{k=0}^{\infty} \lambda^k h_k(x) h_k(y) = \left(\frac{1}{\sinh \gamma} \right)^{\frac{1}{2}} e^{\frac{\gamma}{2}} \exp \left(-\frac{1}{4} \left(\tanh \frac{\gamma}{2} \right) (\xi^2 + \eta^2) - \frac{(\xi - \eta)^2}{4 \sinh \gamma} \right).$$

Proof. Combining the Mehler formula (4) with Proposition 3.1 and Proposition 3.2 we calculate

$$\begin{aligned}
\sum_{k=0}^{\infty} \lambda^k h_k(x) h_k(y) &= \left(\frac{2}{1 - \lambda^2} \right)^{\frac{1}{2}} \exp \left(\frac{2\pi}{\sinh \gamma} \left(-\frac{1}{2} (\cosh \gamma) (x^2 + y^2) + xy \right) \right) \\
&= \left(\frac{2}{1 - e^{-2\gamma}} \right)^{\frac{1}{2}} \exp \left(\frac{1}{2 \sinh \gamma} \left(-\frac{1}{2} (\cosh \gamma) (\xi^2 + \eta^2) + \xi \eta \right) \right) \\
&= \left(\frac{e^\gamma}{e^\gamma - e^{-\gamma}} \right)^{\frac{1}{2}} \exp \left(-\frac{1}{4} \left(\tanh \frac{\gamma}{2} \right) (\xi^2 + \eta^2) - \frac{(\xi - \eta)^2}{4 \sinh \gamma} \right) \\
&= \left(\frac{1}{\sinh \gamma} \right)^{\frac{1}{2}} e^{\frac{\gamma}{2}} \exp \left(-\frac{1}{4} \left(\tanh \frac{\gamma}{2} \right) (\xi^2 + \eta^2) - \frac{(\xi - \eta)^2}{4 \sinh \gamma} \right)
\end{aligned}$$

□

We set

$$\tilde{\mathcal{K}}(\xi, \eta) := \left(\frac{1}{\sinh \gamma} \right)^{\frac{1}{2}} e^{\frac{\gamma}{2}} \exp \left(-\frac{1}{4} \left(\tanh \frac{\gamma}{2} \right) (\xi^2 + \eta^2) - \frac{(\xi - \eta)^2}{4 \sinh \gamma} \right)$$

and note that

$$\mathcal{K}_\beta(\xi, \eta) = \left(\frac{\cosh(\sqrt{\beta}\xi) \cosh(\sqrt{\beta}\eta)}{\pi e^\gamma} \right)^{\frac{1}{2}} \tilde{\mathcal{K}}(\xi, \eta). \quad (5)$$

Lemma 3.4 Let $c \neq 0$ and $a: \mathbb{R} \rightarrow]0, \infty[$ be a smooth function. Then

(i) $R_c: L^2(\mathbb{R}, a(\frac{\xi}{c})d\xi) \rightarrow L^2(\mathbb{R}, a(x)dx)$ defined by

$$(R_c f)(x) := \sqrt{|c|} f(cx)$$

is an isomorphism of Hilbert spaces.

(ii) If an operator K on $L^2(\mathbb{R}, d\xi)$ is given by an integral kernel $K(\xi, \eta)$ via

$$Kf(\xi) = \int_{\mathbb{R}} K(\xi, \eta) f(\eta) d\eta,$$

then the induced operator $K_c := R_c \circ K \circ R_c^{-1}$ on $L^2(\mathbb{R}, dx)$ is given by the kernel

$$K_c(x, y) = cK(cx, cy).$$

Proof.

(i) This follows from the calculation

$$\begin{aligned} \|R_c f\|_{L^2(\mathbb{R}, a(x)dx)}^2 &= \int_{\mathbb{R}} |(R_c f)(x)|^2 a(x) dx \\ &= \int_{\mathbb{R}} |c| |f(cx)|^2 a(x) dx \\ &= \int_{\mathbb{R}} |f(\xi)|^2 a(\frac{\xi}{c}) d\xi \\ &= \|f\|_{L^2(\mathbb{R}, a(\frac{\xi}{c})d\xi)}^2. \end{aligned}$$

(ii) This follows from the calculation

$$\begin{aligned} ((R_c \circ K \circ R_c^{-1})(f))(x) &= \sqrt{|c|} (K(R_c^{-1}f))(cx) \\ &= \sqrt{|c|} \int_{\mathbb{R}} K(cx, \eta) (R_c^{-1}f)(\eta) d\eta \\ &= \sqrt{|c|} \int_{\mathbb{R}} K(cx, \eta) \frac{1}{\sqrt{|c|}} f(\frac{1}{c}\eta) d\eta \\ &= \int_{\mathbb{R}} K(cx, cy) f(y) c dy. \end{aligned}$$

□

Lemma 3.4 yields the following commutative diagram:

$$\begin{array}{ccc} L^2(\mathbb{R}, d\xi) & \xrightarrow{K} & L^2(\mathbb{R}, d\xi) \\ R_c \downarrow & & \downarrow R_c \\ L^2(\mathbb{R}, dx) & \xrightarrow{K_c} & L^2(\mathbb{R}, dx) \end{array}$$

Example 3.5 We consider the Kac kernel

$$\mathcal{K}_\beta(\xi, \eta) = \left(\frac{\cosh(\sqrt{\beta}\xi) \cosh(\sqrt{\beta}\eta)}{\pi \sinh \gamma} \right)^{\frac{1}{2}} \exp \left(-\frac{1}{4} \left(\tanh \frac{\gamma}{2} \right) (\xi^2 + \eta^2) - \frac{(\xi - \eta)^2}{4 \sinh \gamma} \right)$$

and use the constant $c = 2\sqrt{\pi}$. Then, in view of Lemma 3.4 and formula (5), Proposition 3.2 yields

$$\begin{aligned}
\mathcal{K}_{\beta,c}(x,y) &= c\mathcal{K}_{\beta}(cx,cy) \\
&= c\mathcal{K}_{\beta}(\xi,\eta) \\
&= 2\sqrt{\pi} \left(\frac{\cosh(\sqrt{\beta}\xi)\cosh(\sqrt{\beta}\eta)}{\pi e^{\gamma}} \right)^{\frac{1}{2}} \tilde{K}_{\beta}(\xi,\eta) \\
&= 2 \left(\cosh(2\sqrt{\pi\beta}x)\cosh(2\sqrt{\pi\beta}y) \right)^{\frac{1}{2}} \sum_{k=0}^{\infty} e^{-(k+\frac{1}{2})\gamma} h_k(x)h_k(y)
\end{aligned}$$

□

Lemma 3.6 *Let $a: \mathbb{R} \rightarrow [1, \infty[$ be a smooth function and K be a bounded operator on $L^2(\mathbb{R}, dx)$ given by an integral kernel $K(x, y)$ via $Kf(x) = \int_{\mathbb{R}} K(x, y)f(y)dy$. Then*

(i) *The operator $K \circ m_{\sqrt{a}}$ on $L^2(\mathbb{R}, dx)$ is an unbounded integral operator with kernel*

$$K(x, y)\sqrt{a(y)}.$$

(ii) *The operator $m_{\frac{1}{\sqrt{a}}} \circ K$ on $L^2(\mathbb{R}, dx)$ is a bounded integral operator with kernel*

$$\frac{1}{\sqrt{a(x)}}K(x, y).$$

Proof.

(i) This is immediate from

$$(K \circ m_{\sqrt{a}}f)(x) = \int_{\mathbb{R}} K(x, y)\sqrt{a(y)}f(y)dy$$

$$\text{and } \|\sqrt{a}f\|_{L^2(\mathbb{R}, dx)} = \|f\|_{L^2(\mathbb{R}, a(x)dx)} < \infty.$$

(ii) Here we calculate

$$\begin{aligned}
(m_{\frac{1}{\sqrt{a}}} \circ Kf)(x) &= \frac{1}{\sqrt{a(x)}}(Kf)(x) \\
&= \frac{1}{\sqrt{a(x)}} \int_{\mathbb{R}} K(x, y)f(y)dy \\
&= \int_{\mathbb{R}} \frac{1}{\sqrt{a(x)}}K(x, y)f(y)dy
\end{aligned}$$

□

For a smooth map $a: \mathbb{R} \rightarrow [1, \infty[$ Lemma 3.6 yields the following commutative diagram

$$\begin{array}{ccc}
L^2(\mathbb{R}, a(x)dx) & \xrightarrow{K} & L^2(\mathbb{R}, dx) \\
m_{\sqrt{a}} \downarrow & & \downarrow \text{id} \\
L^2(\mathbb{R}, dx) & \xrightarrow{K'} & L^2(\mathbb{R}, dx) \\
\text{id} \downarrow & & \downarrow m_{\frac{1}{\sqrt{a}}} \\
L^2(\mathbb{R}, dx) & \xrightarrow{K''} & L^2(\mathbb{R}, dx)
\end{array}$$

with integral operators K' and K'' given by the kernels $K'(x, y) = K(x, y) \frac{1}{\sqrt{a(y)}}$ and

$$K''(x, y) = \frac{1}{\sqrt{a(x)}} K(x, y) \frac{1}{\sqrt{a(y)}}. \quad (6)$$

Example 3.7 We consider the rescaled Kac kernel

$$\mathcal{K}_{\beta, c}(x, y) = 2 \left(\cosh(2\sqrt{\pi\beta}x) \cosh(2\sqrt{\pi\beta}y) \right)^{\frac{1}{2}} \sum_{k=0}^{\infty} e^{-(k+\frac{1}{2})\gamma} h_k(x) h_k(y)$$

with $c = 2\sqrt{\pi}$ from Example 3.5 and use the function $a(x) = \cosh(2\sqrt{\pi\beta}x)$. Then (6) shows

$$\mathcal{K}_c''(x, y) = 2 \sum_{k=0}^{\infty} e^{-(k+\frac{1}{2})\gamma} h_k(x) h_k(y),$$

where we have omitted the index β since the kernel does not depend on it anymore. Thus the Hilbert basis $(h_k)_{k \in \mathbb{N}_0}$ diagonalizes $\mathcal{K}_c'' : L^2(\mathbb{R}, dx) \rightarrow L^2(\mathbb{R}, dx)$ and the corresponding eigenvalues are given by

$$\mathcal{K}_c'' h_k = 2e^{-(k+\frac{1}{2})\gamma} h_k.$$

Lemma 3.4 now yields the following commutative diagram

$$\begin{array}{ccc} L^2(\mathbb{R}, a(\frac{\xi}{c})d\xi) & \xrightarrow{\mathcal{K}_\beta} & L^2(\mathbb{R}, d\xi) \\ R_c \downarrow & & \downarrow R_c \\ L^2(\mathbb{R}, a(x)dx) & \xrightarrow{\mathcal{K}_{\beta, c}} & L^2(\mathbb{R}, dx) \\ m\sqrt{\pi} \downarrow & & \downarrow \text{id} \\ L^2(\mathbb{R}, dx) & \xrightarrow{\mathcal{K}'_{\beta, c}} & L^2(\mathbb{R}, dx) \\ \text{id} \downarrow & & \downarrow m\frac{1}{\sqrt{a}} \\ L^2(\mathbb{R}, dx) & \xrightarrow{\mathcal{K}_c''} & L^2(\mathbb{R}, dx) \end{array}$$

Note that $e^{\pi x^2} h_k(x)$ is a polynomial of degree k in x (see [Fo89], p.52). In view of the estimate

$$\sqrt{\cosh R\xi} \leq e^{\frac{1}{2}|R\xi|}$$

this shows that the functions $\xi \mapsto \sqrt{\cosh \sqrt{\beta}\xi} h_k(\xi)$ are in $L^2(\mathbb{R}, d\xi)$. Therefore all the h_k are contained in $L^2(\mathbb{R}, \cosh(\sqrt{\beta}\xi)d\xi)$. \square

Proposition 3.8 For $c \in \mathbb{R} \setminus \{0\}$ and $s \in \mathbb{R}$ we have

$$m_{\cosh(sx)} \circ R_c = R_c \circ m_{\cosh(\frac{s}{c}\xi)}.$$

Proof.

$$\begin{aligned} (m_{\cosh(sx)} \circ R_c f)(x) &= \cosh(sx) \sqrt{|c|} f(cx) \\ &= \sqrt{|c|} \cosh\left(\frac{s}{c}cx\right) f(cx) \\ &= (R_c \circ m_{\cosh(\frac{s}{c}\xi)} f)(x) \end{aligned}$$

\square

Proposition 3.9 Set $g(\xi) = \cosh(R'\xi)^t$ for $t \in \mathbb{R}^+$ and $R' \in \mathbb{R}$. Then the Kac operator $\mathcal{K}_\beta: L^2(\mathbb{R}, d\xi) \rightarrow L^2(\mathbb{R}, d\xi)$ satisfies $\mathcal{K}_\beta(L^2(\mathbb{R}, d\xi)) \subseteq L^2(\mathbb{R}, g(\xi)d\xi)$.

Proof. The estimate

$$|g(x)| \leq e^{t|R'\xi|}$$

shows that there exist positive constants c_1 and c_2 such that

$$|g(\xi)\mathcal{K}_\beta(\xi, \eta)| \leq c_1 e^{-c_2(\xi^2 + \eta^2)}.$$

Therefore there exists a positive constant C such that

$$\int_{\mathbb{R}} |\mathcal{K}_\beta(\xi, \eta)| g(\xi) d\xi < C \quad \forall \eta \in \mathbb{R}$$

and

$$\int_{\mathbb{R}} |\mathcal{K}_\beta(\xi, \eta)| d\eta < C \quad \forall \xi \in \mathbb{R}.$$

Now [Fo84], Theorem 6.18, implies that $\mathcal{K}_\beta(\xi, \eta)$ defines a bounded integral operator

$$\mathcal{K}_\beta: L^2(\mathbb{R}, d\eta) \rightarrow L^2(\mathbb{R}, g(\xi)d\xi)$$

with operator norm bounded by C . In particular $\mathcal{K}_\beta f \in L^2(\mathbb{R}, g(\xi)d\xi)$. □

4 Fock Space and Segal-Bargmann Transformation

For $t > 0$ the Fock space $\mathcal{HL}^2(\mathbb{C}, \mu_t)$ consists of all entire functions $F: \mathbb{C} \rightarrow \mathbb{C}$ such that

$$\|F\|_t^2 := \int_{\mathbb{C}} |F(z)|^2 \mu_t(z) dz < \infty$$

where μ_t denotes the weight $\mu_t = t \exp -\pi t |z|^2$. The space $\mathcal{HL}^2(\mathbb{C}, \mu_t)$ with this norm is a Hilbert space (see [Fo89], p.46). Moreover the polynomials

$$\zeta_k^t(z) := \sqrt{\frac{\pi^k}{t^k k!}} z^k \quad \forall k \in \mathbb{N}_0$$

form an orthonormal basis for the space $\mathcal{HL}^2(\mathbb{C}, \mu_t)$. Of special interest for the following discussion is the case $t = 1$. In this case the space $\mathcal{HL}^2(\mathbb{C}, \mu_1)$ will be denoted simply by \mathcal{F} and we call it the Fock space (see [Fo89], p.40):

$$\mathcal{F} = \left\{ F: \mathbb{C} \rightarrow \mathbb{C} \mid F \text{ entire with } \int_{\mathbb{C}} |F(z)|^2 \exp(-\pi z^2) dz < \infty \right\}$$

Its basis ζ_k^1 will be denoted simply by ζ_k with $k = 0, 1, \dots$.

There is an isomorphism $B_t: L^2(\mathbb{R}, d\xi) \rightarrow \mathcal{HL}^2(\mathbb{C}, \mu_t)$, the so called Segal-Bargmann transform, defined by

$$B_t f(z) := \left(\frac{2}{t}\right)^{\frac{1}{4}} \int_{\mathbb{R}} f(\xi) \exp(2\pi\xi z - \frac{\pi}{t}\xi^2 - \frac{\pi t}{2}z^2) d\xi.$$

It is a Hilbert space isomorphism such that $B_t h_k^t = \zeta_k^t$ (see [Fo89], p.40 and p.51), where the Hermite functions h_k^t are generalizations of the Hermite functions $h_k^1 = h_k$.

Recall the operator \mathcal{K}_c'' on $L^2(\mathbb{R}, dx)$ from Example 3.7. Obviously the Segal-Bargmann transform $B := B_1$ induces a linear operator in the space \mathcal{F} . Indeed one finds

Proposition 4.1 For $\lambda = e^{-\gamma}$ the operator $M_\lambda := B \circ \mathcal{K}_c'' \circ B^{-1}: \mathcal{F} \rightarrow \mathcal{F}$ is given by

$$(M_\lambda F)(z) = 2\sqrt{\lambda} F(\lambda z).$$

Proof. Using Example 3.7 and the homogeneity of ζ_k we calculate

$$\begin{aligned} (M_\lambda \zeta_k)(z) &= B(\mathcal{K}_c'' h_k)(z) \\ &= B(2e^{-(k+\frac{1}{2})\gamma} h_k)(z) \\ &= 2e^{-(k+\frac{1}{2})\gamma} \zeta_k(z) \\ &= 2e^{-\frac{1}{2}\gamma} \zeta_k(e^{-\gamma} z). \end{aligned}$$

Since $(\zeta_k)_{k \in \mathbb{N}_0}$ is a Hilbert basis for \mathcal{F} this implies the claim. \square

The functions $\zeta_k(z), k = 0, 1, \dots$ are the eigenfunctions of the operator M_λ with eigenvalue $\rho_k = 2\lambda^{k+\frac{1}{2}}$. Since this operator is nuclear its spectrum consists of these numbers. The above proposition shows that the following diagram commutes:

$$\begin{array}{ccc} L^2(\mathbb{R}, dx) & \xrightarrow{\mathcal{K}_c''} & L^2(\mathbb{R}, dx) \\ \downarrow B & & \downarrow B \\ \mathcal{F} & \xrightarrow{M_\lambda} & \mathcal{F} \end{array}$$

Proposition 4.2 Let $\tau_r: L^2(\mathbb{R}, dx) \rightarrow L^2(\mathbb{R}, dx)$ for $r \in \mathbb{R}$ be the translation defined by

$$\tau_r f(x) = f(x - r),$$

and $\mu_r: L^2(\mathbb{R}, dx) \rightarrow L^2(\mathbb{R}, dx)$ for $r \in \mathbb{R}$ be the multiplication defined by

$$\mu_r f(x) = r f(x).$$

Then we have

$$(Z^*)^k \circ \tau_r = \tau_r \circ (Z^* + \mu_r)^k \quad \forall k \in \mathbb{N}_0.$$

In particular we find

$$(Z^*)^k \circ \tau_r h_0 = \sum_{l=0}^k \binom{k}{l} r^l \tau_r \tilde{h}_{k-l}.$$

Proof. For the first claim it suffices to prove the case $k = 1$. So we calculate

$$\begin{aligned} ((Z^* \circ \tau_r) f)(x) &= x \tau_r f(x) - \frac{1}{2\pi} (\tau_r)'(x) \\ &= x f(x - r) - \frac{1}{2\pi} f'(x - r) \\ &= (x - r) f(x - r) - \frac{1}{2\pi} f'(x - r) + r f(x - r) \\ &= (\tau_r \circ Z^* f)(x) + (\mu_r \circ \tau_r f)(x) \\ &= (\tau_r \circ Z^* f)(x) + (\tau_r \circ \mu_r f)(x) \\ &= (\tau_r \circ (Z^* + \mu_r) f)(x). \end{aligned}$$

Now we calculate

$$\begin{aligned}
(Z^*)^k \circ \tau_r h_0 &= \tau_r \circ (Z^* + \mu_r)^k h_0 \\
&= \tau_r \circ \sum_{l=0}^k \binom{k}{l} \mu_r^l (Z^*)^{k-l} h_0 \\
&= \tau_r \circ \sum_{l=0}^k \binom{k}{l} r^l \tilde{h}_{k-l} \\
&= \sum_{l=0}^k \binom{k}{l} r^l \tau_r \tilde{h}_{k-l}
\end{aligned}$$

□

Proposition 4.3 For each $k \in \mathbb{N}_0$ we have

$$(Z^*)^k \circ m_{e^{sx}} = m_{e^{sx}} \circ (Z^* - \frac{s}{2\pi})^k$$

and

$$m_{e^{sx}} \circ (Z^*)^k = (Z^* + \frac{s}{2\pi})^k \circ m_{e^{sx}}.$$

Proof. It suffices to prove the first equality in the case $k = 1$. So we calculate

$$\begin{aligned}
Z^*(e^{sx} f)(x) &= xe^{sx} f(x) - \frac{1}{2\pi} (e^{sx} f(x))' \\
&= xe^{sx} f(x) - \frac{1}{2\pi} (se^{sx} f(x) + e^{sx} f'(x)) \\
&= e^{sx} (Z^* f)(x) - \frac{s}{2\pi} e^{sx} f(x) \\
&= m_{e^{sx}} \left(Z^* - \frac{s}{2\pi} \right) f(x)
\end{aligned}$$

□

Set

$$\tilde{h}_k := (Z^*)^k h_0.$$

Proposition 4.4 For $s \in \mathbb{R}$ we have

$$m_{e^{sx}} \tilde{h}_k = e^{\frac{s^2}{4\pi}} \sum_{j=0}^k \binom{k}{j} \left(\frac{s}{\pi} \right)^{k-j} \tau_{\frac{s}{2\pi}} \tilde{h}_j.$$

Proof.

$$\begin{aligned}
m_{e^{sx}} h_0(x) &= e^{sx} 2^{\frac{1}{4}} e^{-\pi x^2} \\
&= 2^{\frac{1}{4}} e^{sx - \pi x^2} \\
&= 2^{\frac{1}{4}} e^{-\pi (\frac{s}{2\pi} - x)^2 + (\frac{s}{2\pi})^2 \pi} \\
&= e^{(\frac{s}{2\pi})^2 \pi} 2^{\frac{1}{4}} e^{-\pi (\frac{s}{2\pi} - x)^2} \\
&= e^{(\frac{s}{2\pi})^2 \pi} h_0(x - \frac{s}{2\pi}) \\
&= e^{\frac{s^2}{4\pi}} \tau_{\frac{s}{2\pi}} h_0(x).
\end{aligned}$$

Now we use Proposition 4.3 to calculate

$$\begin{aligned}
m_{e^{sx}} \tilde{h}_k &= m_{e^{sx}} (Z^*)^k h_0 \\
&= \left(Z^* + \frac{S}{2\pi}\right)^k m_{e^{sx}} h_0 \\
&= e^{\frac{s^2}{4\pi}} \left(Z^* + \frac{S}{2\pi}\right)^k \tau_{\frac{s}{2\pi}} h_0 \\
&= e^{\frac{s^2}{4\pi}} \sum_{l=0}^k \binom{k}{l} \left(\frac{S}{2\pi}\right)^{k-l} (Z^*)^l \tau_{\frac{s}{2\pi}} h_0 \\
&= e^{\frac{s^2}{4\pi}} \sum_{l=0}^k \binom{k}{l} \left(\frac{S}{2\pi}\right)^{k-l} \tau_{\frac{s}{2\pi}} (Z^* + \mu_{\frac{s}{2\pi}})^l h_0 \\
&= e^{\frac{s^2}{4\pi}} \sum_{l=0}^k \binom{k}{l} \left(\frac{S}{2\pi}\right)^{k-l} \sum_{j=0}^l \binom{l}{j} \left(\frac{S}{2\pi}\right)^{l-j} \tau_{\frac{s}{2\pi}} (Z^*)^j h_0 \\
&= e^{\frac{s^2}{4\pi}} \sum_{l=0}^k \sum_{j=0}^l \binom{k}{l} \binom{l}{j} \left(\frac{S}{2\pi}\right)^{k-j} \tau_{\frac{s}{2\pi}} \tilde{h}_j \\
&= e^{\frac{s^2}{4\pi}} \sum_{j=0}^k \sum_{l=j}^k \binom{k}{l} \binom{l}{j} \left(\frac{S}{2\pi}\right)^{k-j} \tau_{\frac{s}{2\pi}} \tilde{h}_j
\end{aligned}$$

On the other hand

$$\sum_{l=j}^k \binom{k}{l} \binom{l}{j} = \sum_{n=0}^{k-j} \binom{k}{n+j} \binom{n+j}{j} = \sum_{n=0}^{k-j} \binom{k-j}{n} \binom{k}{j} = 2^{k-j} \binom{k}{j}$$

which inserted in the calculation above gives the claim. \square

Proposition 4.5 For $r \in \mathbb{R}$ we have

$$B \circ \tau_r = m_{e^{\pi r z - \frac{\pi}{2} r^2}} \circ \tau_r \circ B.$$

Here $\tau_r: \mathcal{F} \rightarrow \mathcal{F}$ is also defined by $\tau_r F(z) = F(z - r)$. In other words, for $F \in \mathcal{F}$ we have

$$(B \circ \tau_r \circ B^{-1})F(z) = e^{-\frac{\pi}{2} r^2} e^{\pi r z} F(z - r).$$

Proof.

$$\begin{aligned}
((B \circ \tau_r)f)(z) &= 2^{\frac{1}{4}} \int_{\mathbb{R}} f(\xi - r) \exp(2\pi\xi z - \pi\xi^2 - \frac{\pi}{2} z^2) d\xi \\
&= 2^{\frac{1}{4}} \int_{\mathbb{R}} f(\xi) \exp(2\pi(\xi + r) - \pi(\xi + r)^2 - \frac{\pi}{2} z^2) d\xi \\
&= 2^{\frac{1}{4}} \int_{\mathbb{R}} f(\xi) \exp(2\pi\xi(z - r) - \pi\xi^2 - \frac{\pi}{2}(z - r)^2 + \pi r z - \frac{\pi}{2} r^2) d\xi \\
&= e^{\pi r z - \frac{\pi}{2} r^2} 2^{\frac{1}{4}} \int_{\mathbb{R}} f(\xi) \exp(2\pi\xi(z - r) - \pi\xi^2 - \frac{\pi}{2}(z - r)^2) d\xi \\
&= e^{\pi r z - \frac{\pi}{2} r^2} Bf(z - r) \\
&= e^{\pi r z - \frac{\pi}{2} r^2} \tau_r Bf(z)
\end{aligned}$$

□

Proposition 4.6 For $s \in \mathbb{R}$ we have

$$B \circ m_{e^{sx}} \circ B^{-1} = e^{\frac{1}{8\pi}s^2} m_{e^{\frac{sz}{2}}} \circ \tau_{-s}.$$

In other words, for $F \in \mathcal{F}$ and $z \in \mathbb{C}$ we have

$$(B \circ m_{e^{sx}} \circ B^{-1}F)(z) = e^{\frac{s^2}{8\pi}} e^{\frac{sz}{2}} F\left(z + \frac{s}{2\pi}\right).$$

Proof. Using Proposition 4.4, Proposition 4.5 and the fact that $(B\tilde{h}_j)(z) = z^j$ we calculate

$$\begin{aligned} B \circ m_{\exp sx} \tilde{h}_k &= \exp\left(\frac{s^2}{4\pi}\right) \sum_{j=0}^k \binom{k}{j} \left(\frac{s}{\pi}\right)^{k-j} B \circ \tau_{\frac{s}{2\pi}} \tilde{h}_j \\ &= \exp\left(\frac{s^2}{4\pi}\right) \sum_{j=0}^k \binom{k}{j} \left(\frac{s}{\pi}\right)^{k-j} \exp\left(\frac{sz}{2} - \frac{s^2}{8\pi}\right) \tau_{\frac{s}{2\pi}} B \tilde{h}_j \\ &= \exp\left(\frac{s^2}{8\pi}\right) \exp\left(\frac{sz}{2}\right) \sum_{j=0}^k \binom{k}{j} \left(\frac{s}{\pi}\right)^{k-j} \tau_{\frac{s}{2\pi}} z^j \\ &= \exp\left(\frac{s^2}{8\pi}\right) \exp\left(\frac{sz}{2}\right) \sum_{j=0}^k \binom{k}{j} \left(\frac{s}{\pi}\right)^{k-j} \left(z - \frac{s}{2\pi}\right)^j \\ &= \exp\left(\frac{s^2}{8\pi}\right) \exp\left(\frac{sz}{2}\right) \left(z + \frac{s}{2\pi}\right)^k \end{aligned}$$

and since the \tilde{h}_k form a basis of \mathcal{F} this implies the claim. □

Consider next the unbounded operator $C_s: \mathcal{F} \rightarrow \mathcal{F}$ defined as

$$C_s := B \circ m_{\cosh(sx)} \circ B^{-1}.$$

Then one finds

$$C_s F(z) = \frac{1}{2} e^{\frac{s^2}{8\pi}} \left(e^{\frac{sz}{2}} F\left(z + \frac{s}{2\pi}\right) + e^{-\frac{sz}{2}} F\left(z - \frac{s}{2\pi}\right) \right) \quad (7)$$

and we have the following commutative diagram

$$\begin{array}{ccc} L^2(\mathbb{R}, dx) & \xrightarrow{m_{\cosh(sx)}} & L^2(\mathbb{R}, dx) \\ \downarrow B & & \downarrow B \\ \mathcal{F} & \xrightarrow{C_s} & \mathcal{F} \end{array}$$

Proposition 4.7 For $s \in \mathbb{R}$ and $\lambda = e^{-\gamma}$ we have

$$(C_s \circ M_\lambda F)(z) = \sqrt{\lambda} e^{\frac{1}{8\pi^2}s^2} \left(e^{\frac{sz}{2}} F\left(\lambda z + \lambda \frac{s}{2\pi}\right) + e^{-\frac{sz}{2}} F\left(\lambda z - \lambda \frac{s}{2\pi}\right) \right).$$

Proof.

$$\begin{aligned} (C_s \circ M_\lambda F)(z) &= \frac{1}{2} e^{\frac{s^2}{8\pi}} \left(e^{\frac{sz}{2}} (M_\lambda F)\left(z + \frac{s}{2\pi}\right) + e^{-\frac{sz}{2}} (M_\lambda F)\left(z - \frac{s}{2\pi}\right) \right) \\ &= \sqrt{\lambda} e^{\frac{s^2}{8\pi}} \left(e^{\frac{sz}{2}} F\left(\lambda z + \lambda \frac{s}{2\pi}\right) + e^{-\frac{sz}{2}} F\left(\lambda z - \lambda \frac{s}{2\pi}\right) \right) \end{aligned}$$

□

For $\alpha \neq 0$ consider the map

$$\nu_\alpha : \mathcal{HL}^2(\mathbb{C}, \mu_{\frac{1}{|\alpha|^2}}) \rightarrow \mathcal{F},$$

defined as

$$(\nu_\alpha F)(z) := F(\alpha z).$$

Then one finds for the operator

$$C_s \circ M_\lambda \circ \nu_\alpha : \mathcal{HL}^2(\mathbb{C}, \mu_{\frac{1}{|\alpha|^2}}) \rightarrow \mathcal{F}$$

$$\begin{aligned} (C_s \circ M_\lambda \circ \nu_\alpha F)(z) &= \sqrt{\lambda} e^{\frac{s^2}{8\pi}} \left(e^{\frac{sz}{2}} (M_\lambda \circ \nu_\alpha \circ F)(z + \frac{s}{2\pi}) + e^{-\frac{sz}{2}} (M_\lambda \circ \nu_\alpha \circ F)(z - \frac{s}{2\pi}) \right) \\ &= \sqrt{\lambda} e^{\frac{s^2}{8\pi}} \left(e^{\frac{sz}{2}} F(\alpha\lambda z + \alpha\lambda \frac{s}{2\pi}) + e^{-\frac{sz}{2}} F(\alpha\lambda z - \alpha\lambda \frac{s}{2\pi}) \right) \end{aligned}$$

respectively for the operator

$$\nu_{\alpha^{-1}} \circ C_s \circ M_\lambda \circ \nu_\alpha : \mathcal{HL}^2(\mathbb{C}, \mu_{\frac{1}{|\alpha|^2}}) \rightarrow \mathcal{HL}^2(\mathbb{C}, \mu_{\frac{1}{|\alpha|^2}})$$

$$(\nu_{\alpha^{-1}} \circ C_s \circ M_\lambda \circ \nu_\alpha F)(z) = \sqrt{\lambda} e^{\frac{s^2}{8\pi}} \left(e^{\frac{sz}{2\alpha}} F(\lambda z + \alpha\lambda \frac{s}{2\pi}) + e^{-\frac{sz}{2\alpha}} F(\lambda z - \alpha\lambda \frac{s}{2\pi}) \right)$$

Next chose the parameters s and α as follows:

$$s = 2\sqrt{\pi\beta} \quad \text{and} \quad \alpha = \sqrt{\frac{\pi}{\beta}}.$$

Obviously, $\alpha s = 2\pi$ and therefore one gets for these parameters for the operator

$$\nu_{\sqrt{\frac{\beta}{\pi}}} \circ C_{2\sqrt{\beta\pi}} \circ M_\lambda \circ \nu_{\sqrt{\frac{\pi}{\beta}}} : \mathcal{HL}^2(\mathbb{C}, \mu_{\frac{|\beta|}{\pi}}) \rightarrow \mathcal{HL}^2(\mathbb{C}, \mu_{\frac{|\beta|}{\pi}})$$

defined by

$$(\nu_{\sqrt{\frac{\beta}{\pi}}} \circ C_{2\sqrt{\beta\pi}} \circ M_\lambda \circ \nu_{\sqrt{\frac{\pi}{\beta}}} F)(z) = \sqrt{\lambda} e^{\frac{\beta}{2}} (e^{\beta z} F(\lambda z + \lambda) + e^{-\beta z} F(\lambda z - \lambda)).$$

Proposition 4.8 For real β and $c = 2\sqrt{\pi}$ the operators

$$\mathcal{L}_\beta : \mathcal{HL}^2(\mathbb{C}, \mu_{\frac{|\beta|}{\pi}}) \rightarrow \mathcal{HL}^2(\mathbb{C}, \mu_{\frac{|\beta|}{\pi}})$$

and

$$\frac{1}{\sqrt{\lambda \exp \beta}} m_{\cosh(2\sqrt{\beta\pi}x)} \circ \mathcal{K}_c'' : L^2(\mathbb{R}, dx) \rightarrow L^2(\mathbb{R}, dx)$$

are conjugate operators.

Proof. Indeed one finds after inserting the definitions of the two operators $C_{2\sqrt{\beta\pi}}$ and M_λ

$$C_{2\sqrt{\beta\pi}} \circ M_\lambda = B \circ m_{\cosh 2\sqrt{\beta\pi}x} \circ B^{-1} \circ B \circ \mathcal{K}_c'' \circ B^{-1} = B \circ m_{\cosh 2\sqrt{\beta\pi}x} \circ \mathcal{K}_c'' \circ B^{-1}$$

which shows that $C_{2\sqrt{\beta\pi}} \circ M_\lambda$ is conjugate to the operator $m_{\cosh(2\sqrt{\beta\pi}x)} \circ \mathcal{K}_c''$. Hence $f \in L^2(\mathbb{R}, dx)$ is an eigenfunction of the operator $m_{\cosh(2\sqrt{\beta\pi}x)} \circ \mathcal{K}_c''$ with eigenvalue ρ iff the function Bf is an eigenfunction of the operator $C_{2\sqrt{\beta\pi}} \circ M_\lambda : \mathcal{F} \rightarrow \mathcal{F}$ with the same eigenvalue. In the same way one concludes: f is an eigenfunction of the operator $\frac{1}{\sqrt{\lambda \exp \beta}} (m_{\cosh(2\sqrt{\beta\pi}x)} \circ \mathcal{K}_c'')$ with eigenvalue ρ iff the function $\nu \sqrt{\frac{\beta}{\pi}} \circ Bf$ is an eigenfunction of the operator $\mathcal{L}_\beta : \mathcal{HL}^2(\mathbb{C}, \mu_{\frac{|\beta|}{\pi}}) \rightarrow \mathcal{HL}^2(\mathbb{C}, \mu_{\frac{|\beta|}{\pi}})$ with eigenvalue ρ . \square

Hence given an eigenfunction $f = f(x)$ of the operator $\frac{1}{\sqrt{\lambda \exp \beta}} (m_{\cosh(2\sqrt{\beta\pi}x)} \circ \mathcal{K}_c'')$ the corresponding eigenfunction $F = F(z)$ of the operator $\mathcal{L}_\beta : \mathcal{HL}^2(\mathbb{C}, \mu_{\frac{|\beta|}{\pi}}) \rightarrow \mathcal{HL}^2(\mathbb{C}, \mu_{\frac{|\beta|}{\pi}})$ has the following explicit form:

$$F(z) = 2^{\frac{1}{4}} \int_{\mathbb{R}} f(\xi) \exp \left(2\sqrt{\pi\beta}\xi z - \pi\xi^2 - \frac{\beta}{2}z^2 \right) d\xi.$$

To relate finally the eigenfunctions f of the operator $m_{\cosh(2\sqrt{\beta\pi}x)} \circ \mathcal{K}_c''$ to those of the Kac-Gutzwiller integral operator \mathcal{K}_β with kernel $\mathcal{K}_\beta(\xi, \eta)$ we need

Proposition 4.9 *Let $H : L^2(\mathbb{R}, d\xi) \rightarrow L^2(\mathbb{R}, d\xi)$ be an integral operator with kernel $H(\xi, \eta)$ and $g : \mathbb{R} \rightarrow]0, \infty[$ a smooth function. Then the following statements are equivalent:*

- (1) *f is an eigenfunction of the integral operator $m_{g^2} \circ H$ with kernel $g^2(\xi)H(\xi, \eta)$ for the eigenvalue ρ .*
- (2) *$\frac{f}{g}$ is an eigenfunction of the integral operator H' with kernel $g(\xi)H(\xi, \eta)g(\eta)$ for the eigenvalue ρ .*

Proof. This follows from the calculations

$$\begin{aligned} \rho f(\xi) &= \int_{\mathbb{R}} g^2(\xi)H(\xi, \eta)f(\eta)d\eta \\ &= g(\xi) \int_{\mathbb{R}} g(\xi)H'(\xi, \eta)g(\eta) \frac{f(\eta)}{g(\eta)}d\eta \\ &= g(\xi)H\left(\frac{f}{g}\right)(\xi) \end{aligned}$$

and

$$\begin{aligned} \rho \frac{f(\xi)}{g(\xi)} &= \int_{\mathbb{R}} g(\xi)H(\xi, \eta)g(\eta) \frac{f(\eta)}{g(\eta)}d\eta \\ &= \frac{1}{g(\xi)} \int_{\mathbb{R}} g^2(\xi)H(\xi, \eta)f(\eta)d\eta \\ &= \frac{1}{g(\xi)} m_{g^2} \circ H(f)(\xi). \end{aligned}$$

\square

Theorem 4.10 *For real β the Ruelle operator \mathcal{L}_β and the Kac-Gutzwiller operator \mathcal{G}_β with kernel $(\lambda \exp \beta)^{-\frac{1}{2}} \mathcal{K}_\beta(\xi, \eta)$ have the same eigenvalues (counted with multiplicities) and hence the same spectrum. Their eigenfunctions $F = F(z)$ and $f = f(\xi)$ to the eigenvalue ρ are related as follows:*

(1)

$$F(z) = (8\pi)^{\frac{1}{4}} \int_{\mathbb{R}} \sqrt{\cosh 2\sqrt{\pi\beta\xi}} f(2\sqrt{\pi}\xi) \exp\left(2\sqrt{\pi\beta}\xi z - \pi\xi^2 - \frac{\beta}{2}z^2\right) d\xi.$$

(2)

$$f(\xi) = \left(\frac{1}{2\pi}\right)^{\frac{1}{4}} \frac{1}{\sqrt{\cosh(2\sqrt{\pi\beta}\xi)}} \int_{\mathbb{C}} F\left(\sqrt{\frac{\pi}{\beta}}z\right) \exp\left(\sqrt{\pi}\xi z^* - \frac{1}{4}\xi^2 - \frac{\pi}{2}z^{*2} - \pi z^2\right) dz.$$

Proof. According to Proposition 4.8 the operators $\frac{1}{\sqrt{\lambda \exp \beta}} (m_{\cosh(2\sqrt{\beta\pi}x)} \circ \mathcal{K}_c'')$ with $c = 2\sqrt{\pi}$ and \mathcal{L}_β are conjugate via $\nu_{\sqrt{\frac{\pi}{\beta}}} \circ B$. Proposition 4.9 shows that the operator $m_{\cosh(2\sqrt{\beta\pi}x)} \circ \mathcal{K}_c''$ has the same spectrum as the operator

$$m_{\sqrt{\cosh 2\sqrt{\pi\beta}x}} \circ \mathcal{K}_c'' \circ m_{\sqrt{\cosh 2\sqrt{\pi\beta}x}}$$

with kernel $\sqrt{\cosh 2\sqrt{\pi\beta}x \cosh 2\sqrt{\pi\beta}y} \mathcal{K}_c''(x, y)$. But this operator is conjugate to the operator \mathcal{K}_β with kernel $\mathcal{K}_\beta(\xi, \eta)$ through the map R_c with $c = 2\sqrt{\pi}$. Hence, if $f \in L^2(\mathbb{R}, d\xi)$ is an eigenfunction of the Kac-Gutzwiller operator $\frac{1}{\sqrt{\lambda \exp \beta}} \mathcal{K}_\beta$ with eigenvalue ρ , then

$$(R_{2\sqrt{\pi}}f)(x) = \sqrt{2\sqrt{\pi}} f(2\sqrt{\pi}x)$$

is an eigenfunction of the operator $\frac{1}{\sqrt{\lambda \exp \beta}} \left(m_{\sqrt{\cosh 2\sqrt{\pi\beta}x}} \circ \mathcal{K}_c'' \circ m_{\sqrt{\cosh 2\sqrt{\pi\beta}x}} \right)$ with the same eigenvalue. Hence $\sqrt{\cosh 2\sqrt{\pi\beta}x} (R_{2\sqrt{\pi}}f)(x)$ is an eigenfunction of the operator

$$\frac{1}{\sqrt{\lambda \exp \beta}} \left(m_{\cosh(2\sqrt{\beta\pi}x)} \circ \mathcal{K}_c'' \right)$$

with eigenvalue ρ by Proposition 4.9. But then

$$F(z) = (8\pi)^{\frac{1}{4}} \int_{\mathbb{R}} \sqrt{\cosh 2\sqrt{\pi\beta}\xi} f(2\sqrt{\pi}\xi) \exp\left(2\sqrt{\pi\beta}\xi z - \pi\xi^2 - \frac{\beta}{2}z^2\right) d\xi$$

is an eigenfunction of the operator \mathcal{L}_β with eigenvalue ρ . On the other hand starting with an eigenfunction of the operator \mathcal{L}_β we know that $h(x) = (B^{-1} \circ \nu_{\alpha^{-1}} F)(x)$ is an eigenfunction of the operator

$$\frac{1}{\sqrt{\lambda \exp \beta}} \left(m_{\sqrt{\cosh 2\sqrt{\beta\pi}x}} \circ \mathcal{K}_c'' \circ m_{\sqrt{\cosh 2\sqrt{\beta\pi}x}} \right)$$

which is again conjugate to \mathcal{K}_β via the map R_c^{-1} and hence $R_c^{-1}(\sqrt{\cosh 2\sqrt{\beta\pi}x}^{-1} h)(\xi)$ is an eigenfunction of the operator \mathcal{G}_β . Inserting all the transformations involved we finally get for the corresponding eigenfunction $f = f(\xi)$

$$f(\xi) = \left(\frac{1}{2\pi}\right)^{\frac{1}{4}} \frac{1}{\sqrt{\cosh(2\sqrt{\pi\beta}\xi)}} \int_{\mathbb{C}} F\left(\sqrt{\frac{\pi}{\beta}}z\right) \exp\left(\sqrt{\pi}\xi z^* - \frac{1}{4}\xi^2 - \frac{\pi}{2}z^{*2} - \pi |z|^2\right) dz.$$

□

5 Zeros of the Ruelle zeta function

From our discussion it follows that for real β the eigenvalues of the Ruelle operator \mathcal{L}_β are real. According to (3) the zeros of the Ruelle zeta function $\zeta_R(\beta) := \zeta_R(1, \beta)$ are located at those values of β , where the operator \mathcal{L}_β has eigenvalue $\rho = \lambda^{-1}$ and not at the same time the eigenvalue $\rho = 1$. In the following we will show for $0 < \lambda < \frac{1}{2}$ that there exist infinitely many real β -values such that \mathcal{L}_β has the eigenvalue $\rho = \lambda^{-1}$ and therefore $\zeta_R(\beta)$ presumably has infinitely many “nontrivial” zeros on the real line unless unexpected cancellations take place. For the special case $\lambda = \frac{1}{2}$ we can show furthermore that there exists infinitely many values of β on the line $\operatorname{Re}(\beta) = \ln 2$ such that the operator \mathcal{L}_β has eigenvalue $\rho = \lambda^{-1}$ and hence presumably $\zeta_R(\beta)$ has infinitely many trivial zeros on this line $\operatorname{Re}(\beta) = \ln 2$ unless the aforementioned cancellations take place. We expect this pole and zero structure for Ruelle’s zeta function for the Kac-Baker model to be true indeed for generic $0 < \lambda < 1$.

Consider first the case $\beta = 0$. The spectrum of the operator \mathcal{L}_0 can be determined explicitly and is given by the numbers $2\lambda^k, k = 0, 1, 2, \dots$. The corresponding eigenfunctions are polynomials of degree k . Hence the Ruelle function $\zeta_R(z, \beta)$ at this point can be calculated and turns out to be $(1 - 2z)^{-1}$. This is just the Artin-Mazur zeta function for the subshift of finite type over two symbols. The Ruelle function $\zeta_R(1, \beta)$ hence takes the special value -1 at the point $\beta = 0$ independently of the parameter λ . To show now the existence of infinitely many real β -values such that the operator \mathcal{L}_β has eigenvalue $\rho = \lambda^{-1}$ it is enough to show that for $\beta \rightarrow +\infty$, respectively for $\beta \rightarrow -\infty$, infinitely many eigenvalues become larger than λ^{-1} , since the eigenvalues depend analytically on β . The asymptotic behaviour of the eigenvalues has been determined by B. Moritz in his unpublished diploma thesis [Mo89]. To formulate his result we have to introduce the parity operator $P : B(D) \rightarrow B(D)$ defined via

$$Pf(z) = f(-z).$$

It is straightforward to see that all eigenspaces of the operator \mathcal{L}_β have a basis consisting of eigenfunctions being either even (eigenvalue $+1$) or odd (eigenvalue -1) under the parity operator P . But then the spectrum of this operator is just the union of the spectra of the two following operators $\mathcal{L}_\beta^+ : B(D) \rightarrow B(D)$ and $\mathcal{L}_\beta^- : B(D) \rightarrow B(D)$ defined by

$$\mathcal{L}_\beta^+ f(z) := \exp(\beta z) f(\lambda + \lambda z) + \exp(-\beta z) f(\lambda - \lambda z),$$

respectively

$$\mathcal{L}_\beta^- f(z) := \exp(\beta z) f(\lambda + \lambda z) - \exp(-\beta z) f(\lambda - \lambda z).$$

Obviously the eigenfunctions of these two operators are even, respectively odd. Their eigenvalues will be called even, respectively odd. The main result of Moritz is then the following

Proposition 5.1 *For arbitrary N consider the N even and odd eigenvalues ρ_i of the operator \mathcal{L}_β largest in absolute value. For $0 < \lambda < \frac{1}{2}$ these eigenvalues behave for $\beta \rightarrow \infty$ as $\lambda^i \exp\left(\frac{\lambda}{1-\lambda}\beta\right)$. For $\beta \rightarrow -\infty$ the N even eigenvalues largest in absolute value behave like $(-1)^i \lambda^i \exp\left(-\frac{\lambda}{1+\lambda}\beta\right)$, whereas the odd eigenvalues behave like $(-1)^{i+1} \lambda^i \exp\left(-\frac{\lambda}{1+\lambda}\beta\right)$.*

Proof. Since the arguments for the odd eigenvalues ρ^- are similar to the even eigenvalues ρ^+ we restrict ourselves to the even eigenvalues. Consider the case $\beta \rightarrow \infty$ first. If we write in this case the eigenfunction f with eigenvalue ρ for the operator \mathcal{L}_β^+ in the form $f(z) = \exp\left(\frac{\beta z}{1-\lambda}\right) u(z)$, one finds with $\bar{\rho} = \exp\left(-\frac{\beta \lambda}{1-\lambda}\right) \rho$ and $\bar{\beta} = \frac{2\beta}{1-\lambda}$ by a simple calculation

$$\bar{\rho} u(\lambda + z') = u(\lambda + \lambda^2 + \lambda z') + \exp(-\bar{\beta} \lambda) \exp(-\bar{\beta} z') u(\lambda - \lambda^2 - \lambda z'),$$

where we have also replaced the argument z by $z' + \lambda$. For the function $h(z) := u(\lambda + z)$ one then finds the equation

$$h(\lambda z + \lambda^2) + \exp(-\bar{\beta}\lambda) \exp(-\bar{\beta}z) h(-\lambda^2 - \lambda z) = \bar{\rho}h(z).$$

Hence the function h is an eigenfunction with eigenvalue $\bar{\rho}$ of the operator $\mathcal{T}_{\bar{\beta}} : B(D_R) \rightarrow B(D_R)$ defined as

$$\mathcal{T}_{\bar{\beta}}h(z) = h(\lambda z + \lambda^2) + \exp(-\bar{\beta}\lambda) \exp(-\bar{\beta}z) h(-\lambda^2 - \lambda z),$$

where $D_R = \{z \in \mathbb{C} \mid |z| < R\}$ and $R > \frac{\lambda^2}{1-\lambda}$. This operator is nuclear and its spectrum is closely related to the one of the operator \mathcal{L}_{β}^+ . Writing now

$$\mathcal{T}_{\bar{\beta}} = \mathcal{T}_{\bar{\beta}}^+ + \mathcal{T}_{\bar{\beta}}^-$$

with the obvious definitions of the two operators $\mathcal{T}_{\bar{\beta}}^+$ and $\mathcal{T}_{\bar{\beta}}^-$ one finds for the norm of the operator $\mathcal{T}_{\bar{\beta}}^-$

$$\|\mathcal{T}_{\bar{\beta}}^-\| \leq \exp(-\bar{\beta}\lambda) \exp(\bar{\beta}R).$$

Since for $\lambda < \frac{1}{2}$ one can choose $\lambda > R > \frac{\lambda^2}{1-\lambda}$ we find $\lim_{\bar{\beta} \rightarrow \infty} \|\mathcal{T}_{\bar{\beta}}^-\| = 0$. Hence the operator $\mathcal{T}_{\bar{\beta}}$ approaches for $\bar{\beta} \rightarrow \infty$ in norm the operator $\mathcal{T}_{\bar{\beta}}^+$ and hence also their spectra are identical in this limit. The spectrum of the operator $\mathcal{T}_{\bar{\beta}}^+$, however, is given by the numbers $\{\lambda^i \mid i = 0, 1, \dots\}$.

This shows that for large β the even eigenvalues ρ^+ of the operator \mathcal{L}_{β} behave like $\lambda^i \exp\left(\frac{\lambda\beta}{1-\lambda}\right)$. To find the behaviour of the even eigenvalues of the operator \mathcal{L}_{β} for large negative values of β consider the operator $\tilde{\mathcal{L}}_{\beta}^+$ defined as

$$\tilde{\mathcal{L}}_{\beta}^+ f(z) := \exp(-\beta z) f(\lambda + \lambda z) + \exp(\beta z) f(\lambda - \lambda z)$$

and the behaviour of its eigenvalues for large positive values of β . Writing its eigenfunction f with eigenvalue ρ as $f(z) = \exp\left(\frac{\beta z}{1+\lambda}\right) u(z)$ one finds in this case with $\bar{\rho} := \exp\left(-\frac{\beta\lambda}{1+\lambda}\right)$ and $\bar{\beta} := \frac{2\beta}{1+\lambda}$ the equation

$$h(-\lambda^2 - \lambda z) + \exp(-\bar{\beta}\lambda) \exp(-\bar{\beta}z) h(\lambda^2 + \lambda z) = \bar{\rho}h(z),$$

where we have introduced again the function $h(z) = u(\lambda + z)$. An argument completely analogous to the former case then shows that the large $\bar{\beta}$ behaviour of the eigenvalues $\bar{\rho}$ is determined by the operator $\tilde{\mathcal{T}}_{\bar{\beta}}^+ h(z) = h(-\lambda^2 - \lambda z)$. Its spectrum is given by the numbers $\{(-\lambda)^i \mid i = 0, 1, \dots\}$.

Hence the even eigenvalues of the operator \mathcal{L}_{β} behave for large negative β as $(-\lambda)^i \exp\left(-\frac{\beta\lambda}{1+\lambda}\right)$. This proves the claim. \square

Let us calculate the trace of the Ruelle operator \mathcal{L}_{β} for large positive or negative values of β . Adding up N of the asymptotic eigenvalues one finds for large positive β :

$$\text{trace } \mathcal{L}_{\beta} \underset{\beta \rightarrow \infty}{\sim} \frac{1 - \lambda^{N+1}}{1 - \lambda} \exp\left(\frac{\beta\lambda}{1 - \lambda}\right).$$

For large negative β on the other hand one finds when adding up N asymptotic eigenvalues

$$\text{trace } \mathcal{L}_{\beta} \underset{\beta \rightarrow -\infty}{\sim} O(\lambda^{N+1}).$$

From this we expect that the result of Moritz on the asymptotic behaviour of the eigenvalues indeed is true for general $0 < \lambda < 1$.

Now we can prove the following proposition which determines the location of zeros and poles of the Ruelle zeta function for the Kac model:

Proposition 5.2 *For any λ with $0 < \lambda < 1$ the Fredholm determinant $\det(1 - \lambda \mathcal{L}_\beta)$ has infinitely many zeros on the real line \mathbb{R} . For $\lambda = \frac{1}{2}$ there are infinitely many zeros on the line $\operatorname{Re}(\lambda) = \ln 2$.*

Proof. Since for real β all the eigenvalues $\rho(\beta)$ of the operator \mathcal{L}_β are real and analytic in β , it follows from the asymptotic behaviour of the eigenvalues that infinitely many of them must take the value λ^{-1} for positive and negative values of the parameter β . On the other hand an easy calculation shows that for the special value $\lambda = \frac{1}{2}$ the function $f(z) = \sinh(2\beta z)$ is an eigenfunction of the operator \mathcal{L}_β for all $\beta \in \mathbb{C}$ with eigenvalue $\rho = \exp \beta$. But this eigenvalue takes the value $\lambda^{-1} = 2$ just for $\beta = \beta_n = \ln 2 + 2\pi i n$. \square

The zeros of the Fredholm determinant $\det(1 - \mathcal{L}_\beta)$ on the real line are certainly “nontrivial”, whereas the ones on the line $\operatorname{Re}(\beta) = \ln 2$ which even are equidistant could be called “trivial” ones. Since the Ruelle zeta function $\zeta_R(z)$ has the representation $\zeta_R(z) = \frac{\det(1 - \lambda \mathcal{L}_\beta)}{\det(1 - \mathcal{L}_\beta)}$ this function has infinitely many nontrivial zeros and poles on the real line, whereas for the special value $\lambda = \frac{1}{2}$ infinitely many trivial zeros lie on the line $\operatorname{Re}(z) = \ln 2$. Obviously accidental cancellations with the zeros of $\det(1 - \mathcal{L}_\beta)$ which determine the poles of the Ruelle zeta function could destroy some of these zeros.

Since the location of the nontrivial zeros of the Ruelle zeta function is not known explicitly one has to determine them numerically. Indeed in his paper [Gu82] M. Gutzwiller derived explicit formulas for the matrix elements of more or less the operator $\frac{1}{\sqrt{\lambda \exp \beta}} m_{\cosh(2\sqrt{\beta}\pi x)} \circ \mathcal{K}'_c$ in the basis given by the Hermite functions $h_k(\xi)$ in the space $L^2(\mathbb{R}, d\xi)$. After a conjugation this matrix becomes a symmetric matrix \mathcal{B}_β whose matrix elements $\mathcal{B}_{n,m}(\beta)$ have the following form:

$$\mathcal{B}_{n,m}(\beta) = 2(n!m!)^{-\frac{1}{2}} \exp\left(-\frac{(n+m)\gamma}{2}\right) \frac{M!}{(2\mu)!} \Phi(2\mu - M, 2\mu + 1; -\beta),$$

where Φ denotes the confluent hypergeometric function, $M = \max\{m, n\}$, and the number $2\mu = |m - n|$ must be even. For $|m - n|$ odd the matrix elements $\mathcal{B}_{n,m}(\beta)$ vanish. Since the number $2\mu - M$ is a non-positive integer the above confluent hypergeometric function is just proportional to the Laguerre polynomial according to the formula

$$\Phi(-n, a + 1; x) = \binom{n+a}{n}^{-1} L_n^a(x).$$

Inserting this relation into the expression for the matrix elements $\mathcal{B}_{n,m}$ one finally gets

$$\mathcal{B}_{n,m}(\beta) = 2(n!m!)^{-\frac{1}{2}} \exp\left(-\frac{(n+m)\gamma}{2}\right) \frac{M!}{(2\mu)!} \binom{M}{M-2\mu}^{-1} L_{M-2\mu}^{2\mu}(-\beta).$$

This allows one to calculate the traces of the iterates of the Kac-Gutzwiller Operator as the sum over the diagonal elements of this matrix. For instance for the trace of \mathcal{G}_β one obtains in this way

$$\operatorname{trace} \mathcal{G}_\beta = 2 \sum_{m=0}^{\infty} \exp(-\gamma m) L_m^0(-\beta) = 2 \frac{1}{1-\lambda} \exp\left(\frac{\beta\lambda}{1-\lambda}\right),$$

which is just the definition of the generating function for the Laguerre polynomials $L_m(-\beta) = L_m^0(-\beta)$. In complete analogy one can relate the traces of the iterates of the Kac-Gutzwiller operator \mathcal{G}_β calculated via the matrix $\mathcal{B}(\beta)$ to the expressions calculated via the Ruelle operator

\mathcal{L}_β and its iterates. One arrives thereby at rather complicated formulas which can be interpreted as generating functions for powers of the Laguerre polynomials $L_{M-2\mu}^{2\mu}(-\beta)$. Whether these formulas are known in the literature for the Laguerre polynomials is not known to us.

A detailed numerical study of the zeros and poles of the Ruelle zeta function by the above matrices is under way.

Acknowledgements: This work has been supported by the Deutsche Forschungsgemeinschaft through the DFG Forschergruppe “Zeta Functions and Locally Symmetric Spaces”. D. M. thanks the MPI for Mathematics in Bonn for the kind hospitality extended to him. There the final version of this paper has been prepared.

References

- [AtBo67] M. Atiyah, R. Bott. A Lefschetz fixed point formula for elliptic complexes I. *Ann. Math.* **86**, 374–407 (1967).
- [Ba61] G. Baker. One dimensional order-disorder model which approaches a second order phase transition. *Phys. Rev.* **122**, 1477–1484 (1961).
- [Be86] M. Berry. Riemann’s zeta function, a model of quantum chaos. *Lect. Notes in Physics* **263**, Springer Verlag, Berlin, 1986.
- [Co96] A. Connes. Formule de trace en geometrie non commutative et hypothese de Riemann. *C.R. Acad. Sci. Paris Ser I* **323**, 1231–1236 (1996).
- [Cr46] H. Cramér. *Mathematical Methods in Statistics*. Princeton Univ. Press, 1946.
- [De99] C. Deninger. On dynamical systems and their possible significance for arithmetic geometry. *Progr. Math.* **171**, 29–87 (1999).
- [Fo84] G.B. Folland. *Real Analysis*. Wiley, New York, 1984.
- [Fo89] G.B. Folland. *Harmonic Analysis in Phase Space*. Princeton University Press, 1989.
- [Gu82] M. Gutzwiller. The quantization of a classically ergodic system. *Physica* **5D**, 183–207 (1982).
- [Is01] S. Isola. On the spectrum of Farey and Gauss maps. Preprint University of Camerino, 2001.
- [Ka59] M. Kac. On the partition function of a one-dimensional lattice gas. *Phys. Fluids* **2**, 8–12 (1959).
- [Ka66] M. Kac. The mathematical mechanism of phase transitions. In *Brandeis University Summer Institute in Theoretical Physics*, vol 1, 245–305 eds.: H. Chretien et al. Gordon & Breach, New York, 1966.
- [Ma80] D. Mayer. The Ruelle-Araki transfer operator in classical statistical mechanics. *Lect. Notes in Physics* **123**(1), Springer Verlag, Berlin, 1980.
- [Ma90] D. Mayer. On the thermodynamic formalism for the Gauss map. *Commun. Math. Phys.* **130**, 311–333 (1990).
- [Ma91] D. Mayer. The thermodynamic formalism approach to Selberg’s zeta function for $PSL(2, \mathbb{Z})$. *Bull. Am. Math. Soc.* **25**, 55–60 (1991).
- [Mo89] B. Moritz. *Die Transferoperator-Methode in der Behandlung des Kacschen Spinmodells*. Diplomarbeit, RWTH Aachen, 1989.

- [PTMT94] A. Pentek, Z. Toroczkai, D. Mayer, T. Tel. The Kac model from a dynamical system's point of view. *Phys. Rev. E* 49, 2026–2040 (1994).
- [Ro86] P. Robba. Une introduction naive aux cohomologies de Dwork. *Mem. Soc. Math. France* 23, 61–105 (1986).
- [Ru68] D. Ruelle. Statistical mechanics of a one dimensional lattice gas. *Commun. Math. Phys.* 9, 267–278 (1968).
- [Ru92] D. Ruelle. Dynamical zeta functions: where do they come from and what are they good for? In *Proc. Internat. Congress of Math. Phys. X (Leipzig 1991)*, 43–51, Springer Verlag, Berlin, 1992.
- [Rug94] H.H. Rugh. On the asymptotic form and the reality of spectra of Perron-Frobenius operators. *Nonlinearity* 7, 1055–1066 (1994).
- [ViMa77] K. Vishwanathan, D. Mayer. Statistical mechanics of one-dimensional Ising and Potts models with exponential interactions. *Physica* 89A, 97–112 (1977).