ON ASYMPTOTIC DIMENSION OF GROUPS ACTING ON TREES

G. Bell and A. Dranishnikov

ABSTRACT. We prove the following.

Theorem. Let π be the fundamental group of a finite graph of groups with finitely generated vertex groups G_v having asdim $G_v \leq n$ for all vertices v. Then asdim $\pi \leq n + 1$.

This gives the best possible estimate for the asymptotic dimension of an HNN extension and the amalgamated product.

§0 INTRODUCTION

The notion of the asymptotic dimension asdim X of a metric space X was introduced by Gromov [Gr, pp. 28–30] (asdim₊ X in his notation) for studying asymptotic invariants of discrete groups. He gave three equivalent definitions. The first two definitions deal with coverings of metric spaces and allow us to use topological methods. The third involves mapping into Hilbert space and allows the use of more analytical techniques. We will use all three definitions as well as some characterizations of asdim given in [Dr]. Gromov's first definition is as follows: a metric space X has asymptotic dimension asdim $X \leq n$ if for any (large) number d one can find n + 1 uniformly bounded families $\mathcal{U}^0, \ldots, \mathcal{U}^n$ of d-disjoint sets in X such that the union $\cup_i \mathcal{U}^i$ is a cover of X. Two sets in a metric space are called d-disjoint if every point in the first is at least d away from every point in the second. Asymptotic dimension is a quasi-isometry invariant. Recall that the two metric spaces (X, d_X) and (Y, d_Y) are said to be quasi-isometric if there exist constants λ, ϵ and C and a map $f: X \to Y$ such that $Y \subset N_C(f(X))$ and

$$\frac{1}{\lambda}d_X(z,w) - \epsilon \le d_Y(f(z), f(w)) \le \lambda d_X(z,w) + \epsilon.$$

Typeset by \mathcal{AMS} -T_EX

¹⁹⁹¹ Mathematics Subject Classification. Primary 20H15, Secondary 20E08, 20F69.

Key words and phrases. asymptotic dimension, graph of groups, amalgamated free product, HNN extension.

The second author was partially supported by NSF grant DMS-9971709.

Here, $N_C(f(X))$ denotes the open C-neighborhood of f(X).

A generating set $S, S = S^{-1}$, in a group Γ defines the *word metric* on Γ by the following rule: $d_S(x, y)$ is the minimal length of a presentation of the element $x^{-1}y \in \Gamma$ in the alphabet S. If a group Γ is finitely generated, then all metrics d_S with finite S are quasi-isometric. Thus, one can speak about the asymptotic dimension asdim Γ of a finitely generated group Γ . Gromov proved [Gr, pg. 31] that $\operatorname{asdim} \Gamma < \infty$ for hyperbolic groups Γ . In [DJ] it was shown that Coxeter groups have finite asymptotic dimension. In [BD] we proved that the finiteness of asymptotic dimension is preserved under amalgamated products and HNN extensions of groups. Results on the finiteness of asymptotic dimension became quite important because of a theorem of Yu which states that the Novikov Higher Signature Conjecture holds for manifolds whose fundamental group has a finite asymptotic dimension [Yu].

In this paper we prove a finite asymptotic dimensionality theorem in the case of a general graph of groups. Moreover, we significantly improve the existing estimates for the asymptotic dimension of amalgamated products and HNN extensions. In particular, we show that the asymptotic dimension of the fundamental group of a graph of groups does not exceed the maximum of the asymptotic dimension of the vertex groups plus one.

We note that asymptotic dimension is a coarse invariant, i.e. it is an invariant of the coarse category introduced in [Ro]. We recall that the objects in the coarse category are metric spaces and morphisms are coarsely proper and coarsely uniform (not necessarily continuous) maps. A map $f: X \to Y$ between metric spaces is called *coarsely proper* if the preimage $f^{-1}(B_r(y))$ of every ball in Y is a bounded set in X. A map $f: X \to Y$ is called *coarsely uniform* if there is a function $\rho: \mathbf{R}_+ \to \mathbf{R}_+$, tending to infinity, such that $d_Y(f(x), f(y)) \leq \rho(d_X(x, y))$ for all $x, y \in X$. Thus, quasi-isometric metric spaces are coarsely equivalent. We note that every object in the coarse category is isomorphic to a discrete metric space, i.e. to a space with a metric taking values in the nonnegative integers \mathbf{Z}_+ .

§1 Asymptotic dimension

We recall that the order (or multiplicity) of a cover \mathcal{U} of a space X is the maximum number n such that every $x \in X$ is covered by no more than n elements of \mathcal{U} . If d > 0, the *d*-multiplicity of a cover \mathcal{U} of a metric space X is the maximum number n such that every *d*-ball $B_d(x)$ intersects no more than n elements of \mathcal{U} . Gromov's second definition of asymptotic dimension follows:

asdim $X \leq n$ if for any (large) number d > 0 there is a uniformly bounded cover \mathcal{V} of X with d-multiplicity $\leq n + 1$.

The equivalence of Gromov's first and second definitions was proven in [Gr, pp 29–31]. We recall that the Lebesgue number of a cover \mathcal{U} of a metric space X is $L(\mathcal{U}) =$ inf{max{ $d(x, X \setminus U) \mid U \in \mathcal{U}$ } | $x \in X$ }. Gromov's second definition can be reformulated as follows:

Assertion 1. asdim $X \leq n$ if for any (large) number d > 0 there is a uniformly bounded cover \mathcal{U} of X with multiplicity $\leq n + 1$ and with Lebesgue number $L(\mathcal{U}) > d$.

Proof. Let d be given and let \mathcal{V} be as in Gromov's second definition. We define $\mathcal{U} = \{N_d(V) \mid V \in \mathcal{V}\}$. Clearly, $L(\mathcal{U}) \geq d$. Since $B_d(x)$ intersects no more than n+1 elements of \mathcal{V} , no more than n+1 elements of \mathcal{U} contain x. Thus, \mathcal{U} has multiplicity $\leq n+1$.

On the other hand, let d be given and let \mathcal{U} be an open cover satisfying the conditions of Assertion 1 for 2d. We define $\mathcal{V} = \{U \setminus N_d(X \setminus U) \mid U \in \mathcal{U}\}$. Since $L(\mathcal{U}) \geq 2d$, \mathcal{V} is still a cover of X. The condition $B_d(x) \cap (U \setminus N_d(X \setminus U)) \neq \emptyset$ is equivalent to $B_d(x) \not\subset N_d(X \setminus U)$. The latter is equivalent to the condition $x \notin X \setminus U$, i.e., $x \in U$. Therefore \mathcal{V} has multiplicity $\leq n+1$. \Box

Let \mathcal{U} be an open cover of a metric space X. The canonical projection to the nerve $p: X \to Nerve(\mathcal{U})$ is defined by the partition of unity $\{\phi_U : X \to \mathbf{R}\}_{U \in \mathcal{U}}$, where $\phi_U(x) = d(x, X \setminus U) / \sum_{V \in \mathcal{U}} d(x, X \setminus V)$. The family $\{\phi_U : X \to \mathbf{R}\}_{U \in \mathcal{U}}$ defines a map p to the Hilbert space $l_2(\mathcal{U})$ with basis indexed by \mathcal{U} . The nerve $Nerve(\mathcal{U})$ of the cover \mathcal{U} is realized in $l_2(\mathcal{U})$ by taking every vertex U to the corresponding element of the basis. Clearly, the image of p lies in the nerve.

Proposition 1. For every k and every $\epsilon > 0$ there exists a number $\nu = \nu(\epsilon, k)$ such that for every cover \mathcal{U} of a metric space X of order $\leq k + 1$ with Lebesgue number $L(\mathcal{U}) > \nu$ the canonical projection to the nerve $p_{\mathcal{U}} : X \to Nerve(\mathcal{U})$ is ϵ -Lipschitz.

Proof. We take $\nu \geq \frac{(2k+3)^2}{\epsilon}$. Let $x, y \in X$ and $U \in \mathcal{U}$. The triangle inequality implies

$$|d(x, X \setminus U) - d(y, X \setminus U)| \le d(x, y).$$

Then $|\phi_U(x) - \phi_U(y)| \leq$

$$\frac{1}{\sum_{V \in \mathcal{U}} d(x, X \setminus V)} d(x, y) + d(y, X \setminus U) \left| \frac{1}{\sum_{V \in \mathcal{U}} d(x, X \setminus V)} - \frac{1}{\sum_{V \in \mathcal{U}} d(y, X \setminus V)} \right|$$
$$\leq \frac{1}{L(\mathcal{U})} d(x, y) + \frac{1}{L(\mathcal{U})} \left(\sum_{V \in \mathcal{U}} \left| d(x, X \setminus V) - d(y, X \setminus V) \right| \right) \leq \frac{2k+3}{\nu} d(x, y).$$

Then ||p(x) - p(y)|| =

$$= \left(\sum_{U \in \mathcal{U}} (\phi_U(x) - \phi_U(y))^2\right)^{\frac{1}{2}} \le \left((2k+2)\left(\frac{2k+3}{\nu}d(x,y)\right)^2\right)^{\frac{1}{2}} \le \frac{(2k+3)^2}{\nu}d(x,y) \le \epsilon d(x,y).$$

Every countable simplicial complex K can be realized in the Hilbert space l_2 in such a way that all vertices of K are vectors from some orthonormal basis. As a subspace of the metric space l_2 , K inherits the so-called *uniform metric*. All simplices in this metric are isometric to the standard simplices Δ^n (of different dimensions n). When the simplicial complex K is considered with this uniform metric, we denote it by K_U . Gromov's third definition characterizes asymptotic dimension by means of Lipschitz maps to uniform polyhedra. We note that an additional property of the canonical projection is that the preimage of every (open) simplex $p_{\mathcal{U}}^{-1}(\sigma)$ lies in the union of those $U \in \mathcal{U}$ which correspond to the vertices of σ . Thus, if the cover \mathcal{U} consists of uniformly bounded sets, then the preimages $p_{\mathcal{U}}^{-1}(\sigma)$ are also uniformly bounded. A map $f: X \to K$ to a simplicial complex is called *uniformly cobounded* if the diameters of the preimages of all simplices $f^{-1}(\sigma)$ are uniformly bounded. Since the equivalence of Gromov's third definition to other two is not given in detail in [Gr] we present it here in the form of

Assertion 2. For a metric space X the inequality asdim $X \leq n$ is equivalent to the following condition:

For any (small) $\epsilon > 0$ there is a uniformly cobounded ϵ -Lipschitz map $f : X \to K$ to a uniform simplicial complex of dimension $\leq n$.

Proof. Let K be an n-dimensional simplicial complex. We consider the cover \mathcal{V}_K of K by open stars of its vertices Ost(v, K). Clearly the multiplicity of this cover is n+1. If K is a uniform complex, then there is a uniform lower bound c_n for the Lebesgue number, i.e., $L(\mathcal{V}_K) > c_n$ for all K with dim $K \leq n$. For every uniformly cobounded ϵ -Lipschitz map $f: X \to K$ to a uniform simplicial complex K of dimension $\leq n$ the preimage $f^{-1}(\mathcal{V}_K)$ will be a uniformly bounded cover of X with multiplicity $\leq n+1$ and with Lebesgue number $> c_n/\epsilon$. By Assertion 1 we have asdim $X \leq n$.

On the other hand, let asdim $X \leq n$. Given $\epsilon > 0$ we take $\nu = \nu(\epsilon, n)$ from Proposition 1. By Gromov's second definition of asdim we can take a uniformly bounded cover \mathcal{U} of Xof multiplicity $\leq n+1$ with Lebesgue number $> \nu$. Then by Proposition 1 the projection $p_{\mathcal{U}}: X \to Nerve(\mathcal{U})$ is an ϵ -Lipschitz map to a uniform *n*-dimensional polyhedron. The preimage $p_{\mathcal{U}}^{-1}(\sigma)$ of every simplex is contained in a union of elements of \mathcal{U} having a common point. Hence $p_{\mathcal{U}}$ is uniformly cobounded. \Box

We will use the following theorems proven in [BD].

Finite Union Theorem. Suppose that a metric space is presented as a union $A \cup B$ of subspaces. Then asdim $A \cup B \leq \max\{\text{asdim } A, \text{asdim } B\}$.

A family of metric spaces $\{F_{\alpha}\}$ satisfies the inequality asdim $F_{\alpha} \leq n$ uniformly if for arbitrary large d > 0 there exist R and R-bounded d-disjoint families $\mathcal{U}_{\alpha}^{0} \dots \mathcal{U}_{\alpha}^{n}$ of subsets of F_{α} such that the union $\cup_{i} \mathcal{U}_{\alpha}^{i}$ is a cover of F_{α} .

4

Infinite Union Theorem. Assume that $X = \bigcup_{\alpha} F_{\alpha}$ and asdim $F_{\alpha} \leq n$ uniformly. Suppose that for any r there exists $Y_r \subset X$ with asdim $Y_r \leq n$ and such that the family $\{F_{\alpha} \setminus Y_r\}$ is r-disjoint. Then asdim $X \leq n$.

§2 Simplicial mapping cylinders

An orientation on a simplicial complex K is a linear order on its vertices. Every subcomplex of an oriented complex has the induced orientation.

On the product of two ordered sets $A \times B$ we define a partial ordering called the *product order* by the rule: (a, b) < (a', b') if either a < a' and $b \le b'$ or $a \le a'$ and b < b'.

Let σ be an oriented k-simplex x_0, \ldots, x_k . The product order on $\{x_0, \ldots, x_k\} \times \{0, 1\}$ defines a triangulation on the prism $\sigma \times [0, 1]$ by the following rule: every chain $(x_{i_0}, y_{j_0}) < \cdots < (x_{i_r}, y_{j_r}), y_j = 0, 1$, spans a simplex. We note that the set of vertices for this triangulation equals $\{x_0, \ldots, x_k\} \times \{0, 1\}$.

Proposition 2. For every simplicial map $f : X \to Y$ the mapping cylinder M_f admits a triangulation with the set of vertices equal to the disjoint union of vertices of X and Y.

Proof. We consider an orientation on X. For every simplex σ in X we consider the triangulation of the prism $\sigma \times [0, 1]$ defined above. Since the orientations on simplices σ agree with each other, it defines a triangulation on $X \times [0, 1]$. Let Z = f(X). We consider the image M of the simplicial map $g: X \times [0, 1] \to M$ defined by means of the map f on $X \times \{1\}$. Then M_f is a simplicial complex as it is the union of the simplicial complexes M and Y along the common subcomplex Z. \Box

By the product metric on the cartesian product $X \times Y$ of two metric spaces (X, d_X) and (Y, d_Y) we mean the metric

$$d((x_1, y_1), (x_2, y_2)) = \sqrt{d_X(x_1, x_2)^2 + d_Y(y_1, y_2)^2}$$

where $x_1, x_2 \in X$ and $y_1, y_2 \in Y$.

Proposition 3. For every *n* there is a constant c_n so that for any simplicial map $g : X \to Y$ of an oriented *n*-dimensional simplicial complex X the quotient map $q : X \times [0,1] \to M_g$ of the product $X \times [0,1]$ to the mapping cylinder M_g is c_n -Lipschitz where X and M_g are given the uniform metrics and $X \times [0,1]$ has the product metric.

Proof. We note that the quotient map $q: X \times [0,1] \to M_g$ can be factored through the uniformization $q': X \times [0,1] \to (X \times [0,1])_U$ where the triangulation on $X \times [0,1]$ is defined by means of the orientation on X and $(X \times [0,1])_U$ is a uniform complex. Thus $q = \bar{q} \circ q'$ where $\bar{q}: (X \times [0,1])_U \to M_g$ is a simplicial map between uniform complexes. Since \bar{q} is always 1-Lipschitz, it suffices to show that there is a constant c_n so that for

any oriented *n*-dimensional complex X the uniformization $q': X \times [0,1] \to (X \times [0,1])_U$ is c_n -Lipschitz.

For each k > 0, by a *k*-prism we mean a subset of $X \times [0, 1]$ of the form $\sigma \times [0, 1]$ for some k - 1-simplex σ of X. First, for each k there exists a constant λ_k so that a uniformization of each k-prism is λ_k -Lipschitz.

With the λ_k as above, take $c_n = \max\{\lambda_k \mid 1 \le k \le 2n+2\}$. It remains to show that q' is c_n -Lipschitz. To this end, let x and y in $X \times [0,1]$ be given. Take a simplex σ in X of minimal dimension with $x \in \sigma \times [0,1]$, and a simplex $\tau \in X$ of minimal dimension with $y \in \tau \times [0,1]$.

Since the complex X is considered in the uniform metric, its vertices span a (infinite) simplex Δ_0 . The orientation on X defines an orientation on Δ_0 . Therefore the triangulation on $X \times [0, 1]$ is naturally extendable to a triangulation on $\Delta_0 \times [0, 1]$ with the same set of vertices.

Let \tilde{q} denote the uniformization map of $\Delta_0^{(2n+1)} \times [0,1]$. Then $\tilde{q}|_{X \times [0,1]} = q'$.

Let ρ be a simplex in Δ_0 of minimal dimension so that σ and τ are faces of ρ . Then, dim $\rho \leq \dim \sigma + \dim \tau + 1 \leq 2n+1$. Now, \tilde{q} is c_n -Lipschitz when restricted to the k-prism $\rho \times [0, 1]$. Note that $k \leq 2n+2$. Thus, $d(q(x), q(y)) = d(\tilde{q}(x), \tilde{q}(y)) \leq c_n d(x, y)$. Thus, qis c_n -Lipschitz. \Box

Proposition 4. Let $A \subset W \subset X$ be subsets in a geodesic metric space X such that the r-neighborhood $N_r(A)$ is contained in W and let $f : W \to Y$ be a continuous map to a metric space Y. Assume that the restrictions $f|_{N_r(A)}$ and $f|_{W\setminus N_r(A)}$ are ϵ -Lipschitz. Then f is ϵ -Lipschitz.

Proof. Let $x, y \in W$ be two points. The only possibility for difficulties with checking the Lipschitz condition can occur when x and y lie in different sets $N_r(A)$ and $W \setminus N_r(A)$. Let $x \in N_r(A)$ and $y \in W \setminus N_r(A)$ and let $\gamma : [0, d(x, y)] \to X$ be a geodesic segment joining them in X. Then there is a $t \in [0, d(x, y)]$ such that $\gamma(t) \in \partial N_r(A)$, where $\partial N_r(A) = \{z \in X \mid d(z, A) = r\}$. Then $d_Y(f(x), f(y)) \leq d_Y(f(x), f(\gamma(t))) + d_Y(f(\gamma(t)), f(y)) \leq \epsilon d_X(x, \gamma(t)) + \epsilon d_X(\gamma(t), y) = \epsilon d_X(x, y)$. \Box

We recall that $L(\mathcal{U})$ denotes the Lebesgue number of a cover \mathcal{U} of a metric space. If \mathcal{U} is a cover of a subset $A \subset X$ of a metric space, we define $L(\mathcal{U}) = \inf \{ \sup\{d(x, X \setminus U) \mid U \in \mathcal{U}\} \mid x \in A \}$. Let $b(\mathcal{U})$ denote the least upper bound for the diameters of $U \in \mathcal{U}$ and let $\nu(\epsilon, k)$ be the number defined by Proposition 1.

Lemma 1. Let $r > 8\epsilon$ and let \mathcal{V} and \mathcal{U} be covers of the r-neighborhood $N_r(A)$ of a closed subset A in a geodesic metric space X by open sets such that both \mathcal{U} and \mathcal{V} have order $\leq n + 1$, $Nerve(\mathcal{V})$ is orientable, and $L(\mathcal{U}) > b(\mathcal{V}) > L(\mathcal{V}) \geq \nu(\epsilon/4c_n, n)$, where c_n is the constant from Proposition 3. Then there is an ϵ -Lipschitz map f: $N_r(A) \to M_g$ to the mapping cylinder supplied with the uniform metric of a simplicial map $g: Nerve(\mathcal{V}) \to Nerve(\mathcal{U})$ between the nerves such that f is uniformly cobounded, $f|_{\partial N_r(A)} = q(p_{\mathcal{V}}|_{\partial N_r(A)}, 0), \text{ and } f|_A = p_{\mathcal{U}}|_A, \text{ where } p_{\mathcal{U}} : N_r(A) \to Nerve(\mathcal{U}) \text{ and } p_{\mathcal{V}} : N_r(A) \to Nerve(\mathcal{V}) \text{ are the canonical projections to the nerves.}$

Proof. We recall that the simplicial complexes $Nerve(\mathcal{U})$ and $Nerve(\mathcal{V})$ inherit their metrics from the Hilbert spaces $l_2(\mathcal{U})$ and $l_2(\mathcal{V})$. By Proposition 1 the projections $p_{\mathcal{U}}$: $N_r(A) \to Nerve(\mathcal{U})$ and $p_{\mathcal{V}}: N_r(A) \to Nerve(\mathcal{V})$ are $\epsilon/4c_n$ -Lipschitz. Let $t_x = \frac{2d_X(x,A)}{c_n r}$, we define $\int a(p_{\mathcal{V}}(x), 2-t_n) \qquad if \ d(x,A) > c_n r/2$

$$f(x) = \begin{cases} q(p_{\mathcal{V}}(x), 2 - t_x) & \text{if } d(x, A) > c_n r/\\ t_x g p_{\mathcal{V}}(x) + (1 - t_x) p_{\mathcal{U}}(x) & \text{otherwise.} \end{cases}$$

In the first case $q^{-1}f(x)$ lies in $Nerve(\mathcal{V}) \times [0,1)$. The map $q^{-1}f: N_r(A) \to Nerve(\mathcal{V}) \times [0,1)$ is $\max\{\epsilon\sqrt{2}/4c_n, 2\sqrt{2}/c_nr\}$ -Lipschitz and is therefore ϵ/c_n -Lipschitz. In the second case the linear combination is taken in the Hilbert space $l_2(\mathcal{U})$. It belongs to $Nerve(\mathcal{U})$, since $gp_{\mathcal{V}}(x)$ and $p_{\mathcal{U}}(x)$ lie in the simplex of $Nerve(\mathcal{U})$ spanned by all $U \in \mathcal{U}$ that contain x. Note that the coefficients in the linear combination are $2/c_nr$ -Lipschitz functions. Therefore, they are $\epsilon/4c_n$ -Lipschitz, as $2/r \leq \epsilon/4$.

Proposition 3 implies q is c_n -Lipschitz. Hence the restriction $qq^{-1}f|_{N_{c_nr}(A)\setminus N_{c_nr/2}}$ is ϵ -Lipschitz. Since the coefficients t_x and $1-t_x$ are $\epsilon/4c_n$ -Lipschitz and $||gp_{\mathcal{V}}(x)||, ||p_{\mathcal{U}}(x)|| \leq 1$, it follows from the Leibnitz rule that the restriction of f to $N_{c_nr/2}(A)$ is also ϵ -Lipschitz. In view of Proposition 4 we can conclude that f is an ϵ -Lipschitz map. Note that $f|_{\partial N_r(A)} = p_{\mathcal{V}}|_{\partial N_r(A)}$ and $f|_A = p_{\mathcal{U}}|_A$.

Finally, we show diam $f^{-1}(q(\sigma \times [0,1])) \leq 2b(\mathcal{U})$. For x and y in $f^{-1}(q(\sigma \times [0,1]))$, there are three cases to consider, based on dist(x, A) and dist(y, A). If both dist(x, A)and dist $(y, A) > c_n r/2$, then $f(x) = q(p_{\mathcal{V}}(x), 2 - t_x)$ and $f(y) = q(p_{\mathcal{V}}(y), 2 - t_y)$, so x and y are in σ under the projection to the nerve $p_{\mathcal{V}}$. So, there is some set $V \in \mathcal{V}$ containing x and y. Thus, dist $(x, y) \leq b(\mathcal{V}) < b(\mathcal{U})$. Next, if dist $(x, A) < c_n r/2$ and dist $(y, A) \geq c_n r/2$, then f(x) is the same as above, and f(y) is a linear combination of elements of a simplex in $Nerve(\mathcal{V})$, with coefficients at most 1. The map g takes an open set $V \in \mathcal{V}$ containing x into a set $U \in \mathcal{U}$ containing the elements of the linear combination in the expression for f(y). Thus, dist $(x, y) \leq 2b(\mathcal{U})$. Finally, if both elements lie in the closed $c_n r/2$ neighborhood of A, then both of the elements lie in a linear combination of elements of a simplex in $Nerve(\mathcal{U})$, so dist $(x, y) \leq 2b(\mathcal{U})$. \Box

§3 GROUPS ACTING ON TREES

A norm on a group A is a map $|| || : A \to \mathbb{Z}_+$ such that $||ab|| \le ||a|| + ||b||$, $||a^{-1}|| = ||a||$ and ||x|| = 0 if and only if x is the unit in A. A set of generators $S \subset A$ with $S = S^{-1}$ defines the norm $||x||_S$ as the minimal length of a presentation of x in terms of S. A norm on a group defines a left-invariant metric d by $d(x,y) = ||x^{-1}y||$. If G is a finitely generated group and S and S' are two finite generating sets, then the corresponding metrics d_S and $d_{S'}$ define coarsely equivalent metric spaces (G, d_S) and

 $(G, d_{S'})$. In particular, $\operatorname{asdim}(G, d_S) = \operatorname{asdim}(G, d_{S'})$, and we can speak about the asymptotic dimension $\operatorname{asdim} G$ of a finitely generated group G.

Assume that a group Γ acts on a metric space X. For every R > 0 we define the *R*-stabilizer $W_R(x_0)$ of the point $x_0 \in X$ to be the set of all $g \in \Gamma$ with $g(x_0) \in B_R(x_0)$. Here $B_R(x)$ denotes the closed ball of radius R centered at x.

Lemma 2. Assume that a finitely generated group Γ acts by isometries on a tree X with a base point x_0 . Suppose that asdim $W_R(x_0) \leq k$ for all R. Then asdim $\Gamma \leq k + 1$.

Proof. Let $S = S^{-1}$ be a finite generating set for Γ . We define a map $\pi : \Gamma \to X$ by the formula $\pi(g) = g(x_0)$. Using the uniqueness of a geodesic segment between any two points in X this map can be canonically extended to a Γ -equivariant map $\bar{\pi} : Y \to X$ of the Cayley graph Y of the group Γ . Let $\lambda = \max\{d_X(s(x_0), x_0) \mid s \in S\}$. We show now that π is λ -Lipschitz. Since the metric d_S on Γ is induced from the geodesic metric on the Cayley graph, it suffices to check that $d_X(\pi(g), \pi(g')) \leq \lambda$ for all $g, g' \in \Gamma$ with $d_S(g, g') = 1$. Without loss of generality we assume that g' = gs, where $s \in S$. Then $d_X(\pi(g), \pi(g')) = d_X(g(x_0), gs(x_0)) = d_X(x_0, s(x_0)) \leq \lambda$. Since every edge in the Cayley graph maps via $\bar{\pi}$ to a geodesic segment, it follows that $\bar{\pi}$ is also λ -Lipschitz.

Observe that $W_R(x_0) = \pi^{-1}(B_R(x_0))$, $\gamma B_R(x) = B_R(\gamma(x))$ and $\gamma(\bar{\pi}^{-1}(B_R(x))) = \bar{\pi}^{-1}(B_R(\gamma(x)))$ for all $\gamma \in \Gamma$, $x \in X$ and all R. We note that the spaces $W_R(x_0)$ and $\bar{\pi}^{-1}(B_R(x_0))$ are in a finite Hausdorff distance from each other and therefore, coarsely equivalent. We denote the latter by $\bar{W}_R(x_0)$.

Let c_k be the Lipschitz constant defined by uniformization of the product $K' \times [0, 1]$, for a k-dimensional simplicial complex K' (see Proposition 3).

Given a (small) number $\epsilon > 0$, we will construct an ϵ -Lipschitz, uniformly cobounded map $\psi : Y \to K$ to a uniform k + 1-dimensional simplicial complex. Let $\nu = \nu(\epsilon/4c_k, k)$ be from Proposition 1, and take $r > \max\{\nu, 8/\epsilon\}$. We need only consider the Γ -orbit of x_0 in order to get information about Γ . Since asdim $X \leq 1$, we have asdim $\Gamma x_0 \leq 1$. Thus there is a uniformly bounded cover \mathcal{W} of Γx_0 with multiplicity 2 such that the λr -enlargement $\{N_{\lambda r}(W)\}$ is a cover of Γx_0 with multiplicity 2. Indeed, using Gromov's first definition of asdim, we can find two families of uniformly bounded, d-disjoint sets \mathcal{W}_1 and \mathcal{W}_2 whose union covers Γx_0 with $d \gg \lambda r$; then, put $\mathcal{W} = \mathcal{W}_1 \cup \mathcal{W}_2$. For each $W \in \mathcal{W}$, let \overline{W} denote $\overline{\pi}^{-1}(W)$. As \mathcal{W} is uniformly bounded, there is some R > 0 so that for each $W \in \mathcal{W}$ we can find $x_W \in \Gamma x_0$ with the property that $N_{\lambda r}(W) \subset B_R(x_W)$. As $\overline{\pi}$ is λ -Lipschitz, we have $\overline{\pi}(N_r(\overline{W})) \subset N_{\lambda r}(W)$. Thus, $N_r(\overline{W}) \subset \overline{\pi}^{-1}(B_R(x_W)) = \overline{W}_R(x_W)$. Let $\gamma_W \in \Gamma$ have $\pi(\gamma_W) = x_W$. We use the condition asdim $\overline{W}_R(x_0) \leq k$ to construct two families of uniformly bounded covers \mathcal{V} and \mathcal{U} of the set $\overline{W}_R(x_0)$ both with multiplicity $\leq k + 1$ such that the Lebesgue number $L(\mathcal{V})$ is greater than r and $L(\mathcal{U}) > b(\mathcal{V})$, where $b(\mathcal{V})$ is an upper bound for diameters of the cover \mathcal{V} .

First, we construct a family of uniformly cobounded ϵ -Lipschitz maps $\phi_W : N_r(\overline{W}) \to K_W$ to uniform k-dimensional simplicial complexes as the canonical projections to the

nerve of $\gamma_W \mathcal{V}$ restricted to $N_r(W)$.

Next, let W and W' be elements of W such that $W \cap W' \neq \emptyset$. Thus, in the nerve, Nerve(W), the sets W and W' define an edge e. Put $A_e = W \cap W'$, and let \bar{A}_e denote $\bar{\pi}^{-1}(A_e)$. Observe that $N_r(\bar{A}_e) \subset N_r(\bar{W})$ and $N_r(\bar{W}')$. Let \mathcal{V}_e and \mathcal{U}_e denote the restrictions of the covers \mathcal{V} and \mathcal{U} to $N_r(\bar{A}_e)$. We take the triangulation on the mapping cylinder M_{g_e} of a simplicial map $g_e : Nerve(\mathcal{V}_e) \to Nerve(\mathcal{U}_e)$ as in Proposition 2, and consider M_{g_e} as a uniform simplicial complex. As $r > 8/\epsilon$, we may apply Lemma 1 to $\bar{A}_e \subset Y$ and to the covers \mathcal{V}_e and \mathcal{U}_e to obtain an ϵ -Lipschitz map $f_e: N_r(\bar{A}_e) \to M_{g_e}$. We note that the $N_r(\bar{A}_e)$ are disjoint for distinct edges in the nerve. Indeed, $\bar{\pi}(N_r(\bar{A}_e)) \subset N_{\lambda r}(A_e) \subset N_{\lambda r}(W) \cap N_{\lambda r}(W')$; so, if $N_r(\bar{A}_e) \cap N_r(A_{e'}) \neq \emptyset$, then the λr -enlargement of \mathcal{W} would have multiplicity at least 3. Thus, for each $W \in \mathcal{W}$ define $\psi_W: N_r(\bar{W}) \to K_W \cup_{W \in e} M_{g_e} = L_W$ to the uniform complex L_W , with mapping cylinders attached as the union of the map $\phi_W|_{N_r(\bar{W}) \setminus \cup_{W \in e} N_r(\bar{A}_e)}$ and the restrictions $f_e|_{N_r(\bar{A}_e) \cap N_r(\bar{W})}$, for all edges e in $Nerve(\mathcal{W})$ which contain W as a vertex in the nerve $Nerve(\mathcal{W})$.

We construct K by gluing together the L_W . Clearly, the dimension of K is at most k + 1. The maps $\psi_W : Y \to K$ agree on the common parts \bar{A}_e so they define a map $\psi : Y \to K$. The map ψ is ϵ -Lipschitz by Proposition 4, and uniformly cobounded by the Lemma 1. \Box

In the next section we show (Lemma 3) that if a group Γ acts on a tree with compact quotient and with asdim $\Gamma_x \leq n$ for all stabilizers, then asdim $W_R(x_0) \leq n$ for all R. As a corollary of this and Lemma 2 we obtain the following:

Theorem 1. Assume that a finitely generated group Γ acts on a tree X with a compact orbit space with the stabilizers Γ_x having $\operatorname{asdim} \Gamma_x \leq n$ for all vertices $x \in X$. Then $\operatorname{asdim} \Gamma \leq n+1$.

§4 Graph of groups

We recall some of the basic constructions and known facts about graphs of groups. Our development and notation closely follow that of [S]. Let Y be a non-empty, connected graph. To each $P \in \text{Vert } Y$, associate a group G_P and to each $y \in \text{Edge } Y$, a group $G_y = G_{\bar{y}}$ equipped with two injective homomorphisms, $\phi_y : G_y \to G_{t(y)}$ and $\phi_{\bar{y}} : G_{\bar{y}} \to G_{i(y)}$.

Define the group F(G, Y) to be the group generated by the elements of the G_P and the elements $y \in \text{Edge } Y$ subject to the relations:

$$\bar{y} = y^{-1}$$

and

$$y\phi_y(a)y^{-1} = \phi_{\bar{y}}(a),$$

if $y \in \text{Edge } Y$ and $a \in G_y$. Let c be a path in Y starting at some vertex P_0 . Let y_1, y_2, \ldots, y_n denote the edges associated to c, where $t(y_i) = P_i$. Then the length of c is n, and we write $i(c) = P_0$ and $t(c) = P_n$. A word of type c in F(G, Y) is a pair (c, μ) where c is a path as above and μ is a sequence r_0, r_1, \ldots, r_n , where $r_i \in G_{P_i}$, The associated element of the group F(G, Y) is

$$|c, \mu| = r_0 y_1 r_1 y_2 r_2 \cdots y_n r_n \in F(G, Y).$$

Serre gives two equivalent definitions of the fundamental group of the graph of groups (G, Y). For the first description, let P_0 denote a fixed vertex and G_{P_0} the associated group. Let $\pi = \pi_1(G, Y, P_0)$ be the set of elements of F(G, Y) associated to a path c in Y with $i(c) = t(c) = P_0$. Obviously, $\pi \subset F(G, Y)$ is a subgroup. For the second description, let T be a maximal subtree of Y, and define $\pi = \pi_1(G, Y, T)$ to be the quotient of F(G, Y) by the normal subgroup generated by the elements $t \in \text{Edge } T$. If g_y denotes the image of $y \in \text{Edge } Y$ in π , then the group π is the group generated by the groups G_P and the elements g_y subject to the relations

$$g_{\bar{y}} = g_y^{-1},$$
$$g_y \phi_y(a) g_y^{-1} = \phi_{\bar{y}}(a),$$

and

 $g_t = 1$

where $a \in G_y$, and $t \in \text{Edge } T$. So, in particular, $\phi_t(a) = \phi_{\bar{t}}(a)$ for all $t \in \text{Edge } T$. The equivalence of the descriptions is proven in [S].

Examples. (1) If Y is the graph with two vertices P, Q and one edge y, then $\pi_1(G, Y, P) = \pi_1(G, Y, Q) = G_P *_{G_y} G_Q$, the free product of G_P and G_Q amalgamated over G_y .

(2) If Y is the graph with one vertex P and one edge y then $\phi_{\bar{y}}(G_y)$ is a subgroup of G_P , and $\pi_1(G, Y, P)$ is precisely the HNN extension of G_P over the subgroup $\phi_{\bar{y}}(G_y)$ by means of $\phi_y \phi_{\bar{y}}^{-1}$.

We now describe the construction of the so-called Bass-Serre tree X on which π will act by isometries. Let T be a maximal tree and let π_P denote the canonical image of G_P in π , obtained via conjugation by the path c, where c is the unique path in T from the basepoint P_0 to the vertex P. Similarly, let π_y denote the image of $\phi_y(G_{t(y)})$ in $\pi_{t(y)}$. Then, set

$$\operatorname{Vert} \tilde{X} = \coprod_{P \in \operatorname{Vert} Y} \pi / \pi_P$$

and

$$\operatorname{Edge} \tilde{X} = \coprod_{y \in \operatorname{Edge} Y} \pi/\pi_y.$$

For a more explicit description of the edges, observe that the vertices $x\pi_{i(y)}$ and $xy\pi_{t(y)}$ are connected by an edge for all $y \in \text{Edge } Y$ and all $x \in \pi$. Obviously the stabilizer of the vertices are conjugates of the corresponding vertex groups, and the stabilizer of the edge connecting $x\pi_{i(y)}$ and $xy\pi_{t(y)}$ is $xy\pi_yy^{-1}x^{-1}$, a conjugate of the image of the edge group. This obviously stabilizes the second vertex, and it stabilizes the first vertex since $y\pi_yy^{-1} = \pi_{\bar{y}} \subset \pi_{i(y)}$. It is known (see [S]) that the action of left multiplication on \tilde{X} is isometric.

Now we will assume that the graph Y is finite and that the groups associated to the edges and vertices are finitely generated with some fixed set of generators chosen for each group. We let S denote the disjoint union of the generating sets for the groups, and require that $S = S^{-1}$. By the norm ||x|| of an element $x \in G_P$ we mean the minimal number of generators in the fixed generating set required to present the element x. We endow each of the groups G_P with the word metric given by $dist(x, y) = ||x^{-1}y||$. We extend this metric to the group F(G, Y) and hence to the subgroup $\pi_1(G, Y, P_0)$ in the natural way, by adjoining to S the collection $\{y, y^{-1} \mid y \in Edge Y\}$.

Proposition 5. Let Y be a non-empty, finite, connected graph, and (G, Y) the associated graph of finitely generated groups. Let P_0 be a fixed vertex of Y, then under the action of π on \tilde{X} , the R-stabilizer $W_R(1.P_0)$ is precisely the set of elements of type c in F(G, Y) with $i(c) = P_0$, and $l(c) \leq R$.

Proof. First, every $x = |c, \mu| \in K$ with $l(c) \leq R$ is in $W_R(1.P_0)$ since $\operatorname{dist}_{\tilde{X}}(xG_{P_0}, G_{P_0})$ is the number of edges in the path c which is at most R. For the reverse inclusion, it suffices to show that given a reduced word $x = |c, \mu|$, with l(c) = R, the length of any coset representative xG_{P_0} is also R. Consider the word $xg = r_0y_1r_1\cdots y_nr_ng$, where $y_i \in \operatorname{Edge} Y, r_i \in G_{P_i}$ and $g \in G_{P_0}$. The relations on F(G, Y) allow for only one type of reduction involving edges to occur. It occurs in one of the following two forms:

(1)
$$y\phi_y(a) = \phi_{\bar{y}}(a)y$$
, or
(2) $y\phi_y(a)y^{-1} = \phi_{\bar{y}}(a)$,

where $a \in G_y$ and $y \in \text{Edge } Y$. Clearly only a reduction of type (2) can decrease the (path) length of the word. So, assume that there are edges $y_{j+1}^{-1} = y_j$ and through a sequence of type-(1) reductions, x is translated to

$$r_0y_1\cdots y_jr_j\phi_{y_j}(b)y_{j+1}r'_{j+1}\cdots y_nr'_n.$$

Now, in order for a type-(2) reduction to occur here, we must have $r_j \phi_{y_j}(b) = \phi_{y_j}(d)$ for some $d \in G_{y_j}$. This implies that $r_j = \phi_{y_j}(db^{-1})$, which enables a type-(2) reduction of the original word x. As x was assumed to be reduced, this cannot occur. \Box **Lemma 3.** If a group Γ acts on a tree with compact quotient and finitely generated stabilizers satisfying asdim $\Gamma_x \leq n$ for all vertices x, then asdim $W_R(x_0) \leq n$ for all R.

Proof. Consider the subset $K \subset F(G, Y)$ of all words whose associated path c has $i(c) = P_0$. This set acts on the tree by left multiplication, and the fundamental group is a subset of K. We consider the R-stabilizer $W_R(P_0)$ of the fixed vertex as a subset of K, and show that this set has asymptotic dimension at most n by induction. It will follow then that the stabilizers in the fundamental group will also have asymptotic dimension at most n.

We proceed by induction. The base case is clear since W_0 is precisely G_{P_0} and by assumption, asdim $W_0 \leq n$. By Proposition 5, W_R is precisely the set of all words in Kwith associated path c for which $l(c) \leq R$. Let $H_k \subset K$ be the set of elements whose paths have length precisely k. Then, in the above notation, $W_R = \bigcup_{k \leq R} H_k$, so by the Finite Union Theorem (§1), it suffices to prove that asdim $H_k \leq n$ for all k. Since W_0 and H_0 coincide, we proceed to the induction step on H_k .

Observe that $H_k \subset \bigcup_{y \in Edge Y} H_{k-1}yG_{t(y)}$. Since Y was assumed to be a finite graph, this is a finite union. So, by the Finite Union Theorem, it suffices to show that asdim $H_k \cap H_{k-1}yG_{t(y)} \leq n$, for some fixed $y \in Edge Y$. Next, we let $Y_r = H_{k-1}yN_r(\phi_y(G_y))$, where $N_r(\phi_y(G_y))$ is the r-neighborhood of $\phi_y(G_y)$ in F(G,Y). Now, the set Y_r is coarsely equivalent to $H_{k-1}y\phi_y(G_y)$, and by the relations on F(G,Y), this set is $H_{k-1}\phi_{\bar{y}}(G_y)y =$ $H_{k-1}y$. Finally, $H_{k-1}y$ is coarsely equivalent to H_{k-1} , which, by the inductive hypothesis, has asymptotic dimension not exceeding n. Thus, we conclude that asdim $Y_r \leq n$. Consider the sets $xyG_{t(y)}$, indexed by those $x \in H_{k-1}$, which do not end with an element of $\phi_{\bar{y}}(G_y)$. Notice that this collection does cover $H_k \cap H_{k-1}yG_{t(y)}$ since we may obtain $x\phi_{\bar{y}}(a)yg$ from $xy\phi_y(a)g$, which is of the required form. The map $G_{t(y)} \mapsto xyG_{t(y)}$ is an isometry in the word metric, so $asdim(xyG_{t(y)}) \leq n$ uniformly.

To apply the Infinite Union Theorem from §1, we need only show that the family $\{H_k \cap H_{k-1}yG_{t(y)} \setminus Y_r\}$ is r-disjoint. To this end, let $x \neq x'$ and suppose that z and z' are in $G_{t(y)} \setminus N_r(\phi_y(G_y))$. Consider dist $(xyz, x'yz') = ||z^{-1}y^{-1}x^{-1}x'yz'||$. Write $z = \phi_y(a)s$, and $z' = \phi_y(a')s'$, where ||s|| and ||s'|| > r. Then,

$$\|z^{-1}y^{-1}x^{-1}x'yz'\|$$

= $\|s^{-1}y^{-1}\phi_{\bar{y}}(a^{-1})x^{-1}x'\phi_{\bar{y}}(a')ys'\|.$

Now, a reduction can only occur in the middle, and if $\phi_{\bar{y}}(a^{-1})x^{-1}x'\phi_{\bar{y}}(a')$ is not in $\phi_{\bar{y}}(G_y)$, then

$$||z^{-1}y^{-1}x^{-1}x'yz'|| > ||s|| + ||s'|| > 2r.$$

So, suppose that there is a $b \in G_y$ so that $\phi_{\bar{y}}(b) = \phi_{\bar{y}}(a^{-1})x^{-1}x'\phi_{\bar{y}}(a')$. Then, putting $c = ab(a')^{-1}$, we see that $x' = x\phi_{\bar{y}}(c)$. By construction x' cannot end with a nontrivial element of $\phi_{\bar{y}}$, so that $\phi_{\bar{y}}(c) = e$, which implies x' = x, a contradiction.

We conclude that the families are r-disjoint, and therefore, by the Infinite Union Theorem, asdim $H_k < n$, so that asdim $W_R < n$ for every R. \Box

In view of the Bass-Serre structure theorem [S], every group acting without inversion on a tree with compact quotient is a fundamental group of a graph of groups, so we may reformulate Theorem 1 in terms of graphs of groups.

Theorem 1'. Let (G, Y) be a finite graph of groups with finitely generated vertex groups satisfying the inequality asdim $G_v \leq n$ for all vertex groups. Then for the fundamental group we have asdim $\pi_1(G, Y, v_0) \leq n + 1$ for any vertex v_0 .

The estimate of Theorem 1' is exact for HNN extensions. It is also exact for the general amalgamated product. For example, asdim $A *_C B = 2$, where $A = B = C = \mathbb{Z}$ and the inclusions $C \to A$ and $C \to B$ are equal to the multiplication by 2. Indeed, $A *_C B$ is isomorphic to the fundamental group of the Klein bottle. Hence the group $A *_C B$ is coarsely isomorphic to the universal cover X of the Klein bottle which is equal (topologically) to \mathbb{R}^2 . One can show that X is coarsely isomorphic to the Euclidean plane \mathbb{R}^2 and then asdim $X = \operatorname{asdim} A *_C B = 2$. Another possibility here is to apply Gromov's estimate for the asymptotic dimension of a uniformly contractible manifold X [Gr, pg. 32]: for a uniformly contractible manifold X without boundary, asdim $X = \operatorname{dim} X$.

We believe that the exact formula in the case of the amalgamated product should be asdim $A*_C B = \max\{\operatorname{asdim} A, \operatorname{asdim} B, \operatorname{asdim} C+1\}$ where neither of the monomorphisms $C \to A$ and $C \to B$ is an isomorphism.

References

- [BD] G. Bell and A. Dranishnikov, On asymptotic dimension of groups, Algebraic and Geometric Topology 1 (2001), 57-71.
- [Dr] A. Dranishnikov, Asymptotic topology, Russian math. Surveys 55 (2000), No 6, 71-116.
- [DJ] A. Dranishnikov and T. Januszkiewicz, Every Coxeter group acts amenably on a compact space, Topology Proceedings 24 (1999), 135-141.
- [Gr] M. Gromov, Asymptotic invariants of infinite groups, Geometric Group Theory, vol 2, Cambridge University Press, 1993.
- [Ro] J. Roe, Coarse cohomology and index theory for complete Riemannian manifolds, Memoirs Amer. Math. Soc. No. 497, 1993.
- [S] J.-P. Serre, *Trees*, Springer-Verlag, 1980.
- [Yu] G. Yu, The Novikov conjecture for groups with finite asymptotic dimension, Ann. of Math 147 (1998), no. 2, 325-355.

UNIVERSITY OF LOUISVILLE, DEPARTMENT OF MATHEMATICS, 328 NATURAL SCIENCES BUILDING, LOUISVILLE, KY 40292, USA

UNIVERSITY OF FLORIDA, DEPARTMENT OF MATHEMATICS, P.O. BOX 118105, 358 LITTLE HALL, GAINESVILLE, FL 32611-8105, USA

E-mail address: gcbell01@gwise.louisville.edu, dranish@math.ufl.edu