

# A strange example concerning Property $N_p$

Elena Rubei

## Abstract

We exhibit an example of a line bundle  $M$  on a smooth complex projective variety  $Y$  s.t.  $M$  satisfies Property  $N_p$  for some  $p$ , the  $p$ -module of a minimal resolution of the ideal of the embedding of  $Y$  by  $M$  is nonzero and  $M^2$  does not satisfy Property  $N_p$ .

Let  $M$  be a very ample line bundle on a smooth complex projective variety  $Y$  and let  $\varphi_M : Y \rightarrow \mathbf{P}(H^0(Y, M)^*)$  be the map associated to  $M$ . We recall the definition of Property  $N_p$  of Green-Lazarsfeld (see [Gr1], [Gr2] and also [G-L], [Gr3]):

*let  $Y$  be a smooth complex projective variety and let  $L$  be a very ample line bundle on  $Y$  defining an embedding  $\varphi_L : Y \hookrightarrow \mathbf{P} = \mathbf{P}(H^0(Y, L)^*)$ ; set  $S = S(L) = \text{Sym}^* H^0(L)$ , the homogeneous coordinate ring of the projective space  $\mathbf{P}$ , and consider the graded  $S$ -module  $G = G(L) = \bigoplus_n H^0(Y, L^n)$ ; let  $E_*$*

$$0 \longrightarrow E_n \longrightarrow E_{n-1} \longrightarrow \dots \longrightarrow E_0 \longrightarrow G \longrightarrow 0$$

*be a minimal graded free resolution of  $G$ ; let  $E_p = \bigoplus_{q \geq 0} B_{p,q} \otimes S(-q)$ ; the line bundle  $L$  satisfies Property  $N_p$  ( $p \in \mathbf{N}$ ) if and only if*

$$\begin{aligned} B_{0,0} &= \mathbf{C} \\ B_{0,q} &= 0 \text{ for } q > 0 \\ B_{i,q} &= 0 \text{ for } q \neq i + 1 \text{ and } 1 \leq i \leq p. \end{aligned}$$

(Thus  $L$  satisfies Property  $N_0$  if and only if  $Y \subset \mathbf{P}(H^0(L)^*)$  is projectively normal, i.e.  $L$  is normally generated;  $L$  satisfies Property  $N_1$  if and only if  $L$  satisfies Property  $N_0$  and the homogeneous ideal  $I$  of  $Y \subset \mathbf{P}(H^0(L)^*)$  is generated by quadrics;  $L$  satisfies Property  $N_2$  if and only if  $L$  satisfies Property  $N_1$  and the module of syzygies among quadratic generators  $Q_i \in I$  is spanned by relations of the form  $\sum L_i Q_i = 0$ , where  $L_i$  are linear polynomials; and so on.)

---

**Address:** Elena Rubei, Dipartimento di Matematica "U. Dini", via Morgagni 67/A 50134 Firenze, Italia

**E-mail address:** rubei@math.unifi.it

**2000 Mathematical Subject Classification:** 14C20, 13D02.

In [Ru] we considered the problem to see “how Property  $N_p$  propagates through powers”: let  $M$  be a line bundle on a smooth complex projective variety  $Y$  and suppose  $M$  satisfies Property  $N_p$ ; let  $s \in \mathbf{N}$ ; we wondered for which  $k$  the line bundle  $M^s$  satisfies Property  $N_k$ . We proved: *Let  $Y$  be a smooth complex projective variety and  $M$  a line bundle on  $Y$ ; if  $M$  satisfies Property  $N_p$  then  $M^s$  satisfies Property  $N_p$  if  $s \geq p$ .*

Besides, despite there are several theorems suggesting that there is a “propagation property” for Property  $N_p$ , we observed that surprisingly it is not true that if  $M$  satisfies Property  $N_p$  then any power of  $M$  satisfies Property  $N_p$ ; the counterexample we exhibited was the following: the line bundle  $M = \mathcal{O}_{\mathbf{P}^2}(2)$  satisfies Property  $N_p$  for every  $p$  (see the paper [O-P] and the references given there), but, by Ottaviani-Paoletti’s Theorem in [O-P], for instance the bundle  $M^2 = \mathcal{O}_{\mathbf{P}^2}(4)$  doesn’t satisfy Property  $N_{10}$ .

The example can be considered not convincing by someone because the module  $E_{10}$  of a minimal resolution of  $G(\mathcal{O}_{\mathbf{P}^2}(2))$  on  $S(\mathcal{O}_{\mathbf{P}^2}(2))$  is 0.

The product of projective spaces furnishes the following more convincing example.

**Example.** Let  $r \in \mathbf{N}$  and  $r \geq 1$ .

The line bundle  $\mathcal{O}_{P^r \times P^1}(1, 2)$  satisfies Property  $N_p \forall p$  (see [O-P], Theorem 3.6).

On the contrary  $\mathcal{O}_{P^r \times P^1}(2, 4)$  does not satisfy Property  $N_{10}$ , in fact:

$\mathcal{O}_{P^1 \times P^1}(2, 4)$  does not satisfy Property  $N_{10}$  by the following Gallego-Purnapranja’s Theorem

**Theorem (Gallego–Purnapranja [G-P]).** *Let  $a, b \geq 2$ . The line bundle  $\mathcal{O}_{\mathbf{P}^1 \times \mathbf{P}^1}(a, b)$  satisfies Property  $N_p$  if and only if  $p \leq 2a + 2b - 3$ .*

and the fact that  $\mathcal{O}_{P^1 \times P^1}(2, 4)$  does not satisfy Property  $N_{10}$  implies  $\mathcal{O}_{P^r \times P^1}(2, 4)$  does not satisfy Property  $N_{10}$ , by Remark in Section 2 of [Gr2] (precisely: write  $\mathbf{P}^r = \mathbf{P}(V^*)$  and write the module  $E_p$  of a minimal resolution of  $G(\mathcal{O}_{P^r \times P^1}(2, 4))$  as  $E_p = \bigoplus_{q \geq 0} B_{p,q} \otimes S(-q)$ , as in the definition of Property  $N_p$ ; we have that  $B_{p,q}$  are representations of  $GL(V)$ ; thus, if they are nonzero for  $r = n$  then they are nonzero also for  $r > n$ ) (analogously to the proof of Prop. 1.8 in [O-P]).

If  $r$  is sufficiently great, we have that the 10-th module of a minimal resolution of  $G(\mathcal{O}_{P^r \times P^1}(1, 2))$  is nonzero (in fact, if  $X$  is a projective variety in  $\mathbf{P}^n$  and  $\dim X = m$  with  $m < n - 1$ , then the length  $l$  of a minimal resolution of the ideal  $\mathcal{I}_X$  of  $X$  is  $\geq n - m$ : let  $t$  be s.t.  $H^m(\mathcal{O}_X(t)) \neq 0$ ; thus  $H^{m+1}(\mathcal{I}_X(t)) \neq 0$ ; then by breaking the twist by  $t$  of a minimal resolution of  $\mathcal{I}_X$  into short exact sequences, we get  $l \geq m - n$ ).

**Acknowledgements.** I thank G. Ottaviani for a helpful discussion.

## References

- [G-P] F.J. Gallego, B.P. Purnaprajna *Some results on rational surfaces and Fano varieties* preprint math.AG/0001107
- [Gr1] M. Green *Koszul cohomology and the geometry of projective varieties I* J. Differ. Geom. **20**, 125-171 (1984)
- [Gr2] M. Green *Koszul cohomology and the geometry of projective varieties II* J. Differ. Geom. **20**, 279-289 (1984)
- [Gr3] M. Green *Koszul cohomology and geometry*, in: M. Cornalba et al. (eds), Lectures on Riemann Surfaces, World Scientific Press (1989)
- [G-L] M. Green, R. Lazarsfeld *On the projective normality of complete linear series on an algebraic curve* Invent. math. **83**, 73-90 (1986)
- [O-P] G. Ottaviani, R. Paoletti *Syzygies of Veronese embeddings* preprint math.AG/9811131 to appear in Compositio Mathematica
- [Ru] E. Rubei, *A note on Property  $N_p$* , Manuscripta Mathematica 101 (2000), 449-455.