## THIN GROUPS OF FRACTIONS

### Patrick DEHORNOY

ABSTRACT. A number of properties of spherical Artin groups extend to Garside groups, defined as the groups of fractions of monoids where least common multiples exist, there is no nontrivial unit, and some additional finiteness conditions are satisfied [9]. Here we investigate a wider class of groups of fractions, called *thin*, which are those associated with monoids where minimal common multiples exist, but they are not necessarily unique. Also, we allow units in the involved monoids. The main results are that all thin groups of fractions satisfy a quadratic isoperimetric inequality, and that, under some additional hypotheses, they admit an automatic structure.

#### 1. Introduction

The algebraic theory of braids, as developed in [15] and [14], relies on the existence of Garside's fundamental elements  $\Delta_n$ : for each n, the braid  $\Delta_n$  is an element of the monoid  $B_n^+$  which is a least common multiple of the standard generators  $\sigma_i$ , and the main technical point is that the left divisors of  $\Delta_n$  in  $B_n^+$  coincide with its right divisors. Most of the results established for Artin's braid groups  $B_n$  have been extended to more general groups: spherical Artin groups [12, 4, 5], Garside groups in the sense of [11], and, subsequently, of [9] (also called small or thin Gaussian groups in [11] and [17], respectively). All the considered groups are groups of fractions of monoids in which least common multiples exist, and, in each case, a key rôle is played by some element  $\Delta$  of the associated monoid that satisfies most of the technical properties of Garside's braids  $\Delta_n$ . In particular, it is proved in [9] that the greedy normal form of braids [1, 14, 13] extends to all Garside groups, and that it gives rise to a bi-automatic structure.

The aim of this paper is to consider groups of fractions of monoids where common multiples exist, but *least* common multiples need not exist. In this case, no counterpart of the element  $\Delta$  need exist in general, but a number of properties involving the divisors of  $\Delta$  can still be established when considering subsets of the monoid that are closed under convenient operations. In this way, one can define an extended notion of normal form, which coincides with the greedy normal form when

<sup>2000</sup> Mathematics Subject Classification. 20M05, 20F36, 05C25.

Key words and phrases. Artin group, Gaussian group, Garside group, Garside element, normal form, automatic group.

least common multiples exist. The price to pay for the lack of lcm is a possible non-uniqueness. However, we shall see that, at least in good cases, this non necessarily unique normal form is still associated with an automatic structure. We shall be mainly interested in the *thin* case, defined as the case when a *finite* set of generators with good closure properties exists. The main results we prove are:

Theorem 1.1 (Prop. 4.5). Every thin group of fractions satisfies a quadratic isoperimetric inequality.

Theorem 1.2 (Prop. 9.10). Assume that G is the group of fractions of a thin cancellative monoid M that admits a Garside element  $\Delta$  such that all  $\Delta$ -normal forms in M have the same length. Then G is an automatic group.

These results apply to all thin Gaussian groups, which are the Garside groups of [9], hence in particular to all spherical Artin groups, for which the properties were already known, but they also apply to groups of a completely different flavour, as some simple examples will show.

The possible interest of our approach is double. On the one hand, as we mentioned, new groups are eligible. On the other hand, we hope that extending classical results may help to understand them better and to capture the really important hypotheses: studying Gaussian groups showed in [9] that the fact that Garside's element  $\Delta_n$  is a least common multiple of the generators  $\sigma_i$  of  $B_n$  is useless, and, so, using such a fact gives slightly misleading arguments. Similarly, the approach developed in the current paper shows that a clear distinction should be made between the family of all divisors of  $\Delta$  (the "simple" elements), and a smaller subfamily (the "primitive" elements) which contains the real information: the latter can be extended to the more general framework, while the former cannot, at least if we use the classical definition. This leads us here to an alternative, hopefully improved definition of a simple element. In the current framework, the proof that the Garside groups are automatic reduces to a small number of technical lemmas, each of which is specially easy when lcm's exist (Lemmas 7.8, 7.12, and 7.13).

The organization of the paper is as follows. In Sec. 2, we introduce the notion of a spanning subset of a monoid, which is a generating set satisfying some additional closure property. Then a thin monoid is defined to be a monoid that admits a finite spanning subset. In Sec. 3, we introduce the weaker notion of a quasi-spanning set so as to allow nontrivial units. In Sec. 4, we define thin groups of fractions as those associated with a thin Ore monoid, and we prove Theorem 1.1. In Sec. 5, we show that every thin monoid admits a minimal spanning subset. In Sec. 6, we introduce the notion of an S-simple element associated with a spanning subset S, which is a counterpart for the notion of divisor of  $\Delta_n$  in braid monoids. In Sec. 7, we use S-simple elements to construct a counterpart to the greedy normal form of braids. In Sec. 8, we introduce Garside elements, which are convenient generalizations for the fundamental braids  $\Delta_n$ . Finally, in Sec. 9, we prove Theorem 1.2.

# 2. Spanning subsets of a monoid

We consider in the sequel cancellative monoids. Most of the results until Sec. 4 are valid if we only assume left cancellativity. If M is a monoid, we say that an element of M is a left (right) unit if it admits a left (right) inverse; provided M is left or right cancellative, uv = 1 implies uvu = u and v = vuv, hence vu = 1, so left and right units coincide, and they form a subgroup of M that will be denoted

by  $M^*$ . For  $u \in M^*$ , we denote by  $u^{-1}$  the (unique) left and right inverse of u. As multiplying by a unit on the right is often considered in the sequel, we introduce a notation:

DEFINITION 2.1. Assume that M is a (left) cancellative monoid. For  $x, x' \in M$ , we say that  $x \simeq x'$  holds if we have x' = xu for some u in  $M^*$ . We say that a subset S of M is quasi-finite if it contains finitely many  $\simeq$ -classes.

The relation  $\simeq$  is an equivalence relation which is compatible with left multiplication. For  $S\subseteq M$ , the set  $SM^*$  is the smallest  $\simeq$ -saturated subset of M including S.

If M is a monoid, and x, y lie in M, we say that x is a left divisor of y, written  $x \leq y$ , if y = xz holds for some z. If, in addition, z is not a unit, we say that x is a proper left divisor of y, and write  $x \prec y$ . We have the symmetric notion of a right divisor, but, as left divisors play a distinguished rôle, we shall usually simply say "divisor" for "left divisor". The set of all (left) divisors of x is denoted Div(x). If x is a (left) divisor of y, we equivalently say that y is a (right) multiple of x. Notice that z is compatible with z and z in the sense that z is z (resp. z is equivalent to z is z if z in the sense that z is z in the sense that z in the sense that z is z in the sense that z in the sense that z is z in the sense that z in the sense that z is z in the sense that z in the sense that z is z in the sense that z in the sense that z in the sense that z is z in the sense that z in the sense that z is z in the sense that z is z in the sense that z is z in the sense that z is z in the sense that z in the sense that z is z in the sense that z in the sense that z is z in the sense that z in the sense that z is z in the sense that z in the sense that z is z in the sense that z is z in the sense that z in the sense that z is z.

The central notion of this paper is that of a spanning subset of a monoid; it is defined by means of some closure properties involving left divisors:

DEFINITION 2.2. Assume that M is a (left) cancellative monoid, and S is a subset of M. We say that S spans M if S generates M, it contains 1, it is  $\simeq$ -saturated, i.e.,  $SM^* \subseteq S$  holds, and

(2.1) If we have 
$$x \leq z$$
 and  $y \leq z$  with  $x, y \in S$ , then there exist  $x', y'$  in  $S$  satisfying  $xy' = yx' \leq z$ .

We say that M is thin (resp. quasi-thin) if it admits a finite (resp. quasi-finite) spanning subset.

By definition, a thin monoid is finitely generated, but the converse need not be true, as a spanning subset is more than a generating subset. Note that the converse of Implication (2.1) always holds:  $xy' = yx' \leq z$  trivially implies  $x \leq z$  and  $y \leq z$ . Spanning subsets always exist: if M is a monoid, then M is a spanning subset of itself. Actually, we shall be mainly interested in the case when small spanning subsets exist, so, typically, in the thin case. For a monoid with no nontrivial unit, or, more generally, with finitely many units, being quasi-thin is equivalent to being thin.

EXAMPLE 2.3. Let M be a spherical Artin monoid, i.e., one associated with a finite Coxeter group W. For  $x, y \in M$ , define  $x \setminus y$  to be the unique element z such that xz is the least common multiple of x and y. Then the closure of the standard generators  $\sigma_i$  under the operation  $\setminus$  is a finite spanning subset of M, in one-to-one correspondence with some subset of W [11]. So spherical Artin monoids are thin.

More generally, if M is a Gaussian monoid in the sense of [11, 9], *i.e.*, a cancellative monoid in which least common multiples exist and division has no infinite descending chain, then the closure of the set of atoms under operation  $\setminus$  is a minimal spanning subset of M. Thus the monoid M is thin if and only if the




FIGURE 2.1. Spanning subset

latter closure is finite, *i.e.*, if M is thin in the sense of [17] (or small in the sense of [11]), so the terminologies are compatible.<sup>1</sup>

Lemma 2.4. Assume that M is a (left) cancellative monoid, and S is subset of M. Then Condition (2.1) for S is equivalent to

(2.2) If we have 
$$xy'' = yx''$$
 with  $x, y \in S$ , then there exist  $x', y'$  in  $S$  and  $z$  in  $M$  satisfying  $xy' = yx', x'' = x'z$ , and  $y'' = y'z$  (Fig. 2.1).

PROOF. It is clear that (2.2) implies (2.1); conversely, assuming xy'' = yx'', Condition (2.1) implies that there exist x', y' in S and z in M satisfying xy' = yx', xy'' = xy'z, and yx'' = yx'z, hence x'' = x'z and y'' = y'z if we can cancel x and y on the left.

If S, T are subsets of a monoid M, we put  $ST = \{xy : x \in S, y \in T\}$ . In particular,  $S^p$  is the set of all elements that can be written as  $x_1 \cdots x_p$  with  $x_1, \ldots, x_p \in S$ . Notice that  $1 \in S$  implies  $S \subseteq S^p$  for  $p \ge 2$ . We put  $S^0 = \{1\}$ .

LEMMA 2.5. Assume that M is a monoid, and S is a subset of M satisfying Condition (2.2). Then, if we have xy'' = yx'' with  $x \in S^p$  and  $y \in S^q$ , there exist an element z of M and two sequences  $x_{i,j}$ ,  $y_{i,j}$ ,  $0 \le i \le p$ ,  $0 \le j \le q$ , of elements of S satisfying  $x_{i,j-1}y_{i,j} = y_{i-1,j}x_{i,j}$  for all i, j, and  $x = \prod x_{i,0}$ ,  $y = \prod y_{0,j}$ ,  $x'' = \prod x_{i,q} z$ , and  $y'' = \prod y_{p,j} z$ . So, in particular, there exist x' in  $S^p$ , y' in  $S^q$  and z in M satisfying xy' = yx', x'' = x'z, and y'' = y'z.

PROOF. (Fig. 2.2) First, the condition is sufficient, as the local equalities  $x_{i,j}y_{i,j+1}=y_{i,j}x_{i+1,j}$  imply  $\prod_i x_{i,0}\prod_j y_{p,j}=\prod_j y_{0,j}\prod_i x_{i,q}$ , hence xy''=yx'' when x,y,x'',y'' have the above specified values.

We prove now that the condition is necessary. The result is trivial for p=0 or q=0. Indeed, p=0 means x=1: then the hypothesis  $y\in S^q$  allows us to write  $y=\prod_j y_{0,j}$  with  $y_{0,1},\ldots,y_{0,q}\in S$ , and the hypothesis y''=yx'' then gives  $y''=\prod_j y_{0,j}z$  with z=x''. Then we use induction on p+q. By the remark above, the first nontrivial case is p=q=1, and, then, the result is true by Condition (2.2). Assume now  $p+q\geq 3$ , with  $p,q\geq 1$ . Then at least one of p,q is greater than 1. Assume for instance  $q\geq 2$ . Write  $y=y_1y_2$  with  $y_1\in S^{q_1},y_2\in S^{q_2}$  and  $1\leq q_1,q_2< q$ . Applying the induction hypothesis to  $x\in S^p,y_1\in S^{q_1}$  and  $xy''=y_1(y_2x'')$  gives  $z_1$  in M and  $x_{i,j},y_{i,j},0\leq i\leq p,0\leq j\leq q_1$  in S satisfying  $x_{i,j-1}y_{i,j}=y_{i-1,j}x_{i,j},y_2x''=x_1z_1,y''=y_1'z_1$  with

$$x = \prod_{1}^{p} x_{i,0}, \quad y_1 = \prod_{1}^{q_1} y_{0,j}, \quad x_1 = \prod_{1}^{p} x_{i,q_1}, \quad y_1' = \prod_{1}^{q_1} y_{p,j}.$$

<sup>&</sup>lt;sup>1</sup>In order to uniformize terminology with other authors, we use *Garside monoid* as a synonym for thin Gaussian monoid, and *Garside group* for the group of fractions of a Garside monoid [9].




FIGURE 2.2. Power of a spanning subset

Applying the induction hypothesis to  $x_1 \in S^p$ ,  $y_2 \in S^{q_2}$  and  $x_1z_1 = y_2x''$  gives z in M and  $x_{i,j}, y_{i,j}, 0 \le i \le p$ ,  $q_1 < q_2$  in S satisfying  $x_{i,j-1}y_{i,j} = y_{i-1,j}x_{i,j}$ ,  $x'' = x'z, z_1 = y_2'z$  with

$$x = \prod_{1}^{p} x_{i,0}, \quad y_2 = \prod_{q_1+1}^{q_2} y_{0,j}, \quad x' = \prod_{1}^{p} x_{i,q}, \quad y'_2 = \prod_{q_1+1}^{q_2} y_{p,j}.$$

Putting  $y' = y_1'y_2'$  gives the expected result.

Finally, with the previous notation, put  $x' = \prod_i x_{i,q}$  and  $y' = \prod_j y_{p,j}$ . By construction, x' lies in  $S^p$ , y' lies in  $S^q$ , and we have xy' = yx', x'' = x'z, and y'' = y'z.

Applying Lemma 2.5 with q = p, we obtain

Proposition 2.6. Assume that M is a (left) cancellative monoid. If S spans M, so does  $S^p$  for every positive p.

Cancellativity is not used in the proof of Lemma 2.5, so, at the expense of using (2.2) instead of (2.1) in the definition of a spanning subset, we could state Prop. 2.6 for a general monoid.

PROPOSITION 2.7. Assume that M is a (left) cancellative monoid, and S spans M. Then every right divisor of an element of S lies in S, and we have  $M^*S \subseteq S$ .

PROOF. Assume  $y=xz\in S$ . As S generates M, we have  $x\in S^p$  for some p, so, applying Lemma 2.5 to the equality  $x\cdot z=y\cdot 1$ , we find x in  $S^p$ , y' in S, and z' in M satisfying xy'=yx', z=y'z', and 1=x'z'. The latter relation shows that x' and z' are units and z=y'z' then implies  $z\in SM^*$ , hence  $z\in S$  as S is supposed to be closed under right multiplication by a unit.

Assume  $x \in S$  and  $u \in M^*$ . Then we have  $x = u^{-1}(ux)$ , so ux is a right divisor of x, and, by the previous result, it belongs to S.

One of the interests of spanning subsets is that they completely determine the monoid in the sense below. In the sequel, if S is a set (of letters), and R is a set of relations over S, *i.e.*, of equalities of the form  $x_1 \cdots x_p = y_1 \cdots y_q$  with  $x_1, \ldots, y_q \in S$ , we denote by  $\langle S; R \rangle^+$  the monoid so presented, and by  $\langle S; R \rangle$  the group with the same presentation.




FIGURE 2.3. Quadratic isoperimetric inequality

DEFINITION 2.8. Assume that M is a (left) cancellative monoid, and S spans M. We denote by  $R_S$  the set of all relations xy' = yx' with  $x, y, x', y' \in S$ .

PROPOSITION 2.9. Assume that M is a (left) cancellative monoid, and S spans M. Then  $\langle S; R_S \rangle^+$  is a presentation of M, and every equality  $x_1 \cdots x_p = y_1 \cdots y_q$  with  $x_1, \ldots, y_q \in S$  can be proved using  $O((p+q)^2)$  relations of  $R_S$ .

PROOF. (Fig. 2.3) The set S generates M by definition. Assume  $x_1 \cdots x_p = y_1 \cdots y_q$  with  $x_1, \ldots, y_q \in S$ . By Lemma 2.5, there exist  $x_1', \ldots, x_p', y_1', \ldots, y_q' \in S$  and  $z \in M$  satisfying

$$x_1 \cdots x_p y_1' \cdots y_q' = y_1 \cdots y_q x_1' \cdots x_p', \quad x_1' \cdots x_p' z = 1, \text{ and } y_1' \cdots y_q' z = 1,$$

and, moreover, the first equality can be established using pq relations in  $R_S$ . As for the other ones, we know by Prop. 2.7 that each of the elements  $x'_i \cdots x'_p z$  and  $y'_j \cdots y'_q z$  belongs to S, and, therefore, the equality  $x'_1 \cdots x'_p z = 1$  can be established using p relations of  $R_S$  (of the special form xy = 1), and, similarly,  $y'_1 \cdots y'_q z = 1$  can be established using q relations of  $R_S$ . So, finally, the equality  $x_1 \cdots x_p = y_1 \cdots y_q$  can be established using at most  $(p+q)^2/4 + (p+q)$  relations of  $R_S$ .

Applying the previous result to the case of a finite spanning subset, we obtain:

Proposition 2.10. Every thin cancellative monoid satisfies a quadratic isoperimetric inequality.

In order to construct new examples of thin monoids, Prop. 2.9 suggests that we consider presentations where the relations are of the form xy' = yx', i.e., involve words of length 2 at most. Every monoid admits a presentation of this type, and the question arises of recognizing spanning subsets. Typically, if  $\langle S;R\rangle^+$  is a presentation of the type above for a monoid M, there is no obvious reason why S should span M, as some equalities in M may follow from the relations of R but not decompose into such relations using the scheme of Fig. 2.3. In particular, there is no reason why the equality  $R = R_S$  should hold. Here, we shall refer to [10], where the notion of a complete presentation is defined. The idea is that a presentation is complete if the relations have no hidden consequence, i.e., if enough relations have been displayed to avoid any such hidden consequence. Then the result is that, if  $\langle S;R\rangle^+$  is a complete presentation for the monoid M, then a sufficient condition for S to span M is that each relation in R has length 2 at most.




FIGURE 2.4. Characteristic graph associated with a spanning subset

EXAMPLE 2.11. With this method, we can exhibit thin monoids that do not resemble the Gaussian monoids of Example 2.3, namely monoids where least common multiples do not exist. The following three examples are typical, and they will be considered throughout the paper:

$$M_1 = \langle a, b \; ; \; a^2 = b^2, ab = ba \rangle^+,$$
  
 $M_2 = \langle a, b, c \; ; \; a^2 = b^2 = c^2, ab = bc = ca, ac = ba = cb \rangle^+,$   
 $M_3 = \langle a, b, c \; ; \; ac = ca = b^2, ab = bc, cb = ba \rangle^+.$ 

Applying the criterion of [10], one checks that the above presentations are complete, and that, in each case, the involved set of generators completed with 1 is a spanning subset. Thus the monoids  $M_i$  are thin. Moreover, there is no relation xy = xy' or yx = y'x with  $y \neq y'$  in the above presentations, so, according to [10] again, the monoids  $M_i$  are cancellative.

If S spans a monoid M, then M is determined by S and  $R_S$ . Especially when the considered set S is finite, i.e., in the thin case, it is natural to introduce the subgraph of the Cayley graph of M displaying the relations of  $R_S$ : by the remarks above, such a (finite) graph completely determines the monoid. In the Gaussian case, i.e., when least common multiples exist, the graph is a lattice, in the sense that any two vertices admit a unique immediate common successor. In the general case, this need not be true. For instance, we display in Fig. 2.4 the graphs associated with the braid monoid  $B_3^+$  and the spanning subset  $\{1, \sigma_1, \sigma_2, \sigma_1\sigma_2, \sigma_2\sigma_1\}$ , and with the thin monoids  $M_1, M_2, M_3$  of Example 2.11.

## 3. Quasi-spanning subsets

Many results for the thin case extend to the quasi-thin case, and, to deal with the latter, it is convenient to introduce the notion of a quasi-spanning subset of a monoid, of which a typical example is a  $\simeq$ -selector through a spanning set, *i.e.*, a subset that picks one element in each equivalence class.

DEFINITION 3.1. Assume that M is a (left) cancellative monoid, and S is a subset of M. We say that S quasi-spans M if  $SM^*$  spans M.

By definition, a spanning subset is quasi-spanning, and both notions coincide if there is no nontrivial unit.

PROPOSITION 3.2. Assume that M is a (left) cancellative monoid, and S spans M. Then every  $\simeq$ -selector through S quasi-spans M. Conversely, if S is a minimal quasi-spanning subset of M, then S is a  $\simeq$ -selector.

PROOF. If  $\Sigma$  is a  $\simeq$ -selector through S, we have  $\Sigma M^* = S$ , so  $\Sigma$  is quasi-spanning. On the other hand, assume that S is a minimal quasi-spanning subset of M. Assume  $y \simeq x$ , with  $x, y \in S$  and  $x \neq y$ . Then  $(S - \{y\})M^*$  is equal to  $SM^*$ , and, therefore,  $S - \{y\}$  quasi-spans M, contradicting the minimality of S.  $\square$ 

Corollary 3.3. A cancellative monoid is quasi-thin if and only if it admits a finite quasi-spanning subset.

Lemma 3.4. Assume that M is a (left) cancellative monoid, and S is a subset of M. Then S quasi-spans M if and only if S contains 1,  $SM^*$  generates M,  $M^*S \subseteq SM^*$  holds, and

(3.1) If we have 
$$x \leq z$$
 and  $y \leq z$  with  $x, y \in S$ , then there exist  $x', y'$  in  $S$  satisfying  $xy' \simeq yx' \leq z$ .

PROOF. Assume that S satisfies the above conditions. Then  $SM^*$  contains 1, it generates M by hypothesis, and it is  $\simeq$ -saturated by construction. Assume  $xu \leq z$  and  $yv \leq z$  with  $x, y \in S$ , and  $u, v \in M^*$ . Then we have  $x \leq z$  and  $y \leq z$ , so, by (3.1), there exist x', y' in S satisfying  $xy' \simeq yx' \leq z$ , say  $xy' = yx'w \leq z$  with  $w \in M^*$ . Then we have also  $(xu)(u^{-1}y') = (yv)(v^{-1}x'w) \leq z$ , and the elements  $u^{-1}y'$  and  $v^{-1}x'w$  belong to  $M^*SM^*$ , hence to  $SM^*$ , which therefore satisfies Condition (2.1), and spans M.

Conversely, assume that S quasi-spans M. Then  $1 \in M^*$  implies  $S \subseteq SM^*$ , and restricting Condition (2.1) for  $SM^*$  to S yields Condition (3.1). Moreover, we have  $M^*SM^* \subseteq SM^*$  by Prop. 2.7, hence, a fortiori,  $M^*S \subseteq SM^*$ . So all conditions of Lemma 3.4 are satisfied.

If S quasi-spans a monoid M, then we have  $M^*S \subseteq SM^*$  by Lemma 3.4, and a straightforward induction then implies

$$(3.2) (SM^*)^n = S^n M^*$$

for every positive n. As  $SM^*$  is supposed to generate M, it follows that every element of M admits a decomposition of the form  $x = x_1 \cdots x_n u$  with  $x_1, \ldots, x_n \in S$  and  $u \in M^*$ .

Proposition 3.5. Assume that M is a (left) cancellative monoid and S quasispans M.

- (i) The subset  $S^p$  quasi-spans M for every positive p.
- (ii) If we have  $x \leq z$  and  $y \leq z$  with  $x \in S^p$  and  $y \in S^q$ , then there exist  $x' \in S^p$  and  $y' \in S^q$  satisfying  $xy' \simeq yx' \leq z$ .
- (iii) If we have  $x_i \leq z$  with  $x_i \in S$  for  $1 \leq i \leq n$ , then there exists x in  $S^n$  satisfying  $x_i \leq x \leq z$  for  $1 \leq i \leq n$ .

PROOF. (i) By Lemma 3.4,  $SM^*$  spans M, so, by Prop. 2.6,  $(SM^*)^p$  spans M as well for every positive p. By (3.2), the latter is  $S^pM^*$ , and, by Lemma 3.4 again,  $S^pM^*$  spanning M implies  $S^p$  quasi-spanning M.

(ii) Applying Lemma 2.5 to the spanning subset  $SM^*$  of M, we obtain x'' in  $(SM^*)^p$  and y'' in  $(SM^*)^q$  satisfying  $xy'' = yx'' \leq z$ . By (3.2), we have  $(SM^*)^p = xy'' \leq z$ .

 $S^pM^*$  and  $(SM^*)^q = S^qM^*$ , so we deduce that there exist x' in  $S^p$ , y' in  $S^q$ , and u, v in  $M^*$  satisfying x'' = x'u and y'' = y'v, hence  $xy' \simeq yx' \preceq z$ .

(iii) We use induction on  $n \ge 1$ . For n = 1, we take  $x = x_1$ . Assume  $n \ge 2$ . By induction hypothesis, some y in  $S^{n-1}$  satisfies  $x_i \le y \le z$  for  $i \le n-1$ . Applying (ii) to  $x_n$  and y, we obtain x' in S and y' in  $S^{n-1}$  satisfying  $x_n y' \simeq y x' \le z$ . Putting x = y x' gives the result.

PROPOSITION 3.6. Assume that M is a (left) cancellative monoid and S is a minimal quasi-spanning subset of M. Then, for every pair (x, u) in  $S \times M^*$ , there exists a unique pair (x', u') in  $S \times M^*$  satisfying ux = x'u'. The mapping  $x \mapsto x'$  defines an action of the group  $M^*$  on S. If x is fixed under this action, the mapping  $u \mapsto u'$  is an endomorphism of  $M^*$  for every x.

PROOF. By Lemma 3.4, we have  $M^*S \subseteq SM^*$ , so  $ux \in SM^*$ , i.e., there exist x' in S and u' in  $M^*$  satisfying ux = x'u'. The uniqueness of x' follows from Prop. 3.2, and that of u' then follows from left cancellativity. Writing ux for x', we have  $1x \simeq x$  for every x in S, and, therefore 1x = x, and  $ux = u^vx = u^vx$ , hence  $ux = u^vx = u^vx$ .

Assume now  ${}^{u}x=x$ . Writing  $u^{x}$  for u', we find  $(uv)x=uxv^{x}=xu^{x}v^{x}$ , hence  $(uv)^{x}=u^{x}v^{x}$ .

The previous result gives a way for constructing quasi-thin monoids with a prescribed group of units. Assume that M is a cancellative monoid with no non-trivial unit, S spans M, and G is a group with a left action on M that preserves S globally. Consider the semi-direct product  $M \rtimes G$  where (x,u)(y,v) is defined to be (xu(y),uv). The set of units in  $M \rtimes G$  is  $\{1\} \times G$ , and the set  $S \times \{1\}$  is a quasi-spanning subset of  $M \rtimes G$ . More generally, instead of a semidirect product, we could also use a crossed product as in [17].

In the same way as a spanning subset determines a monoid, a quasi-spanning subset together with the units determine a monoid.

DEFINITION 3.7. Assume that M is a (left) cancellative monoid, and S quasispans M. We denote by  $R_S^*$  the set consisting of

- (i) all relations xy' = yx'u with  $x, y, x', y' \in S$  and  $u \in M^*$ ,
- (ii) all relations ux = x'u' with  $x, x' \in S$  and  $u, u' \in M^*$ ,
- (iii) all relations uv = w with  $u, v, w \in M^*$ .

PROPOSITION 3.8. Assume that M is a (left) cancellative monoid, and S quasispans M. Then  $(S \cup M^*; R_S^*)^+$  is a presentation of M, and every equality  $x_1 \cdots x_p = y_1 \cdots y_q$  with  $x_1, \ldots, y_q \in S$  can be proved using  $O((p+q)^2)$  relations of  $R_S^*$ .

PROOF. By Lemma 3.4, the set  $SM^*$  spans M, so, by Prop. 2.9,  $\langle SM^*; R_{SM^*} \rangle^+$  is a presentation of M, and every equality  $x_1 \cdots x_p = y_1 \cdots y_q$  with  $x_1, \ldots, y_q \in SM^*$  can be established using at most  $O((p+q)^2)$  relations in  $R_{SM^*}$ . So, it suffices to prove that every relation in  $R_{SM^*}$  can be decomposed into a uniformly bounded number of relations in  $R_S^*$ . By construction, every element in  $SM^*$  can be expressed as xu with  $x \in S$  and  $u \in M^*$ . So assume xuy'v' = yvx'u' with  $x, y, x', y' \in S$  and  $u, v, u', v' \in M^*$ . There exist x'', y'' in S and u'', v'' in  $M^*$  satisfying the type (ii) relations uy' = y''u'' and vx' = x''v'': then xuy'v' = yvx'u' implies the type (i) relation xy''w = yx'' with  $w = u''v'u'^{-1}v''^{-1}$ , and the latter equality follows from three type (iii) relations (Fig. 3.1). Thus every relation of  $R_{SM^*}$  can be decomposed into at most six relations in  $R_S^*$ .




Figure 3.1. Decomposition of a relation

Instead of using all units in the presentation, we can replace  $M^*$  with any set that generates it (as a monoid). We obtain:

COROLLARY 3.9. Assume that M is a (left) cancellative monoid, S quasispans M, and  $\langle S'; R' \rangle^+$  is a presentation of  $M^*$  as a monoid. Let R consist of (i) all relations  $xy' = yx'u_1 \cdots u_p$  with  $x, y, x', y' \in S, u_1, \ldots, u_p \in S'$ , (ii) all relations  $ux = x'u'_1 \cdots u'_p$  with  $x, x' \in S, u, u'_1, \ldots, u'_p \in S'$ . Then  $\langle S \cup S'; R \cup R' \rangle^+$  is a presentation of M.

The question of whether an isoperimetric inequality is satisfied when the previous presentation is finite is open: even if we assume that the presentation of the group  $M^*$  satisfies such a condition, there is no easy way to conclude for M as we know nothing about the elements denoted u'', v'', and w in Fig. 3.1.

#### 4. Thin groups of fractions

By a well known result of Ore [7], a cancellative monoid M embeds in a group of (right) fractions if and only if any two elements of M admit at least one common multiple. A monoid satisfying such conditions will be called a  $Ore\ monoid$  in the sequel. It is then natural to consider those groups of fractions that are associated with thin monoids:

DEFINITION 4.1. We say that a group G is a *thin* (resp. *quasi-thin*) group of fractions if there exists a thin (resp. quasi-thin) Ore monoid M such that G is the group of fractions of M.

Thus, the braid groups  $B_n$ , the spherical Artin groups, and, more generally, the Garside groups of [9] are thin groups of fractions. We shall give more examples below.

A nice point is that, when a quasi-spanning subset is known in a monoid M, then it is easy to study the possible existence of common multiples in M. Indeed, we have the following criterion:

Proposition 4.2. Assume that M is a (left) cancellative monoid, and S is a quasi-spanning subset in M. Then any two elements of M admit a common multiple if and only the following condition holds:

(4.1) For all x, y in S, there exist x', y' in S satisfying  $xy' \simeq yx'$ .

PROOF. Assume  $x, y \in S$ . If common multiples always exist in M, there exists z satisfying  $x \leq z$  and  $y \leq z$ , and, therefore, since S is quasi-spanning, there exist x', y' in S satisfying  $xy' \simeq yx' (\leq z)$ . So (4.1) holds.

Conversely, assume (4.1). First, we claim that, for all  $x, y \in SM^*$ , there exist  $x', y' \in SM^*$  satisfying xy' = yx'. Indeed, assume  $x \simeq x_0, y \simeq y_0$  with  $x, y \in S$ . By hypothesis, there exists z satisfying  $x_0 \leq z$  and  $y_0 \leq z$ . Then  $x \leq z$  and  $y \leq z$  hold as well. As  $SM^*$  spans M, we deduce that there exist x', y' in  $SM^*$  satisfying  $xy' = yx'(\leq z)$ .

We prove now that, if x belongs to  $(SM^*)^p$  and y belongs to  $(SM^*)^q$ , then x and y admit a common multiple in M. The result is trivial for p=0 and q=0, and the case p=q=1 has been treated above. Then we use a recurrence on p+q: the principle is to construct a diagram like the one in Fig. 2.2 starting from the left and the top edges. By hypothesis, each small square can be closed, so, inductively, the full diagram can be completed.

It follows that, if M is a cancellative monoid and S is a quasi-spanning subset in M that satisfies (4.1), then M is a Ore monoid, and it embeds in a group of right fractions. Moreover, the latter admits the presentation  $\langle S; R_S \rangle$ , where  $R_S$  is as in Prop. 2.9. In particular, we can state:

COROLLARY 4.3. Assume that M is a thin cancellative monoid and S spans M and satisfies (4.1). Then the group  $\langle S; R_S \rangle$  is a thin group of fractions.

EXAMPLE 4.4. For i = 1, ..., 3, the monoid  $M_i$  of Example 2.11 satisfies Condition (4.1), as we can check on the graph of Fig. 2.4. Thus, any two elements in  $M_i$  admit a common multiple, and, therefore,  $M_i$  embeds in a group of fractions. So the groups

$$G_1 = \langle a, b \; ; \; a^2 = b^2, ab = ba \rangle$$

$$G_2 = \langle a, b, c \; ; \; a^2 = b^2 = c^2, ab = bc = ca, ac = ba = cb \rangle$$

$$G_3 = \langle a, b, c \; ; \; ac = ca = b^2, ab = bc, cb = ba \rangle$$

are thin groups of fractions.

Observe that the monoid  $\langle a, b, c \,; \, ab = bc = ca \rangle^+$  is the Birman-Ko-Lee monoid for the braid group  $B_3$  [2]; then  $M_2$  is the quotient of the latter monoid under the additional relation  $a^2 = b^2$ , and, therefore, the group  $G_2$  is the quotient of  $B_3$  obtained by adding the relation  $\sigma_1^2 = \sigma_2^2$ , thus an intermediate group between  $B_3$  and the symmetric group  $S_3$ .

We can now state our first general result about thin groups of fractions:

Proposition 4.5. Every thin group of fractions satisfies a quadratic isoperimetric inequality.

PROOF. Assume that G is the group of fractions of the Ore monoid M, and S is a finite spanning subset of M. By Prop. 2.9,  $\langle S; R_S \rangle^+$  is a presentation of M, and, therefore,  $\langle S; R_S \rangle$  is a presentation of G, which is finite by construction. Assume that  $x_1^{e_1} \cdots x_n^{e_n} = 1$  holds in G, with  $x_1, \ldots, x_n \in S$  and  $e_1, \ldots, e_n = \pm 1$ . First, an easy induction on m shows that, if the sequence  $(e_1, \ldots, e_n)$  contains p times +1, q times -1, and m subpairs (-1, +1), then there exist  $y_1, \ldots, y_p$  and  $z_1, \ldots, z_q$  in M such that  $x_1^{e_1} \cdots x_n^{e_n} = y_1 \cdots y_p z_q^{-1} \cdots z_1^{-1}$  holds in G, and the equality can be established using m relations of  $R_S$ . Then  $x_1^{e_1} \cdots x_n^{e_n} = 1$  in G implies  $y_1 \cdots y_p = 1$ 

 $z_1 \cdots z_q$  in G, hence in M. By Prop. 2.9 again, the latter equality, if true, can be established using at most  $(p+q)^2$  relations of  $R_S$ . We have  $m \leq pq \leq (p+q)^2/4$  and p+q=n, so, alltogether, we need  $5n^2/4$  relations of  $R_S$  at most to establish  $x_1^{e_1} \cdots x_n^{e_n} = 1$ .

So, for instance, all Garside groups satisfy a quadratic isoperimetric inequality—as was already proved in [11]—and so do the groups  $G_i$  of Example 4.4.

If we consider a Ore monoid M which is quasi-thin only, there is no clear result, as the complexity of the group of units is involved. If the group  $M^*$  is finite, then the presentation of Prop. 3.8 is finite, and it gives rise to a quadratic isoperimetric inequality: however, in this case, M is thin, and Prop. 4.5 applies.

### 5. Primitive elements

The rest of the paper is centered on the possible existence of an automatic structure for a thin group of fractions, a strengthening of the existence of a quadratic isoperimetric inequality. In this section, we show that every thin monoid admits a minimal spanning subset, whose elements will be called primitive. Such primitive elements are themselves connected with atoms, and we begin with some observations about such elements which extend earlier results of [11, 9] so as to allow the existence of nontrivial units.

DEFINITION 5.1. Assume that M is a cancellative monoid. We say that an element x of M is an atom if x is not a unit but x = yz implies that either y or z is a unit. The set of all atoms in M is denoted by  $A_M$ .

Lemma 5.2. If M is a cancellative monoid, then the atoms of M are closed under left and right multiplication by a unit, and, therefore,  $\simeq$ -saturated.

PROOF. Assume  $x \in A_M$  and  $u \in M^*$ . First,  $xu \in M^*$  would imply  $x \in M^*$ , so we have  $xu \notin M^*$ . Then assume xu = yz. We have  $x = y(zu^{-1})$ , hence either y or  $zu^{-1}$  is a unit, and, in the latter case, so z. So xu is an atom. The case of left multiplication by a unit is similar.

Definition 5.3. Assume that M is a monoid. For  $x \in M$ , we put  $\|x\| = 0$  if x is a unit, and

$$||x|| = \sup\{n : (\exists x_1, \dots, x_n \in A_M)(x = x_1 \cdots x_n)\}\$$

otherwise, if such a decomposition exists and the involved supremum is finite. Then we say that M is *quasi-atomic* if ||x|| exists for each x in M.

Thus, M is quasi-atomic if and only if it is generated by its atoms and its units, and, moreover, for each x in M, the maximal number of atoms occurring in a decomposition of x is finite. If there is no nontrivial unit in the monoid M, then M is quasi-atomic if and only if it is atomic in the sense of [11, 9].

Lemma 5.4. Assume that M is a quasi-atomic monoid. Then we have

$$||x|| = 0 \Leftrightarrow x \in M^*,$$

$$||xy|| \ge ||x|| + ||y||,$$

$$(5.3) u \in M^* \Rightarrow ||xu|| = ||ux|| = ||x||,$$

for all x, y in M. So  $x \simeq x'$  implies ||x|| = ||x'||.

PROOF. For (5.1),  $x \in M^*$  implies ||x|| = 0 by definition. Conversely, for  $x \notin M^*$ , the hypothesis that ||x|| exists means that x can be expressed as a finite product of atoms, so, by definition,  $||x|| \ge 1$  holds in this case.

Assume now  $\|x\| = p$  and  $\|y\| = q$ . For p = q = 0, both x and y are units, so is xy, and we have  $\|xy\| = 0 = p + q$ . For p > 0 and q = 0, y is a unit, while x admits a decomposition  $x = x_1 \cdots x_p$  with  $x_1, \ldots, x_p \in A_M$ . Then  $x_p y$  is an atom, and we obtain  $xy = x_1 \cdots x_{p-1}(x_p y)$ , hence  $\|xy\| \ge p = p + q$ . The argument is similar for p = 0 and q > 0. Assume now p > 0 and q > 0. Then x and y admit decompositions  $x = x_1 \cdots x_p$ ,  $y = y_1 \cdots y_q$  with  $x_1, \ldots, y_q \in A_M$ , and we deduce  $xy = x_1 \cdots x_p y_1 \cdots y_q$ , hence  $\|xy\| \ge p + q$ . This shows (5.2).

Assume  $u \in M^*$ . Then (5.2) gives  $||xu|| \ge ||x||$ . Applying this with xu instead of x and  $u^{-1}$  instead of u, we obtain  $||(xu)u^{-1}|| \ge ||xu||$ , i.e.,  $||x|| \ge ||xu||$ , whence ||xu|| = ||x||. The argument is similar for ux.

PROPOSITION 5.5. A monoid M is quasi-atomic if and only if there exists a mapping  $l: M \to \mathbb{N}$  satisfying, for all x, y in M,

$$(5.4) l(x) = 0 \Rightarrow x \in M^*,$$

$$(5.5) l(xy) \ge l(x) + l(y).$$

PROOF. Lemma 5.4 shows that the mapping  $||\cdot||$  satisfies (5.4) and (5.5) when M is quasi-atomic, so the condition is necessary.

Conversely, assume that l is a mapping satisfying (5.4) and (5.5). Assume  $x \in M-M^*$ , and let  $x=x_1\cdots x_p$  be a decomposition of x into non-invertible elements. By (5.4), we have  $l(x_i) \geq 1$  for each i, and, by (5.5),  $l(x) \geq l(x_1) + \cdots + l(x_p)$ , hence  $l(x) \geq p$ . So, the supremum n of the lengths of the decompositions of x into a product of non-invertible elements satisfies  $n \leq l(x)$ , and, therefore, it is finite. Now, let  $x = x_1 \cdots x_n$  be such a decomposition with maximal length. We claim that  $x_1, \ldots, x_n$  are atoms. Indeed, if  $x_i$  is not an atom, it can be decomposed as  $x_i = x_i'x_i''$  with neither  $x_i'$  nor  $x_i''$  in  $M^*$ , and replacing  $x_i$  with  $x_i'x_i''$  gives a decomposition of x of length  $x_i$  atoms. This shows that  $x_i$  is quasi-atomic, and that  $||x|| \leq l(x)$  holds for every x in  $x_i'$  in  $x_i'$  in  $x_i'$  is quasi-atomic, and that

EXAMPLE 5.6. The monoids  $M_i$  of Example 2.11 all are quasi-atomic, and even atomic as they contain no nontrivial unit: as the defining relations preserve the length, the latter induces a well defined mapping l of  $M_i$  to  $\mathbb{N}$  that satisfies (5.4) and (5.5).

The previous situation is general, as we have:

Proposition 5.7. Every thin cancellative monoid is quasi-atomic.

PROOF. Assume that M is a thin cancellative monoid, and S is a finite spanning subset of M. Let  $x \in M$ . As S generates M, x belongs to  $S^p$  for some p. Now, let  $x = x_1 \cdots x_n$  be an arbitrary decomposition of x with  $x_1, \ldots, x_n \in S - M^*$ . By Prop. 2.7,  $S^p$  spans M, so the element  $x_i \cdots x_n$ , which is a right divisor of x, belongs to  $S^p$  as well. Assume  $n \geq \operatorname{card}(S^p)$ . Then there exist i, j with  $0 \leq i < j \leq n$  satisfying  $x_i \cdots x_n = x_j \cdots x_n$ , which implies  $x_i \cdots x_{j-1} = 1$  and contradicts the assumption  $x_i \notin M^*$ . Thus we must have  $n \leq \operatorname{card}(S^p) \leq \operatorname{card}(S)^p$ . Let us define l(x) to be the maximal possible value of n in a decomposition as above. It is clear

that l(x) = 0 implies  $x \in M^*$ , and that  $l(xy) \ge l(x) + l(y)$  always holds, as shows concatenating a maximal decomposition for x and a maximal decomposition for y. So the mapping l satisfies the conditions (5.4) and (5.5), and, by Prop. 5.5, M is quasi-atomic.

When we only assume that a finite quasi-spanning set exists, the situation is more complicated. However, we can still recognize quasi-atomicity as follows:

PROPOSITION 5.8. Assume that M is a quasi-thin cancellative monoid. Then M is quasi-atomic if and only if the group  $M^*$  is closed under conjugation in M, in the sense that, if xu = u'x holds, then  $u \in M^*$  implies  $u' \in M^*$ .

PROOF. Assume that M is quasi-atomic and we have xu = u'x with  $u \in M^*$ . By Lemma 5.4, we have  $||x|| = ||xu|| = ||u'x|| \ge ||u'|| + ||x||$ , hence ||u'|| = 0, and  $u' \in M^*$  by (5.1). So the condition is necessary.

Conversely, assume that  $M^*$  is closed under conjugation, and S is a finite quasispanning set in M. We adapt the argument of the proof of Prop. 5.7. Let  $x \in M$ . Then x belongs to  $S^pM^*$  for some p. Let  $x = x_1 \cdots x_n u$  be any decomposition of x with with  $x_1, \ldots, x_n \in S - M^*$  and  $u \in M^*$ . The set  $S^pM^*$  spans M, so, by Prop. 2.7,  $x_i \cdots x_n$ , which is a right divisor of x, belongs to  $S^pM^*$  for each i. If  $n \geq \operatorname{card}(S^p)$  holds, there exist i, j with  $0 \leq i < j \leq n$  satisfying  $x_i \cdots x_n \simeq x_j \cdots x_n$ , so we have  $(x_i \cdots x_{j-1})(x_j \cdots x_n) = (x_j \cdots x_n)u$  for some unit u. This, by hypothesis, implies  $x_i \cdots x_{j-1} \in M^*$ , contradicting the hypothesis  $x_i \notin M^*$ . So we must have  $n \leq \operatorname{card}(S^p) \leq \operatorname{card}(S)^p$ . If we define l(x) to be the maximal possible value of n in a decomposition as above, then l satisfies (5.4) and (5.5), and M is quasi-atomic.

If M is a quasi-atomic cancellative monoid, then the relation  $\prec$  is a strict partial ordering with no infinite descending chain (and so is its right counterpart). Indeed, by (5.2),  $x \prec y$  implies ||x|| < ||y||, so  $\prec$  may admit no cycle, hence it is a strict partial ordering, and it admits no infinite descending chain since ( $\mathbb{N}$ , <) does. In such a framework, we can introduce the notion of a minimal common (right) multiple ("mcm"), which extends the notion of a least common multiple ("lcm"):

DEFINITION 5.9. Assume that M is a monoid. For  $x, y \in M$ , we say that z is an mcm of x and y if both  $x \leq z$  and  $y \leq z$  hold, but  $x \leq z'$  and  $y \leq z'$  hold for no proper divisor z' of z. We denote by C(x,y) the set of all elements y' such that xy' is an mcm of x and y (if any).

An mcm is like a lcm, except that we require no uniqueness. For instance, in the monoid  $M_1$  of Example 2.11, the elements a and b admit two mcm's, namely  $a^2$  and ab, but they admit no lcm, as we have neither  $a^2 \leq ab$  nor  $ab \leq a^2$ .

LEMMA 5.10. Assume that M is a quasi-atomic cancellative monoid. Then, for all x, y in M, every common multiple of x and y (if any) is a multiple of some mcm of x and y.

PROOF. Assume  $x \leq z$  and  $y \leq z$ . Let  $Z = \{z' \leq z : x \leq z', y \leq z'\}$ . Then any element z' of Z such that ||z'|| has the least possible value is an mcm of x and y.  $\square$ 

We are now ready to introduce the notion of a primitive element:

DEFINITION 5.11. Assume that M is a quasi-atomic cancellative monoid. We say that an element x of M is *primitive* if x belongs to the smallest subset S of M

that contains the atoms and is such that, if x and y belong to S, so does y' whenever xy' is an mcm of x and y. The set of all primitive elements of M is denoted by  $P_M$ .

In other words,  $P_M$  is the closure of  $A_M$  under operation C.

EXAMPLE 5.12. If any two elements admitting a common multiple admit a lcm, the set C(x,y) is either empty, or it consists of a single element  $x \setminus y$ , so the primitive elements are the closure of the atoms under operation  $\setminus$ .

Consider now the monoid  $M_1$  of Example 2.11. There are two atoms, namely a and b. We observed above that a and b admit two right mcm's, namely  $a^2$  and ab, so C(a,b) consist of the two elements a and b, and so does C(b,a). It follows that the closure of  $A_M$  under C is the set  $\{1,a,b\}$ , *i.e.*, there are three primitive elements in  $M_1$ .

The reader can easily check that there are four primitive elements in the monoids  $M_2$  and  $M_3$ , namely 1 and the atoms a, b, and c.

Lemma 5.13. Assume that M is a quasi-atomic cancellative monoid. Then the set  $P_M$  is closed under right multiplication by a unit, and, therefore, it is  $\simeq$ -saturated.

PROOF. Assume  $x \in P_M$ , and  $u \in M^*$ . If x is an atom, then xu is an atom as well, so it is primitive. Otherwise, there exist y, z in  $P_M$  such that yx is an mcm of y and z. In this case, yxu is also an mcm of y and z, so xu also belongs to C(y,z), and, therefore, it is primitive. So, we have  $P_MM^* \subseteq P_M$ , and, therefore,  $P_M$  is  $\simeq$ -saturated.

We can now prove:

Proposition 5.14. Assume that M is a quasi-atomic cancellative monoid. Then  $P_M$  spans M, and every spanning subset of M includes  $P_M$ .

PROOF. First,  $P_M$  is  $\simeq$ -saturated by Lemma 5.13. Then, if x is an atom of M and u is a unit, xu is an mcm of x and x, and, therefore, u is primitive. Thus  $P_M$  includes  $A_M$  and  $M^*$ , and, therefore, it generates M. Next, assume  $x \leq z$  and  $y \leq z$  with  $x, y \in P_M$ . By Lemma 5.10, there exist x', y' such that  $xy' = yx' \leq z$  holds and xy' is an mcm of x and y. This implies  $x' \in P_M$  and  $y' \in P_M$  by definition. Hence  $P_M$  satisfies Condition (2.1), *i.e.*, it spans M.

Let S be an arbitrary spanning subset of M. As S generates M, it necessarily includes  $A_M$ . Then, S has to be closed under C. Indeed, assume that x, y lie in S, and xy'' is an mcm of x and y, say xy'' = yx''. As S spans M, there exist y' in S and z satisfying y'' = y'z. The hypothesis that xy'' is an mcm implies ||xy'|| = ||xy''||, hence ||z|| = 0, and  $y' \simeq y''$ . As S is  $\simeq$ -saturated by definition, we deduce  $y'' \in S$ . So, S includes  $A_M$  and it is closed under C, hence it includes the closure  $P_M$  of  $A_M$  under C.

COROLLARY 5.15. Assume that M is a quasi-atomic cancellative monoid. Then M is thin (resp. quasi-thin) if and only the set  $P_M$  is finite (resp. quasi-finite).

Once we know that primitive elements span M, we can apply Prop. 2.7 and we deduce that, if M is a quasi-atomic cancellative monoid, then the set  $P_M$  is closed under left multiplication by a unit, and every right divisor of a primitive element is primitive.

Another consequence of Prop. 5.14 is that, if M is a quasi-atomic cancellative monoid, and S is a  $\simeq$ -selector through  $P_M$ , then S is a minimal quasi-spanning subset of M, and  $S \cap A_M$  is a  $\simeq$ -selector through  $A_M$ . Indeed, by construction,  $SM^*$  is equal to  $P_M$ , so S quasi-spans M. If S' is a proper subset of S,  $S'M^*$  is a proper subset of  $P_M$ , so it cannot span M, and S' cannot quasi-span M. Finally, every atom x is primitive, so it belongs to  $SM^*$ , and, therefore, x is  $\simeq$ -equivalent to one element of S.

### 6. Simple elements

A crucial feature in Garside's and Thurston's analysis of the braid monoids and its subsequent extensions is the existence of a finite subset that is closed both under lcm and right divisor: in the current framework, this means that there exists a finite spanning subset S that is closed under lcm, *i.e.*, the lcm of two elements of S belongs to S. The least such set S happens to be the closure of primitive elements under lcm, and its elements, called minimal in [5, 6], or simple in [9], play a prominent rôle. In particular, there exists a maximal simple element  $\Delta$  which enjoys most of the properties of Garside's fundamental braids  $\Delta_n$  [15].

So, in the current approach, a natural idea would be to look for finite spanning subsets closed under mcm. Unfortunately, when least common multiples do not exist, more precisely when common multiples exist but some elements admit at least two non  $\simeq$ -equivalent mcm's, no such set may exist:

PROPOSITION 6.1. Assume that M is a quasi-atomic cancellative monoid, any two elements of M admit a common multiple, and S is a finite spanning subset of M that is closed under mcm. Assume in addition that  $x \leq y \in S$  implies  $x \in S$ . Then any two elements of M admit a lcm.

PROOF. As S is finite and closed under right mcm, there exists  $\Delta$  in S such that  $x \leq \Delta$  holds for every x in S, i.e., there exists  $x^*$  satisfying  $xx^* = \Delta$ ; as M is left cancellative,  $x^*$  is unique, and, as S spans M, every right divisor of an element of S belongs to S, so  $x^*$  belongs to S. The mapping  $x \mapsto x^*$  is injective, and, therefore, it is a permutation of S. Assume  $x, y \in S$ , and let xy' and xy'' be two right mcm's of x and y. By hypothesis, xy' belongs to S, so, by the previous remark, there exists z in S satisfying  $xy' = z^*$ , i.e.,  $zxy' = \Delta$ . By hypothesis, S is closed under left divisors, so zx and, similarly, zy belong to S, and so does their right mcm zxy''. So we must have  $zxy'' \leq \Delta$ , hence  $y'' \leq y'$ . In other words, xy' is a lcm of x and y. Finally, as S generates M, the existence of a lcm for each pair of elements of S inductively implies the existence of a lcm for each pair of elements of S.

Thus, we must find a more subtle definition. The following one is convenient, in the sense that it will prove appropriate for the construction of a normal form.

DEFINITION 6.2. Assume that M is a cancellative monoid, and S quasi-spans M. We say that an element x of M is S-simple if  $y \prec x$  implies  $\mathrm{Div}(y) \cap S \neq \mathrm{Div}(x) \cap S$ . If M is quasi-atomic, we say simple for  $P_M$ -simple.

The elements of S always are S-simple. Indeed, for  $x \in S$ , we have  $x \in \text{Div}(x) \cap S$ , but  $y \prec x$  implies  $x \not\prec y$ , *i.e.*,  $x \notin \text{Div}(y) \cap S$ . So, in particular, a primitive element is always simple. By definition, an element is S-simple if and only if it is an mcm of its divisors lying in S. In particular, in the Gaussian case,

an element is S-simple if and only if it is the lcm of its divisors lying in S, and, therefore, a S-simple element x is completely determined by the set  $\mathrm{Div}(x) \cap S$ . This need not be true in the general case.

EXAMPLE 6.3. Let M be a free commutative monoid based on  $\{a_1, \ldots, a_n\}$ . Then the atoms of M are  $a_1, \ldots, a_n$ , there are n+1 primitive elements, namely 1 and the atoms, and there are  $2^n$  simple elements, namely the elements  $a_I = \prod_{i \in I} a_i$  for  $I \subseteq \{1, \ldots, n\}$ . Indeed,  $a_i \preceq a_I$  is equivalent to  $i \in I$  in this case, and, for every x, the element  $a_I$  with  $I = \{i : a_i \preceq x\}$  is a divisor of x with the same divisors in  $P_M$ , so no element not of the form  $a_I$  may be simple.

On the other hand, there are three primitive elements, namely 1, a, and b in the monoid  $M_1$  of Example 2.11. These elements are simple, and there are two more simple elements, namely  $a^2$  and ab. Here, we have  $\text{Div}(a^2) \cap P_M = \text{Div}(ab) \cap P_M = P_M$ , which gives an example where a simple element is not determined by the family of its primitive divisors.

Similarly, there are seven simple elements in  $M_3$ , namely the four primitive elements 1, a, b, c, and, in addition, the three elements ab, ba, and  $b^2$ : the sets of primitive divisors of the latter elements are  $\{a,b\}$ ,  $\{b,c\}$ , and  $\{a,b,c\}$  respectively, so, here, a simple element happens to be determined by its primitive divisors (although the monoid admits no lcm).

In the Gaussian case, *i.e.*, when least common multiples exist, the current definition of a simple element is equivalent to that of [9]. In particular, in the thin case, the simple elements of M have a natural characterization extending that given for a free commutative monoid in Example 6.3.

Proposition 6.4. Assume that M is a thin Gaussian monoid, i.e., a Garside monoid. Then the simple elements of M are finite in number, and they coincide with the divisors of the  $lcm \Delta$  of  $P_M$ .

PROOF. Let  $\{x_i; i=1,\ldots,n\}$  be an enumeration of  $P_M$ . For  $I\subseteq\{1,\ldots,n\}$ , let  $x_I$  be the lcm of the  $x_i$ 's with  $i\in I$ . Then  $x_I$  is simple, and, conversely, every simple element must be of this form. Let  $\Delta$  be the lcm of  $P_M$ . Then, by construction, every simple element  $x_I$  is a divisor of  $\Delta$ . The computation rules for lcm's then imply that simple elements span M [9], and, as a consequence, that every divisor of  $\Delta$  is simple.

It is well known that, if M is a spherical Artin monoid, then the simple elements are in one-to-one correspondence with the elements of the associated finite Coxeter group [4, 12]: for instance, the n! simple elements in the braid monoid  $B_n^+$  are in one-to-one correspondence with the permutations of n objects. More generally, it is shown in [9] that the simple elements of a Garside monoid make a finite lattice with a unique maximal element, the lcm  $\Delta$  of the primitive element. As shows the case of the monoid  $M_1$ , such a property need not be true in the general case.

For future use, we gather now some general results about S-simple elements.

LEMMA 6.5. Assume that M is a cancellative monoid, and S quasi-spans M. (i) An element of M is S-simple if and only if it is  $SM^*$ -simple.

- (ii) The set of all S-simple elements is closed under left and right multiplication by a unit, and, therefore, it is  $\simeq$ -saturated.
- PROOF. (i) Assume that x is S-simple, and  $y \prec x$  holds. By definition, we have  $\mathrm{Div}(y) \cap S \neq \mathrm{Div}(x) \cap S$ , so, a fortiori,  $\mathrm{Div}(y) \cap SM^* \neq \mathrm{Div}(x) \cap SM^*$ , and x

is  $SM^*$ -simple. Conversely, assume that x is  $SM^*$ -simple, and  $y \prec x$  holds. Then we have  $z \preceq x$  and  $z \not\preceq y$  for some z in  $SM^*$ . By definition, we have z = z'u for some z' in S and u in  $M^*$ . Then  $z' \preceq x$  and  $z' \not\preceq y$  hold, and x is S-simple.

(ii) Assume that x is S-simple and u is a unit. Then we have  $\mathrm{Div}(xu) = \mathrm{Div}(x)$ , and  $y \prec xu$  is equivalent to  $y \prec x$ . Hence  $y \prec x$  implies  $\mathrm{Div}(y) \cap S \neq \mathrm{Div}(xu) \cap S$ , and xu is S-simple. The argument is similar for left multiplication by u, as  $\mathrm{Div}(ux) = \mathrm{Div}(x)$  holds.

In the general case, as shows the example of  $M_1$ , simple elements need not span the monoid. However, we still have the following closure property:

Lemma 6.6. Assume that M is a cancellative monoid, and S quasi-spans M. Then every right divisor of a S-simple element of M is S-simple.

PROOF. Assume that xy is S-simple. We wish to show that y is S-simple. Assume  $y' \prec y$ . Then we have  $xy' \prec xy$ , so, by definition, there exists z in S satisfying  $z \preceq xy$  and  $z \not\preceq xy'$ . As  $SM^*$  spans M, there exists z' in  $SM^*$  and x' in M satisfying  $xz' = zx' \preceq xy$ , hence  $z' \preceq y$ . Thus, z' belongs to  $\mathrm{Div}(y) \cap SM^*$ . On the other hand,  $z' \preceq y'$  would imply  $zx' = xz' \preceq xy'$ , hence  $z \preceq xy'$ , contradicting the hypothesis. So  $z' \preceq y'$  is impossible. We have  $\mathrm{Div}(y') \cap SM^* \neq \mathrm{Div}(y) \cap SM^*$ , so y is  $SM^*$ -simple, hence S-simple.

Proposition 6.4 implies that, in the Gaussian case, there exist at most  $2^n$  simple elements when there are n primitive elements, a bound which we have seen is nearly reached in the case of a free commutative monoid. The result extends to the general case as follows:

PROPOSITION 6.7. Assume that M is a cancellative monoid, and S is a quasi-spanning subset of M with n elements. Then every S-simple element belongs to  $S^nM^*$ , and, therefore, there are at most  $n^n \simeq$ -equivalence classes of S-simple elements in M.

PROOF. Assume that x is S-simple, and let  $x_1, \ldots, x_p$  be an enumeration of  $\mathrm{Div}(x) \cap S$ . By Proposition 3.5(iii), there exists x' in  $S^p$  such that  $x_i \leq x'$  holds for  $1 \leq i \leq p$ . Then, we have  $\mathrm{Div}(x') \cap S \supseteq \mathrm{Div}(x) \cap S$ , hence  $\mathrm{Div}(x') \cap S = \mathrm{Div}(x) \cap S$ . By definition of a S-simple element,  $x' \prec x$  is impossible, so  $x' \simeq x$  is the only possibility, which shows that x belongs to  $S^pM^*$ , and, therefore, to  $S^nM^*$  since 1 is primitive and  $p \leq n$  holds.

COROLLARY 6.8. If M is a thin cancellative monoid, then the simple elements of M are finite in number. More precisely, if M contains n primitive elements, it contains at most  $n^n$  simple elements.

If M is a quasi-atomic cancellative monoid and S is a  $\simeq$ -selector through simple elements, then  $S \cap P_M$  is a  $\simeq$ -selector through  $P_M$ , and  $S \cap A_M$  is a  $\simeq$ -selector through  $A_M$ . Conversely, every  $\simeq$ -selector through  $A_M$  can be extended into a selector through  $P_M$ , and, then, through simple elements.

Finally, let us observe that simple elements, as atoms and primitive elements, are defined intrinsically, and, therefore, they are preserved by automorphisms:

PROPOSITION 6.9. Assume that M is a quasi-atomic cancellative monoid, and  $\phi$  is an automorphism of M. Then  $\phi$  globally preserves  $A_M$ ,  $P_M$ , and the set of all simple elements in M.

PROOF. As  $\phi$  maps units to units, then it maps non-atoms to non-atoms, and, therefore, it maps atoms to atoms. Then, it maps every mcm of two elements to an mcm of their images, and, therefore, it maps every primitive element to a primitive element. Finally,  $\phi$  preserves the relations  $\leq$  and  $\prec$ , so it maps simple elements to simple elements.

#### 7. Normal forms

The main interest of simple elements in the Gaussian case, *i.e.*, when least common multiples exist, is that they can be used to construct good normal forms. In particular, the greedy normal form originally defined for the braid monoids [?, 18, 13, 14] extends to every Gaussian monoid, and, subsequently, to the corresponding group of fractions [9]. The principle is that, for  $x \neq 1$  in the considered monoid M, there exists a maximal simple divisor  $x_1$  of x, namely the gcd of x and the maximal simple element  $\Delta$ , so we can write  $x = x_1x'$ , and, applying the process to x', we inductively obtain a decomposition  $x = x_1x_2 \cdots$  in terms of simple elements. This decomposition enjoys good properties, and, in particular, it gives rise to a bi-automatic structure on the associated group of fractions.

A crucial technical point in the above construction is that simple elements happen to span the monoid, in the Gaussian case. We shall see now that a similar construction is still possible in the general case when we start with an arbitrary spanning set S and use the derived notion of a S-simple element. The price to pay for the generalization is that a given element possibly may have more than one normal decomposition, but, this fact excepted, the results remain similar, and the proofs are extremely easy.

As in the Gaussian case, we start from the fact that, for every element x, there exists a maximal S-simple divisor of x:

Lemma 7.1. Assume that M is a quasi-atomic cancellative monoid, S quasi-spans M, and  $x_0$  is a S-simple element. Then, for every x in M, there exists a S-simple divisor  $x_1$  of x satisfying  $\mathrm{Div}(x) \cap S = \mathrm{Div}(x_1) \cap S$ ; moreover, we may assume  $x_0 \leq x_1$  whenever  $x_0 \leq x$  holds.

If M is Gaussian, then  $x_1$  is a lcm of  $Div(x) \cap S$ , and, so, it is unique.

PROOF. Assume  $x_0 \leq x$ . Let Y be the set of all S-simple elements y satisfying  $x_0 \leq y \leq x$ , and let  $x_1$  be an element of Y such that  $\|x_1\|$  has the maximal possible value: such an element exists since  $y \in Y$  implies  $\|y\| \leq \|x\|$ . Write  $x = x_1x''$ . Assume  $z \in \operatorname{Div}(x) \cap S$ . As S quasi-spans M, there must exist z' in S, and  $x'_1$  in M satisfying  $x_1z' = x'_1z' \leq x$ . So we have  $x_0 \leq x_1 \leq x_1z' \leq x$ . Moreover, provided z' has been chosen so that  $\|x_1z'\|$  has the least possible value, no proper divisor of  $x_1z'$  is a multiple of z, which implies that  $x_1z'$  is S-simple, and, therefore, it belongs to Y. The definition of  $x_1$  then implies  $\|x_1z'\| = \|x_1\|$ , hence  $z' \in M^*$ , and then  $z \leq x_1$ . So we have  $\operatorname{Div}(x_1) \cap S = \operatorname{Div}(x) \cap S$ . Take  $x_0 = 1$  for the general result.

In the Gaussian case, the lcm of  $\mathrm{Div}(x) \cap S$  is a S-simple element satisfying the requirements, and it divides every other element satisfying them, so it must be the only solution.

DEFINITION 7.2. Assume that M is a cancellative monoid, and S quasi-spans M. We say that a sequence  $(x_1, \ldots, x_n)$  in M is S-prenormal if, for each i, we have  $\mathrm{Div}(x_i) \cap S = \mathrm{Div}(x_i \cdots x_n) \cap S$ . We say that  $(x_1, \ldots, x_n)$  is S-normal if it is

S-prenormal, and, in addition, each factor  $x_i$  is S-simple. If M is quasi-atomic, we say (pre)normal for  $P_M$ -(pre)normal.

Say that a sequence  $(x_1, \ldots, x_n)$  is a decomposition for x if  $x = x_1 \cdots x_n$  holds. Iterating Lemma 7.1, we find:

PROPOSITION 7.3. Assume that M is a quasi-atomic cancellative monoid, S quasi-spans M, and  $x_0$  is a S-simple element. Then every element x of M satisfying  $x_0 \leq x$  admits a S-normal decomposition  $(x_1, \ldots, x_n)$  with  $x_0 \leq x_1$ .

PROOF. Lemma 7.1 gives a S-simple element  $x_1$  satisfying both  $x_0 \leq x_1$  and  $\operatorname{Div}(x_1) \cap S = \operatorname{Div}(x) \cap S$ . Write  $x = x_1 x'$ . If x' is a unit, then x is S-simple by Lemma 6.5(ii), and we are done. Otherwise, we have ||x'|| < ||x||: inductively, we find a S-normal decomposition  $(x_2, \ldots, x_n)$  of x', and concatenating  $x_1$  with the latter gives a S-normal decomposition of x.

By Lemma 7.1, the normal form of Prop. 7.3 is unique in the Gaussian case, and it coincides with the greedy normal form of [14, 9]. More generally, the S-normal form is unique whenever distinct S-normal elements never admit the same divisors in S. Now, the latter condition need not be true, and the normal form need not be unique in general.

EXAMPLE 7.4. Consider once again the monoid  $M_1$  of Example 2.11. Then  $a^2$  and ab are simple elements, and  $(a^2, a^2)$  and (ab, ab) are two normal decompositions for  $a^4$  in M. On the other hand, we observed that, in the case of  $M_3$ , the simple elements are uniquely determined by their primitive divisors. So, in this case, the normal decomposition is unique.

When nontrivial units exist, we can replace the family of all S-simple elements by a  $\simeq$ -selector, at the expense of keeping a unit at the end of the decomposition.

COROLLARY 7.5. Assume that M is a quasi-atomic cancellative monoid, S quasi-spans M, and  $\Sigma$  is a  $\simeq$ -selector through S-simple elements in M. Then every element x of M admits a decomposition  $(x_1, \ldots, x_n, u)$  with  $x_1, \ldots, x_n \in \Sigma$ ,  $u \in M^*$  and  $x_i \triangleright_S x_{i+1}$  for each i.

PROOF. Let  $x \in M$ . By Prop. 7.3, x admits a S-normal decomposition  $(x'_1, \ldots, x'_n)$ , where  $x'_1, \ldots, x'_n$  are S-simple and  $x'_i \triangleright_S x'_{i+1}$  holds for every i. Using (3.2), we find  $x_1, \ldots, x_n$  in  $\Sigma$  and u in  $M^*$  satisfying  $x_i \simeq x'_i$  for every i and  $x_1 \cdots x_n u = x'_1 \cdots x'_n = x$ . By Lemma 7.7,  $x'_i \triangleright_S x'_{i+1}$  implies  $x_i \triangleright_S x_{i+1}$ .

The interest of the current construction lies in that S-normal sequences admit a purely local characterization.

DEFINITION 7.6. Assume that M is a monoid and S is a subset of M. For  $x, y \in M$ , we say that x covers y w.r.t. S, denoted  $x \triangleright_S y$ , if  $\mathrm{Div}(xy) \cap S = \mathrm{Div}(x) \cap S$  holds, *i.e.*, if every element of S dividing xy already divides x. If M is quasi-atomic, we write  $\triangleright$  for  $\triangleright_{P_M}$ .

Lemma 7.7. Assume that M is a cancellative monoid and S is a subset of M. Then the relations  $\triangleright_S$  and  $\triangleright_{SM^*}$  coincide.

PROOF. It is obvious that  $x \triangleright_{SM^*} y$  implies  $x \triangleright_S y$ . Conversely, assume  $x \triangleright_S y$  and  $z' \simeq z \in S$ . Then  $z' \preceq xy$  (resp.  $z' \preceq x$ ) is equivalent to  $z \preceq xy$  (resp.  $z \preceq x$ ). So  $z' \preceq xy$  implies  $z \preceq xy$ , hence  $z \preceq x$ , hence  $z' \preceq x$ , and we have  $x \triangleright_{SM^*} y$ .




Figure 7.1. Covering relation

All properties of the normal form relie on the following basic observations:

Lemma 7.8. Assume that M is a cancellative monoid, and S quasi-spans M. Then, for x, y, z in M:

- (i) The relation  $y \triangleright_{S} z$  implies  $xy \triangleright_{S} z$ ;
- (ii) The conjunction of  $x \triangleright_S y$  and  $y \triangleright_S z$  implies  $x \triangleright_S yz$ .

PROOF. (i) (Fig. 7.1) Assume  $s \in S$  and  $s \leq xyz$ . As  $SM^*$  spans M, there exist s' in  $SM^*$ , and x' in M satisfying  $sx' = xs' \leq xyz$ , hence  $s' \leq yz$ . By Lemma 7.7,  $y \triangleright_S z$  implies  $y \triangleright_{SM^*} z$ , so  $s' \leq yz$  implies  $s' \leq y$ , and, therefore, we have  $s \leq xs' \leq xy$ .

(ii) Assume  $s \in S$  and  $s \leq xyz$ . By (i),  $y \triangleright_S z$  implies  $xy \triangleright_S z$ , so we deduce  $s \leq xy$ . Then the hypothesis  $x \triangleright_S y$  implies  $s \leq x$ .

We can know establish the expected local characterization of normal sequences, a necessary first step toward a possible automatic structure:

PROPOSITION 7.9. Assume that M is a cancellative monoid, and S quasi-spans M. Then a sequence  $(x_1, \ldots, x_n)$  in  $M^n$  is S-prenormal if and only if  $x_i \triangleright_S x_{i+1}$  holds for each i.

PROOF. By definition, the sequence  $(x_1, \ldots, x_n)$  is S-prenormal if and only if  $x_i \triangleright_S x_{i+1} \cdots x_n$  holds for each i. By definition, the latter relation always implies  $x_i \triangleright_S x_{i+1}$ . By Lemma 7.8(ii), the converse implication is also true: a descending induction on j shows that  $(\forall i \geq j)(x_i \triangleright_S x_{i+1})$  implies  $x_i \triangleright_S x_{i+1} \cdots x_p$ . Indeed, the conjunction of  $x_j \triangleright_S x_{j+1}$  and  $x_{j+1} \triangleright_S x_{j+2} \cdots x_n$  implies  $x_j \triangleright_S x_{j+1} \cdots x_n$ .

REMARK 7.10. Instead of using S-simple elements, we could think of simply considering elements of S, and constructing a normal form of x starting with a maximal divisor of x in S. But, then, the normal sequences would not necessarily admit the local characterization of Prop. 7.9. For instance, in the monoid  $M_1$ , if we take  $S = \text{Div}(a^2)$  (a spanning subset that we shall consider in Sec. 8 below), the two sequences (a, b) and (b, a) would be S-normal, as a is a maximal divisor of ab in S, and b is a maximal divisor of ab in S, but the concatenated sequence (a, b, a) would not, as we have  $a^2 \leq aba$ , and therefore a is not a maximal divisor of aba in S.

We have seen that the normal form of Prop. 7.3 need not be unique in general. We shall need in Sec. 9 below the following refinement of Prop. 7.3 that connects the various normal decompositions of an element:

PROPOSITION 7.11. Assume that M is a quasi-atomic cancellative monoid, S quasi-spans M, and  $x_1, \ldots, x_n$  are S-simple elements of M. Then  $x_1 \cdots x_n$  admits a S-normal decomposition  $(x'_1, \ldots, x'_m)$  such that  $m \leq n$  holds and, for each i, we have  $x_1 \cdots x_{f(i)} \leq x'_1 \cdots x'_i$  for some increasing mapping f of  $\{1, \ldots, m\}$  into  $\{1, \ldots, n\}$  with f(m) = n.

PROOF. The result is trivial for n=1. Assume n=2. Applying Prop. 7.3 to  $x_1x_2$ , we find a S-normal decomposition of  $x_1x_2$  that begins with some multiple  $x_1'$  of  $x_1$ . Two cases may happen. Either  $x_1x_2$  is S-simple, and  $(x_1')$  is the expected decomposition. Or we have  $x_1' = x_1y$  with  $y \prec x_2$ , and, therefore,  $x_1x_2 = x_1'x_2'$  with  $x_2 = yx_2'$ . By Lemma 6.6, the latter relation forces  $x_2'$  to be S-simple, and, therefore,  $(x_1', x_2')$  is a S-normal decomposition of the expected form.

For  $n \geq 3$ , we use induction on n. Applying the induction hypothesis, we find a S-normal decomposition  $(y_2,\ldots,y_p)$  for  $x_2\cdots x_n$  and an increasing mapping g of  $\{2,\ldots,p\}$  into  $\{2,\ldots,n\}$  satisfying  $x_2\cdots x_{g(i)} \preceq y_2\cdots y_i$  for  $2\leq i\leq p$ . If p< n holds, we can apply the induction hypothesis to  $x_1,y_2,\ldots,y_p$ , and get the result directly. So, assume p=n. Then g must be the identity mapping. Applying the result with n=2 to  $x_1y_2$ , we obtain a S-normal decomposition of length 2 or 1. In the latter case, we resort to the induction hypothesis directly. So, assume that we have obtained  $(x_1',y_2')$  with  $x_1\preceq x_1'$  and  $x_1'y_2'=x_1y_2$ . We apply the induction hypothesis another time to  $y_2'y_3\cdots y_n$ , obtaining a S-normal decomposition  $(x_2',\ldots,x_m')$ . Then  $(x_1',x_2',\ldots,x_m')$  satisfies our requirements. Indeed, by construction, we have  $y_2 \rhd_S y_3\cdots y_n$ , hence, by Lemma 7.8,  $x_1'y_2'=x_1y_2 \rhd_S y_3\cdots y_n$ , and, as  $x_1' \rhd_S y_2'$  holds by construction,  $x_1' \rhd_S y_2'y_3\cdots y_n$ , hence  $x_1' \rhd_S x_2'$ . So the sequence  $(x_1',\ldots,x_m')$  is S-normal. The relations  $x_1\cdots x_{f(i)} \preceq x_1'\cdots x_i'$  follow from the induction hypothesis.

Although natural, the previous result was not obvious: putting in normal form a product of two simple elements might have required say three simple elements, since the conditions for being normal discard some decompositions.

We consider now the effect of multiplication on normal forms, *i.e.*, we try to connect the normal form(s) of an element x with those of yx and xy, especially when y is S-simple. As one can expect, such results will be crucial for constructing an automatic structure.

LEMMA 7.12. Assume that M is a quasi-atomic cancellative monoid, and S quasi-spans M. Let x, y be arbitrary elements of M, and  $(x_1, \ldots, x_n)$  be a S-prenormal decomposition of x. Put  $y_0 = y$ , and, inductively, let  $(x'_i, y_i)$  be any S-prenormal decomposition of  $y_{i-1}x_i$ . Then  $(x'_1, \ldots, x'_n, y_n)$  is a S-prenormal decomposition of yx.

If, in addition, y is S-simple and  $(x_1, \ldots, x_n)$  is S-normal, then we may assume that each element  $y_i$  is S-simple, and then  $(x'_1, \ldots, x'_n, y_n)$  is a S-normal decomposition of yx.

PROOF. (Fig. 7.2) We have  $x'_n \triangleright_S y_n$  by construction, so the point is to show  $x'_i \triangleright_S x'_{i+1}$  for each i. Assume  $z \in S$  and  $z \preceq x'_i x'_{i+1}$ . This implies  $z \preceq x'_i x'_{i+1} y_{i+1}$ , i.e.,  $z \preceq y_{i-1} x_i x_{i+1}$ . By hypothesis, we have  $x_i \triangleright_S x_{i+1}$ , hence, by Lemma 7.8(i),  $y_{i-1} x_i \triangleright_S x_{i+1}$ . So  $z \preceq y_{i-1} x_i x_{i+1}$  implies  $z \preceq y_{i-1} x_i$ , i.e.,  $z \preceq x'_i y_i$ . Now, by hypothesis, we have  $x'_i \triangleright_S y_i$ , so we deduce  $z \preceq x'_i$ , hence  $x'_i \triangleright_S x'_{i+1}$ , and the sequence  $(x'_1, \ldots, x'_n, y_n)$  is S-prenormal.

If y and each  $x_i$  are S-simple, we can inductively assume that  $y_{i-1}$  and  $x'_i$  are S-simple: indeed, in this case, Prop. 7.11 guarantees that  $y_{i-1}x_i$  admits a S-normal form of length 2 at most, and, if we define  $(x'_i, y_i)$  to be such a S-normal sequence (with possibly  $y_i = 1$ ), then induction continues.




FIGURE 7.2. Comparing normal forms of x and yx

Let us finally consider multiplication on the right. A similar argument is possible, but it works in the Gaussian case only.

LEMMA 7.13. Assume that M is a Gaussian monoid, S quasi-spans M. Let x, y be arbitrary elements of M, and  $(x_1, \ldots, x_n)$  be a S-prenormal decomposition of xy. Put  $y_n = y$ , and, inductively, define  $x_i'$  and  $y_{i-1}$  so that  $y_{i-1}x_i = x_i'y_i$  holds and the latter is a left lcm of  $x_i$  and  $y_i$ . Then  $(x_1', \ldots, x_n')$  is a S-prenormal decomposition of x.

PROOF. (Fig. 7.2 again) Let us show that  $x_i \triangleright_S x_{i+1}$  implies  $x_i' \triangleright_S x_{i+1}'$ . Assume  $z \in S$  and  $z \preceq x_i' x_{i+1}'$ . As in the proof of Lemma 7.12, we deduce  $z \preceq y_{i-1} x_i x_{i+1}$ , hence, as  $x_i \triangleright_S x_{i+1}$  implies  $y_{i-1} x_i \triangleright_S x_{i+1}$ ,  $z \preceq y_{i-1} x_i$ , i.e.,  $z \preceq x_i' y_i$ . By construction,  $y_i$  and  $x_{i+1}'$  have no common divisor; in the Gaussian case, this implies that every common divisor of  $x_i' y_i$  and  $x_i' x_{i+1}'$  is a divisor of  $x_i'$ . So, we have  $z \preceq x_i'$ , and  $x_i' \triangleright_S x_{i+1}'$ .

Finally, we observe that  $y_0 = 1$  necessarily holds, as, by the results of [9],  $y_0x_1 \cdots x_n$  has to be a left lcm of  $x_1 \cdots x_n$  and y, hence to equal  $x_1 \cdots x_n$ .

EXAMPLE 7.14. When lcm's need not exist, the previous argument fails, and so does the result itself. For instance, let us consider  $M_1$  again. Choose  $x=a^3$ , y=b. Then  $(a^2,a)$  is a normal decomposition of x. One possibility according to Lemma 7.13 is to define  $y_0=1$ ,  $y_1=b$ ,  $x_1'=b$ ,  $x_2'=a$ . Indeed, ab=ba is a left mcm of a and b, and  $1a^2=bb$  is a left mcm of  $a^2$  and b. Now, (b,a) is not a (pre)normal sequence, as we have  $a \in S$ ,  $a \leq ba$ , and  $a \not\leq b$ , hence  $b \not\geqslant_S a$ .

### 8. Garside elements

As was recalled above, if M is a thin Gaussian monoid, *i.e.*, a Garside monoid, then the lcm  $\Delta$  of all primitive elements plays an important rôle. Technically, the point is that the left divisors of  $\Delta$  coincide with its right divisors, which implies in particular that conjugation by  $\Delta$  gives an automorphism of M, and that some power of  $\Delta$  belongs to the center of M. Conversely, it is proved in [9] that, if M is a Gaussian monoid and  $\Delta$  is an element of M such that the left divisors of  $\Delta$  coincide with its right divisors and they generate M, then these divisors of  $\Delta$  span M, and, therefore, M is thin, and, therefore, it is a Garside monoid.

In the general case, there seems to be no reason why the existence of a finite spanning set should imply the existence of an element  $\Delta$  with similar properties. Even worse, Prop. 6.1 shows that the existence of such an element is impossible in the non-Gaussian case if we require both closure under mcm and left divisors.

However, we shall see now how to define an appropriate notion of a Garside element which may exist in the non-Gaussian case, and extends the usual notion in the Gaussian case. We shall then prove in the general case a large part of the results established in the Gaussian case.

DEFINITION 8.1. Assume that M is a cancellative monoid. We say that an element  $\Delta$  of M is a Garside element if  $Div(\Delta)$  is a finite spanning subset of M.

Notice that, if  $\Delta$  is a Garside element in M, then M must be thin by definition, hence quasi-atomic by Prop. 5.7, and every primitive element of M must divide  $\Delta$ , since, by Prop. 5.14, the family  $P_M$  is the least spanning subset of M, and, therefore, it must be included in  $\mathrm{Div}(\Delta)$ . Let us mention that most of the subsequent results could be extended to a *quasi-Garside* element, the latter being defined as an element  $\Delta$  such that  $\mathrm{Div}(\Delta)$  spans M and is quasi-finite.

Lemma 8.2. Assume that M is a thin cancellative monoid, and  $\Delta$  is a Garside element in M. Then, for every element x in  $\mathrm{Div}(\Delta)$ , there exists a unique element  $x^*$  in  $\mathrm{Div}(\Delta)$  satisfying  $xx^* = \Delta$ ; the mapping  $x \mapsto x^*$  is a permutation of  $\mathrm{Div}(\Delta)$ ; for  $x, y \in \mathrm{Div}(\Delta)$ , x being a left divisor of y is equivalent to  $y^*$  being a right divisor of  $x^*$ .

PROOF. (The argument was already used for Prop. 6.1.) By definition  $x \leq \Delta$  means that  $xx^* = \Delta$  holds for some right divisor  $x^*$  of  $\Delta$ , which is unique as M is assumed to be (left) cancellative. By hypothesis, the family  $\mathrm{Div}(\Delta)$  spans M which contains  $\Delta$ , so, by Prop. 2.7, it also contains every right divisor of  $\Delta$ , so, in particular,  $x^*$  belongs to  $\mathrm{Div}(\Delta)$ . Then  $x^* = y^*$  implies  $xx^* = \Delta = yy^* = yx^*$ , hence x = y, as M is cancellative. This proves that  $x \mapsto x^*$  is an injection of  $\mathrm{Div}(\Delta)$  into itself, hence a bijection as  $\mathrm{Div}(\Delta)$  is assumed to be finite. Finally y = xz implies  $xx^* = \Delta = yy^* = xzy^*$ , hence  $x^* = zy^*$ .

We deduce that, in the Gaussian case, our current notion of a Garside element coincides with that considered in [9]:

Lemma 8.3. Assume that M is a thin cancellative monoid.

- (i) If  $\Delta$  is a Garside element in M, then the left and the right divisors of  $\Delta$  coincide and they generate M.
- (ii) Conversely, if M is Gaussian and  $\Delta$  is an element of M such that the left and the right divisors of  $\Delta$  coincide and they generate M, then  $\Delta$  is a Garside element in M.
- PROOF. (i) Assume that  $\Delta$  is Garside. By Prop. 2.7, every right divisor of  $\Delta$  belongs to  $\mathrm{Div}(\Delta)$ , hence is a left divisor of  $\Delta$ , while, by Lemma 8.2, every element of  $\mathrm{Div}(\Delta)$  belongs to the range of the mapping  $x \mapsto x^*$ , hence it is a right divisor of  $\Delta$ : so the left and the right divisors of  $\Delta$  coincide.
- (ii) Assume now that M is Gaussian and the left and the right divisors of  $\Delta$  coincide and they generate M. Assume  $x, y \leq \Delta$  and xy'' = yx''. Let xy' = yx' be the lcm of x and y. By definition, we have  $x' \leq x''$  and  $y' \leq y''$ . Moreover,  $x, y \leq \Delta$  implies  $xy' \leq \Delta$ . So xy' and yx' are left divisors of  $\Delta$ , hence they are right divisors of  $\Delta$  as well, and so are y' and x'. Finally, x' and y' belong to  $\text{Div}(\Delta)$ , and the latter spans M. So  $\Delta$  is a Garside element.

In the thin Gaussian case, *i.e.*, in a Garside monoid, there always exists a unique minimal Garside element, namely the lcm of all primitive elements. In the general case, we have no such result, but the following examples show that Garside elements may still exist.

EXAMPLE 8.4. Consider again the monoid  $M_1$  of Example 2.11. Let  $\Delta_1 = a^2$  and  $\Delta_2 = ab$ . Then  $\Delta_1$  and  $\Delta_2$  both are minimal Garside elements. For instance,

we have  $\text{Div}(\Delta_1) = \{1, a, b, a^2\}$ , a spanning subset of  $M_1$ , and the left and the right divisors of  $\Delta_1$  coincide. Observe that, in this case, the divisors of  $\Delta_1$  properly include the primitive elements.

The reader can check similarly that the monoid  $M_2$  contains three minimal Garside elements, namely  $a^2$ , ab, and ac, while  $M_3$  contains one minimal Garside element only, namely  $b^2$ .

Proposition 8.5. Assume that M is a thin cancellative monoid, and  $\Delta$  is a Garside element in M. The mapping  $x \mapsto x^{**}$  extends into an automorphism  $\phi_{\Delta}$  of M and we have

$$(8.1) x\Delta = \Delta\phi_{\Delta}(x)$$

for every x in M. The automorphism  $\phi_{\Delta}$  globally preserves  $\mathrm{Div}(\Delta)$ , the units, the atoms, the primitive elements, and the simple elements of M. The order of  $\phi_{\Delta}$  is a finite integer e, and the element  $\Delta^e$  belongs to the center of M, which therefore is not trivial.

PROOF. By Lemma 8.2, the mapping  $x \mapsto x^{**}$  is a permutation of  $\operatorname{Div}(\Delta)$ , and it has a finite order say e. By definition, we have  $x\Delta = x(x^*x^{**}) = (xx^*)x^{**} = \Delta x^{**}$  for every x in  $\operatorname{Div}(\Delta)$ . Assume  $x_1 \cdots x_p = y_1 \cdots y_q$  with  $x_1, \ldots, y_q \in \operatorname{Div}(\Delta)$ . Using the previous remark, we obtain

$$\Delta x_1^{**} \cdots x_p^{**} = x_1 \cdots x_p \Delta = y_1 \cdots y_q \Delta = \Delta y_1^{**} \cdots y_q^{**},$$

hence  $x_1^{**} \cdots x_p^{**} = y_1^{**} \cdots y_q^{**}$  by cancelling  $\Delta$ . Thus putting  $\phi_{\Delta}(x_1 \cdots x_p) = x_1^{**} \cdots x_p^{**}$  yields a well defined mapping. As  $\operatorname{Div}(\Delta)$  generates M, the mapping  $\phi_{\Delta}$  is defined everywhere on M, and, by construction, it is an endomorphism and (8.1) is satisfied. Then,  $\phi_{\Delta}^e$  is also an endomorphism, and it is the identity on  $\operatorname{Div}(\Delta)$ , so it is the identity everywhere. Hence  $\phi_{\Delta}$  must be an automorphism. Moreover, (8.1) inductively implies  $x\Delta^k = \Delta^k \phi_{\Delta}^k(x)$  for every positive k and every x, so, in particular,  $x\Delta^e = \Delta^e x$  for every x, i.e.,  $\Delta^e$  commutes with every element of M. Finally, we apply Prop. 6.9.

EXAMPLE 8.6. Different Garside elements may give rise to different automorphisms. For instance, in  $M_2$ , the automorphism  $\phi_{a^2}$  is the identity, while  $\phi_{ab}$  and  $\phi_{ac}$  have order 3, and they correspond to the cyclic permutations (a, c, b) and (a, b, c) of the atoms respectively.

PROPOSITION 8.7. Assume that M is a thin cancellative monoid, and  $\Delta$  is a Garside element in M. Then any two elements of M admit a common multiple; more precisely, for  $x \in \text{Div}(\Delta)^p$  and  $y \in \text{Div}(\Delta)^q$ , we have xy' = yx' for some x' in  $\text{Div}(\Delta)^p$  and y' in  $\text{Div}(\Delta)^q$ .

PROOF. The proof of Prop. 4.2 shows that, if S spans M and any two elements of S admit a common multiple, then two elements x of  $S^p$  and y of  $S^q$  admit a common multiple xy' = yx' with  $x' \in S^p$  and  $y' \in S^q$ . Here we apply the result to the spanning subset  $\text{Div}(\Delta)$ . The only point to check is the result in the case p = q = 1. Now, for  $x \leq \Delta$  and  $y \leq \Delta$ , we can take  $x' = y^*$  and  $y' = x^*$ .

For a while let us write  $Div_r(x)$  for the set of all right divisors of x.




Figure 8.1. Divisors of  $\Delta^k$ 

Lemma 8.8. Assume that M is a thin cancellative monoid, and  $\Delta$  is a Garside element in M. Then, for every positive integer k, we have  $\mathrm{Div}(\Delta^k) = \mathrm{Div}_r(\Delta^k) = \mathrm{Div}(\Delta)^k$ , and, therefore,  $\Delta^k$  is a Garside element.

PROOF. We prove the three relations  $\mathrm{Div}_r(\Delta^k) \subseteq \mathrm{Div}(\Delta)^k \subseteq \mathrm{Div}(\Delta^k) \subseteq \mathrm{Div}_r(\Delta^k)$ . First, by Prop. 2.6, the set  $\mathrm{Div}(\Delta)^k$  spans M, and it contains  $\Delta^k$ , so, by Prop. 2.7, it also contains every right divisor of  $\Delta^k$ .

The second inclusion is proved using induction on k. The result is trivial for k=1. Assume  $k\geq 2$ , and let  $x\in \mathrm{Div}(\Delta)^k$ , say  $x=x_1x'$  with  $x_1\preceq \Delta$  and  $x'\in \mathrm{Div}(\Delta)^{k-1}$  (Fig. 8.1). By construction,  $x_1^*$  belongs to  $\mathrm{Div}(\Delta)$ , so, by Proposition 8.7, we have  $x'y=x_1^*x''$  for some  $y\in \mathrm{Div}(\Delta)$  and  $x''\in \mathrm{Div}(\Delta)^{k-1}$ . By induction hypothesis, we have  $x''\preceq \Delta^{k-1}$ , and, therefore,

$$x \leq xy = x_1 x' y = x_1 x_1^* x'' = \Delta x'' \leq \Delta \Delta^{k-1} = \Delta^k$$
.

For the third inclusion, assume  $x \leq \Delta^k$ , say  $xy = \Delta^k$ . We find

$$\phi_{\Delta}^{-k}(y)xy = \phi_{\Delta}^{-k}(y)\Delta^k = \Delta^k\phi_{\Delta}^k(\phi_{\Delta}^{-k}(y)) = \Delta^ky,$$

hence  $\phi_{\Delta}^{-k}(y)x = \Delta^k$ , which shows that x is a right divisor of  $\Delta^k$ .

Proposition 8.9. Assume that M is a thin cancellative monoid, and  $\Delta$  is a Garside element in M. Then any two elements of M admit a common left multiple.

PROOF. Assume  $x, y \in M$ . Then both x and y belong to  $Div(\Delta)^k$  for k large enough. By Lemma 8.8, this implies that  $\Delta^k$  is both a common right multiple, and a common left multiple of x and y.

If a thin cancellative monoid M contains a Garside element, then, by Proposition 8.7, it is a Ore monoid, and, therefore, it embeds in a thin group of (right) fractions G. Using the Garside element, we can also express every element of G as a left fraction whose denominator is a power of  $\Delta$ .

PROPOSITION 8.10. Assume that M is a thin cancellative monoid, and  $\Delta$  is a Garside element in M. Then M embeds in a group of fractions G; every element of G admits a unique decomposition  $\Delta^{-k}x$  with  $k \in \mathbb{Z}$  and  $x \in M$  satisfying  $\Delta \not\preceq x$ .

PROOF. Let  $z=xy^{-1}$  be an element of G. As y belongs to  $\mathrm{Div}(\Delta)^\ell$  for some positive  $\ell$ , we have  $y \leq \Delta^\ell$  by Lemma 8.8, say  $yx' = \Delta^\ell$ . Then we find

$$z = xx'x'^{-1}y^{-1} = xx'\Delta^{-\ell} = \Delta^{-\ell}\phi_{\Delta}^{-\ell}(xx'),$$

i.e.,  $z = \Delta^{-\ell} z_0$  for some  $z_0$  in M. Assume  $p \leq \ell$  and  $y = \Delta^p z \in M$ . Then, in M, we have  $z_0 = \Delta^{\ell-p} y$ , hence  $\ell - p \leq ||z_0||$ . Thus the set  $\{p \in \mathbb{Z} : \Delta^p z \in M\}$  must have a least element, say k. Then, by construction, z can be expressed as

 $\Delta^{-k}x$  for some x in M. As any relation  $x = \Delta x'$  in M would imply  $z = \Delta^{-k+1}x'$  and contradict the definition of k, we have  $\Delta \not \preceq x$ . Finally,  $\Delta^{-k}x = \Delta^{-k'}x'$  with k' > k implies  $\Delta^{-k+k'}x = x'$  in M, hence  $\Delta \preceq x'$ , showing the uniqueness of the decomposition  $\Delta^{-k}x$  when  $\Delta \not \preceq x$  is required.

If  $\Delta$  is a Garside element in a monoid M, then, by hypothesis, the set  $\mathrm{Div}(\Delta)$  spans M, and, therefore, there exist the associated notions of a  $\mathrm{Div}(\Delta)$ -simple element and a  $\mathrm{Div}(\Delta)$ -normal sequence. For simplicity, we call them  $\Delta$ -simple and  $\Delta$ -normal respectively, and we write  $\triangleright_{\Delta}$  for  $\triangleright_{\mathrm{Div}(\Delta)}$ . Using Prop. 8.10 and the results of Sec. 7, we obtain:

PROPOSITION 8.11. Assume that M is a thin cancellative monoid,  $\Delta$  is a Garside element in M, and G is the group of fractions of M. Then every element of G admits a decomposition  $\Delta^{-k}x_1\cdots x_p$  where k is a uniquely determined integer and  $(x_1,\ldots,x_p)$  is a  $\Delta$ -normal sequence with  $x_1 \not\simeq \Delta$ .

PROOF. The only point to establish is that, if  $(x_1, \ldots, x_p)$  is a  $\Delta$ -normal sequence, then  $\Delta \not\preceq x_1 \cdots x_p$  is equivalent to  $x_1 \not\simeq \Delta$ . The condition is obviously necessary. Conversely, as  $\Delta$  belongs to  $\mathrm{Div}(\Delta)$ , the relation  $\Delta \preceq x_1 \cdots x_p$  implies  $\Delta \preceq x_1$  by definition of a  $\Delta$ -normal sequence. Now, as  $\Delta$  is divisible by every element of  $\mathrm{Div}(\Delta)$ , no proper multiple of  $\Delta$  may be  $\Delta$ -simple, and  $\Delta \preceq x_1$  implies  $\Delta \simeq x_1$  when  $x_1$  is  $\Delta$ -simple.

EXAMPLE 8.12. Even if we use a minimal Garside element, the  $\Delta$ -normal form need not coincide with the  $(P_M)$ -normal form in general. For instance, consider once more the monoid  $M_1$  of Example 2.11. We have seen that  $\Delta_1 = a^2$  is a minimal Garside element in  $M_1$ . Then the  $\Delta_1$ -simple elements coincide with the simple elements: there are five of them, namely 1, a, b,  $a^2$ , and ab. Now, the relations  $\triangleright$  and  $\triangleright_{\Delta_1}$  do not coincide, because we have  $\mathrm{Div}(\Delta_1) = P_{M_1} \cup \{\Delta_1\}$ . It follows that the  $\Delta_1$ -simple elements are determined by their divisors in  $\mathrm{Div}(\Delta_1)$ , while they are not determined by their primitive divisors: both  $a^2$  and ab are divisible by 1, a, b, but only  $a^2$  is divisible by  $a^2$ . As a consequence, the  $\Delta_1$ -normal form is unique, while we have seen the normal form is not.

#### 9. Automatic structure

In the Gaussian case, *i.e.*, when lcm exist, thinness implies the existence of a Garside element, and the latter implies the existence of an automatic structure for the associated group of fractions. We shall show now that the latter result extends to more general cases: indeed, we shall prove that, under suitable hypotheses, the normal form of Prop. 8.11 is associated with an automatic structure.

The first steps, namely proving that the normal decompositions make a regular language, are easy.

PROPOSITION 9.1. Assume that M is a thin cancellative monoid,  $\Delta$  is a Garside element in M, and G is the group of fractions of M. Let  $\Sigma_{\Delta}$  denote the set of all  $\Delta$ -simple elements in M. Then the language consisting of all normal sequences in the sense of Prop. 8.11 is regular.

PROOF. By Prop. 6.7, there are finitely many  $\Delta$ -simple elements, *i.e.*, the set  $\Sigma_{\Delta}$  is finite. Put  $A = \Sigma_{\Delta} \cup \{\Delta^{-1}\}$ . A word over A, *i.e.*, a finite sequence

 $(x_1, \ldots, x_n)$  of letters, is a normal form if and only if the following requirements are obeyed:

- a letter  $\Delta^{-1}$  or  $\Delta$  cannot follow any other letter;
- a letter x in  $\Sigma_{\Delta} \{\Delta\}$  may follow only  $\Delta^{-1}$  or one of those (finitely many) letters y in  $\Sigma_{\Delta}$  that satisfy  $y \triangleright_{\!\!\! \Delta} x$ .

Define a state set Q to be  $A \cup \{1, \bot\}$ , where 1 is an initial state and  $\bot$  is a failure state, and a transition function  $F: Q \times A \to Q$  by

$Q \downarrow A \rightarrow$	$x \neq \Delta, \Delta^{-1}$	Δ	$\Delta^{-1}$
$y \neq \Delta, \Delta^{-1}$		1	Т
Δ	x	Δ	
$\Delta^{-1}$	x		$\Delta^{-1}$
1	x	Δ	$\Delta^{-1}$

Then the finite state automaton  $(Q, A, F, 1, Q - \{\bot\})$  recognizes the language of  $\Delta$ -normal forms (see for instance [14] for definitions).

Provided all  $\Delta$ -normal forms have the same length, we can readily apply the method of [3] or [19], and deduce:

Corollary 9.2. Assume that G is the group of fractions of a cancellative monoid M that admits a Garside element  $\Delta$  such that all  $\Delta$ -normal forms of an element have the same length. Then G has rational growth, i.e., the number of elements of G with a  $\Delta$ -normal form of length n is a rational function of n.

If G is a group generated by a family A, we denote by  $\Gamma_A(G)$  the Cayley graph of G with respect to A, *i.e.*, the labelled graph whose vertices are the elements of G and there exists a z-labelled edge from x to y if y = xz holds in G. For  $x, y \in G$ , the distance  $\operatorname{dist}_{A,G}(x,y)$  between x and y in  $\Gamma_A(G)$  is the minimal length of an unoriented path from x to y.

DEFINITION 9.3. Assume that G is a group generated by A. The *synchronous distance* between two words on A, *i.e.*, two sequences of letters in A, say  $(x_1, \ldots, x_p)$  and  $(y_1, \ldots, y_q)$ , is defined to be the supremum of the numbers

$$\operatorname{dist}_{A,G}(x_1\cdots x_{\inf(i,p)},y_1\cdots y_{\inf(i,q)})$$

for  $1 \le i \le \sup(p, q)$ .

By the results of [14], the  $\Delta$ -normal form of Prop. 8.11 is associated with a (left) automatic structure if and only if the fellow traveller property (FTP) is satisfied, *i.e.*, for every x in the group and every y in  $\Sigma_{\Delta} \cup \{\Delta^{-1}\}$ ,

- the synchronous distance between any two  $\Delta\text{-normal}$  decompositions of x is uniformly bounded, and
- the synchronous distance between a  $\Delta$ -normal decomposition of x and one of yx is uniformly bounded.

We shall see that such conditions are satisfied in good cases. To this end, we shall first establish a bound for the distance between the various normal forms of an element in the monoid. (The notion of the synchronous distance is extended to the case of the monoid in the obvious way.)

Lemma 9.4. Assume that M is a quasi-atomic cancellative monoid, S is a quasi-spanning subset of M of cardinality k and, for every x in M, the following condition holds:

(9.1) All S-normal decompositions of x have the same length.

Then the synchronous distance between any two S-normal decompositions of an element of M is uniformly bounded by 2(k-1).

We begin with two auxiliary results.

LEMMA 9.5. Assume that M is a (left) cancellative monoid, and S quasispans M. Then  $x_1 \triangleright_S \cdots \triangleright_S x_k \triangleright_S x$  implies  $x_1 \cdots x_k \triangleright_{S^k} x$ .

PROOF. We use induction on  $k \geq 0$ . Assume  $z \in S^k$  and  $z \leq x_1 \cdots x_k x$ . For k = 0, *i.e.*, for z = 1, the result is vacuously true. Otherwise, write  $z = z_1 z'$ , with  $z_1 \in S$  and  $z' \in S^{k-1}$ . By Lemma 7.8(ii),  $x_1 \triangleright_S \cdots \triangleright_S x_k \triangleright_S x$  implies  $x_1 \triangleright_S x_2 \cdots x_k x$ . By hypothesis, we have  $z_1 \leq x_1 \cdots x_k x$ , hence  $z_1 \leq x_1$ , say  $x_1 = z_1 x'_1$ . Then, by Lemma 7.8(i), we have  $x'_1 x_2 \triangleright_S x_3 \triangleright_S \cdots \triangleright_S x_k \triangleright_S x$ , and, as M is (left) cancellative,  $z' \leq (x'_1 x_2) x_3 \cdots x_k x$ . By induction hypothesis, this implies  $z' \leq (x'_1 x_2) x_3 \cdots x_k$ , hence  $z = z_1 z' \leq z_1 (x'_1 x_2) x_3 \cdots x_k$ , i.e.,  $z \leq x_1 \cdots x_k$ .

LEMMA 9.6. Under the hypotheses of Lemma 9.4, if  $(x_1, \ldots, x_n)$  is a S-normal decomposition for x, and  $x'_1$  is a maximal S-simple divisor of x, then there exist  $x'_2, \ldots, x'_k$  such that  $(x'_1, \ldots, x'_k, x_{k+1}, \ldots, x_n)$  is another S-normal decomposition of x.

PROOF. As  $x_1'$  is S-simple, it belongs to  $S^k$  by Prop. 6.7. By Lemma 9.5, we have  $x_1 \cdots x_k \triangleright_{S^k} x_{k+1} \cdots x_n$ , so  $x_1' \leq x$  implies  $x_1' \leq x_1 \cdots x_k$ , say  $x_1 \cdots x_k = x_1'y$ . Let  $(x_2', \dots, x_{k'}')$  be a S-normal decomposition of y. By hypothesis, we have  $\operatorname{Div}(x_1') \cap S = \operatorname{Div}(x) \cap S$ , hence  $x_1' \triangleright_S x_2' \cdots x_{k'}'$ , so  $(x_1', \dots, x_{k'}')$  is a S-normal sequence, hence another S-normal decomposition for  $x_1 \cdots x_k$ . Then Condition (9.1) implies k' = k.

Let us now consider the S-covering relation between  $x'_k$  and  $x_{k+1}$ . As in the proof of Prop. 7.11, let x' be a maximal S-simple divisor of  $x'_k x_{k+1}$  satisfying  $x'_k \leq x'$ . Write  $x' = x'_k z$ . Then  $x'_1 \cdots x'_k z$  equals  $x'_1 \cdots x'_{k-1} x'$ , so it belongs to  $S^k$ , and, therefore, by Prop. 7.11, it must admit at least one normal form of length k at most. On the other hand, we have  $x_k \triangleright_S x_{k+1}$  and  $z \leq x_{k+1}$ , hence  $x_k \triangleright_S z$ , so, if z is not invertible,  $(x_1, \ldots, x_k, z)$  is another S-normal decomposition of  $x'_1 \cdots x'_{k-1} x'$ . Condition (9.1) discards this possibility. Hence, z must be invertible, i.e., we must have  $x'_k \triangleright_S x_{k+1}$ . So the sequence  $(x'_1, \ldots, x'_k, x_{k+1})$  is S-normal, and, trivially, so is  $(x'_1, \ldots, x'_k, x_{k+1}, \ldots, x_n)$ .

PROOF OF LEMMA 9.4. Let  $(x_1, \ldots, x_n)$  and  $(x'_1, \ldots, x'_n)$  be two S-normal decomposition of an element x of M. Applying Lemma 9.6 to  $(x_1, \ldots, x_n)$  and to  $x'_1$ , we find  $x_{2,1}, \ldots, x_{k,1}$  so that  $(x'_1, x_{2,1}, \ldots, x_{k,1}, x_{k+1}, \ldots, x_n)$  is another S-normal decomposition of x. Then, applying Lemma 9.6 to the latter sequence and to  $x'_2$ , we find  $x_{3,2}, \ldots, x_{k+1,2}$  so that  $(x'_1, x'_2, x_{3,2}, \ldots, x_{k+1,2}, x_{k+2}, \ldots, x_n)$  is a S-normal decomposition of x. Similarly, having found a S-normal form  $(x'_1, \ldots, x'_i, x_{i+1,i}, \ldots, x_{i+k-1,i}, x_{i+k}, \ldots, x_n)$  for x, applying Lemma 9.6 to this sequence and to  $x'_{i+1}$  yields a new S-normal decomposition  $(x'_1, \ldots, x'_{i+1}, x_{i+2,i+1}, \ldots, x_{i+k,i+1}, x_{i+k+1}, \ldots, x_n)$ . Now, we read on Fig. 9.1 that, for each i, the distance between  $x_1 \cdots x_i$  and  $x'_1 \cdots x'_i$ 




FIGURE 9.1. Comparing two normal forms of x (here k=3)

is bounded above by 2(k-1), as  $x_1 \cdots x_{i+k-1}$  is a common multiple of these elements.

Applying the previous result to the case of monoids with a Garside element, we deduce:

Proposition 9.7. Assume that M is a thin cancellative monoid,  $\Delta$  is a Garside element in M with k divisors, and G is the group of fractions of M. Assume moreover that, for every x in M, the following condition holds:

(9.2) All  $\Delta$ -normal decompositions of x have the same length.

Then the synchronous distance between any two  $\Delta$ -normal decompositions of an element of G is uniformly bounded by 2(k-1).

PROOF. We have seen that, if  $(\Delta^{-1}, \ldots, \Delta^{-1}, x_1, \ldots, x_n)$ , k times  $\Delta^{-1}$ , and  $(\Delta^{-1}, \ldots, \Delta^{-1}, x'_1, \ldots, x'_{n'})$ , k' times  $\Delta^{-1}$ , are two  $\Delta$ -normal decompositions for some element z of G, then, necessarily, k = k' holds, and, therefore,  $(x_1, \ldots, x_n)$  and  $(x'_1, \ldots, x'_{n'})$  are two  $\Delta$ -normal decompositions for some element of M. Then we apply Lemma 9.4 with  $S = \text{Div}(\Delta)$ .

The case of left multiplication in the monoid has already been treated in Lemma 7.12, which gives:

Lemma 9.8. Assume that M is a quasi-atomic cancellative monoid, S quasi-spans M, and y is a S-simple element of M. Then, for every element x of M, and every S-normal decomposition of x, there exists a S-normal decomposition of yx at synchronous distance at most 1.

It remains to extend the result to the group of fractions.

Proposition 9.9. Assume that M is a thin cancellative monoid,  $\Delta$  is a Garside element in M with k divisors, G is the group of fractions of M, and y is a  $\Delta$ -simple element of M. Then, for every element z of G, and every  $\Delta$ -normal decomposition of z, there exists a  $\Delta$ -normal decomposition of yz at synchronous distance at most 3k.

PROOF. Assume first  $y \in \text{Div}(\Delta)$ . Assume  $z = \Delta^{-k}x$ , with  $x \in M$  and  $\Delta \not\preceq x$ . Then we have  $yz = \Delta^{-k}y'x$  with  $y' = \phi_{\Delta}^{-k}(y)$ . By Prop. 8.5, we have  $y' \preceq \Delta$ , so, in particular, y' is  $\Delta$ -simple, and we can apply Lemma 9.8 to y' and any  $\Delta$ -normal decomposition  $(x_1, \ldots, x_n)$  of x to obtain a  $\Delta$ -normal decomposition  $(x'_1, \ldots, x'_n, y'_n)$  of y'x. There remains one point to check: if it contains at least one  $\Delta^{-1}$ , the sequence  $(\Delta^{-1}, \ldots, \Delta^{-1}, x'_1, \ldots, x'_n, y'_n)$  is  $\Delta$ -normal only if  $x'_1$  is not  $\Delta$ : if  $x'_1 = \Delta$  holds, we must cancel  $x'_1$  with the last  $\Delta^{-1}$ , and repeat the

reduction until we possibly find  $x_i' \neq \Delta$ . As each such reduction increases the synchronous distance by 2, there could be a problem here. Actually, we shall prove that  $x_1' \simeq x_2' \simeq \Delta$  implies  $\Delta \leq x_1$ , hence  $x_1 \simeq \Delta$ . Here we use the hypothesis that y' is not only  $\Delta$ -simple, but also it is a divisor of  $\Delta$ . First,  $x_1' \simeq x_2' \simeq \Delta$  implies  $x_1'x_2' \simeq \Delta^2$ . Indeed, for  $u \in M^*$ , we have  $u\Delta = xv$  for some x and v satisfying  $x \in \text{Div}(\Delta)$  and  $v \in M^*$ , and  $||x|| = ||\Delta||$  implies  $x \simeq \Delta$ . So we deduce  $\Delta^2 \leq y'x_1x_2$ , i.e.,  $y'y'^*\Delta \leq y'x_1x_2$ , hence  $y'^*\Delta \leq x_1x_2$ , i.e.,  $\Delta\phi_{\Delta}(y'^*) \leq x_1x_2$  which implies  $\Delta \leq x_1x_2$ , and, finally,  $\Delta \leq x_1$  as  $x_1 \triangleright_{\Delta} x_2$  holds by hypothesis. So, at most one reduction  $\Delta^{-1}\Delta$  may occur, and the synchronous distance between the  $\Delta$ -normal form of x and that of yx is at most 3.

The result for an arbitrary  $\Delta$ -simple element y follows, as, by Prop. 6.7, any such element is the product of at most k elements of  $Div(\Delta)$ .

Putting Propositions 9.1, 9.7, and 9.9 together, we deduce

PROPOSITION 9.10. Assume that G is the group of fractions of a cancellative monoid M that admits a Garside element  $\Delta$  such that all  $\Delta$ -normal forms of an element have the same length. Then G is an automatic group.

The previous result applies in particular to every thin Gaussian group, i.e., to every Garside group—hence in particular to every spherical Artin group. But non-Gaussian groups are also eligible:

EXAMPLE 9.11. Consider once more the groups  $G_1$  and  $G_3$  of Example 4.4. We have seen in Example 8.12 that the monoid  $M_1$  contains a Garside element  $\Delta$  such that the  $\Delta$ -simple elements are determined by their divisors in  $\text{Div}(\Delta)$ . So the associated  $\Delta$ -normal form is unique, and, therefore, the length requirement is satisfied. The argument is similar for  $M_3$ . So the groups  $G_1$  and  $G_3$  are automatic.

The case of  $G_2$  is slightly different. Indeed, in the monoid  $M_2$ ,  $a^2$  is a Garside element, but ab and ac are  $a^2$ -simple elements with the same divisors in  $Div(a^2)$ , namely 1, a, b, c. Now, we have the following sufficient condition:

PROPOSITION 9.12. Assume that M is a thin cancellative monoid with no non-trivial unit,  $\Delta$  is a Garside element in M, and the following condition holds in M: If x and x' are distinct  $\Delta$ -simple elements with the same divisors in  $\mathrm{Div}(\Delta)$ , then every common multiple of x and x' is a multiple of some  $\Delta$ -simple common multiple of x and x'. Then the  $\Delta$ -normal form is unique, and, therefore, the group of fractions of M is automatic.

PROOF. It suffices to show that, for every x in M, there exists a unique  $\Delta$ -simple element  $x_1$  with the same divisors as x in  $\mathrm{Div}(\Delta)$ . Now, assume that  $x_1$  and  $x_1'$  satisfy these conditions and are distinct. Then, by hypothesis, there exists a  $\Delta$ -simple element  $x_1''$  satisfying  $x_1 \preceq x_1'' \preceq x$  and  $x_1' \preceq x_1'' \preceq x$ . Then  $x_1' = x_1''$  would imply  $x_1 \prec x_1'$ , contradicting the  $\Delta$ -simplicity of  $x_1'$ . So we must have  $x_1 \prec x_1''$ , and, therefore,  $\mathrm{Div}(x_1) \cap \mathrm{Div}(\Delta) \neq \mathrm{Div}(x_1') \cap \mathrm{Div}(\Delta) \subseteq \mathrm{Div}(x) \cap \mathrm{Div}(\Delta)$ , which contradicts  $\mathrm{Div}(x_1) \cap \mathrm{Div}(\Delta) = \mathrm{Div}(x) \cap \mathrm{Div}(\Delta)$ .

EXAMPLE 9.13. The previous criterion applies to the monoid  $M_2$ : indeed, for  $\Delta_1 = a^2$ , the only problem with  $\Delta_1$  occurs with the  $\Delta_1$ -simple elements ab and ac. Now, every common multiple of ab and ac is a multiple of  $a^2$ , i.e., of  $\Delta_1$ . We deduce that  $G_2$  is automatic.

Let us conclude with some open questions.

QUESTION 9.14. If  $\Delta$  is a Garside element in a thin cancellative monoid M, do all  $\Delta$ -normal decompositions of a given element of M necessarily have the same length, *i.e.*, is the additional assumption of Prop. 9.10 superfluous?

In the Gaussian case, Lemma 7.13 gives a uniform bound for the synchronous distance between the normal form of x and that of xy when y is simple. It is then easy to deduce that the  $\Delta$ -normal form of Prop. 9.10 gives rise to a bi-automatic structure—alternatively, we can also replace in this case the dissymmetric form  $\Delta^{-k}x_1\cdots y_n$  with a symmetric one  $y_p^{-1}\cdots y_1^{-1}x_1\cdots x_n$  [18, 9]. In the general case, the argument fails, the behaviour of  $\Delta$ -normal form with respect to right multiplication remains unknown, and so does the existence of an automatic structure involving a symmetric fractionary decomposition (defining the latter in the non-Gaussian case seems to require a uniform bound for the distance between the possible various mcm's of two elements in the monoid).

QUESTION 9.15. Under the hypotheses of Prop. 9.10, is the group G bi-automatic?

(In the case of the groups  $G_1$ ,  $G_2$ ,  $G_3$  of Example 4.4, a simple specific argument gives a positive answer.)

By Prop. 8.7, common multiples must exist in every thin cancellative monoid admitting a Garside element. In the Gaussian case, i.e., when we assume not only that common multiples exist, but even that least common multiples exist, then the lcm of all primitive elements is a Garside element.

QUESTION 9.16. Does every thin cancellative monoid admitting common multiples contain a Garside element? More precisely, need every mcm of the primitive elements be a Garside element?

Finally, let us mention an open problem dealing with the Gaussian case:

QUESTION 9.17. Is every finitely generated Gaussian group thin, *i.e.*, is every finitely generated Gaussian group necessarily a Garside group?

### References

- S.I. Adyan, Fragments of the word Delta in a braid group, Mat. Zam. Acad. Sci. SSSR 36-1 (1984) 25-34; translated Math. Notes of the Acad. Sci. USSR; 36-1 (1984) 505-510.
- [2] J. Birman, K.H. Ko & S.J. Lee, A new approach to the word problem in the braid groups, Advances in Math. 139-2 (1998) 322-353.
- [3] M. Brazil, Monoid growth functions for braid groups, Int. J. Algebra & Comput. 1-2 (1991) 201–205.
- [4] E. Brieskorn & K. Saito, Artin-Gruppen und Coxeter-Gruppen, Invent. Math. 17 (1972) 245–271.
- [5] R. Charney, Artin groups of finite type are biautomatic, Math. Ann. 292-4 (1992) 671-683.
- [6] R. Charney, Geodesic automation and growth functions for Artin groups of finite type, Math. Ann. 301-2 (1995) 307-324.
- [7] A.H. Clifford & G.B. Preston, The Algebraic Theory of Semigroups, vol. 1, Amer. Math. Soc. Surveys 7, (1961).
- [8] P. Dehornoy, Groups with a complemented presentation, J. Pure Appl. Algebra 116 (1997) 115–137.
- [9] P. Dehornoy, Groupes de Garside, Ann. Sci. Ec. Norm. Sup., to appear; ArXiv math.GR/0111157.
- [10] P. Dehornoy, Complete positive group presentations, Preprint; ArXiv math.GR/0111275.

- [11] P. Dehornoy & L. Paris, Gaussian groups and Garside groups, two generalizations of Artin groups, Proc. London Math. Soc. 79-3 (1999) 569-604.
- [12] P. Deligne, Les immeubles des groupes de tresses généralisés, Invent. Math. 17 (1972) 273–302.
- [13] E. A. Elrifai & H. R. Morton, Algorithms for positive braids, Quart. J. Math. Oxford 45-2 (1994) 479-497.
- [14] D. Epstein & al., Word Processing in Groups, Jones & Bartlett Publ. (1992).
- [15] F. A. Garside, The braid group and other groups, Quart. J. Math. Oxford 20 No.78 (1969) 235–254.
- [16] R. C. Lyndon & P. E. Schupp, Combinatorial group theory, Springer (1977).
- [17] M. Picantin, The center of thin Gaussian groups, J. of Algebra 245-1 (2001) 92–122.
- [18] W. Thurston, Finite state algorithms for the braid group, Circulated notes (1988).
- [19] P. Xu, Growth of the positive braid groups, J. Pure Appl. Algebra 80 (1992) 197–215.

LABORATOIRE SDAD, MATHÉMATIQUES, UNIVERSITÉ DE CAEN BP 5186, 14032 CAEN, FRANCE  $E\text{-}mail\ address$ : dehornoy@math.unicaen.fr URL: //www.math.unicaen.fr/~dehornoy