

# THE DEFINITION OF A DIFFERENTIAL FORM ON A TANGENT STRUCTURE

Carlos A. Torre

Escuela de Matemática, Universidad de Costa Rica,  
San José, Costa Rica

*ABSTRACT.* Three definitions of a differential form on a tangent structure are considered. It is proved that the (covariant) definition given by Souriau (as a collection of forms indexed by the plaques) is equivalent to a smooth section of the corresponding vector bundle if the space does not have transverse points.

## 1. Introduction.

A diffeology on any set  $X$  was defined by Souriau [3,4] as a collection of plaques  $p: U \subset \mathbf{R}^n \rightarrow X$ , for any  $n \in \mathbf{N}$  which covers  $X$  and is closed under composition with smooth maps  $\phi: V \subset \mathbf{R}^m \rightarrow \mathbf{R}^n$  where  $U, V$  are open sets. In this context differential forms were defined as a collection of forms on open subsets of  $\mathbf{R}^n$ , indexed by the plaques. They were used to construct a picture of quantization on a class of coadjoint orbits of diffeomorphisms groups [1,2]. A tangent structure was defined in [7] for coadjoint orbits of diffeomorphisms groups, allowing a geometric definition of a differential form as a section of the corresponding vector bundle [5,6], and in this way allowing also the construction of a Poisson structure and precuantization on coadjoint orbits of diffeomorphisms groups. In this paper we consider the relation between the two definitions of a differential form mentioned above, and the more algebraic one as a multilinear alternating map on the  $C^\infty(X)$  module of vector fields on  $X$ .

In section 2 we review four definitions of a differential form on a finite dimensional manifold and recall the proof about the equivalence between them. In section 3 we recall the definition of a tangent structure and three definitions of a differential form on it: the global definition given algebraically, the local definition, as a collection of forms indexed by plaques, as defined by Souriau, and as a smooth section on the vector bundle of exterior forms. It is proved that the last two are equivalent if the space does not have transverse points.

## 2. A review of differential forms on manifolds.

An exterior  $k$ -form on a vector space  $V$  is an alternating  $k$ -linear map

$$\omega: V^k \rightarrow \mathbf{R}$$

The set of them is denoted  $\Lambda^k V^*$ , forming a vector space with the ordinary scalar multiplication and sum of functions and the set  $\Lambda V^* := \cup_{k \in \mathbf{N}} \Lambda^k V^*$  is an associative algebra with the wedge product:

$$\omega^k \wedge \omega^l(\xi_1, \dots, \xi_{k+l}) := \sum (-1)^\sigma \omega^k(\xi_{i_1}, \dots, \xi_{i_k}) \omega^l(\xi_{j_1}, \dots, \xi_{j_l})$$

where  $\omega^i \in \Lambda^i V^*$  for  $i = k, l$  and the sum is over all permutations  $(i_1, \dots, i_k, j_1, \dots, j_l)$  of  $(1, \dots, k+l)$  and  $\sigma$  is the order of the permutation. This operation is skew commutative:

$$\omega^k \wedge \omega^l = (-1)^{kl} \omega^l \wedge \omega^k$$

In particular if  $k = l = 1$  then

$$\omega_1 \wedge \omega_2(\xi, \eta) = \omega_1(\xi)\omega_2(\eta) - \omega_2(\xi)\omega_1(\eta)$$

and more generally

$$\omega_1 \wedge \cdots \wedge \omega_k(\xi_1, \dots, \xi_k) = \det(\omega_j(\xi_i))$$

In  $V = \mathbf{R}^n$  every exterior 1-form  $\omega$  is a scalar multiple of a projection along some vector  $y \in V$ :  $\omega(\xi) = (\xi, y)$ . In particular the projection along the  $i$ -th coordinate vector is denoted  $x_i$ . More generally, if  $P: V \rightarrow F$  is a projection along a  $k$ -dimensional subspace  $F$  with ortonormal base  $\{\beta_1, \dots, \beta_k\}$ , then the map

$$\omega_P^k(\xi_1, \dots, \xi_k) = \det(a_{ij})$$

where  $P(\xi_i) = \sum_{j=1}^k a_{ij}\beta_j$  defines a  $k$ -form on  $V$ . This is the oriented  $k$ -volume of the projection along  $F$  of the parallelepiped generated by  $\xi_1, \dots, \xi_k$ . Conversely, using linearity it is found that every exterior  $k$ -form is a linear combination of forms  $\omega_P^k$  where  $P$  runs over all projections along the  $k$ -dimensional coordinate subspaces and therefore the set

$$\{x_{i_1} \wedge \cdots \wedge x_{i_k} / \{i_1, \dots, i_k\} \subset \{1, \dots, n\}\}$$

is a base of  $\Lambda^k V^*$ . In particular if  $k = n$ , every exterior  $n$ -form is a scalar multiple of

$$\nu(\xi_1, \dots, \xi_n) = \det(\xi_{ij})$$

where  $\xi_i = \sum_{j=1}^n \xi_{ij}e_j$  and  $\{e_1, \dots, e_n\}$  is the standard base. This is the oriented volume of the parallelepiped with edges  $\xi_1, \dots, \xi_n$ . Every  $k$ -form with  $k > n$  is zero.

If  $\omega_1, \omega_2$  are projections along  $v_1, v_2$  then  $\omega_1 \wedge \omega_2(\xi_1, \xi_2)$  is the oriented area of the projection of the parallelepiped generated by  $\xi_1, \xi_2$  over the plane generated by  $v_1, v_2$ .

The *pullback* of  $\omega \in \Lambda^k V^*$  by a linear map  $L: W \rightarrow V$  is the exterior  $k$ -form  $L^*\omega \in \Lambda^k W^*$  defined by

$$L^*\omega(\xi_1, \dots, \xi_k) = \omega(L\xi_1, \dots, L\xi_k)$$

The pullback defines an algebra homomorphism and preserves composition of linear maps (in the appropriate order).

A *differential  $k$ -form* on an open set  $U \subset \mathbf{R}^n$  is an alternating  $C^\infty(U)$   $k$ -linear map

$$\omega: [\Gamma(U)]^k \rightarrow C^\infty(U)$$

where  $\Gamma(U)$  denotes the  $C^\infty(U)$ -module of smooth vector fields on  $U$ . Therefore, each  $x \in U$  defines  $\omega_x \in \Lambda^k V^*$  by

$$\omega_x(\xi_1, \dots, \xi_k) := \omega(\bar{\xi}_1, \dots, \bar{\xi}_k)(x)$$

where  $\bar{\xi}_1, \dots, \bar{\xi}_k$  are smooth vector fields such that  $\bar{\xi}_i(x) = \xi_i$  (they exist:  $\bar{\xi}_i(x) = \sum \xi_i \frac{\partial}{\partial x_i}$ ).

$\omega_x$  is well defined: if  $\xi(x_0) = 0$ , then using the  $C^\infty(U)$ -linearity and the expression  $\xi = \sum_j b_j \frac{\partial}{\partial x_j}$ , with  $b_j \in C^\infty(U)$  and  $b_j(x_0) = 0$  it follows that  $\omega(\xi)(x_0) = 0$ .

Conversely, by evaluation on the vector fields  $\frac{\partial}{\partial x_j}$ , it follows that an exterior  $k$ -form

$$\omega_x = \sum_{|I|=k} a_{i_1 \dots i_k}(x) x_{i_1} \wedge \cdots \wedge x_{i_k}$$

for each  $x \in U$  defines a differential  $k$ -form on  $U$  iff each  $a_{i_1 \dots i_k} \in C^\infty(U)$ . In particular, the  $k$ -forms defined by  $x_{i_1} \wedge \cdots \wedge x_{i_k}$  for each  $x \in U$  is denoted  $dx_{i_1} \wedge \cdots \wedge dx_{i_k}$ , since  $dx_i$  denoted the 1-form

dual of the vector field  $\frac{\partial}{\partial x_i}$  and  $dx_i$  is also the differential of the projection along the  $i$ -axis (denoted  $x_i$  previously). Therefore, we have two equivalent definitions of a  $k$ -form on  $U$ .

If  $f: V \subset \mathbf{R}^m \rightarrow U$  is smooth, the *pullback* of a  $k$ -form  $\omega$  on  $U$  is defined using the differential of  $f$ :

$$f^*\omega(\xi_1, \dots, \xi_k)(x) = \omega_{f(x)}(df(\xi_1)(x), \dots, df(\xi_k)(x))$$

**Definition 2.1.** A  $k$ -form on a smooth manifold  $M$  given by an atlas  $\mathcal{A} = \{(\alpha_i, U_i)/i \in I\}$  is defined in four different ways (at least):

1. It is an exterior  $k$ -form  $\omega_x$  on  $T_x M$  for each  $x \in M$  such that if  $\xi_i \in \Gamma(M)$  for  $i = 1, \dots, k$ , then

$$\omega(\xi_1, \dots, \xi_k)(x) := \omega_x(\xi_1(x), \dots, \xi_k(x))$$

is smooth.

2. It is a  $C^\infty(M)$   $k$ -linear alternating map

$$\omega: [\Gamma(M)]^k \rightarrow C^\infty(M)$$

3. It is a collection  $(\omega_i)_{i \in I}$  of  $k$ -forms, where  $\omega_i$  is a  $k$ -form on  $U_i \subset \mathbf{R}^n$  such that

$$\omega_j = (\alpha_i^{-1} \circ \alpha_j)^* \omega_i$$

for each  $i, j \in I$ .

4. It is a smooth section on the vector bundle of exterior  $k$ -forms on  $M$ .

The four definitions are equivalent:  $1 \Rightarrow 2$  follows from evaluation at a point.  $2 \Rightarrow 1$  follows as follows: from a vector  $\xi_i = \sum \xi_{ij} \frac{\partial}{\partial x_j} \in T_{x_0} M$  construct a vector field  $\bar{\xi}_i(x) = \sum \xi_{ij} \frac{\partial}{\partial x_j}$  in a neighborhood of  $x_0$  and extend globally through multiplication by a smooth function  $g$  such that  $x_0 \in V \subset g < U$  (that is:  $\text{Supp } g \subset U$ ,  $g = 0$  on  $V$  and  $V$  is a neighborhood of  $x_0$  such that  $\bar{V} \subset U$ ) and defining  $\bar{\xi}_i(x) = 0$  outside  $U$ . Then  $\omega_x$  is defined using the vector fields  $\bar{\xi}_i$ . In order to prove that the definition is independent of this particular choice of  $\bar{\xi}_i$ , assume that  $\xi(x_0) = 0$ , take  $U, g$  as before and a local expression  $\xi(x) = \sum b_j(x) \frac{\partial}{\partial x_j}$  on  $U$ . Then

$$\omega(\xi)(x_0) = g(x_0) \sum b_j(x_0) \frac{\partial}{\partial x_j} = 0$$

$1 \Rightarrow 3$  is obtained as follows: define  $\omega_i = \alpha_i^* \omega$ , use  $\alpha_j = \alpha_i \circ (\alpha_i^{-1} \circ \alpha_j)$  and the fact that the pullback preserves composition and  $d\alpha_i$  preserves smoothness of vector fields. In order to prove  $3 \Rightarrow 1$ , for each  $x \in M$  take  $i \in I$  such  $x \in U_i$  and define  $\omega_x = (\alpha_i^{-1})^* \omega_i(x)$  which does not depend on the chart chosen. Finally  $1 \iff 4$ : the vector bundle is given by  $\wedge^k T^* M \rightarrow M$  where

$$\wedge^k T^* M = \{(x, \xi)/x \in M, \xi \in \wedge^k T_x^* M\}$$

$\pi(x, \xi) = x$  and with the atlas

$$\{(\pi^{-1}(U_i), (\alpha_i \times id) \circ \phi_i)/i \in I\}$$

where  $\phi_i: \pi^{-1}(U_i) \rightarrow U_i \times \mathbf{R}^k$  is given by  $\phi_i(x, \xi) := (x, \xi_{i_1 \dots i_k}(x))$  such that  $\xi = \sum \xi_{i_1 \dots i_k} dx_{i_1} \wedge \dots \wedge dx_{i_k}$ . Now, given  $\omega$  under definition 1, define  $\bar{\omega}$  under definition 4 (and viseversa) by

$$\bar{\omega}(x)(\xi_1(x), \dots, \xi_k(x)) = (x, \omega_x(\xi_1(x), \dots, \xi_k(x)))$$

The smoothness of  $\bar{\omega}$  and  $\omega$  are equivalent using the definition of smoothness with the local expression of these forms. This proves the equivalence of the four definitions.

The space of  $k$ -forms on  $M$  is denoted  $\epsilon^k(M)$ . It is a  $C^\infty(M)$ -module and  $\epsilon(M) := \cup_{k \geq 0} \epsilon^k(M)$  is an associative algebra with the wedge product, where  $\epsilon^0(M)$  is defined as  $C^\infty(M)$ .

For example, if  $f: M \rightarrow \mathbf{R}$  is a smooth function, then  $df$  defined by  $df_x(\xi_x) = \frac{d}{dt}|_0 d(\alpha(t))$  where  $\alpha'(0) = \xi_x$  defines a 1-form. For every 1-form  $\omega$  on a Riemannian manifold  $M$  there exists a smooth vector field  $\eta$  such that  $\omega(\xi)(x) = (\xi_x, \eta_x)$ . In particular, if  $\omega = df$  then  $\eta$  is called the gradient of  $f$ .

### 3. Tangent structures.

An  $n$ -plaque on a set  $X$  is a map  $p: U \subset \mathbf{R}^n \rightarrow X$  with  $U$  open. A *diffeology* on  $X$  is a collection  $P(X)$  of  $n$ -plaques for each  $n \in \mathbf{N}$  which covers  $X$  and such that if  $p \in P(X)$  and  $\phi: U \subset \mathbf{R}^m \rightarrow \mathbf{R}^n$  is smooth, then  $p \circ \phi \in P(X)$ . A *tangent structure* on  $(X, P(X))$  is a collection of equivalence relations  $\sim_F^n$  on the set of  $n$ -plaques at  $F \in X$  (that is,  $p \in P(X), p(0) = F$ ) denoted  $P_F^n(X)$ , for each  $n \in \mathbf{N}, F \in X$ , satisfying the following two consistency conditions:

1. If  $p_1 \sim_F^n p_2$  and  $\phi: U \subset \mathbf{R}^m \rightarrow \mathbf{R}^n$ , then  $p_1 \circ \phi \sim_F^m p_2 \circ \phi$ .
2. If  $p: U \rightarrow X$  is a plaque and  $V \subset U$  is an open neighborhood of 0, then  $p \sim p|_V$ .

The class of  $p$  is denoted  $[p]$  or  $[p(t)]_t$ . A diffeological space with a tangent structure is called a *TDS*. The tangent structure is called *linear* if  $P_F^n(X) / \sim_F^n$  carries a vector space structure for each  $F \in X, n \in \mathbf{N}$  (this set is called the  *$n$ -th tangent space at  $F$*  and is denoted  $T_F^n X$ ) satisfying

- a. If  $p_{12} \in [p_1] + [p_2]$  and  $\phi: U \subset \mathbf{R}^m \rightarrow \mathbf{R}^n$  is smooth, then

$$p_{12} \circ \phi \in [p_1 \circ \phi] + [p_2 \circ \phi]$$

- b. If  $p \in c[p_1] \Rightarrow p \circ \phi \in c[p_1 \circ \phi]$ .

The tangent structure is called *continuous* if given two  $(n+m)$ -plaques  $p_i(r, s), i = 1, 2$  such that  $p_1(r, 0) = p_2(r, 0)$  for each  $r$ , there exists a  $(n+m)$ -plaque  $p_{12}(r, s)$  such that

$$[p_{12}(r, s)]_s = [p_1(r, s)]_s + [p_2(r, s)]_s \quad \forall r$$

If  $(X, P(X), \sim), (Y, P(Y), \equiv)$  are two *TDS*, a map  $f: X \rightarrow Y$  is called *smooth* if it preserves plaques, directions, and the vector space structure, that is:

1.  $p \in P(X) \Rightarrow f \circ p \in P(Y)$
2.  $p_1 \sim_F^n p_2 \Rightarrow f \circ p_1 \equiv_{f(F)} f \circ p_2$
3.  $d^k f: T^k X \rightarrow T^k Y$  given by  $[p] \rightarrow [f \circ p]$  is linear if the tangent structure is linear.

The set of smooth maps is denoted  $C^\infty(X, Y)$  and the set of maps satisfying only 1. is denoted  $c^\infty(X, Y)$ .

Any manifold  $M$  has a *standard TDS* for which  $P(M)$  is formed by the smooth maps  $p: U \subset \mathbf{R}^n \rightarrow M$  (which contains the atlas) and

$$p_1 \sim_F^n p_2 \iff D^i p_1|_0 = D^i p_2|_0 \quad \forall i = 0, 1, \dots, n.$$

The tangent spaces and the smooth maps with this *TDS* coincide with those given by the manifold. In particular the standard manifold structure on  $\mathbf{R}^n$  provides a standard *TDS*. Using this, any subalgebra  $\mathcal{F}$  of  $c^\infty(X)$  on any diffeology  $(X, P(X))$  allows the construction of a tangent structure on  $(X, P(X))$ , called the *standard TDS defined by  $\mathcal{F}$* :

$$p_1 \sim_F^n p_2 \iff D^i f \circ p_1|_0 = D^i f \circ p_2|_0 \quad \forall f \in \mathcal{F}$$

in which we are using the standard *TDS* on  $\mathbf{R}$ .

The tangent spaces defines the *tangent bundles*  $T^n X$ . A *vector field* is a section of this bundle such that  $\forall f \in C^\infty(X, \mathbf{R})$  the map  $df(\xi): X \rightarrow \mathbf{R}$  given by  $F \rightarrow df(\xi(F))$  is smooth. The set of smooth vector fields is denoted  $\Gamma(X)$ . In every linear *TDS* the set  $\Gamma(X)$  is a  $C^\infty(X)$  module.

A vector field is called *locally integrable* if for every  $n$ -plaque  $p: U \subset \mathbf{R}^n \rightarrow X$  and every  $r_0 \in U$  there exists a neighborhood  $V$  of  $r_0$  and a  $(n+1)$ -plaque  $q$  such that  $q(r, 0) = p(r)$  and

$$\xi(p(r)) = [q(r, t)]_t \quad \forall r \in V_0$$

Notice that every smooth vector field on a manifold is locally integrable. If the tangent structure is linear and continuous, the set of locally integrable vector fields is a  $C^\infty(X)$  module, denoted  $\Gamma_T(X)$ .

A point  $F \in X$  is called *weakly (strongly resp.) transverse of degree  $(n, m)$*  if there exists  $p_1: U_1 \subset \mathbf{R}^n \rightarrow X$ ,  $p_2: U_2 \subset \mathbf{R}^m \rightarrow X$  at  $F$  such that there does not exist  $V_1 \subset U_1$ ,  $V_2 \subset U_2$  open at  $F$  and a  $(n+m)$ -plaque  $q: V_1 \times V_2 \rightarrow X$  satisfying  $q(r, 0) = p_1(r), \forall r$  and  $q(0, s) = p_2(s), \forall s$  (or  $[q(r, 0)]_r = [p_1(r)]_r, [q(0, s)]_s$  respectively)

### Examples 3.1.

- [1] Manifolds modelled on a locally convex vector space  $E$  do not have transverse points: if  $\{(\alpha_i, U_i)/i \in I\}$  is an atlas of  $M$  and  $p_1, p_2$  are  $n, m$ -plaques respectively, at  $F \in M$  then  $p_i = \alpha_i \circ \phi_i, i = 1, 2$  where  $\phi_i$  is smooth. We may assume that  $\alpha_1 = \alpha_2, \phi_1: V_1 \subset \mathbf{R}^n \rightarrow U \subset E, \phi_2: V_2 \subset \mathbf{R}^m \rightarrow U$ . Consider  $V'_i \subset V_i$  such that

$$\phi_1(V'_1) + \phi_2(V'_2) \subset U$$

Let  $\phi: V'_1 \times V'_2 \rightarrow U$  defined by  $\phi(u, v) = \phi_1(u) + \phi_2(v)$ , then  $q = \alpha_1 \circ \phi$  satisfies the required condition.

- [2] Let  $X$  be the group of diffeomorphisms of a finite dimensional manifold  $M$ . Let  $P(X)$  be the set of plaques  $p: U \subset \mathbf{R}^n \rightarrow X$  such that  $\bar{p}: U \times M \rightarrow M$  given by  $\bar{p}(r, m) = p(r)(m)$  is smooth and consider the standard tangent structure defined by  $c^\infty(X)$ . Then  $X$  does not have transverse points: take  $q(r, s) = F \circ p_1(r)p_2(s) \circ F^{-1}$ .
- [3] Let  $X = \mathbf{R}^2$ , let  $P(X)$  be the set of smooth maps  $p: U \subset \mathbf{R}^n \rightarrow X$  whose image is a line (under the standard  $TDS$ ), then  $c^\infty(X)$  consists of maps which are smooth along lines. With the standard  $TDS$  the tangent structure is linear and continuous, each  $T_F X$  is  $\mathbf{R}^2$ . The only locally integrable vector field is the zero vector field. Notice that every point is strongly transverse of degree  $(1, 1)$ . More generally we consider a manifold  $M$  (or a collection of manifolds  $(M_i)_{i \in I}$ ) and a set of smooth maps  $p: U \subset \mathbf{R}^n \rightarrow M$ , for different  $n \in \mathbf{N}$ , then consider the diffeology generated by these maps. For example, we take  $M = S^2$  and consider the maps whose image is a segment of a parallel (notice that the intersection of 2 plaques is either a point of another plaque). The poles are strongly transverse points of degree  $(1, 1)$ . A vector field is locally integrable  $\iff$  it is zero at poles. However, the union of two tangent surfaces (in this context) at only one point  $F$  do not have transverse points and every vector field is locally integrable.

Notice that every strongly transverse point is weakly transverse. Every smooth vector field is zero at a strongly transverse point.

### Definition 3.1.

Given a linear  $TDS$   $(X, P(X), \sim)$ , a  $k$ -form on  $X$  is defined in one of the following ways:

1. It is an exterior  $k$ -form  $\omega_F, \forall F \in X$  such that if  $\xi_1, \dots, \xi_k$  are vector fields then

$$\omega(\xi_1, \dots, \xi_k)(F) := \omega_F(\xi_1(F), \dots, \xi_k(F))$$

is a smooth map.

2. It is a  $C^\infty(X, \mathbf{R})$  alternating  $k$ -linear map

$$\omega: (\Gamma(X))^k \rightarrow C^\infty(X, \mathbf{R})$$

3. It is a collection  $(\omega_p)_{p \in P(X)}$  where  $\omega_p$  is a  $k$ -form on  $U \subset \mathbf{R}^n$  if  $p: U \rightarrow X$  such that

- a.  $\omega_{p \circ \phi} = \phi^* \omega_p, \forall \phi: U' \subset \mathbf{R}^m \rightarrow \mathbf{R}^n$  smooth.  
 b. If  $p_1, p_2$  are two plaques satisfying  $p_1(r_1) = p_2(r_2) = F$  which are tangent at  $F$  along directions  $v_1, \dots, v_k \in \mathbf{R}^n$  (this means that  $[p_1(tv_i + r_1)]_t = [p_2(tv_i + r_2)]_t$  for all  $i = 1, \dots, k$ ) then

$$\omega_{p_1(r_1)}(v_1, \dots, v_k) = \omega_{p_2(r_2)}(v_1, \dots, v_k)$$

and

$$d\omega_{p_1(r_1)}(v_1, \dots, v_{k+1}) = d\omega_{p_2(r_2)}(v_1, \dots, v_{k+1})$$

for all  $v_{k+1} \in \mathbf{R}^n$ . This is called the *tangent condition* for further reference.

The set of  $k$ -forms under definition  $i$  is denoted  $\epsilon_i^k(X)$  for  $i = 1, 2, 3$ . Each of them is a  $C^\infty(X)$  module and  $\epsilon_i^k(X)$ , is also an associative algebra with the wedge product defined as usual under each of the three definitions.

The pullback of  $\omega \in \epsilon_1^k(X_2)$  under a smooth map  $h: X_1 \rightarrow X_2$  is defined by

$$(h^* \omega)_F(\eta_1(F), \dots, \eta_k(F)) = \omega_{\eta(F)}(dh(F)\eta_1(F), \dots, dh(F)\eta_k(F))$$

If  $\omega \in \epsilon_3^k(X_2)$  then define

$$(h^* \omega) = (\omega_{p \circ h})_{p \in P(X_1)}$$

This collection satisfies the tangent condition. If  $\omega \in \epsilon_2^k(X_2)$  we need  $dh(F)$  to be surjective  $\forall F \in X$  and define

$$h^* \omega(\eta_1, \dots, \eta_k) = \omega(dh(\eta_1), \dots, dh(\eta_k)) \circ h$$

**Proposition 3.1.**

If  $(X, P(X), \sim)$  does not have transverse points then  $\epsilon_1^k(X) = \epsilon_3^k(X)$  for any  $k \in \mathbf{N}$

*Proof:* First we define a map  $\Psi: \epsilon_1^k(X) \rightarrow \epsilon_3^k(X)$  by

$$\Psi(\omega) = (p^* \omega)_{p \in P(X)}$$

This map is linear and  $\Psi(f\omega) = f\Psi(\omega)$  if  $f \in C^\infty(X)$ . We prove that  $\Psi$  is 1-1: Assume that  $\Psi(\omega) = 0$  and that  $\omega_F \neq 0$  for some  $F \in X$ , then  $\exists v_1, \dots, v_k \in \underline{T_F X}$  such that  $\omega_F(v_1, \dots, v_k) \neq 0$ . Let  $p_1, \dots, p_k$  be 1-plaques such that  $[p_i] = v_i$ . Since  $F$  is not strongly transverse, and using induction, there exists a neighborhood  $I$  of 0 in  $\mathbf{R}$  and an  $n$ -plaque  $p: I^k \rightarrow X$  such that  $[p(te_i)] = [p_i(t)]$ , therefore

$$\begin{aligned} \omega_{p(0)}(e_1, \dots, e_k) &= \omega_{p(0)}([p(te_1)], \dots, [p(te_k)]) \\ &= \omega_{p(0)}([p_1(t)], \dots, [p_k(t)]) \\ &= \omega_F(v_1, \dots, v_k) \neq 0 \end{aligned}$$

Therefore  $\omega$  is 1-1. Now we prove that  $\Psi$  is onto: Let  $(\omega_p)_{p \in P(X)} \in \epsilon_3^k(X)$ , define  $\underline{\omega}$  as follows: let  $F \in X$  and let  $v_1, \dots, v_k \in \underline{T_F X}$ , let  $p \in P(X)$  such that  $[p(te_i)] = v_i$  (exists because  $F$  is not strongly transverse), define

$$\underline{\omega}_F(v_1, \dots, v_k) = (\omega_p)(0)(e_1, \dots, e_k)$$

This value is independent of  $p$  because if  $p'$  is another plaque as before then

$$(\omega_p)(0)(e_1, \dots, e_k) = (\omega_{p'})(0)(e_1, \dots, e_k)$$

because of the tangent condition. Next we prove that  $\underline{\omega}$  is smooth: Let  $\xi_1, \dots, \xi_k \in \Gamma_I(X)$  and let  $p: U \subset \mathbf{R}^n \rightarrow X$  be a plaque. We shall prove that

$$\underline{\omega}(p(r))(\xi_1(p(r)), \dots, \xi_k(p(r))): U \rightarrow \mathbf{R}$$

is smooth. Let  $q_1, \dots, q_n$  be  $(n+1)$ -plaques such that  $[q_i(r, t)]_t = \xi_i(p(r)) \forall r$ . Let  $q: V \subset \mathbf{R}^{n+k} \rightarrow X$  be a plaque as in the definition of transverse point such that

$$p(r, 0, \dots, t_i, 0, \dots, 0) = q_i(r, t_i)$$

then  $q(r, 0, \dots, 0) = p(r), \forall r$  and

$$\begin{aligned} [q(r, 0, \dots, 0, t_i, 0, \dots, 0)]_{t_i} &= [q_i(r, t_i)]_{t_i} \\ \underline{\omega}_{p(r)}(\xi_1(p(r)), \dots, \xi_k(p(r))) &= \underline{\omega}_{q(r, 0)}([q_1(r, t)]_t, \dots, [q_k(r, t)]_t) \\ &= (\omega_q)(r, 0)([(r, t_1, 0, \dots, 0)]_{t_1}, [(r, 0, t_2, \dots, 0)]_{t_2}, \dots, [(r, \dots, t_k)]_{t_k}) \\ &= (\omega_q)(r, 0)(v_1(r), \dots, v_k(r)) \end{aligned}$$

and  $v_i(r) = [(r, 0, \dots, 0, t_i, 0, \dots, 0)]_{t_i}$  is a smooth vector field on  $U \subset \bar{U}$ , therefore  $\underline{\omega}$  is smooth. Notice that if  $p: U \subset \mathbf{R}^n \rightarrow X$  is a plaque then

$$\begin{aligned} (p^* \underline{\omega})(v_1, \dots, v_k) &= \underline{\omega}_{p(r)}([p(tv_1)], \dots, [p(tv_k)]) \\ &= (\omega_p)(r)(v_1, \dots, v_k) \end{aligned}$$

It follows from the definition that  $\Psi$  is onto since  $\Psi(\underline{\omega}) = \omega$  and this proves the proposition.

### Remarks 3.1.

1. The theorem remains still valid under a weaker condition than non transversality: for every pair of  $(n+1)$ -plaques  $p_1, p_2: U \times V \subset \mathbf{R}^n \times \mathbf{R} \rightarrow X$  such that  $p_1(r, 0) = p_2(r, 0), \forall r \in U$  there exists a  $(n+2)$ -plaque

$$q: U' \times V' \times V'' \subset \mathbf{R}^n \rightarrow \mathbf{R} \times \mathbf{R} \rightarrow X$$

such that

$$\begin{aligned} [q(r, t_1, 0)]_{t_1} &= [p_1(r, t_1)]_{t_1} \quad \forall r \in U' \\ [q(r, 0, t_2)]_{t_2} &= [p_2(r, t_2)]_{t_2} \quad \forall r \in U' \end{aligned}$$

2. For any  $X$  we have  $\epsilon_1^k(X) \subset \epsilon_2^k(X)$ .
3. In general  $\epsilon_2^k(X)$  is larger than  $\epsilon_1^k(X)$ : consider the example

$$X = \{(x, 0)/x \in \mathbf{R}\} \cup \{(0, y)/y \in \mathbf{R}\}$$

where  $P(X)$  is the set of smooth maps  $p: U \subset \mathbf{R}^n \rightarrow X$  whose image is a subset of the x-axis only, or a subset of the y-axis only. Then  $c^\infty(X)$  is the set of pairs  $(f_1, f_2)$  such that  $f_1, f_2: \mathbf{R} \rightarrow \mathbf{R}$  is smooth and  $f_1(0) = f_2(0)$ . With the standard diffeology  $T_{(0,0)}X = \mathbf{R} \sqcup \mathbf{R}$  and  $\underline{T}_{(0,0)}X = \{0\}$ , otherwise,  $T_F X = \mathbf{R} = \underline{T}_F(X)$ . The point  $(0, 0)$  is transverse and  $\Gamma(X)$  is the set of pairs  $(f_1 \frac{\partial}{\partial x}, f_2 \frac{\partial}{\partial y})$  such that  $f_1, f_2: \mathbf{R} \rightarrow \mathbf{R}$  is smooth and  $f_1(0) = f_2(0) = 0$ . Let

$$\xi_1(x, y) = x \frac{\partial}{\partial x} \quad \xi_2(x, y) = y \frac{\partial}{\partial y}$$

Since every smooth function  $h: \mathbf{R} \rightarrow \mathbf{R}$  such that  $h(0) = 0$  satisfies  $h(t) = tg(t)$  where  $g$  is also smooth then every field is expressed as

$$\xi(x, y) = h_1(x)\xi_1(x, y) + h_2(y)\xi_2(x, y)$$

where  $h_1, h_2: \mathbf{R} \rightarrow \mathbf{R}$  and conversely, define

$$\omega(\xi(x, y)) = h_1(x)(x^2 + 1) + h_2(y)(2y^2 + 1)$$

then  $\omega \in \epsilon_2(X)$  and satisfies  $\omega(\xi_1(0, 0)) = 1$ , however  $\xi_1(0, 0) = 0$ , therefore there is not  $\underline{\omega} \in \epsilon_1(X)$  which correspond to  $\omega$ . Similar constructions may be given with any  $k > 1$ .

4. A construction similar to the above may be done for any  $X$  such that  $\Gamma(X)$  is a free module and there exists a base  $\{\xi_i/i \in I\}$  such that at some point  $F$  the set of vectors

$$\{\xi_i(F)/i \in I\} \subset T_F X$$

is linearly dependent (this occurs if  $\dim T_F X \neq \text{Card} I$  for some  $F$ ). This occurs if  $\Gamma(X)$  is free and  $X$  has transverse points.

5.  $\epsilon_1(X) = \epsilon_2(X)$  if  $\Gamma(X)$  is a free module and there exists a base  $\{\xi_i/i \in I\}$  of  $\Gamma(X)$  such that  $\{\xi_i(F)/i \in I\}$  is linearly independent  $\forall F \in X$  (this occurs for example with  $\mathbf{R}^M$ ,  $M \in \mathbf{N} \cup \{\infty\}$  and manifolds with trivial tangent bundles such as  $S^1$ ), because if  $\omega(\xi_i) = h_i$  then

$$\omega(\xi = \sum_i f_i \xi_i) = \sum_i h_i f_i$$

and  $\xi(F) = 0$  implies  $f_i(F) = 0, \forall i \in I$

**References**

- [1] P. Donato, Géométrie des orbites coadjointes des groupes de difféomorphismes, in: C. Albert, ed., *Géométrie Symplectique et Mécanique*, Lecture Notes in Mathematics **1416** (Springer, Berlin, 1988) 84–104.
- [2] P. Donato, Les difféomorphismes du cercle comme orbit symplectique dans les moments de Virasoro, Preprint CPT-92/P.2681, CNRS-Luminy, 1992.
- [3] J. M. Souriau, Un algorithme générateur de structures quantiques, *Astérisque*, hors série (1985) 341–399.
- [4] J. M. Souriau, Groupes différentiels, *Lecture notes in math.*, 838 (1980) 91-128.
- [6] C. Torre, A Tangent bundle on diffeological spaces, math/9801046 (1998).
- [5] C. Torre, Examples of smooth diffeological spaces, Preprint (1999).
- [7] C. Torre and A. Banyaga, A symplectic structure on coadjoint orbits of diffeomorphism subgroups, *Ciencia y Tecnologia* 17 No 2, (1993) 1–14.