

Action-angle coordinates for time-dependent completely integrable Hamiltonian systems

GIOVANNI GIACHETTA¹, LUIGI MANGIAROTTI^{†2} AND GENNADI SARDANASHVILY^{‡3}

[†] Department of Mathematics and Physics, University of Camerino, 62032 Camerino (MC), Italy

[‡] Department of Theoretical Physics, Physics Faculty, Moscow State University, 117234 Moscow, Russia

Abstract. A time-dependent completely integrable Hamiltonian system is extended to the autonomous one, and time-dependent action-angle coordinates are introduced around an invariant manifold diffeomorphic to the product of a torus and the time axis. Written relative to these coordinates, a Hamiltonian and first integrals of a time-dependent completely integrable system become time-independent.

MSC (2000): 70H06

1 Introduction

A time-dependent Hamiltonian system of m degrees of freedom is said to be completely integrable if it admits m independent first integrals in involution. They are constant on trajectories of this system. Therefore, one may hope to find coordinates such that a Hamiltonian and first integrals of a time-dependent completely integrable system become time-independent. They are time-dependent action-angle coordinates. To introduce them, we use the fact that, in the framework of homogeneous Hamiltonian formalism, a time-dependent completely integrable Hamiltonian system of m degrees of freedom can be extended to an autonomous completely integrable system of $m + 1$ degrees of freedom where the time is treated as a dynamic variable [3, 4, 8]. In accordance with the well-known theorem [2, 7], this autonomous system admits the action-angle coordinates around a compact invariant manifold. The problem is that invariant manifolds of a time-dependent completely integrable Hamiltonian system are not compact because of the time axis. Therefore, we extend the above mentioned theorem to an invariant manifold which need not be compact, but obeys a certain condition (see Theorem 3 below). This invariant manifold is diffeomorphic to the product $\mathbb{R} \times T^m$ of the time axis \mathbb{R} and an m -dimensional torus T^m , and admits an open neighbourhood equipped with the time-dependent action-angle coordinates (I_i, t, ϕ^i) , $i = 1, \dots, m$, where t is the Cartesian coordinate on \mathbb{R} and ϕ^i

¹E-mail address: giovanni.giachetta@unicam.it

²E-mail address: luigi.mangiarotti@unicam.it

³E-mail address: sard@grav.phys.msu.su; URL: <http://webcenter.ru/~sardan/>

are cycle coordinates on T^m . Written with respect to these coordinates, a Hamiltonian and first integrals of a time-dependent completely integrable system are proved to be functions of action variables I_i only.

2 Time-dependent completely integrable Hamiltonian systems

Recall that the configuration space of a time-dependent mechanical system is a fibre bundle $Q \rightarrow \mathbb{R}$ over the time axis \mathbb{R} equipped with the bundle coordinates (t, q^k) , $k = 1, \dots, m$. The corresponding momentum phase space is the vertical cotangent bundle V^*Q of $Q \rightarrow \mathbb{R}$ endowed with holonomic bundle coordinates (t, q^k, p_k) [9, 10, 11].

The cotangent bundle T^*Q of Q is coordinated by

$$(q^0 = t, q^k, p_0 = p, p_k), \quad p' = p + \frac{\partial q^k}{\partial t} p_k, \quad (1)$$

and plays the role of the homogeneous momentum phase space of time-dependent mechanics. It is provided with the canonical Liouville form $\Xi = p_\lambda dq^\lambda$, the canonical symplectic form $\Omega_T = d\Xi$, and the corresponding Poisson bracket

$$\{f, f'\}_T = \partial^\lambda f \partial_\lambda f' - \partial_\lambda f \partial^\lambda f', \quad f, f' \in C^\infty(T^*Q). \quad (2)$$

There is the canonical one-dimensional fibre bundle

$$\zeta : T^*Q \rightarrow V^*Q, \quad (3)$$

whose kernel is the annihilator of the vertical tangent bundle $VQ \subset TQ$. A glance at the transformation law (1) shows that it is a trivial affine bundle. Indeed, given a global section h of ζ , one can equip T^*Q with the fibre coordinate $r = p - h$ possessing the identity transition functions.

The fibre bundle (3) provides the vertical cotangent bundle V^*Q with the canonical Poisson structure $\{\cdot, \cdot\}_V$ such that

$$\zeta^* \{f, f'\}_V = \{\zeta^* f, \zeta^* f'\}_T, \quad (4)$$

$$\{f, f'\}_V = \partial^k f \partial_k f' - \partial_k f \partial^k f', \quad (5)$$

for all $f, f' \in C^\infty(V^*Q)$ [8, 9]. The corresponding symplectic foliation coincides with the fibration $V^*Q \rightarrow \mathbb{R}$.

However, the Poisson structure (5) fails to define any dynamic equation on the momentum phase space V^*Q because Hamiltonian vector fields

$$\vartheta_f = \partial^k f \partial_k - \partial_k f \partial^k, \quad \vartheta_f \lrcorner df' = \{f, f'\}_V, \quad f, f' \in C^\infty(V^*Q),$$

of functions on V^*Q are vertical. Hamiltonian dynamics of time-dependent mechanics is described in a different way as a particular Hamiltonian dynamics on fibre bundles [5, 6].

A Hamiltonian on the momentum phase space $V^*Q \rightarrow \mathbb{R}$ of time-dependent mechanics is defined as a global section

$$h : V^*Q \rightarrow T^*Q, \quad p \circ h = -\mathcal{H}(t, q^j, p_j),$$

of the affine bundle ζ (3). It yields the pull-back Hamiltonian form

$$H = h^*\Xi = p_k dq^k - \mathcal{H}dt \quad (6)$$

on V^*Q . Given H (6), there exists a unique vector field γ_H on V^*Q such that

$$\gamma_H \rfloor dt = 1, \quad \gamma_H \rfloor dH = 0. \quad (7)$$

This vector field reads

$$\gamma_H = \partial_t + \partial^k \mathcal{H} \partial_k - \partial_k \mathcal{H} \partial^k. \quad (8)$$

It defines the Hamilton equation

$$\dot{t} = 1, \quad \dot{q}^k = \partial^k \mathcal{H}, \quad \dot{p}_k = -\partial_k \mathcal{H} \quad (9)$$

on V^*Q , where $(t, q^k, p_k, \dot{t}, \dot{q}^k, \dot{p}_k)$ are holonomic coordinates on the tangent bundle TV^*Q . Solutions of this equation are trajectories of the vector field γ_H . They assemble into a (regular) foliation of V^*Q .

A first integral of the Hamilton equation (9) is defined as a smooth real function F on V^*Q whose Lie derivative

$$\mathbf{L}_{\gamma_H} F = \gamma_H \rfloor dF = \partial_t F + \{\mathcal{H}, F\}_V$$

along the vector field γ_H (8) vanishes, i.e., the function F is constant on trajectories of the vector field γ_H . A time-dependent Hamiltonian system (V^*Q, H) on V^*Q is said to be completely integrable if the Hamilton equation (9) admits m first integrals F_k which are in involution with respect to the Poisson bracket $\{\cdot, \cdot\}_V$ (5) and whose differentials dF_k are linearly independent almost everywhere. This system can be extended to an autonomous completely integrable Hamiltonian system on T^*Q as follows.

Let us consider the pull-back ζ^*H of the Hamiltonian form $H = h^*\Xi$ onto the cotangent bundle T^*Q . It is readily observed that the difference $\Xi - \zeta^*h^*\Xi$ is a horizontal one-form on $T^*Q \rightarrow \mathbb{R}$ and that

$$\mathcal{H}^* = \partial_t \rfloor (\Xi - \zeta^*h^*\Xi) = p + \mathcal{H} \quad (10)$$

is a function on T^*Q [11]. Let us regard \mathcal{H}^* (10) as a Hamiltonian of an autonomous Hamiltonian system on the symplectic manifold (T^*Q, Ω_T) . The Hamiltonian vector field of \mathcal{H}^* on T^*Q reads

$$\gamma_T = \partial_t - \partial_t \mathcal{H} \partial^0 + \partial^k \mathcal{H} \partial_k - \partial_k \mathcal{H} \partial^k. \quad (11)$$

It is projected onto the vector field γ_H (8) on V^*Q , and the relation

$$\zeta^*(\mathbf{L}_{\gamma_H}f) = \{\mathcal{H}^*, \zeta^*f\}_T, \quad \forall f \in C^\infty(V^*Q).$$

holds. An immediate consequence of this relation is the following.

Proposition 1. Let $(V^*Q, H; F_k)$ be a time-dependent completely integrable Hamiltonian system with first integrals $\{F_k\}$ on V^*Q . Then $(T^*Q; \mathcal{H}^*, \zeta^*F_k)$ is an autonomous completely integrable Hamiltonian system on T^*Q whose first integrals $\{\mathcal{H}^*, \zeta^*F_k\}$ are in involution with respect to the Poisson bracket $\{\cdot, \cdot\}_T$ (2). Furthermore, let N be a connected invariant manifold of the time-dependent completely integrable Hamiltonian system $(V^*Q, H; F_k)$. Then $h(N) \subset T^*Q$ is a connected invariant manifold of the completely integrable Hamiltonian system $(T^*Q; \mathcal{H}^*, \zeta^*F_k)$ on T^*Q . If N contains no critical points of first integrals F_k , then $\{\mathcal{H}^*, \zeta^*F_k\}$ have no critical points in $h(N)$.

Hereafter, we restrict our consideration to complete vector fields γ_H (8). This means that the Hamilton equation (9) admits a unique global solution through each point of the momentum phase space V^*Q , and the trajectories of γ_H define a trivial fibre bundle

$$\xi : V^*Q \rightarrow V_t^*Q \quad (12)$$

of V^*Q over any fibre V_t^*Q of $V^*Q \rightarrow \mathbb{R}$. Without a loss of generality, we choose the fibre $i_0 : V_0^* \rightarrow V^*Q$ at $t = 0$. Since N is an invariant manifold, the fibration (12) also yields the fibration of N onto

$$N_0 = N \cap V_0^*Q \quad (13)$$

which makes N to the trivial fibre bundle

$$N \cong \xi^{-1}(N_0) = \mathbb{R} \times N_0.$$

It is readily observed that N_0 (13) is an invariant manifold of the completely integrable system $\{i_0^*F_k\}$ on the symplectic leaf V_0^*Q .

3 Time-dependent action-angle coordinates

Recall the following theorem [2, 7].

Theorem 2. Let (Z, Ω) be a $2m$ -dimensional symplectic manifold and f_1, \dots, f_m smooth real functions in involution on Z whose differentials df_k are linearly independent almost everywhere. Let M be a connected compact invariant manifold of the completely integrable system $\{f_k\}$. If M contains no critical points of f_k , there exists an open neighbourhood of M which is isomorphic to the symplectic annulus $V \times T^m$, where $V \subset \mathbb{R}^m$ is a

non-empty (open contractible) domain. It is equipped with the action-angle coordinates $(I_k, \phi^k \bmod 2\pi)$ such that the symplectic form on $V \times T^m$ reads

$$\Omega = dI_k \wedge d\phi^k,$$

and all f_k are functions of action coordinates (I_k) only.

Let us introduce time-dependent action-angle coordinates around an invariant manifold N of a time-dependent completely integrable Hamiltonian system (V^*Q, H) as those induced by the action-angle coordinates around the invariant manifold $h(N)$ of the autonomous completely integrable system (T^*Q, \mathcal{H}^*) in Proposition 1. Since N and, consequently, $h(N)$ are non-compact, we start from the following generalization of Theorem 2.

Theorem 3. Let M' be a connected invariant manifold of an autonomous completely integrable system (F_λ) , $\lambda = 1, \dots, n$, on a symplectic manifold (Z, Ω_Z) , and let the Hamiltonian vector fields of first integrals F_λ on M' be complete. Let there exist a neighbourhood U of M' such that F_λ have no critical points in U and the submersion

$$\times F_\lambda : U \rightarrow \mathbb{R}^n$$

is a trivial fibre bundle over a domain $V' \subset \mathbb{R}^n$. Then U is isomorphic to the symplectic annulus

$$W = V' \times (\mathbb{R}^{n-m} \times T^m) \tag{14}$$

provided with the generalized action-angle coordinates

$$(I_1, \dots, I_n; x^1, \dots, x^{n-m}; \phi^1, \dots, \phi^m) \tag{15}$$

such that the symplectic form on W reads

$$\Omega_Z = dI_i \wedge x^i + dI_{n-m+k} \wedge d\phi^k,$$

and the first integrals F_λ are functions of the action coordinates (I_α) only.

Proof. In accordance with the well-known theorem [1, 2], the invariant manifold M' is diffeomorphic to the product $\mathbb{R}^{n-m} \times T^m$, provided with coordinates

$$(y^\lambda) = (x^1, \dots, x^{n-m}; \phi^1, \dots, \phi^m)$$

along the integral curves of Hamiltonian vector fields of first integrals F_λ on M' . Let (J_λ) be coordinates on $V' \subset \mathbb{R}^n$ which are values of first integrals F_λ . Since $W' \rightarrow V'$ is a

trivial bundle, (J_λ, y^λ) are bundle coordinates on the annulus W' (14). The symplectic form Ω_Z on W' written with respect to these coordinates reads

$$\Omega_Z = a_\beta^\alpha dJ_\alpha \wedge dy^\beta + \frac{1}{2} b_{\alpha\beta} dy^\alpha \wedge dy^\beta,$$

where a_β^α is a nowhere degenerate matrix function on W' . By the definition of coordinates (y^λ) , the Hamiltonian vector fields ϑ_λ of first integrals F_λ take the coordinate form $\vartheta_\lambda = \vartheta_\lambda^\alpha(J_\mu) \partial_\alpha$ and obey the relations

$$\vartheta_\lambda \rfloor \Omega_Z = -dJ_\lambda, \quad a_\beta^\alpha \vartheta_\lambda^\beta = \delta_\lambda^\alpha, \quad \vartheta_\lambda^\alpha b_{\alpha\beta} = 0.$$

It follows that $v_\lambda^\alpha = (a^{-1})_\lambda^\alpha$ (i.e., the matrix function a_β^α depends only on coordinates J_μ), and that $b_{\alpha\beta} = 0$. Since Ω_Z is closed, one obtains the relation

$$\partial^\lambda a_\beta^\alpha dJ_\lambda \wedge dJ_\alpha = 0.$$

Then, by virtue of the relative Poincaré lemma, $a_\beta^\alpha = \partial^\alpha I_\beta$ where I_β are functions on the domain V' . Thus, we obtain the desired coordinates (15).

In particular, let M' be a compact invariant manifold of a completely integrable system $\{F_\lambda\}$, $\lambda = 1, \dots, n$, on a symplectic manifold (Z, Ω_Z) which does not contain critical points of the first integrals F_λ . One can show that the condition of Theorem 3 always holds [7], and we come to Theorem 2.

Now, let us apply Theorem 3 to completely integrable Hamiltonian systems in Proposition 1.

Theorem 4. Let the vector field γ_H (8) be complete. Let a connected invariant manifold N of a time-dependent completely integrable Hamiltonian system $(V^*Q, H; F_k)$ contain no critical points of first integrals F_k , and let its projection N_0 (13) onto the fibre V_0^*Q along trajectories of γ_H be compact. Then the invariant manifold $h(N)$ of the completely integrable Hamiltonian system $(T^*Q; \mathcal{H}^*, \zeta^*F_k)$ has an open neighbourhood U which obeys the condition of Theorem 3.

Proof. We first show that the invariant manifold N_0 of a completely integrable system on the fibre V_0^*Q does not contain critical points of the first integrals $i_0^*F_k$. On the contrary, let us suppose that $di_0^*F_k = i_0^*dF_k$ vanishes at some point $y \in N_0$. Then $u \rfloor dF_k(y) = 0$ for all vertical tangent vectors u to $V^*Q \rightarrow \mathbb{R}$ at $y \in V^*Q$. If $\partial_t \rfloor dF_k(y)$ also vanishes, then $dF_k(y) = 0$. If $\partial_t \rfloor dF_k(y) \neq 0$, then $\gamma_H(y) \rfloor dF_k(y) \neq 0$. The both results contradict the condition that F_k is a first integral without critical points in N . Since N_0 is compact and does not contain critical points of $i_0^*F_k$, Theorem 2 states that there is an open neighbourhood of N_0 in V_0^*Q isomorphic to $V \times N_0$ where $V \subset \mathbb{R}^m$ is an open domain, and that $\{v\} \times N_0$, $v \in V$, are also invariant manifolds in V_0^*Q . Then

$$W = \xi^{-1}(V \times N_0) \cong V \times N \tag{16}$$

is an open neighbourhood in V^*Q of the invariant manifold N foliated by invariant manifolds $\xi^{-1}(\{v\} \times N_0)$, $v \in V$, of the time-dependent completely integrable Hamiltonian system on V^*Q . Since $F_k = \xi^*(i_0^*F_k)$, we obtain that $dF_k = \xi^*d(i_0^*F_k) \neq 0$ on W , i.e., the first integrals F_k of the time-dependent completely integrable system have no critical points in the neighbourhood W of the invariant manifold N . For any real number $r \in (-\varepsilon, \varepsilon)$, let us consider a section

$$h_r : V^*Q \rightarrow T^*Q, \quad p \circ h_r = -\mathcal{H}(t, q^j, p_j) + r,$$

of the affine bundle ζ (3). Then the images $h_r(W)$, $r \in (-\varepsilon, \varepsilon)$, of the open neighbourhood W (16) of N in V^*Q make up an open neighbourhood U of $h(N) = h_0(N)$ in T^*Q . Because $\zeta(U) = W$, the pull-backs ζ^*F_k of first integrals F_k have no critical points in U , and so is obviously the function \mathcal{H}^* (10). Since the coordinate $r = p - h$ provides a trivialization of the affine bundle ζ , the open neighbourhood U of $h(N)$ is diffeomorphic to the product

$$(-\varepsilon, \varepsilon) \times h(W) \cong (-\varepsilon, \varepsilon) \times V \times h(N)$$

which is a trivialization of the fibration

$$\mathcal{H}^* \times (\times \zeta^*F_k) : U \rightarrow (-\varepsilon, \varepsilon) \times V.$$

Now, it remains to prove that the Hamiltonian vector fields of \mathcal{H}^* and ζ^*F_k on U are complete. It is readily observed that the Hamiltonian vector field γ_T (11) of \mathcal{H}^* is tangent to the manifolds $h_r(W)$, and is the image

$$\gamma_T = Th_r \circ \gamma_H \circ \zeta$$

of the vector field γ_H (8) on W (16). The latter is complete on W , and so is γ_T on U . Similarly, the Hamiltonian vector field

$$\gamma_k = -\partial_t F_k \partial^0 + \partial^i F_k \partial_i - \partial_i F_k \partial^i$$

of the function ζ^*F_k on T^*Q with respect to the Poisson bracket $\{, \}_T$ (2) is tangent to the manifolds $h_r(W)$, and is the image

$$\gamma_k = Th_r \circ \vartheta_k \circ \zeta$$

of the Hamiltonian vector field

$$\vartheta_k = \partial^i F_k \partial_i - \partial_i F_k \partial^i$$

of the first integral F_k on W with respect to the Poisson bracket $\{, \}_V$ (5). The vector fields ϑ_k on W are vertical relative to the fibration $W \rightarrow \mathbb{R}$, and are tangent to compact

manifolds. Therefore, they are complete, and so are the vector fields γ_k on U . Thus, U is the desired open neighbourhood of the invariant manifold $h(N)$.

In accordance with Theorem 3, the open neighbourhood U of the invariant manifold $h(N)$ of the completely integrable Hamiltonian system $(T^*Q; \mathcal{H}^*, \zeta^* F_k)$ in Theorem 4 is isomorphic to the symplectic annulus

$$W' = V' \times (\mathbb{R} \times T^m), \quad V' = (-\varepsilon, \varepsilon) \times V, \quad (17)$$

provided with the generalized action-angle coordinates

$$(I_0, \dots, I_m; t, \phi^1, \dots, \phi^m). \quad (18)$$

Moreover, running the proof of Theorem 3, we find that $J_0 = r$, $a_0^\alpha = \delta_0^\alpha$ and, as a consequence,

$$a_0^0 = \frac{\partial I_0}{\partial J_0} = 1, \quad a_i^0 = \frac{\partial I_i}{\partial J_0} = 0,$$

i.e., the action coordinate I_0 is linear in the coordinate r , while I_i are independent of r . With respect to the coordinates (18), the symplectic form on W' reads

$$\Omega_T = dI_0 \wedge dt + dI_k \wedge d\phi^k,$$

the Hamiltonian \mathcal{H}^* is an affine function

$$\mathcal{H}^* = I_0 + \mathcal{H}'(I_j) \quad (19)$$

of the action coordinate I_0 , while the first integrals $\zeta^* F_k$ depends only on the action coordinates I_i . The Hamiltonian vector field of the Hamiltonian (19) is

$$\gamma_T = \partial_t + \partial^i \mathcal{H}' \partial_i. \quad (20)$$

Since the action coordinates I_i are independent on the coordinate r , the symplectic annulus W' (17) inherits the composite fibration

$$W' \xrightarrow{\zeta} V \times (\mathbb{R} \times T^m) \longrightarrow \mathbb{R}. \quad (21)$$

Therefore, one can regard

$$W = V \times (\mathbb{R} \times T^m) \quad (22)$$

as a momentum phase space of the time-dependent Hamiltonian system in question around the invariant manifold N . It is coordinated by $(I_i; t, \phi^i)$, which we agree to call the time-dependent action-angle coordinates. By the relation similar to (4), W can be equipped with the Poisson structure

$$\{f, f'\}_W = \partial^i f \partial_i f' - \partial_i f \partial^i f',$$

while the global section

$$h' : W \rightarrow W', \quad I_0 \circ h' = -\mathcal{H}',$$

of the trivial bundle ζ (21) provides W with the Hamiltonian form

$$H' = I_i d\phi^i - \mathcal{H}'(I_j) dt.$$

The associated vector field γ_H (7) is exactly the projection onto W of the Hamiltonian vector field γ_T (20), and takes the same coordinate form. It defines the Hamilton equation

$$I_i = \text{const}, \quad \dot{\phi}^i = \partial^i \mathcal{H}'(I_j) \quad (23)$$

on W . One can think of this equation (23) as being the Hamilton equation of a time-dependent completely integrable Hamiltonian system around the invariant manifold N relative to time-dependent action-angle coordinates.

References

- [1] V.Arnold, *Russ. Math. Surv.* **18**, No.5, 13 (1963).
- [2] V.Arnold (Ed.), *Dynamical Systems III* (Springer-Verlag, Berlin, 1988).
- [3] S.Bouquet and A.Bourdier, *Phys. Rev. E* **57**, 1273 (1998).
- [4] A.Dewisme and S.Bouquet, *J. Math. Phys.* **34**, 997 (1993).
- [5] G.Giachetta, L.Mangiarotti and G.Sardanashvily, *New Lagrangian and Hamiltonian Methods in Field theory* (World Scientific, Singapore, 1997)
- [6] G.Giachetta, L.Mangiarotti and G.Sardanashvily, *J. Phys. A* **32**, 6629 (1999).
- [7] V.Lazutkin, *KAM Theory and Semiclassical Approximations to Eigenfunctions* (Springer-Verlag, Berlin, 1993).
- [8] A.Lichtenberg and M.Liebertmann, *Regular and Stochastic Motion* (Springer-Verlag, Berlin, 1983).
- [9] L.Mangiarotti and G.Sardanashvily, *Gauge Mechanics* (World Scientific, Singapore, 1998).
- [10] G.Sardanashvily, *J. Math. Phys.* **39**, 2714 (1998).
- [11] G.Sardanashvily, *J. Math. Phys.* **41**, 5245 (2000).