

# Action-angle coordinates for time-dependent completely integrable Hamiltonian systems

GIOVANNI GIACHETTA<sup>1</sup>, LUIGI MANGIAROTTI<sup>†2</sup> AND GENNADI SARDANASHVILY<sup>‡3</sup>

<sup>†</sup> Department of Mathematics and Physics, University of Camerino, 62032 Camerino (MC), Italy

<sup>‡</sup> Department of Theoretical Physics, Physics Faculty, Moscow State University, 117234 Moscow, Russia

**Abstract.** A time-dependent completely integrable Hamiltonian system is proved to admit the action-angle coordinates around any regular instantly compact invariant manifold. Written relative to these coordinates, its Hamiltonian and first integrals are functions only of action coordinates.

**MSC (2000):** 70H06

## 1 Introduction

A time-dependent Hamiltonian system of  $m$  degrees of freedom is called a completely integrable system (henceforth CIS) if it admits  $m$  independent first integrals in involution. We address the representation of its Hamiltonian in terms of first integrals (see [7, 9] for the low dimension case). By virtue of the classical Arnold–Liouville theorem [2, 8], an autonomous CIS admits the action-angle coordinates around a regular compact invariant manifold. In the framework of homogeneous Hamiltonian formalism, a time-dependent CIS of  $m$  degrees of freedom can be extended to an autonomous CIS of  $m + 1$  degrees of freedom where the time is regarded as a dynamic variable [3, 4, 10]. Using this fact, we aim to introduce action-angle coordinates for a time-dependent CIS. The problem is that its invariant manifolds are not compact because of the time axis. Generalizing the above mentioned theorem, we prove that there is a system of action-angle coordinates on an open neighbourhood  $U$  of a regular invariant manifold  $M$  of an autonomous CIS if Hamiltonian vector fields of first integrals on  $M$  are complete and the foliation of  $U$  by invariant manifolds is trivial. If  $M$  is a compact invariant manifold, these conditions always hold [8]. Then we show that, if a regular connected invariant manifold of a time-dependent CIS is compact at each instant, it is diffeomorphic to the product of the time axis  $\mathbb{R}$  and an  $m$ -dimensional torus  $T^m$ , and admits an open neighbourhood equipped with the time-dependent action-angle coordinates  $(I_i, t, \phi^i)$ ,  $i = 1, \dots, m$ , where  $t$  is the

---

<sup>1</sup>E-mail address: giovanni.giachetta@unicam.it

<sup>2</sup>E-mail address: luigi.mangiarotti@unicam.it

<sup>3</sup>E-mail address: sard@grav.phys.msu.su; URL: <http://webcenter.ru/~sardan/>

Cartesian coordinate on  $\mathbb{R}$  and  $\phi^i$  are cycle coordinates on  $T^m$ . Written with respect to these coordinates, a Hamiltonian and first integrals of a time-dependent CIS are functions only of action coordinates  $I_i$ .

Note that the choice of time-dependent action-angle coordinates by no means is unique. We show that, whenever a smooth function  $\mathcal{F}$  on  $\mathbb{R}^m$ , there exist the action-angle coordinates such that the Hamiltonian of a time-dependent CIS equals  $\mathcal{F}$  relative to these coordinates.

## 2 Time-dependent completely integrable Hamiltonian systems

Recall that the configuration space of a time-dependent mechanical system is a fibre bundle  $Q \rightarrow \mathbb{R}$  over the time axis  $\mathbb{R}$  equipped with the bundle coordinates  $(t, q^k)$ ,  $k = 1, \dots, m$ . The corresponding momentum phase space is the vertical cotangent bundle  $V^*Q$  of  $Q \rightarrow \mathbb{R}$  endowed with holonomic coordinates  $(t, q^k, p_k = \dot{q}_k)$  [11-13]. The cotangent bundle  $T^*Q$ , coordinated by  $(q^\lambda, p_\lambda) = (t, q^k, p_0, p_k)$ , is the homogeneous momentum phase space of time-dependent mechanics. It is provided with the canonical Liouville form  $\Xi = p_\lambda dq^\lambda$ , the canonical symplectic form  $\Omega = d\Xi$ , and the corresponding Poisson bracket

$$\{f, f'\}_T = \partial^\lambda f \partial_\lambda f' - \partial_\lambda f \partial^\lambda f', \quad f, f' \in C^\infty(T^*Q). \quad (1)$$

There is the one-dimensional trivial affine bundle

$$\zeta : T^*Q \rightarrow V^*Q. \quad (2)$$

Given its global section  $h$ , one can equip  $T^*Q$  with the fibre coordinate  $r = p_0 - h$  possessing the identity transition functions.

The fibre bundle (2) provides the vertical cotangent bundle  $V^*Q$  with the canonical Poisson structure  $\{, \}_V$  such that

$$\zeta^* \{f, f'\}_V = \{\zeta^* f, \zeta^* f'\}_T, \quad \forall f, f' \in C^\infty(V^*Q), \quad (3)$$

$$\{f, f'\}_V = \partial^k f \partial_k f' - \partial_k f \partial^k f', \quad (4)$$

[10, 11]. Its characteristic symplectic foliation coincides with the fibration  $V^*Q \rightarrow \mathbb{R}$ . However, the Poisson structure (4) fails to define any dynamic equation on the momentum phase space  $V^*Q$  because Hamiltonian vector fields

$$\vartheta_f = \partial^k f \partial_k - \partial_k f \partial^k, \quad \vartheta_f \lrcorner df' = \{f, f'\}_V, \quad f, f' \in C^\infty(V^*Q),$$

of functions on  $V^*Q$  are vertical.

The dynamics of Hamiltonian time-dependent mechanics is described as the particular Hamiltonian dynamics on fibre bundles [5, 6]. A Hamiltonian of time-dependent mechanics is defined as a global section

$$h : V^*Q \rightarrow T^*Q, \quad p_0 \circ h = -\mathcal{H}(t, q^j, p_j),$$

of the affine bundle  $\zeta$  (2). It yields the pull-back Hamiltonian form

$$H = h^*\Xi = p_k dq^k - \mathcal{H}dt \quad (5)$$

on  $V^*Q$  [11,12]. Then there exists a unique vector field  $\gamma_H$  on  $V^*Q$  such that

$$\gamma_H]dt = 1, \quad \gamma_H]dH = 0, \quad (6)$$

$$\gamma_H = \partial_t + \partial^k \mathcal{H} \partial_k - \partial_k \mathcal{H} \partial^k. \quad (7)$$

Its trajectories obey the Hamilton equation

$$\dot{q}^k = \partial^k \mathcal{H}, \quad \dot{p}_k = -\partial_k \mathcal{H} \quad (8)$$

on  $V^*Q$ . They assemble into a (regular) foliation of  $V^*Q$ .

A first integral of the Hamilton equation (8) is a smooth real function  $F$  on  $V^*Q$  whose Lie derivative

$$\mathbf{L}_{\gamma_H} F = \gamma_H]dF = \partial_t F + \{\mathcal{H}, F\}_V$$

along the vector field  $\gamma_H$  (7) vanishes, i.e.,  $F$  is constant on trajectories of  $\gamma_H$ . A time-dependent Hamiltonian system  $(V^*Q, H)$  is said to be completely integrable if the Hamilton equation (8) admits  $m$  first integrals  $F_k$  which are in involution with respect to the Poisson bracket  $\{\cdot, \cdot\}_V$  (4), and whose differentials  $dF_k$  are linearly independent almost everywhere. One can associate to this CIS an autonomous CIS on  $T^*Q$  as follows.

Let us consider the pull-back  $\zeta^*H$  of the Hamiltonian form  $H$  (5) onto the cotangent bundle  $T^*Q$ . It is readily observed that

$$\mathcal{H}^* = \partial_t](\Xi - \zeta^*h^*\Xi) = p + \mathcal{H} \quad (9)$$

is a function on  $T^*Q$  [13]. Let us regard  $\mathcal{H}^*$  as a Hamiltonian of an autonomous Hamiltonian system on the symplectic manifold  $(T^*Q, \Omega)$ . Its Hamiltonian vector field

$$\gamma_T = \partial_t - \partial_t \mathcal{H} \partial^0 + \partial^k \mathcal{H} \partial_k - \partial_k \mathcal{H} \partial^k \quad (10)$$

is projected onto the vector field  $\gamma_H$  (7) on  $V^*Q$  so that

$$\zeta^*(\mathbf{L}_{\gamma_H} f) = \{\mathcal{H}^*, \zeta^* f\}_T, \quad \forall f \in C^\infty(V^*Q).$$

An immediate consequence of this relation is the following.

*Theorem 1.* (i) Given a time-dependent CIS  $(V^*Q, H; F_k)$  on  $V^*Q$ , the functions  $\{\mathcal{H}^*, \zeta^*F_k\}$  are first integrals of an autonomous CIS on  $T^*Q$ . (ii) Let  $N$  be a connected regular invariant manifold of the time-dependent CIS  $(V^*Q, H; F_k)$ . Then  $h(N) \subset T^*Q$  is a connected regular invariant manifold of the autonomous CIS  $(T^*Q; \mathcal{H}^*, \zeta^*F_k)$  on  $T^*Q$ .

Hereafter, we assume that the vector field  $\gamma_H$  (7) is complete. In this case, the Hamilton equation (8) admits a unique global solution through each point of the momentum phase space  $V^*Q$ , and trajectories of  $\gamma_H$  define a trivial fibre bundle  $V^*Q \rightarrow V_t^*Q$  over any fibre  $V_t^*Q$  of  $V^*Q \rightarrow \mathbb{R}$ . Without a loss of generality, we choose the fibre  $i_0 : V_0^*Q \rightarrow V^*Q$  at  $t = 0$ . Since  $N$  is an invariant manifold, the fibration

$$\xi : V^*Q \rightarrow V_0^*Q \quad (11)$$

also yields the fibration of  $N$  onto  $N_0 = N \cap V_0^*Q$  such that

$$N = \xi^{-1}(N_0) \cong \mathbb{R} \times N_0$$

is a trivial fibre bundle. It is readily observed that  $N_0$  is an invariant manifold of the CIS  $\{i_0^*F_k\}$  on the symplectic leaf  $V_0^*Q$ .

### 3 Time-dependent action-angle coordinates

Let us introduce the action-angle coordinates around an invariant manifold  $N$  of a time-dependent CIS on  $V^*Q$  by use of the action-angle coordinates around the invariant manifold  $h(N)$  of the autonomous CIS on  $T^*Q$  in Theorem 1. Since  $N$  and, consequently,  $h(N)$  are non-compact, we first prove the following.

*Theorem 2.* Let  $M$  be a connected invariant manifold of an autonomous CIS  $\{F_\lambda\}$ ,  $\lambda = 1, \dots, n$ , on a symplectic manifold  $(Z, \Omega_Z)$ , and let the Hamiltonian vector fields of the first integrals  $F_\lambda$  on  $M$  be complete. Let  $U$  be a neighbourhood  $U$  of  $M$  such that  $F_\lambda$  have no critical points in  $U$  and the submersion  $\times F_\lambda : U \rightarrow \mathbb{R}^n$  is a trivial fibre bundle over a domain  $V' \subset \mathbb{R}^n$ . Then  $U$  is isomorphic to the symplectic annulus

$$W' = V' \times (\mathbb{R}^{n-m} \times T^m), \quad (12)$$

provided with the generalized action-angle coordinates

$$(I_1, \dots, I_n; x^1, \dots, x^{n-m}; \phi^1, \dots, \phi^m) \quad (13)$$

such that the symplectic form on  $W'$  reads

$$\Omega_Z = dI_i \wedge dx^i + dI_{n-m+k} \wedge d\phi^k,$$

and the first integrals  $F_\lambda$  depend only on the action coordinates  $(I_\alpha)$ .

*Proof.* In accordance with the well-known theorem [1, 2], the invariant manifold  $M$  is diffeomorphic to the product  $\mathbb{R}^{n-m} \times T^m$ , provided with coordinates

$$(y^\lambda) = (x^1, \dots, x^{n-m}; \phi^1, \dots, \phi^m)$$

along the integral curves of Hamiltonian vector fields of first integrals  $F_\lambda$  on  $M'$ . Let  $(J_\lambda)$  be coordinates on  $V' \subset \mathbb{R}^n$  which are values of first integrals  $F_\lambda$ . Since  $W' \rightarrow V'$  is a trivial bundle,  $(J_\lambda, y^\lambda)$  are bundle coordinates on the annulus  $W'$  (12). Written relative to these coordinates, the symplectic form  $\Omega_Z$  on  $W'$  reads

$$\Omega_Z = a_\beta^\alpha dJ_\alpha \wedge dy^\beta + \frac{1}{2} b_{\alpha\beta} dy^\alpha \wedge dy^\beta,$$

where  $a_\beta^\alpha$  is a nowhere degenerate matrix function on  $W'$ . By the definition of coordinates  $(y^\lambda)$ , the Hamiltonian vector fields  $\vartheta_\lambda$  of first integrals  $F_\lambda$  take the coordinate form  $\vartheta_\lambda = \vartheta_\lambda^\alpha (J_\mu) \partial_\alpha$  and obey the relations

$$\vartheta_\lambda \lrcorner \Omega_Z = -dJ_\lambda, \quad a_\beta^\alpha \vartheta_\lambda^\beta = \delta_\lambda^\alpha, \quad \vartheta_\lambda^\alpha b_{\alpha\beta} = 0.$$

It follows that  $\vartheta_\lambda^\alpha = (a^{-1})_\lambda^\alpha$  (i.e., the matrix function  $a_\beta^\alpha$  depends only on coordinates  $J_\mu$ ), and that  $b_{\alpha\beta} = 0$ . Since  $\Omega_Z$  is closed, the relations

$$\partial^\lambda a_\beta^\alpha dJ_\lambda \wedge dJ_\alpha = 0$$

hold. Then, by virtue of the relative Poincaré lemma,  $a_\beta^\alpha = \partial^\alpha I_\beta$  where  $I_\beta$  are functions on the domain  $V'$ . Thus, we obtain the desired coordinates (13).

Now, we apply Theorem 2 to the CISs in Theorem 1.

*Theorem 3.* Let  $N$  be a connected regular invariant manifold of a time-dependent CIS  $(V^*Q, H; F_k)$ , and let its projection  $N_0$  onto the fibre  $V_0^*Q$  along trajectories of  $\gamma_H$  be compact. Then the invariant manifold  $h(N)$  of the autonomous CIS  $(T^*Q; \mathcal{H}^*, \zeta^* F_k)$  has an open neighbourhood  $U$  obeying the condition of Theorem 2.

*Proof.* We first show that the invariant manifold  $N_0$  of an autonomous CIS on the symplectic leaf  $V_0^*Q$  does not contain critical points of the first integrals  $i_0^* F_k$ . On the contrary, let us suppose that  $i_0^* dF_k$  vanishes at some point  $y \in N_0$ . Then  $u \lrcorner dF_k(y) = 0$  for all vertical tangent vectors  $u$  to  $V^*Q \rightarrow \mathbb{R}$  at  $y \in V^*Q$ . If  $\partial_t \lrcorner dF_k(y)$  also vanishes, then  $dF_k(y) = 0$ . If  $\partial_t \lrcorner dF_k(y) \neq 0$ , then  $\gamma_H(y) \lrcorner dF_k(y) \neq 0$ . The both results contradict the condition that  $F_k$  has no critical points in  $N$ . Since  $N_0$  is compact and regular, there is an open neighbourhood of  $N_0$  in  $V_0^*Q$  isomorphic to  $V \times N_0$  where  $V \subset \mathbb{R}^m$  is a domain, and that  $\{v\} \times N_0$ ,  $v \in V$ , are also invariant manifolds in  $V_0^*Q$  [8]. Then

$$W = \xi^{-1}(V \times N_0) \cong V \times N \tag{14}$$

is an open neighbourhood in  $V^*Q$  of the invariant manifold  $N$  foliated by invariant manifolds  $\xi^{-1}(\{v\} \times N_0)$ ,  $v \in V$ , of the time-dependent CIS on  $V^*Q$ . Since  $F_k = \xi^*(i_0^*F_k)$ , we obtain that  $dF_k = \xi^*d(i_0^*F_k) \neq 0$  on  $W$ , i.e., the first integrals  $F_k$  have no critical points in  $W$ . For any real number  $r \in (-\varepsilon, \varepsilon)$ , let us consider a section

$$h_r : V^*Q \rightarrow T^*Q, \quad p \circ h_r = -\mathcal{H}(t, q^j, p_j) + r,$$

of the affine bundle  $\zeta$  (2). Then the images  $h_r(W)$  of  $W$  (14) in  $V^*Q$  make up an open neighbourhood  $U$  of  $h(N)$  in  $T^*Q$ . Because  $\zeta(U) = W$ , the pull-backs  $\zeta^*F_k$  of first integrals  $F_k$  are free from critical points in  $U$ , and so is obviously the function  $\mathcal{H}^*$  (9). Since the coordinate  $r = p - h$  provides a trivialization of the affine bundle  $\zeta$ , the open neighbourhood  $U$  of  $h(N)$  is diffeomorphic to the product

$$(-\varepsilon, \varepsilon) \times h(W) \cong (-\varepsilon, \varepsilon) \times V \times h(N)$$

which is a trivialization of the fibration

$$\mathcal{H}^* \times (\times \zeta^*F_k) : U \rightarrow (-\varepsilon, \varepsilon) \times V.$$

It remains to prove that the Hamiltonian vector fields of  $\mathcal{H}^*$  and  $\zeta^*F_k$  on  $U$  are complete. It is readily observed that the Hamiltonian vector field  $\gamma_T$  (10) of  $\mathcal{H}^*$  is tangent to the manifolds  $h_r(W)$ , and it is the image  $\gamma_T = Th_r \circ \gamma_H \circ \zeta$  of the vector field  $\gamma_H$  (7) on  $W$  (14). The latter is complete on  $W$ , and so is  $\gamma_T$  on  $U$ . Similarly, the Hamiltonian vector field

$$\gamma_k = -\partial_t F_k \partial^0 + \partial^i F_k \partial_i - \partial_i F_k \partial^i$$

of the function  $\zeta^*F_k$  on  $T^*Q$  with respect to the Poisson bracket  $\{\cdot, \cdot\}_T$  (1) is tangent to the manifolds  $h_r(W)$ , and is the image  $\gamma_k = Th_r \circ \vartheta_k \circ \zeta$  of the Hamiltonian vector field  $\vartheta_k$  of the first integral  $F_k$  on  $W$  with respect to the Poisson bracket  $\{\cdot, \cdot\}_V$  (4). The vector fields  $\vartheta_k$  on  $W$  are vertical relative to the fibration  $W \rightarrow \mathbb{R}$ , and are tangent to compact manifolds. Therefore, they are complete, and so are the vector fields  $\gamma_k$  on  $U$ . Thus,  $U$  is the desired open neighbourhood of the invariant manifold  $h(N)$ .

In accordance with Theorem 2, the open neighbourhood  $U$  of the invariant manifold  $h(N)$  of the autonomous CIS  $(T^*Q; \mathcal{H}^*, \zeta^*F_k)$  in Theorem 3 is isomorphic to the symplectic annulus

$$W' = V' \times (\mathbb{R} \times T^m), \quad V' = (-\varepsilon, \varepsilon) \times V, \quad (15)$$

provided with the generalized action-angle coordinates

$$(I_0, \dots, I_m; t, \phi^1, \dots, \phi^m). \quad (16)$$

Moreover, running the proof of Theorem 2, we find that  $J_0 = r$  and, as a consequence,

$$a_0^0 = \frac{\partial I_0}{\partial J_0} = 1, \quad a_i^0 = \frac{\partial I_i}{\partial J_0} = 0,$$

i.e., the action coordinate  $I_0$  is linear in the coordinate  $r$ , while  $I_i$  are independent of  $r$ . With respect to the coordinates (16), the symplectic form on  $W'$  reads

$$\Omega = dI_0 \wedge dt + dI_k \wedge d\phi^k,$$

the Hamiltonian  $\mathcal{H}^*$  is an affine function

$$\mathcal{H}^* = I_0 + \mathcal{H}'(I_j) \tag{17}$$

of the action coordinate  $I_0$ , while the first integrals  $J_k = \zeta^* F_k$  depend only on the action coordinates  $I_i$ .

Since the action coordinates  $I_i$  are independent of the coordinate  $r$ , the symplectic annulus  $W'$  (15) inherits the fibration

$$W' \xrightarrow{\zeta} W'' = V \times (\mathbb{R} \times T^m). \tag{18}$$

Therefore, one can regard  $W'' \cong W$  (18), coordinated by  $(I_i; t, \phi^i)$ , as the momentum phase space of the time-dependent Hamiltonian system in question around the invariant manifold  $N$ . Its Hamiltonian relative to these coordinates is  $\mathcal{H}'(I_j)$  (17). The corresponding Hamilton equation (8) on  $W''$  reads

$$\dot{I}_i = 0, \quad \dot{\phi}^i = \partial^i \mathcal{H}'(I_j).$$

Its form motivates us to regard  $(I_i; t, \phi^i)$  as action-angle coordinates for a time-dependent CIS.

As was mentioned above, the choice of time-dependent action-angle coordinates by no means is unique. In particular, given the fibration  $\xi$  (11), let us provide  $V \times N_0 \subset V_0^*Q$  in Theorem 3 with the initial date action-angle coordinates  $(\bar{I}_k; \bar{\phi}^k)$  for an autonomous CIS  $\{i_0^* F_k\}$  on the symplectic leaf  $V_0^*Q$ . Then, it is readily observed that  $(\bar{I}_k; t, \bar{\phi}^k)$  are time-dependent action-angle coordinates on  $W''$  (18) such that the Hamiltonian  $\mathcal{H}'$  of a time-dependent CIS relative to these coordinates vanishes. Now, let  $\mathcal{F}(\bar{I}_i)$  be an arbitrary smooth function of action coordinates  $\bar{I}_i$ . Let us introduce new coordinates  $(\bar{I}_k; t, \phi'^k = \bar{\phi}^k + t\partial^k \mathcal{F})$ . These are time-dependent action-angle coordinates such that  $\mathcal{H}' = \mathcal{F}$ .

## References

- [1] V.Arnold, *Russ. Math. Surv.* **18**, No.5, 13 (1963).
- [2] V.Arnold (Ed.), *Dynamical Systems III* (Springer-Verlag, Berlin, 1988).
- [3] S.Bouquet and A.Bourdier, *Phys. Rev. E* **57**, 1273 (1998).
- [4] A.Dewisme and S.Bouquet, *J. Math. Phys.* **34**, 997 (1993).
- [5] G.Giachetta, L.Mangiarotti and G.Sardanashvily, *New Lagrangian and Hamiltonian Methods in Field theory* (World Scientific, Singapore, 1997)
- [6] G.Giachetta, L.Mangiarotti and G.Sardanashvily, *J. Phys. A* **32**, 6629 (1999).
- [7] R.Kaushal, *Int. J. Theor. Phys.* **37** (1998) 1793.
- [8] V.Lazutkin, *KAM Theory and Semiclassical Approximations to Eigenfunctions* (Springer-Verlag, Berlin, 1993).
- [9] H.Lewis, P.Leach, S.Bouquet and M.Feix, *J. Math. Phys.* **33** (1992) 591.
- [10] A.Lichtenberg and M.Liebermann, *Regular and Stochastic Motion* (Springer-Verlag, Berlin, 1983).
- [11] L.Mangiarotti and G.Sardanashvily, *Gauge Mechanics* (World Scientific, Singapore, 1998).
- [12] G.Sardanashvily, *J. Math. Phys.* **39**, 2714 (1998).
- [13] G.Sardanashvily, *J. Math. Phys.* **41**, 5245 (2000).