

Action-angle coordinates for time-dependent completely integrable Hamiltonian systems

Giovanni Giachetta¹, Luigi Mangiarotti² and Gennadi Sardanashvily³

^y Department of Mathematics and Physics, University of Camerino, 62032 Camerino (MC), Italy

^z Department of Theoretical Physics, Physics Faculty, Moscow State University, 117234 Moscow, Russia

Abstract. A time-dependent completely integrable Hamiltonian system is proved to admit the action-angle coordinates around any instantly compact regular invariant manifold. Written relative to these coordinates, its Hamiltonian and first integrals are functions only of action coordinates.

1 Introduction

A time-dependent Hamiltonian system of m degrees of freedom is called a completely integrable system (henceforth CIS) if it admits m independent first integrals in involution. In the framework of homogeneous Hamiltonian formalism, a time-dependent CIS of m degrees of freedom can be extended to an autonomous CIS of $m + 1$ degrees of freedom where the time is regarded as a dynamic variable [3, 4, 8]. Using this fact, we aim to introduce action-angle coordinates for a time-dependent CIS.

By virtue of the classical Arnold-Liouville theorem [2, 6], an autonomous CIS admits the action-angle coordinates around a regular compact invariant manifold. The problem is that its invariant manifolds are not compact because of the time axis. Generalizing the above mentioned theorem, we prove that there is a system of action-angle coordinates on an open neighbourhood U of a regular invariant manifold M of an autonomous CIS if Hamiltonian vector fields of first integrals on M are complete and the foliation of U by invariant manifolds is trivial. If M is a compact invariant manifold, these conditions always hold [6]. Then we show that, if a regular connected invariant manifold of a time-dependent CIS is compact at each instant, it is diffeomorphic to the product of the time axis R and an m -dimensional torus T^m , and admits an open neighbourhood equipped with the time-dependent action-angle coordinates $(I_i; t; \theta^i)$, $i = 1; \dots; m$, where t is the Cartesian coordinate on R and θ^i are cyclic coordinates on T^m . Written with respect to these coordinates, a Hamiltonian and first integrals of a time-dependent CIS are functions only of action coordinates I_i .

Thus, action-angle coordinates for a time-dependent CIS provide a particular solution of the problem of a representation of its Hamiltonian in terms of first integrals. However, this representation need not hold with respect to any coordinate system because a Hamiltonian fails to be a scalar under time-dependent canonical transformations [5, 7].

¹E-mail address: giovannigiachetta@unicam.it

²E-mail address: luigimangiarotti@unicam.it

³E-mail address: sard@grav.phys.msu.su; URL: <http://webcenter.ru/~sardan/>

2 Time-dependent completely integrable Hamiltonian systems

Recall that the configuration space of a time-dependent mechanical system is a fibre bundle $Q \rightarrow \mathbb{R}$ over the time axis \mathbb{R} equipped with the bundle coordinates $(t; q^k)$, $k = 1, \dots, m$. The corresponding momentum phase space is the vertical cotangent bundle VQ of $Q \rightarrow \mathbb{R}$ endowed with holonomic coordinates $(t; q^k; p_k = \dot{q}_k)$ [9-11]. The cotangent bundle TQ , coordinated by $(q; p) = (t; q^k; p_0; p_k)$, is the homogeneous momentum phase space of time-dependent mechanics. It is provided with the canonical Liouville form $\omega = p dq$, the canonical symplectic form $\Omega = d\omega$, and the corresponding Poisson bracket

$$\{f, g\}_{TQ} = \frac{\partial f}{\partial q^k} \frac{\partial g}{\partial p_k} - \frac{\partial f}{\partial p_k} \frac{\partial g}{\partial q^k}; \quad f, g \in C^1(TQ); \quad (1)$$

There is the one-dimensional trivial affine bundle

$$\pi: TQ \rightarrow VQ; \quad (2)$$

Given its global section h , one can equip TQ with the fibre coordinate $r = p_0 - h$ possessing the identity transition functions. The fibre bundle (2) provides the vertical cotangent bundle VQ with the canonical Poisson structure $\{f, g\}_V$ such that

$$\{f, g\}_V = \frac{\partial f}{\partial q^k} \frac{\partial g}{\partial p_k} - \frac{\partial f}{\partial p_k} \frac{\partial g}{\partial q^k}; \quad f, g \in C^1(VQ); \quad (3)$$

$$\{f, g\}_V = \frac{\partial f}{\partial q^k} \frac{\partial g}{\partial p_k} - \frac{\partial f}{\partial p_k} \frac{\partial g}{\partial q^k}; \quad (4)$$

Its characteristic symplectic foliation coincides with the fibration $VQ \rightarrow \mathbb{R}$.

A Hamiltonian of time-dependent mechanics is defined as a global section

$$h: VQ \rightarrow TQ; \quad p_0 - h = H(t; q; p_j);$$

of the affine bundle (2) [9, 10]. It yields the pull-back Hamiltonian form

$$H = h^* \omega = p_k dq^k - H dt \quad (5)$$

on VQ [9, 10]. Then there exists a unique vector field X_H on VQ such that

$$X_H \lrcorner \omega = 1; \quad X_H \lrcorner H = 0; \quad (6)$$

$$X_H = \frac{\partial}{\partial t} + \frac{\partial H}{\partial q^k} \frac{\partial}{\partial p_k} - \frac{\partial H}{\partial p_k} \frac{\partial}{\partial q^k}; \quad (7)$$

Its trajectories obey the Hamilton equation

$$\dot{q}^k = \frac{\partial H}{\partial p_k}; \quad \dot{p}_k = - \frac{\partial H}{\partial q^k}; \quad (8)$$

A first integral of the Hamilton equation (8) is a smooth real function F on VQ whose Lie derivative

$$L_{X_H} F = X_H \lrcorner dF = \frac{\partial F}{\partial t} + \frac{\partial F}{\partial q^k} \dot{q}^k + \frac{\partial F}{\partial p_k} \dot{p}_k$$

along the vector field X_H (7) vanishes, i.e., F is constant on trajectories of X_H . A time-dependent Hamiltonian system $(VQ; H)$ is said to be completely integrable if the Hamilton

equation (8) admits m first integrals F_k which are in involution with respect to the Poisson bracket $\{f, g\}_V$ (4), and whose differentials dF_k are linearly independent almost everywhere, i.e., the set of points where this condition fails is nowhere dense in $V \times Q$. One can associate to this CIS an autonomous CIS on $T \times Q$ as follows.

Let us consider the pullback H of the Hamiltonian form H (5) onto the cotangent bundle $T \times Q$. It is readily observed that

$$H = \partial_t C(\quad h) = p_0 + H \quad (9)$$

is a function on $T \times Q$. Let us regard H as a Hamiltonian of an autonomous Hamiltonian system on the symplectic manifold $(T \times Q; \quad)$ [11]. Its Hamiltonian vector field

$$X_H = \partial_t + \partial_t H \partial^0 + \partial^k H \partial_k - \partial_k H \partial^k \quad (10)$$

is projected onto the vector field X_H (7) on $V \times Q$ so that

$$(L_{X_H} f) = f X_H; \quad f g_T; \quad 8 f^2 C^1(V \times Q):$$

An immediate consequence of this relation is the following.

Theorem 1. (i) Given a time-dependent CIS $(V \times Q; H; F_k)$ on $V \times Q$, the Hamiltonian system $(T \times Q; H; F_k)$ on $T \times Q$ is completely integrable. (ii) Let N be a connected regular invariant manifold of $(V \times Q; H; F_k)$. Then $h(N) \subset T \times Q$ is a connected regular invariant manifold of the autonomous CIS $(T \times Q; H; F_k)$.

Hereafter, we assume that the vector field X_H (7) is complete. In this case, the Hamilton equation (8) admits a unique global solution through each point of the momentum phase space $V \times Q$, and trajectories of X_H define a trivial fibre bundle $V \times Q \rightarrow V_t \times Q$ over any fibre $V_t \times Q$ of $V \times Q \rightarrow \mathbb{R}$. Without loss of generality, we choose the fibre $i_0: V_0 \times Q \rightarrow V \times Q$ at $t=0$. Since N is an invariant manifold, the fibration

$$: V \times Q \rightarrow V_0 \times Q \quad (11)$$

also yields the fibration of N onto $N_0 = N \cap V_0 \times Q$ such that $N \rightarrow N_0$ is a trivial fibre bundle.

3 Time-dependent action-angle coordinates

Let us introduce the action-angle coordinates around an invariant manifold N of a time-dependent CIS on $V \times Q$ by use of the action-angle coordinates around the invariant manifold $h(N)$ of the autonomous CIS on $T \times Q$ in Theorem 1. Since N and, consequently, $h(N)$ are non-compact, we first prove the following.

Theorem 2. Let M be a connected invariant manifold of an autonomous CIS $(\mathbb{Z}^n; \quad = 1; \dots; n)$, on a symplectic manifold $(\mathbb{Z}; \quad_z)$, and let the Hamiltonian vector fields of the first integrals F on M be complete. Let U be a neighbourhood of M such that F have no

critical points in U and the submersion $F : U \rightarrow \mathbb{R}^n$ is a trivial bundle of Lagrangian invariant manifolds over a domain $V^0 \subset \mathbb{R}^n$. Then U is isomorphic to the symplectic annulus

$$W^0 = V^0 \times (\mathbb{R}^{n-m} \times T^m); \quad (12)$$

provided with the generalized action-angle coordinates

$$(I_1; \dots; I_n; x^1; \dots; x^{n-m}; \varphi^1; \dots; \varphi^m) \quad (13)$$

such that the symplectic form on W^0 reads

$$\omega_Z = dI_i \wedge dx^i + dI_{n-m+k} \wedge d\varphi^k;$$

and the first integrals F depend only on the action coordinates (I) .

Proof. In accordance with the well-known theorem [1, 2], the invariant manifold M is diffeomorphic to the product $\mathbb{R}^{n-m} \times T^m$, provided with coordinates $(y) = (s^a; \varphi^i)$ where φ^i are linear functions of parameters s along the integral curves of Hamiltonian vector fields of first integrals F on M . Let (J) be coordinates on V^0 which are values of first integrals F . Since $W^0 \rightarrow V^0$ is a trivial bundle, $(J; y)$ are bundle coordinates on the annulus W^0 (12). It should be emphasized that, since group parameters are given up to a shift, the coordinates y on W^0 are determined up to a shift by functions of coordinates J . Written relative to these coordinates, the symplectic form ω_Z on W^0 reads

$$\omega_Z = dJ^a \wedge dJ_a + dJ^i \wedge d\varphi_i; \quad (14)$$

By the definition of coordinates (y) , the Hamiltonian vector fields $\#$ of first integrals take the coordinate form $\# = \#(J) \partial_y$ where

$$\#_a = \partial_a + \#_a^i(J) \partial_i; \quad \#_i = \#_i^k(J) \partial_k; \quad (15)$$

and they obey the relations

$$\#^a \lrcorner \omega_Z = dJ^a; \quad \#^i = 0; \quad (16)$$

It follows that ω_Z is a nondegenerate matrix and $\# = (\#^a)$, i.e., the matrix functions depend only on coordinates J . Let us apply the relation (16) to the Hamiltonian vector fields $\#_a$ and $\#_i$ (15). We obtain

$$\#_b^a = \delta_b^a; \quad \#_a^i = 0; \quad (17)$$

$$\#_i^k \partial_k = \partial_i; \quad \#_i^a = 0; \quad (18)$$

The first of the equalities (18) shows that the matrix $\#_i^j$ is nondegenerate, and so is $\#_i^k$. Then the second one results in $\#_k^a = 0$.

Using the well-known Kunneth formula for the de Rham cohomology of a product, one can easily justify that the closed form ω_Z on W^0 (12) is exact. Moreover, $\omega_Z = d\varphi$ where φ takes the form

$$\varphi = (J^a; \varphi^i) dJ_a + \varphi_i(J) d\varphi^i;$$

Of course, α_i is determined up to an exact form. Using the fact that components of $d\alpha = \sum \alpha_i dz_i$ are independent of y and obey the equalities (17) { (18), we obtain the following.

- (i) $\alpha_i^a = \partial_i \alpha^a + \partial^a \alpha_i = 0$. It follows that α_i^a is independent of y , i.e., α^a is a one-form in y and, consequently, is independent of y since y is a cyclic coordinate. Hence, $\partial^a \alpha_i = 0$, i.e., α_i is a function only of coordinates J_j .
- (ii) $\alpha_i^k = \partial_i \alpha^k + \partial^k \alpha_i$. Similarly, one shows that α^k is independent of y and $\alpha_i^k = \partial^k \alpha_i$, i.e., $\partial^k \alpha_i$ is a nondegenerate matrix.
- (iii) $\alpha_b^a = \partial_b \alpha^a = \alpha_b^a$. Hence, $\alpha^a = D^a(J) + s^a$.
- (iv) $\alpha_b^i = \partial_b \alpha^i$, i.e., α^i is a one-form in s^a .

In view of items (i) { (iv), the Liouville form reads

$$\alpha = x^a dJ_a + \mathbb{D}^i(J) + B_a^i(J) s^a dJ_i + \alpha_i(J_j) dy^i;$$

where $x^a = s^a + D^a(J)$. Since the matrix $\partial^k \alpha_i$ is nondegenerate, one can introduce new coordinates $I_i = \alpha_i(J_j)$, $I_a = J_a$. We obtain

$$\alpha = x^a dI_a + \mathbb{D}^{0i}(I) + B_a^{0i}(I) s^a dI_i + I_i dy^i;$$

Finally, put

$$I_i = y^i - \mathbb{D}^{0i}(I) + B_a^{0i}(I) s^a$$

in order to obtain the desired coordinates

$$I_a = J_a; \quad I_i(J_k); \quad x^a = s^a; \quad \alpha^a(s^b; y^i; J): \quad (19)$$

Of course, the generalized action-angle coordinates (13) by no means are unique. For instance, let F_a , $a = 1; \dots; n-m$ be an arbitrary smooth function on R^m . Let us consider the canonical coordinate transformation

$$I_a^0 = I_a - F_a(I_j); \quad I_k^0 = I_k; \quad x^{0a} = x^a; \quad \alpha^0 = \alpha + x^a \partial^a F_a(I_j): \quad (20)$$

Then $(I_a^0; I_k^0; x^{0a}; \alpha^0)$ are generalized action-angle coordinates on the symplectic annulus which differs from W^0 (12) in another trivialization.

Now, we apply Theorem 2 to the CISs in Theorem 1.

Theorem 3. Let N be a connected regular invariant manifold of a time-dependent CIS $(V \times Q; H; F_k)$, and let the image N_0 of its projection (11) be compact. Then the invariant manifold $h(N)$ of the autonomous CIS $(T \times Q; H; F_k)$ has an open neighbourhood U obeying the condition of Theorem 2.

Proof. (i) We first show that functions $i_0 F_k$ make up a CIS on the symplectic leaf $(V_0 \times Q; \omega_0)$ and N_0 is its invariant manifold without critical points. Clearly, the functions $i_0 F_k$ are in involution, and N_0 is their connected invariant manifold. Let us show that the set of critical points of $i_0 F_k$ is nowhere dense in $V_0 \times Q$ and N_0 has none of these points. Let $V_0 \times Q$ be equipped with some coordinates $(\bar{q}^k; \bar{p}_k)$. Then the trivial bundle (11) is provided with the bundle coordinates $(t; \bar{q}^k; \bar{p}_k)$ which play a role of the initial data coordinates on the

momentum phase space $V \times Q$. Written with respect to these coordinates, the first integrals F_k become time-independent. It follows that

$$dF_k(y) = d_{i_0}F_k(y) \quad (21)$$

for any point $y \in V \times Q$. In particular, if $y_0 \in V_0 \times Q$ is a critical point of $f_{i_0}F_k$, then the trajectory $\pi^{-1}(y_0)$ is a critical set for the first integrals fF_k . The desired statement at once follows from this result.

(ii) Since N_0 obeys the condition in item (i), there is an open neighbourhood of N_0 in $V_0 \times Q$ isomorphic to $V \times N_0$ where $V \subset \mathbb{R}^m$ is a domain, and $f_v g|_{N_0}$, $v \in V$, are also invariant manifolds in $V_0 \times Q$ [6]. Then

$$W = \pi^{-1}(V \times N_0) = V \times N \quad (22)$$

is an open neighbourhood in $V \times Q$ of the invariant manifold N foliated by invariant manifolds $\pi^{-1}(f_v g|_{N_0})$, $v \in V$, of the time-dependent CIS on $V \times Q$. By virtue of the equality (21), the first integrals fF_k have no critical points in W . For any real number $r \in (0, \infty)$, let us consider a section

$$h_r : V \times Q \rightarrow T \times Q; \quad p_0 \mapsto h_r = H(t; q^j; p_j) + r;$$

of the affine bundle (2). Then the images $h_r(W)$ of W (22) make up an open neighbourhood U of $h(N)$ in $T \times Q$. Because $h_r^{-1}(U) = W$, the pullbacks F_k of first integrals F_k are free from critical points in U , and so is the function H (9). Since the coordinate $r = p_0 - h$ provides a trivialization of the affine bundle, the open neighbourhood U of $h(N)$ is diffeomorphic to the product

$$(0, \infty) \times h(W) = (0, \infty) \times V \times h(N)$$

which is a trivialization of the fibration

$$H : (0, \infty) \times F_k : U \rightarrow (0, \infty) \times V :$$

(iii) It remains to prove that the Hamiltonian vector fields of H and F_k on U are complete. It is readily observed that the Hamiltonian vector field X_H (10) of H is tangent to the manifolds $h_r(W)$, and is the image $X_H = Th_r X_H$ of the vector field X_H (7). The latter is complete on W , and so is X_H on U . Similarly, the Hamiltonian vector field

$$X_k = \partial_t F_k \partial^0 + \partial^i F_k \partial_i - \partial_i F_k \partial^i$$

of the function F_k on $T \times Q$ with respect to the Poisson bracket $f; g_T$ (1) is tangent to the manifolds $h_r(W)$, and is the image $X_k = Th_r X_k$ of the Hamiltonian vector field X_k of the first integral F_k on W with respect to the Poisson bracket $f; g_V$ (4). The vector fields X_k on W are vertical relative to the fibration $W \rightarrow \mathbb{R}$, and are tangent to compact manifolds. Therefore, they are complete, and so are the vector fields X_k on U . Thus, U is the desired open neighbourhood of the invariant manifold $h(N)$.

In accordance with Theorem 2, the open neighbourhood U of the invariant manifold $h(N)$ of the autonomous CIS $(T^*Q; H; F_k)$ in Theorem 3 is isomorphic to the symplectic annulus

$$W^0 = V^0 \cap (R \times T^m); \quad V^0 = (t; \theta) \in V; \quad (23)$$

provided with the generalized action-angle coordinates

$$(I_0; \dots; I_m; t; \theta^1; \dots; \theta^m): \quad (24)$$

A glance at the Hamiltonian vector field $\#_0 = \tau$ (10) and the relation (16) – (19), shows that $I_0 = J_0 = H$ and the corresponding generalized angle coordinate is $x^0 = t$, while the first integrals $J_k = F_k$ depend only on the action coordinates I_i .

Since the action coordinates I_i are independent of the coordinate r , the symplectic annulus W^0 (23) inherits theibration

$$W^0 \cap W^\infty = V \cap (R \times T^m): \quad (25)$$

By the relation similar to (3), the product $W^\infty = W$ (25), coordinated by $(I_i; t; \theta^i)$, is provided with the Poisson structure

$$\{f, g\}_W = \partial_i f \partial_i g^0 - \partial_i g \partial_i f^0; \quad f, g \in C^1(W^\infty):$$

Therefore, one can regard W^∞ as the momentum phase space of the time-dependent CIS in question around the invariant manifold N .

It is readily observed that the Hamiltonian vector field τ of the autonomous Hamiltonian $H = I_0$ is $\tau = \partial_t$, and so is its projection τ_H (7) on W^∞ . Consequently, the Hamilton equation (8) of a time-dependent CIS with respect to the action-angle coordinates take the form $\tau_H = 0$, $\dot{\theta}^i = 0$. Consequently, $(I_i; t; \theta^i)$ are the initial date coordinates. One can introduce such coordinates as follows. Given theibration (11), let us provide $N_0 \subset V \subset V_0 \subset Q$ in Theorem 3 with action-angle coordinates $(\bar{I}_i; \bar{\theta}^i)$ for the CIS $(f_0 F_k g)$ on the symplectic leaf $V_0 \subset Q$. Then, it is readily observed that $(\bar{I}_i; t; \bar{\theta}^i)$ are time-dependent action-angle coordinates on W^∞ (25) such that the Hamiltonian $H(\bar{I}_j)$ of a time-dependent CIS relative to these coordinates vanishes, i.e., $\tau_H = \bar{I}_0$. Using the canonical transformations (20), one can consider time-dependent action-angle coordinates besides the initial date ones. Given a smooth function H on R^m , one can provide W^∞ with the action-angle coordinates

$$I_0 = \bar{I}_0 + H(\bar{I}_j); \quad I_i = \bar{I}_i; \quad \theta^i = \bar{\theta}^i + t \partial_i H(\bar{I}_j)$$

such that $H(I_i)$ is a Hamiltonian of time-dependent CIS on W^∞ .

References

- [1] Arnold V 1963 Russ. Math. Surv. 18 No.5 13
- [2] Arnold V (Ed.) 1998 Dynamical Systems III (Berlin: Springer-Verlag)
- [3] Bouquet S and Bourdier A 1998 Phys. Rev. E 57 1273

- [4] Dewism e A and Bouquet S 1993 J. M ath. Phys. 34 997
- [5] Kaushal R 1998 Int. J. Theor. Phys. 37 1793
- [6] Lazutkin V 1993 KAM Theory and Sem iclassical Approxim ations to Eigenfunctions (Berlin: Springer-Verlag)
- [7] Lew is H , Leach O , Bouquet S and Feix M 1992 J. M ath. Phys. 33 591
- [8] Lichtenberg A and Lieberman M 1983 Regular and Stochastic Motion (Berlin: Springer-Verlag)
- [9] Mangiarotti L and Sardanashvily G 1998 Gauge M echanics (Singapore: W orld Scien-
ti c)
- [10] Sardanashvily G 1998 J. M ath. Phys. 39 2714
- [11] Sardanashvily G 2000 J. M ath. Phys. 41 5245