

RIEMANN-ROCH FOR QUOTIENTS AND TODD CLASSES OF SIMPLICIAL TORIC VARIETIES

DAN EDIDIN AND WILLIAM GRAHAM

Department of Mathematics, University of Missouri, Columbia, MO
65211.

E-mail address: edidin@math.missouri.edu

Department of Mathematics, University of Georgia, Athens, GA 30602

E-mail address: wag@math.uga.edu

Dedicated to Steven L. Kleiman on the occasion of his 60th birthday.

ABSTRACT. In this paper we give an explicit formula for the Riemann-Roch map for singular schemes which are quotients of smooth schemes by diagonalizable groups. As an application we obtain a simple proof of a formula for the Todd class of a simplicial toric variety. An equivariant version of this formula was previously obtained for complete simplicial toric varieties by Brion and Vergne [BV] using different techniques.

1. INTRODUCTION

Suppose G is a linear algebraic group acting properly (hence with finite stabilizers) on a scheme X , with a geometric quotient $Y = X/G$. There is [BFM] a Riemann-Roch map $\tau_Y : G(Y)_{\mathbb{Q}} \rightarrow CH^*(Y)_{\mathbb{Q}}$, where $G(Y)_{\mathbb{Q}}$ denotes the rational Grothendieck group of coherent sheaves on Y and $CH^*(Y)_{\mathbb{Q}}$ denotes the rational Chow groups of Y . Similarly, there is an equivariant Riemann-Roch map $\tau_X^G : G^G(X)_{\mathbb{Q}} \rightarrow CH_G^*(X)_{\mathbb{Q}}$, where $G^G(X)$ is the Grothendieck group of G -equivariant coherent sheaves on X , and $CH_G^*(X)$ denotes the equivariant Chow groups of X .

The purpose of this paper is to obtain a formula (Theorem 3.1) for τ_Y in the case where X is smooth, G is a diagonalizable group (for example, a torus), and the ground field is \mathbb{C} . This is of interest because Riemann-Roch maps for non-smooth schemes are in general difficult to compute. The formula we give implies Kawasaki's Riemann-Roch formula for orbifolds

Both authors were partially supported by N.S.F. grants.

2000 Mathematics Subject Classification: Primary 14C40; Secondary 19L47, 14L32

which are global quotients of smooth spaces by diagonalizable groups. For quasi-projective quotients of quasi-projective varieties, Theorem 3.1 can be deduced from Toen's Riemann-Roch theorem for Deligne-Mumford stacks [To, Thm. 4.10]. Our proof, however, does not require quasi-projectivity hypotheses; moreover, it is more elementary than Toen's proof and completely avoids the use of stacks. An application of this result is a simple proof of a formula of Brion and Vergne [BV] for the Todd class of a simplicial toric variety. Related results for equivariant Todd classes of toric varieties were also obtained by Guillemin [Gu] and Ginzburg-Guillemin-Karshon [GGK] using an equivariant version of Kawaski's Riemann-Roch theorem.

The ingredients of our proof are the localization theorem in equivariant K -theory and the equivariant Riemann-Roch theorem of our previous paper [EG2]. The localization theorem relates the equivariant K -theory of X to the equivariant K -theory of X^g , the fixed locus of $g \in G$. It is necessary to consider the fixed loci for all $g \in G$, and this leads to an expression for τ_Y in terms of τ_X^G which is more subtle than might be naively expected.

In particular, there is a natural map $\text{Inv} : G^G(X) \rightarrow G(Y)$ taking the class of an equivariant coherent sheaf \mathcal{F} to the sheaf of invariant sections \mathcal{F}^G . Also, there is an isomorphism $\phi : CH_G^*(X)_{\mathbb{Q}} \rightarrow CH^*(Y)_{\mathbb{Q}}$, taking the class of an invariant cycle Z to the class $\frac{1}{e_Z}[Z/G]$, where (in characteristic 0) e_Z is the order of the stabilizer group in G of a general point of Z . These maps fit into a diagram

$$\begin{array}{ccc} K^G(X) & \xrightarrow{\tau_X^G} & CH_G^*(X)_{\mathbb{Q}} \\ \text{Inv} \downarrow & & \phi \downarrow \\ G(Y) & \xrightarrow{\tau_Y} & CH^*(Y)_{\mathbb{Q}}. \end{array}$$

The naive formula for τ_Y is the formula (for $\alpha \in G^G(X)$)

$$(1) \quad \tau_Y(\alpha^G) = \phi \circ \tau_X^G(\alpha).$$

This would be true if the above diagram commuted. However, in general it does not, unless the action of G is free. The correct formula for $\tau_Y(\alpha^G)$ is a sum of terms coming from the fixed point loci X^g as g runs over the (finite) set of elements of G such that X^g is nonempty. The right hand side of (1) is the contribution corresponding to the fixed point set of the identity in G , which is of course X . If G acts freely on X then the X^g are empty for $g \neq 1$, which explains why the naive formula is true in that case.

In a subsequent paper we will work out a more general formula for actions of reductive groups on algebraic spaces of arbitrary characteristic. We will also give an equivariant version of the the formula in Theorem 3.1.

2. BACKGROUND

2.1. Group actions and quotients. In this paper all schemes are assumed to be of finite type and separated over the ground field \mathbb{C} . All algebraic groups are assumed to be linear. Because we are working over \mathbb{C} , every *reductive* group (i.e., a group with trivial unipotent radical) is *linearly reductive* (i.e., every representation splits as a direct sum of irreducibles). A group G is *diagonalizable* if it is isomorphic to a closed subgroup of D_n , the group of diagonal matrices in $\mathrm{GL}(n)$ [B, Prop. 8.4]. Since \mathbb{C} is algebraically closed, [B, Prop. 8.7] implies that a group is diagonalizable if and only if it is isomorphic to $(\mathbb{C}^*)^n \times F$ for some finite abelian group F .

If G is an algebraic group acting on a scheme X then we say that G acts *properly* if the map $G \times X \rightarrow X \times X$ given by $(g, x) \mapsto (x, gx)$ is proper. This implies that the stabilizer of any point is proper, hence finite. However, a group action can have finite stabilizers without being proper.

If G acts properly on a scheme X a geometric quotient $Y = X/G$ always exists in the category of algebraic spaces, but in general Y may not be a scheme. To avoid the technicalities of working with algebraic spaces, in this paper when we refer to a quotient $Y = X/G$ we assume that the quotient is a scheme. If G is reductive, the orbits of the G action on X are closed¹, and X can be covered by G -invariant affines then a geometric quotient exists by [MFK, Proposition 1.9]. If X is normal and G is a torus then Sumihiro's theorem [Su] states that X can be covered by G -invariant affines. The properness of the action on X implies that the quotient is a separated scheme [MFK, Lemma 0.6] and that the quotient map $X \rightarrow X/G$ is an affine morphism [MFK, Proposition 0.7].

An important fact about geometric quotients is the following.

Proposition 2.1. *Suppose that a reductive group G acts properly on a scheme X and $X \rightarrow Y$ is a geometric quotient. Let $p: X' \rightarrow X$ be a finite G -equivariant map. Then:*

- (1) G acts properly on X' .
- (2) There is a geometric quotient $X' \rightarrow Y'$ and a finite map $q: Y' \rightarrow Y$ such that the following diagram commutes:

$$\begin{array}{ccc} X' & \xrightarrow{p} & X \\ \downarrow & & \downarrow \\ Y' & \xrightarrow{q} & Y. \end{array}$$

Proof. (1) Since G acts properly on X , the map $G \times X \rightarrow X \times X$ given by $(g, x) \mapsto (x, gx)$ is proper. The fiber product $(G \times X)_{X \times X}(X' \times X')$ is

¹This is automatic if G acts with finite stabilizers, since all orbits have dimension equal $\dim G$.

isomorphic to the product $G \times (X' \times_X X')$ via the map $(g, x, x'_1, x'_2) \mapsto (g, x'_1, g^{-1}x'_2)$. Hence the map $G \times (X' \times_X X') \rightarrow X' \times X'$ given by $(g, x'_1, x'_2) \mapsto (x'_1, gx'_2)$ is proper. The map $G \times X' \rightarrow X' \times X'$ factors as

$$G \times X' \xrightarrow{1 \times \Delta_p} G \times (X' \times_X X') \rightarrow X' \times X'.$$

The first map is proper (in fact, it is a closed embedding, since $p: X' \rightarrow X$ is separated). We have shown that the second map is proper, so the composition is as well. Thus G acts properly on X' . (Note that this argument only requires that p be separated.)

Since G acts properly on X the quotient map $X \rightarrow Y$ is affine. Hence, X is covered by G -invariant affines X_1, \dots, X_k where $X_i = \text{Spec } A_i$. Since $X' \rightarrow X$ is finite, X'_1, \dots, X'_k is a cover of X' by G -invariant affines, where $X'_i = p^{-1}(X_i) = \text{Spec}(B_i)$ and the B_i 's are finite A_i -algebras. As noted above, [MFK, Proposition 1.9] implies that there is a geometric quotient $X' \rightarrow Y'$. Since geometric quotients are categorical quotients ([MFK, Proposition 0.1]) the G -invariant map $X' \rightarrow X \rightarrow Y$ factors through the quotient $X' \rightarrow Y'$; i.e., the diagram

$$\begin{array}{ccc} X' & \xrightarrow{p} & X \\ \downarrow & & \downarrow \\ Y' & \xrightarrow{q} & Y \end{array}$$

commutes.

It remains to show that the induced map $Y' \rightarrow Y$ is finite. To do this we will work locally on Y and assume that $Y = \text{Spec } C$. Since the quotient map $X \rightarrow Y$ is affine it follows that $X = \text{Spec } A$. Since $\text{Spec } A \rightarrow \text{Spec } A^G$ is a categorical quotient [MFK, Theorem 1.1], and categorical quotients are unique up to isomorphism (cf. [MFK, Proposition 0.1]), it follows that $C \simeq A^G$.

The map p is affine (being finite) so $X' = \text{Spec } B$, which again implies that $Y' = \text{Spec } B^G$. Because p is finite, B is a finite A -algebra; we must prove that B^G is a finite A^G -algebra. Since G is reductive, A^G and B^G are both of finite type over \mathbb{C} [MFK, Theorem 1.1]. Thus B^G is of finite type over A^G . Now if $b \in B^G$ then, since B is a finite A -algebra, b is integral over A . Let $p(t) \in A[t]$ be a monic polynomial for b . Let $q(t) = Ep(t)$ where $E: A[t] \rightarrow A[t]$ is the Reynolds operator [MFK, Definition 1.5]. (Here $A[t]$ is made into a G -module by taking the G -action on t to be trivial.) Since $1 \in A$ is G -invariant, $q(t)$ is a monic polynomial with coefficients in A^G . By functoriality of E , Eb is a root of $q(t)$. But $b \in B^G$ so $Eb = b$. Therefore, B^G is integral over A^G . \square

2.2. Invariants and equivariant K -theory. If X is a scheme with a G -action, denote by $K^G(X)$ the Grothendieck group of G -equivariant locally

free sheaves; this is a ring, with the product given by the tensor product of equivariant vector bundles. Let $G^G(X)$ denote the Grothendieck group of G -equivariant coherent sheaves; this is a module for the ring $K^G(X)$. Identify the representation ring $R(G)$ with $K^G(pt)$; then pullback by the projection $X \rightarrow pt$ gives a natural map $R(G) \rightarrow K^G(X)$. In this way, $K^G(X)$ becomes an $R(G)$ -algebra and $G^G(X)$ an $R(G)$ -module.

If X is smooth and G is reductive then every G -equivariant coherent sheaf has a finite resolution by G -equivariant locally free sheaves. This implies that $G^G(X) \simeq K^G(X)$ [Th1, Theorem 2.18].

If $X \xrightarrow{\pi} Y$ is a geometric quotient and \mathcal{F} is a G -equivariant coherent sheaf on X , write \mathcal{F}^G for the sheaf of \mathcal{O}_Y -modules $(\pi_*(\mathcal{F}))^G$.

Proposition 2.2. *If a reductive group G acts properly on a scheme X and $X \rightarrow Y$ is a geometric quotient, then the assignment $\mathcal{F} \mapsto \mathcal{F}^G$ is an exact functor from the category of G -equivariant coherent \mathcal{O}_X -modules to the category of coherent \mathcal{O}_Y -modules.*

Proof. Since the quotient map is affine we may assume that $X = \text{Spec } A$ and $Y = \text{Spec } A^G$.

We first show that the assignment $\mathcal{F} \mapsto \mathcal{F}^G$ is exact. Since taking invariants is left exact for any group G , it suffices to prove that any surjection $M \twoheadrightarrow N$ of G -equivariant finitely generated A -modules restricts to a surjection $M^G \twoheadrightarrow N^G$ of invariant submodules. Any G -module is a union of its finite-dimensional G -submodules ([MFK, Lemma p. 25]) so we may reduce to the case that M, N are finite-dimensional. In this case, since G is reductive, the map $M^G \rightarrow N^G$ is surjective by Schur's lemma.

To show that the assignment takes coherent sheaves to coherent sheaves, we must show that if G is reductive and N is a finitely generated G -equivariant A -module, then N^G is finitely generated as an A^G -module. This fact is well known; see [Gr, Theorem 16.8] for a proof. \square

Proposition 2.3. *Suppose that G acts properly on X and $X' \xrightarrow{p} X$ is a finite G -equivariant map. Let $Y' \xrightarrow{q} Y$ be the induced finite map of quotients (Proposition 2.1). If \mathcal{F} is a G -equivariant coherent sheaf on X' then $q_*(\mathcal{F}^G) = (p_*\mathcal{F})^G$ as coherent sheaves on Y .*

Proof. As in the proof of Proposition 2.2 we can reduce to the case $X = \text{Spec } A$, $X' = \text{Spec } B$, $Y = \text{Spec } A^G$, $Y' = \text{Spec } B^G$ and $\mathcal{F} = \tilde{M}$ where M is a G -equivariant B -module. In this case both $q_*(\mathcal{F}^G)$ and $(p_*\mathcal{F})^G$ correspond to the A^G -module M^G , where the A^G -module structure is obtained via $A^G \rightarrow B^G \hookrightarrow B$. \square

As a consequence of Propositions 2.2 and 2.3, the assignment $\mathcal{F} \mapsto \mathcal{F}^G$ induces a map of Grothendieck groups $\text{Inv}: G^G(X) \rightarrow G^G(Y)$ which is covariant for finite morphisms. If $\alpha \in G^G(X)$, we will write α^G for $\text{Inv}(\alpha)$.

2.3. The localization theorem for diagonalizable groups in equivariant K -theory. For simplicity, throughout the rest of the paper all Grothendieck and Chow groups are taken with complex coefficients.

In this section G will denote a diagonalizable group. Let $R = R(G) \otimes \mathbb{C}$ denote the representation ring of G tensored with \mathbb{C} . Then R coincides with the coordinate ring $\mathcal{O}(G)$ of G [B, Proposition p.119]. If $h \in G$, let $\mathfrak{m}_h \subset R$ denote the corresponding maximal ideal.

Suppose that G acts on X . If $h \in G$, let $i_h: X^h \hookrightarrow X$ denote the inclusion of the fixed locus X^h of h into X . The localization theorem says that the proper push-forward $i_{h*}: G^G(X^h)_{\mathfrak{m}_h} \rightarrow G^G(X)_{\mathfrak{m}_h}$ is an isomorphism [Th2, Theorem 2.1]; here $G^G(X)_{\mathfrak{m}_h}$ is the localization of $G^G(X)$ at the maximal ideal \mathfrak{m}_h of R .

If N is an equivariant vector bundle of rank r , we write $\lambda_{-1}(N)$ for the class $\sum_{i=0}^r (-1)^i [\Lambda^i N]$ in $K^G(X)$. If $N = \bigoplus_{i=1}^r L_i$ splits as a direct sum of equivariant line bundles then $\lambda_{-1}(N) = \prod_{i=1}^r (1 - [L_i])$.

If X is smooth, then i_h is a regular embedding (note that X^h need not be connected or equidimensional). Let N_h denote the normal bundle to this embedding, and N_h^* the conormal bundle. Then [Th2, Lemma 3.3] says $\lambda_{-1}(N_h^*)$ is a unit in the ring $K^G(X^h) \simeq G^G(X^h)$ and the inverse to i_{h*} is the map

$$\frac{i_h^*}{\lambda_{-1}(N_h^*)}: G^G(X)_{\mathfrak{m}_h} \rightarrow G^G(X^h)_{\mathfrak{m}_h}.$$

Proposition 2.4. *Suppose that G acts on X with finite stabilizers. Then the natural map*

$$G^G(X) \rightarrow \bigoplus_h G^G(X)_{\mathfrak{m}_h}$$

is an isomorphism, where the sum is over the finite set of $h \in G$ such that X^h is nonempty.

Proof. If G acts with finite stabilizers, the proof of [EG2, Proposition 5.1] shows that there is an ideal $J \subset R$ with R/J supported at the points $h \in G$ for which X^h is nonempty, such that $JR^G(X) = 0$. Hence

$$G^G(X) = G^G(X)/JR^G(X) = G^G(X) \otimes_R R/J.$$

Now R/J is Artinian, so the map

$$R \rightarrow \bigoplus_{\mathfrak{m}_h \in \text{Spec } R/J} (R/J)_{\mathfrak{m}_h} / \mathfrak{m}_h^{k_h} = \bigoplus_{\mathfrak{m}_h \in \text{Spec } R/J} (R/J)_{\mathfrak{m}_h}$$

is an isomorphism; here k_h is an integer such that $\mathfrak{m}_h^{k_h} R/J = \mathfrak{m}_h^{k_h+1} R/J$. Tensoring with $G^G(X)$ yields the proposition. \square

Henceforth, if G acts on X with finite stabilizers, we will view $G^G(X)_{\mathfrak{m}_h}$ as a subset of $G_G(X)$. If $f : G_G(X) \rightarrow V$ is any map from $G^G(X)$ to any vector space V , we use the same letter f to denote the induced map $G^G(X)_{\mathfrak{m}_h} \rightarrow V$. In particular, we obtain $\text{Inv} : G^G(X)_{\mathfrak{m}_h} \rightarrow G(Y)$.

Lemma 2.5. *Suppose $p : X' \rightarrow X$ is a finite G -equivariant map, where G acts with finite stabilizers. Then $p_*(G^G(X')_{\mathfrak{m}_h}) \subset G^G(X)_{\mathfrak{m}_h}$.*

Proof. Suppose that $\alpha \in G^G(X')_{\mathfrak{m}_h}$. By definition, this means that α is in the kernel of $G^G(X') \rightarrow G^G(X')_{\mathfrak{m}_g}$ for all $g \neq h$. This is equivalent to saying that for all $g \neq h$ there exists $r \in R \setminus \mathfrak{m}_g$ such that $r\alpha = 0$. The equivariant projection formula implies that $rp_*(\alpha) = p_*(r\alpha) = 0$. Hence $p_*(\alpha) \in G^G(X)_{\mathfrak{m}_h}$. \square

2.4. Equivariant Chow groups and quotients. If X is an equidimensional scheme we will use the notation $CH_G^i(X)$ to denote the ‘‘codimension i ’’ equivariant Chow group $A_{\dim X - i}^G(X)$ defined in [EG1] (tensored with \mathbb{C}). If G acts with finite stabilizers then $CH_G^i(X)$ is generated by classes of G -invariant subvarieties [EG2, Prop. 5.2]. We reserve the notation $A_G^*(X)$ for the G -equivariant operational Chow ring defined in [EG1]. When X is smooth we can identify $A_G^*(X)$ with $CH_G^*(X)$ [EG1, Proposition 4].

Now suppose that G acts properly on X , with a geometric quotient $X \rightarrow Y$. By [EG1, Theorem 3], there is an isomorphism

$$\phi : CH_G^*(X) \rightarrow CH^*(Y),$$

defined as follows. Let $W \subset X$ be a G -invariant subvariety with image $Z \subset Y$. Set e_W to be the order of the stabilizer of the generic point of W . Then $\phi([W]_G) = \frac{1}{e_W}[Z]$; this map descends to give the isomorphism of Chow groups. The proof of this theorem (cf. [EG1, Proposition 11(a)]) shows that if $p : X' \rightarrow X$ is a finite equivariant morphism and $q : Y' \rightarrow Y$ is the corresponding morphism of quotients, then $\phi_X \circ p_* = q_* \circ \phi_{X'}$.

2.5. The equivariant Riemann-Roch map. By [EG2], there is an equivariant Riemann-Roch map $\tau_X^G : G^G(X) \rightarrow \prod_{i=0}^{\infty} CH_G^i(X)$, with the same functorial properties as the non-equivariant Riemann-Roch map of [BFM]. If G acts with finite stabilizers, then $CH_G^i(X) = 0$ for $i > \dim X$ so the Riemann-Roch map has target $CH_G^*(X)$.

Proposition 2.6. *Suppose that G is a diagonalizable group acting on X with finite stabilizers. Then $\tau_X^G(G^G(X)_{\mathfrak{m}_h}) = 0$ for $h \neq 1$, and $\tau_X^G : G^G(X)_{\mathfrak{m}_1} \rightarrow CH_G^*(X)$ is an isomorphism.*

Proof. Let $\widehat{G^G(X)}$ denote the completion of $G^G(X)$ with respect to the ideal \mathfrak{m}_1 . By [EG2, Theorem 3.1(a), Theorem 4.1] $\tau_X^G : G^G(X) \rightarrow CH_G^*(X)$

factors through an isomorphism $\widehat{G^G(X)} \rightarrow CH_G^*(X)$. The proof of Proposition 2.4 shows that there is an integer k_1 such that $\mathfrak{m}_1^{k_1} G^G(X) = \mathfrak{m}_1^{k_1+1} G^G(X)$. Thus

$$\widehat{G^G(X)} = G^G(X)/\mathfrak{m}_1^{k_1} G^G(X) = G^G(X)_{\mathfrak{m}_1}$$

and the map $G^G(X) \rightarrow \widehat{G^G(X)}$ is the projection $G^G(X) \rightarrow G^G(X)_{\mathfrak{m}_1}$. The proposition follows. \square

There is an equivariant Chern character map $\text{ch}^G : K^G(X) \rightarrow A_G^*(X)$. Suppose that X is smooth. If X is covered by G -invariant affines (for example, if G acts properly and a geometric quotient $X \rightarrow Y$ exists), then, identifying $G^G(X)$ with $K^G(X)$, the proof of [EG2, Theorem 3.1d] implies that $\tau_X^G(\alpha) = \text{ch}^G(\alpha) \text{Td}^G(T_X)$; here $\alpha \in G^G(X)$ and T_X is the tangent bundle of X .²

Proposition 2.7. *Suppose that X is smooth (so we identify $G^G(X)$ with $K^G(X)$) and that G acts on X with finite stabilizers. Let $\gamma \in G^G(X)$. Suppose that γ is a unit in $G^G(X)_{\mathfrak{m}_1}$. Then $\text{ch}^G(\gamma)$ is a unit in $CH_G^*(X)$. If $\frac{\alpha}{\gamma} \in G^G(X)_{\mathfrak{m}_1}$, then*

$$\tau_X^G\left(\frac{\alpha}{\gamma}\right) = \frac{1}{\text{ch}^G(\gamma)} \tau_X^G(\alpha).$$

Proof. By [EG2, Theorems 5.1 and 6.1], there is an isomorphism $\widehat{G^G(X)} \rightarrow G^G(X)_{\mathfrak{m}_1}$ where $\widehat{G^G(X)}$ is the completion along the augmentation ideal I of bundles of virtual rank 0 in $K^G(X) = G^G(X)$. Since γ is invertible after localizing at \mathfrak{m}_1 , it is invertible in the completion. Hence $\gamma \notin IG^G(X)$, so it has non-zero virtual rank. Hence $\text{ch}^G(\gamma) = \text{rk } \gamma + A$ where $A \in CH_G^{>0}(X)$. Thus $\text{ch}^G(\gamma)$ has an inverse in the completion $\prod_i CH_G^i(X)$. However, since G acts with finite stabilizers, $CH_G^i(X) = 0$ for $i > \dim X$, so the completion is just $CH_G^*(X)$. Then

$$\tau_X^G(\alpha) = \tau_X^G\left(\frac{\alpha}{\gamma} \cdot \gamma\right) = \tau_X^G\left(\frac{\alpha}{\gamma}\right) \cdot \text{ch}^G(\gamma).$$

Multiplying both sides by the inverse of $\text{ch}^G(\gamma)$ gives the result. \square

²This formula does not follow from the *statement* of Theorem 3.1d if G is not connected. However, the proof only requires that if X is a scheme then there is a system of pairs (U, V) such that the mixed spaces $X \times^G U$ are schemes. This is true if X is covered by a finite number of invariant affines.

2.6. Subgroup actions on equivariant K -theory for diagonalizable group actions. In this section we again assume that G is diagonalizable and let $R = R(G) \otimes \mathbb{C}$.

Let $H \subset G$ be a finite subgroup and set $\overline{G} = G/H$, $\overline{R} = R(\overline{G}) \otimes \mathbb{C}$. Define an H -action on R by

$$(2) \quad (h \cdot r)(g) = r(h^{-1}g).$$

Then $h \cdot \mathfrak{m}_g = \mathfrak{m}_{hg}$, and $R^H = \overline{R}$.

Assume that H acts trivially on X . If \mathcal{E} is a G -equivariant sheaf on X , and $U \subseteq X$ is any open subset, then any $h \in H$ induces a map (also denoted h)

$$h : \mathcal{E}(U) \rightarrow \mathcal{E}(U).$$

We can decompose $\mathcal{E}(U)$ into a sum of weight spaces for H :

$$\mathcal{E}(U) = \bigoplus_{\chi \in \hat{H}} \mathcal{E}_\chi(U),$$

where \hat{H} is the character group of H , and $\mathcal{E}_\chi(U)$ is the χ -weight space of $\mathcal{E}(U)$. This yields a decomposition of \mathcal{E} into H -eigensheaves:

$$\mathcal{E} = \bigoplus_{\chi \in \hat{H}} \mathcal{E}_\chi.$$

Each H -eigensheaf is G -equivariant, so we have $[\mathcal{E}] = \sum [\mathcal{E}_\chi]$ in $G^G(X)$. Define an H -action on $G^G(X)$ by the rule

$$(3) \quad h \cdot \sum [\mathcal{E}_\chi] = \sum \chi(h^{-1}) [\mathcal{E}_\chi].$$

If X is a point, then $G^G(X) = R$ and the action (3) agrees with the action (2). For general X (with H acting trivially), the H -actions on R and on $G^G(X)$ are compatible, in the sense that, for $r \in R$, $\alpha \in G^G(X)$,

$$h \cdot (r\alpha) = (h \cdot r)(h \cdot \alpha).$$

This implies that $h \cdot \mathfrak{m}_g G^G(X) = \mathfrak{m}_{hg} G^G(X)$, and, in particular,

$$h^{-1} \cdot \mathfrak{m}_h G^G(X) = \mathfrak{m}_1 G^G(X).$$

For simplicity of notation, we will sometimes write $\alpha(h)$ for $h^{-1} \cdot \alpha$.

We remark that there is a decomposition

$$G^G(X) \simeq G^{\overline{G}}(X) \otimes_{\overline{R}} R,$$

and the H -action on $G^G(X)$ defined in (3) coincides with the H -action on $G^{\overline{G}}(X) \otimes_{\overline{R}} R$ induced by the H -action on R .

The following lemma will be used in proving the Riemann-Roch theorem for quotients.

Lemma 2.8. *Suppose that G acts on X , and $H \subset G$ acts trivially on X . If $\alpha \in G^G(X)$, $h \in H$, then $\alpha^G = (h \cdot \alpha)^G$.*

Proof. We may assume that $\alpha = [\mathcal{E}] = \sum[\mathcal{E}_\chi]$. Then (with χ_1 denoting the trivial character of H), $[\mathcal{E}_\chi]^G = 0$ for $\chi \neq \chi_1$, so $[\mathcal{E}]^G = [\mathcal{E}_{\chi_1}]^G$ and $(h \cdot [\mathcal{E}])^G = \sum \chi(h)[\mathcal{E}_\chi]^G = \chi_1(h)[\mathcal{E}_{\chi_1}]^G$. The result follows since $\chi_1(h) = 1$. \square

3. THE RIEMANN-ROCH THEOREM FOR QUOTIENTS OF DIAGONALIZABLE GROUP ACTIONS

The main theorem of the paper is the following Riemann-Roch theorem for quotients.

Theorem 3.1. *Let G be a diagonalizable group acting properly on a smooth scheme X and let $X \rightarrow Y$ be a geometric quotient. If $h \in G$, let $i_h : X^h \hookrightarrow X$ denote the inclusion of the fixed locus of h , with normal bundle N_h ; let $j_h : Y^h = X^h/G \hookrightarrow Y$ denote the induced inclusion. If $\alpha \in K^G(X)$, then*

$$(4) \quad \tau_Y(\alpha^G) = \sum_{h \in \text{Supp } \alpha} j_{h*} \circ \phi_{X^h} \circ \tau_{X^h}^G(h^{-1} \cdot \frac{i_h^* \alpha}{\lambda_{-1}(N_h^*)}),$$

where $\text{Supp } \alpha$ is the (finite) set of points $h \in G$ such that $\alpha_h \neq 0$. Equivalently,

$$(5) \quad \tau_Y(\alpha^G) = \sum_{h \in \text{Supp } \alpha} \phi_X \circ i_{h*} \left(\frac{\text{ch}^G(i_h^* \alpha(h))}{\text{ch}^G(\lambda_{-1}(N_h^*)(h))} \text{Td}^G(T_{X^h}) \right).$$

Remark 3.2. When X and Y are quasi-projective, Theorem 3.1 follows from the Riemann-Roch theorem for Deligne-Mumford stacks proved in [To]. Our proof avoids the quasiprojectivity hypothesis. Note that the quotient of a quasi-projective variety need not be quasi-projective. For example, a simplicial toric variety whose fan is non-polytopal has no ample line bundles. In particular, our theorem can be applied to compute the Todd class of such toric varieties (Theorem 4.2).

Remark 3.3. In the right hand side of (4), the element $\frac{i_h^* \alpha}{\lambda_{-1}(N_h^*)}$ is viewed as an element of $G^G(X^h)_{\mathfrak{m}_h} \subset G^G(X^h)$. Since h^{-1} acts trivially on X^h , the results of the previous subsection imply that $h^{-1} \cdot \frac{i_h^* \alpha}{\lambda_{-1}(N_h^*)}$ is an element of $G^G(X^h)_{\mathfrak{m}_1} \subset G^G(X^h)$.

Remark 3.4. If $\alpha = [\mathcal{E}]$ where \mathcal{E} is an equivariant coherent sheaf then the sums in equations (4) and (5) are in $CH^*(Y) \otimes \mathbb{Q}$ even though the individual terms are in $CH^*(Y) \otimes \mathbb{C}$.

The remainder of the section will be devoted to the proof of this theorem. Observe that (5) follows from (4) by Section 2.5.

Let $\alpha = \sum \alpha_h$ be the decomposition corresponding to Proposition 2.4. We will prove that for all $h \in G$,

$$(6) \quad \tau_Y((\alpha_h)^G) = j_{h*} \circ \phi_{X^h} \circ \tau_{X^h}^G(h^{-1} \cdot \frac{i_h^* \alpha}{\lambda_{-1}(N_h^*)}).$$

Step 1. Suppose that G acts freely on a (not necessarily smooth) scheme X' with quotient scheme Y' . We claim that for $\beta \in G^G(X')$,

$$(7) \quad \tau_Y(\beta^G) = \phi_X \circ \tau_X^G(\beta).$$

Indeed, this follows from [EG2, Theorem 3.1(e)], since we identify $G^G(X')$ with $G(Y')$ by the map $\beta \mapsto \beta^G$, and $CH_G^*(X')$ with $CH^*(Y')$ by the map $\phi_{X'}$.

Note that the freeness of the action implies that $\beta_h = 0$ for $h \neq 1$; so $\beta = \beta_1$.

Step 2. We prove (6) for $h = 1$. In this case, we want to show that

$$(8) \quad \tau_Y((\alpha_1)^G) = \phi_X \circ \tau_X^G(\alpha).$$

By Proposition 2.6, $\tau_X^G(\alpha_h) = 0$ for $h \neq 1$, so $\tau_X^G(\alpha) = \tau_X^G(\alpha_1)$.

Lemma 3.5. *Let $p: X' \rightarrow X$ be a finite surjective G -equivariant morphism of (not necessarily smooth) G -schemes, where G acts properly on X (and hence also on X' by Proposition 2.1). Then $p_*: G^G(X')_{\mathfrak{m}_1} \rightarrow G^G(X)_{\mathfrak{m}_1}$ is surjective.*

Proof. By [EG2, Theorem 3.1], we have a commutative diagram

$$\begin{array}{ccc} G^G(X')_{\mathfrak{m}_1} & \xrightarrow{\tau_{X'}^G} & CH_G^*(X') \\ \downarrow p_* & & \downarrow p_* \\ G^G(X)_{\mathfrak{m}_1} & \xrightarrow{\tau_X^G} & CH_G^*(X). \end{array}$$

The horizontal arrows are isomorphisms by Proposition 2.6. Since G acts properly, the rational equivariant Chow groups are generated by G -invariant cycles ([EG1, Proposition 13(a)]), so the right hand p_* is surjective. Therefore the left hand p_* is surjective. This proves the lemma. \square

By [Se, Theorem 6.1] there exists a finite surjective morphism $p: X' \rightarrow X$ of G -schemes such that G acts freely on X' (we do not require that X' be smooth). By Proposition 2.1 there is a geometric quotient $X' \rightarrow Y'$ such that the induced map $q: Y' \rightarrow Y$ is also finite. The preceding lemma implies that there exists $\beta_1 \in G^G(X')_{\mathfrak{m}_1}$ such that $p_*\beta_1 = \alpha_1$. By Lemma 2.5, $q_*((\beta_1)^G) = \alpha_1^G$. Therefore,

$$\tau_Y((\alpha_1)^G) = \tau_Y(q_*((\beta_1)^G)) = q_*\tau_{Y'}((\beta_1)^G).$$

By Step 1, this equals $q_* \circ \phi_{X'} \circ \tau_{X'}^G(\beta_1)$. As noted in Section 2.4, $q_* \circ \phi_{X'} = \phi_X \circ p_*$. Hence

$$q_* \circ \phi_{X'} \circ \tau_{X'}^G(\beta_1) = \phi_X \circ p_* \circ \tau_{X'}^G(\beta_1) = \phi_X \circ \tau_X^G \circ p_*(\beta_1) = \phi_X \circ \tau_X^G(\alpha_1).$$

This completes Step 2.

Step 3. We prove (6) for arbitrary h . As usual, we view $G^G(X)_{\mathfrak{m}_h}$ and $G^G(X^h)_{\mathfrak{m}_h}$ as summands in $G^G(X)$ and $G^G(X^h)$, respectively. Let

$$\beta_h = \frac{i_h^* \alpha}{\lambda_{-1}(N_h^*)} \in G^G(X^h)_{\mathfrak{m}_h}.$$

By Proposition 2.3, $(i_{h*} \beta_h)^G = j_{h*}((\beta_h)^G)$. Hence

$$(9) \quad \tau_Y((\alpha_h)^G) = \tau_Y((i_{h*} \beta_h)^G) = \tau_Y \circ j_{h*}((\beta_h)^G) = j_{h*} \circ \tau_{Y^h}((\beta_h)^G),$$

Now, $h^{-1} \cdot \beta_h \in G^G(X^h)_{\mathfrak{m}_1}$, and by Lemma 2.8, $(h^{-1} \cdot \beta_h)^G = (\beta_h)^G$. By Step 2,

$$\tau_{Y^h}((h^{-1} \cdot \beta_h)^G) = \phi_{X^h} \circ \tau_{X^h}^G(h^{-1} \cdot \beta_h) = \phi_{X^h} \circ \tau_{X^h}^G(h^{-1} \cdot \frac{i_h^* \alpha}{\lambda_{-1}(N_h^*)}).$$

Substituting this into (9) yields (6). This completes the proof of Theorem 3.1.

4. TODD CLASSES OF SIMPLICIAL TORIC VARIETIES

In this section we apply the Riemann-Roch formula of Theorem 3.1 to give a formula for the Todd class, $\tau(\mathcal{O}_X)$, of a simplicial toric variety X . When X is complete this is the same as the formula of Brion and Vergne [BV, Theorem 4.1].

4.1. Global coordinates on toric varieties. We begin by recalling some basic facts about toric varieties, following notation as in [F] and [BV], as well as the construction of global coordinates in [C].

Let T be a d -dimensional torus. Denote the the lattice of one-parameter subgroups by $N = \text{Hom}(\mathbb{C}^*, T)$ and the character group by $M = \text{Hom}(T, \mathbb{C}^*)$. A toric variety X is a normal variety on which T acts, with a dense orbit isomorphic to T . Such a variety is determined by its fan Σ in $N_{\mathbb{R}}$. The toric variety is said to be *simplicial* if every cone in the fan is generated by linearly independent vectors. In this case the toric variety has finite quotient singularities. If each cone is generated by elements of part of a basis for N , then the toric variety is smooth. The toric variety X is complete if and only if the support of Σ is $N_{\mathbb{R}}$.

Denote by $\Sigma(k)$ the set of k -dimensional cones of Σ . If σ is a cone, let $\sigma(k)$ denote its k -dimensional faces. A k -dimensional cone $\tau \in \Sigma(k)$ determines a $d - k$ -dimensional T -invariant subvariety $V(\tau) \subset X$.

If $\tau \in \Sigma(1)$ is a 1-dimensional cone we will refer to the primitive lattice vector generating the cone by n_τ . Then $A_{d-1}(X)$ is generated by $\{[V(\tau)]\}_{\tau \in \Sigma(1)}$.

Define a map $(\mathbb{C}^*)^{\Sigma(1)} \rightarrow T$ by $(t_\tau)_{\tau \in \Sigma(1)} \rightarrow \prod_{\tau \in \Sigma(1)} n_\tau(t_\tau)$. Let G be the kernel of this map of tori. Let $Z(\Sigma)$ be the subscheme of $\mathbb{A}^{\Sigma(1)}$ defined by the monomial ideal $B(\Sigma) = \langle \hat{x}_\sigma : \sigma \in \Sigma \rangle$, where $\hat{x}_\sigma = \prod_{\tau \notin \sigma(1)} x_\tau$ and $\{x_\tau\}_{\tau \in \Sigma(1)}$ are coordinates on $\mathbb{A}^{\Sigma(1)}$. The group G acts on $W = \mathbb{A}^{\Sigma(1)} - Z(\Sigma)$ by the restriction of the diagonal $(\mathbb{C}^*)^{\Sigma(1)}$ action. Let $a_\tau : (\mathbb{C}^*)^{\Sigma(1)} \rightarrow \mathbb{C}^*$ be the projection to the τ -th factor. We will also refer to the restriction of this character to any subgroup of $(\mathbb{C}^*)^{\Sigma(1)}$ as a_τ .

Theorem 4.1. [C] *X is a simplicial toric variety if and only if X is a geometric quotient W/G .*

Under the quotient map $W \rightarrow X$, the coordinate hyperplane x_τ maps to the invariant Weil divisor $V(\tau)$. More generally, if σ is the k -dimensional cone generated by τ_1, \dots, τ_k , then $V(\sigma)$ is the image under the quotient map of the linear subspace W_σ defined by the ideal $(x_{\tau_1}, x_{\tau_2}, \dots, x_{\tau_k})$. Let G_σ be the stabilizer of the linear subspace W_σ . By definition,

$$\begin{aligned} G_\sigma &= \{g \in G \mid a_\tau(g) = 1 \text{ if } \tau \notin \sigma(1)\} \\ &= \{(t_\tau)_{\tau \in \sigma(1)} \mid t_\tau \in \mathbb{C}^*, \prod_{\tau \in \sigma(1)} n_\tau(t_\tau) = 1\}. \end{aligned}$$

This implies that G_σ is a finite group whose order is the order of $N_\sigma / \sum_{\tau \in \sigma(1)} \mathbb{Z}n_\tau$, the multiplicity of the cone σ . Let G_Σ be the union of the subgroups G_σ of G .

For any $g \in G$, set

$$g(1) = \{\tau \in \Sigma(1) \mid a_\tau(g) \neq 1\}.$$

The fixed locus of g in $\mathbb{A}^{\Sigma(1)}$ is the linear subspace cut out by the ideal generated by $\{x_\tau\}_{\tau \in g(1)}$. Thus, W^g is the intersection of this linear subspace with the open set $W \subset \mathbb{A}^{\Sigma(1)}$. Observe that, by definition, the linear space defined by the ideal $(x_{\tau_1}, x_{\tau_2}, \dots, x_{\tau_k})$ is in $Z(\Sigma)$ if and only if the rays τ_1, \dots, τ_k do not all lie in a single cone of the fan Σ . This implies that W^g is nonempty if and only if $g \in G_\sigma$ for some cone σ . Hence the support of the R -module $K^G(W)$ is $G_\Sigma \subset G = \text{Spec } R$.

4.2. Todd classes of simplicial toric varieties. The Todd class of the toric variety X is by definition $\text{Td}(X) := \tau(\mathcal{O}_X)$. Since $\mathcal{O}_X = (\mathcal{O}_W)^G$, we can apply Theorem 3.1 to obtain a formula for $\text{Td}(X)$. When X is complete this formula was previously obtained by Brion and Vergne [BV, Theorem 4.1].

For $\tau \in \Sigma(1)$ let $z_\tau \in CH_G^1(W)$ be the equivariant fundamental class of the invariant divisor $\{x_\tau = 0\}$. The generic stabilizer of this divisor is G_τ , which by the preceding subsection is trivial. Hence $F_\tau := \phi(z_\tau)$ is the fundamental class of the Weil divisor $V(\tau)$.

Theorem 4.2.

$$(10) \quad \mathrm{Td}(X) = \tau(\mathcal{O}_X) = \sum_{g \in G_\Sigma} \prod_{\tau \in \Sigma(1)} \frac{F_\tau}{1 - a_\tau(g)e^{-F_\tau}}.$$

Proof. By Theorem 3.1

$$(11) \quad \mathrm{Td}(X) = \tau(\mathcal{O}_W^G) = \sum_{g \in G_\Sigma} \phi \circ i_{g*} \left(\frac{\mathrm{Td}^G(T_{W^g})}{\mathrm{ch}^G(\lambda_{-1}(N_{W^g}^*(g)))} \right).$$

As noted above, W^g is the intersection of W with the linear subspace cut out by the ideal generated by $\{x_\tau\}_{\tau \in g(1)}$. Hence

$$(12) \quad [W^g] = \prod_{\tau \in g(1)} z_\tau \in CH_G^*(W).$$

The equivariant normal bundle to $W^g \subset W$ is the pullback to W^g of the direct sum of equivariant line bundles

$$\bigoplus_{\tau \in g(1)} \mathcal{O}(x_\tau).$$

Observe that this direct sum decomposition is also a decomposition into eigenbundles for the finite subgroup $H = \langle g \rangle$, where the weight on the τ -th factor is the character a_τ . Since $c_1(\mathcal{O}(x_\tau)) = z_\tau$, we obtain

$$(13) \quad \mathrm{ch}^G(\lambda_{-1}(N_{W^g}^*(g))) = i_g^* \left(\prod_{\tau \in g(1)} 1 - a_\tau^{-1}(g)e^{-z_\tau} \right).$$

Likewise, the tangent bundle to W^g splits as

$$\bigoplus_{\tau \notin g(1)} \mathcal{O}(x_\tau).$$

Hence,

$$(14) \quad \mathrm{Td}^G(T_{W^g}) = i_g^* \left(\prod_{\tau \notin g(1)} \frac{z_\tau}{1 - e^{-z_\tau}} \right).$$

If $\tau \notin g(1)$, then $a_\tau(g) = 1$, so

$$(15) \quad \mathrm{Td}^G(T_{W^g}) = i_g^* \left(\prod_{\tau \notin g(1)} \frac{z_\tau}{1 - a_\tau(g^{-1})e^{-z_\tau}} \right).$$

Substituting (12)–(15) into equation (11) shows that the contribution of W^g to the Todd class is

$$(16) \quad \phi \left(\prod_{\tau \in \Sigma(1)} \frac{z_\tau}{1 - a_\tau(g^{-1})e^{-z_\tau}} \right)$$

Summing over all $g \in G_\Sigma$ and noting that $\phi(z_\tau) = F_\tau$ yields the Brion-Vergne formula. (Note that we are using the fact that $g \in G_\Sigma$ if and only if $g^{-1} \in G_\Sigma$.) \square

Remark 4.3. When X is complete, then the image of formula (10) in cohomology is the same as the restriction from T -equivariant cohomology of the formula in [BV, Theorem 4.1] to ordinary cohomology. The methods of this paper can also be used to yield a formula for the T -equivariant Todd class in $CH_T^*(Y)$, but we do not pursue this here.

Acknowledgments. The authors are grateful to Michel Brion for helpful comments on an earlier version of the paper.

REFERENCES

- [BFM] Baum, P., Fulton, W. and MacPherson, R., *Riemann-Roch for singular varieties*, Publ. Math. I.H.E.S. **45** (1975), 101–145.
- [B] Borel, A., *Linear Algebraic Groups* (2nd enlarged edition), G.T.M **126**, Springer-Verlag (1991).
- [BV] Brion, M. and Vergne M., *An equivariant Riemann-Roch theorem for complete simplicial toric varieties*, J. Reine. Agnew. Math. **482** (1997), 67–92.
- [C] Cox, D., *The homogeneous coordinate ring of a toric variety*, J. Algebraic Geom. **4** (1995), 17–50.
- [EG1] Edidin, D. and Graham, W., *Equivariant intersection theory*, Invent. Math., **131** (1998), 595–634.
- [EG2] Edidin, D. and Graham, W., *Riemann-Roch for equivariant Chow groups*, Duke Math J., **102** (2000), 567–594.
- [F] Fulton, W., *Introduction to Toric Varieties*. Annals of Mathematics Studies, **131**, Princeton University Press (1993).
- [GGK] Ginzburg, V., Guillemin, V. and Karshon, E., *Cobordism techniques in differential geometry*, monograph, to be published by the American Mathematical Society.
- [Gr] Grosshans, F., *Algebraic Homogeneous Spaces and Invariant Theory*, Lecture Notes in Math, **1673**, Springer-Verlag 1997.
- [Gu] Guillemin, V., *Riemann-Roch for toric orbifolds*, J. Diff. Geom. **45** (1997), 53–73.
- [MFK] Mumford, D., Fogarty, J. and Kirwan, F., *Geometric Invariant Theory: 3rd enlarged edition*, Springer-Verlag 1994.
- [Se] Seshadri, C. S., *Quotient spaces modulo algebraic groups*, Ann. Math. **95** (1972), 511–556.
- [Su] Sumihiro, H., *Equivariant completion*, J. Math. Kyoto Univ. **14** (1974), 1–28.
- [Th1] Thomason, R., *Equivariant resolution, linearization, and Hilbert’s fourteenth problem over arbitrary base schemes*, Adv. in Math. **65** (1987), 16–34.
- [Th2] Thomason, R., *Une formule de Lefschetz en K -théorie équivariante algébrique*, Duke Math. J. **68** (1992), 447–462.
- [To] Toen, B., *Théorèmes de Riemann-Roch pour les champs de Deligne-Mumford*. K -Theory **18** (1999), 33–76.