

NONCOMMUTATIVE RICCI CURVATURE AND DIRAC OPERATOR ON $\mathbb{C}_q[SL_2]$ AT ROOTS OF UNITY

Shahn Majid¹

School of Mathematical Sciences
Queen Mary, University of London, Mile End Rd
London E1 4NS, UK

12 June – revised August, 2002

Abstract We find a unique torsion free Riemannian spin connection for the natural Killing metric on the quantum group $\mathbb{C}_q[SL_2]$, using a recent frame bundle formulation. We find that its covariant Ricci curvature is essentially proportional to the metric (i.e. an Einstein space). We compute the Dirac operator \not{D} and find for q an odd r 'th root of unity that its eigenvalues are given by q -integers $[m]_q$ for $m = 0, 1, \dots, r - 1$ offset by the constant background curvature. We fully solve the Dirac equation for $r = 3$.

Keywords: noncommutative geometry, quantum groups, Riemann tensor, gravity, roots of unity, Jones index, spinor.

1 Introduction

This is a follow-up to our work [1] where we covered the electromagnetic theory on the quantum group $\mathbb{C}_q[SL_2]$ at a root of unity. We now compute the canonical Riemannian geometry induced by the Killing form using the frame-bundle formulation of noncommutative geometry on algebras in [2][3]. Just as compact Lie groups are natural examples of Riemannian manifolds, so quantum groups such as $\mathbb{C}_q[SL_2]$ *should* be natural examples of noncommutative Riemannian geometry. We will see that this is indeed the case if we use the frame-bundle formulation coming out of quantum group theory.

There are already a lot of attempts to q -deform geometric structures, including so-called Levi-Civita connections on specific quantum groups, for example [4], but on a largely ad-hoc or case-by-case basis. The problem remains how definitive or complete any of these various q -formulae might be. Our goal in the present paper is exactly the opposite: the frame bundle formulation is defined generally at the level of (in principle) any unital algebra, including matrix algebras, finite group function algebras and algebras not connected in any way to q -deformations. In other words a ‘functorial theory’ at an algebraic level. Our goal is to

¹Royal Society University Research Fellow

take this existing theory and specialise it to $\mathbb{C}_q[SL_2]$ as a test of the general theory. We find reasonable answers including a unique spin connection, a constant Ricci curvature and an interesting Dirac operator \mathcal{D} . We study its modes at roots of unity where all constructions are finite. The root of unity theory is perhaps the most interesting both mathematically and physically. It is the case occurring naturally in the WZW model and hence in some approaches to quantum gravity[5]. The geometry in this case is little studied, however; aside from [1], we note [6] in connection with quantum planes at roots of unity.

We refer to [3] for the frame-bundle formulation but will recall the key formulae for torsion, metric compatibility, Ricci tensor etc. when we need them. The main ingredients are a (possibly noncommutative) algebra \mathcal{A} of ‘functions’ for the base space, a quantum group H for the functions on the frame group, a quantum principal bundle with some algebra P and fixed subalgebra \mathcal{A} and of course compatible cotangent bundles $\Omega^1(\mathcal{A}), \Omega^1(H), \Omega^1(P)$. Then we need a representation of the frame quantum group and a framing isomorphism of $\Omega^1(\mathcal{A})$ with the corresponding associated vector bundle. In the case of interest in the present paper, we take $\Omega^1(H) = H \otimes \Lambda^1$, where Λ^1 is the dual of the (braided) Lie algebra of the frame group, and we take $P = \mathcal{A} \otimes H$ a trivial bundle. The framing isomorphism just amounts to a left \mathcal{A} -basis $\{e_i\}$ of $\Omega^1(\mathcal{A})$ or ‘vielbein’. A spin connection just amounts to a Lie-algebra valued 1-form or a map $A : \Lambda^1 \rightarrow \Omega^1(\mathcal{A})$. In our case of a quantum group \mathcal{A} we take $H = \mathcal{A}$ i.e. reduce the frame quantum group to \mathcal{A} itself. Then there is a natural choice of vielbein as explained in [3], given by the Maurer-Cartan form. So the new result of the present paper is to obtain a unique compatible spin connection and thereby compute the entire Riemannian geometry at least for $\mathbb{C}_q[SL_2]$. We expect similar results for other standard $\mathbb{C}_q[G]$. Let us mention one interesting phenomenon; as for the finite group S_3 in [3] we find that the Ricci tensor is essentially proportional to the Killing metric (an ‘Einstein space’) but shifted by an invariant element $\theta \otimes \theta$. Here θ is a ‘time-direction’ induced by the noncommutative geometry and not part of the 3-geometry of $\mathbb{C}_q[SL_2]$. The shift by $\theta \otimes \theta$ turns out to be exactly such that the Ricci tensor has zero modes.

It is also a very interesting question how such examples as $\mathbb{C}_q[SL_2]$ could fit into the mathematically more sophisticated K-theory and operator algebra approach of A. Connes and others[7]. There the geometry is defined by a spectral triple or ‘axiomatic Dirac operator’ obeying some axioms suggested by operator theory, including self-adjointness. By contrast we find that our constructed Dirac operator on $\mathbb{C}_q[SL_2]$ at an odd root of unity has eigenvalues on a complex circle with properties perhaps reminiscent of the discrete series Jones index of type II_1 subfactors.

2 Preliminaries

Here we fix notations in the conventions that we will use and do some preliminary computations, in Section 2.2. We let $q^2 \neq 1$. The quantum group $\mathcal{A} = \mathbb{C}_q[SL_2]$ has a matrix of generators $\{t^\alpha_\beta\} = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$ with relations

$$ba = qab, \quad ca = qac, \quad db = qbd, \quad dc = qcd, \quad cb = bc, \quad da - ad = q\mu bc, \quad ad - q^{-1}bc = 1,$$

where $\mu = 1 - q^{-2}$. The coproduct Δ and counit ϵ have the usual matrix coalgebra form. We denote the antipode or ‘matrix inverse’ by S .

We will also work with the dimensional Hopf algebra $\mathcal{A} = \mathbb{C}_q[SL_2]$ reduced at q a primitive r 'th root of unity. This has the further relations

$$c^r = b^r = 0, \quad a^r = d^r = 1$$

where a^r, b^r, c^r, d^r generate an undeformed $\mathbb{C}[SL_2]$ central sub-Hopf algebra of the original $\mathbb{C}_q[SL_2]$. Note also that in the reduced case $a = (1+q^{-1}bc)d^{-1}$ is redundant and $\dim(\mathcal{A}) = r^3$. In this case a basis of \mathcal{A} is $\{c^k b^n d^m\}$ for $0 \leq m, n, k \leq r - 1$. Explicit computations are done via Mathematica for $r = 3, 5, 7$ for concreteness, but we expect identical results for all odd r . Equations involving only the invariant differential forms do not directly involve the function algebra and are solved for all $q^2 \neq -1$.

2.1 Exterior algebra

We take the standard bicovariant exterior algebra on $\mathbb{C}_q[SL_2]$ which has the lowest dimensional (4d) space of 1-forms[8]. This is the same set-up as in [1] except that we carefully switch to Λ^1 the space of left-invariant 1-forms as used in [3] (rather than right-invariant). Thus, we take a basis $\{e_\alpha^\beta\} = \begin{pmatrix} e_a & e_b \\ e_c & e_d \end{pmatrix}$, where $e_1^2 = e_b$, etc., and form a right crossed module with right multiplication and the right coaction

$$\Delta_R(e_\alpha^\beta) = e_\gamma^\delta \otimes t^\gamma_\alpha S t^\beta_\delta.$$

$\Omega^1 = \mathcal{A} \otimes \Lambda^1$ is spanned by the left-invariant forms as a free left module over $\mathbb{C}_q[SL_2]$. These also generate the invariant exterior algebra Λ and $\Omega = \mathcal{A} \otimes \Lambda$. Here e_a, e_b, e_c behave like usual forms or Grassmann variables and

$$e_a \wedge e_d + e_d \wedge e_a + \mu e_c \wedge e_b = 0, \quad e_d \wedge e_c + q^2 e_c \wedge e_d + \mu e_a \wedge e_c = 0$$

$$e_b \wedge e_d + q^2 e_d \wedge e_b + \mu e_b \wedge e_a = 0, \quad e_d^2 = \mu e_c \wedge e_b. \tag{1}$$

It is useful to define $e_z \equiv q^{-2}e_a - e_d$, which obeys

$$e_z \wedge e_c + q^2 e_c \wedge e_z = 0, \quad e_b \wedge e_z + q^2 e_z \wedge e_b = 0, \quad e_z \wedge e_z = (1 - q^{-4})e_c \wedge e_b. \quad (2)$$

The relations among 1-forms are obtained by setting to zero the kernel of $\Psi - \text{id}$, where Ψ is the crossed-module braiding. Its specific form can be obtained by converting the right-invariant formulae in [1] to left ones by $e_a \leftrightarrow e_d$ and left-right reversal of all products including tensor products. We will need the eigenvectors of Ψ in Section 4.

The right module structure on 1-forms is defined via the commutation relations

$$e_a \begin{pmatrix} a & b \\ c & d \end{pmatrix} = \begin{pmatrix} qa & q^{-1}b \\ qc & q^{-1}d \end{pmatrix} e_a, \quad [e_c, b] = [e_c, d] = [e_b, a] = [e_b, c] = 0$$

$$[e_c, a] = q\mu b e_a, \quad [e_c, c] = q\mu d e_a, \quad [e_b, b] = q\mu a e_a, \quad [e_b, d] = q\mu c e_a$$

$$[e_d, a]_{q^{-1}} = \mu b e_b, \quad [e_d, b]_q = \mu a e_c + q\mu^2 b e_a, \quad [e_d, c]_{q^{-1}} = \mu d e_b, \quad [e_d, d]_q = \mu c e_c + q\mu^2 d e_a$$

where $[x, y]_q = xy - qyx$ and $a, b, c, d \in \mathbb{C}_q[SL_2]$.

2.2 Exterior derivative and Lie bracket structure constants

The differential on the exterior algebra structure is defined by graded anticommutator $d = \mu^{-1}[\theta, \]$ where $\theta = e_a + e_d$. In particular

$$de_a = -e_c \wedge e_b, \quad de_d = e_c \wedge e_b, \quad de_c = q^2 e_c \wedge e_z, \quad de_b = -e_b \wedge e_z, \quad de_z = -(q^{-2} + 1)e_c \wedge e_b. \quad (3)$$

Lemma 2.1 *cf.[1] For all invertible $q^2 \neq 1$,*

$$\begin{aligned} e_a.c^k b^n d^m &= q^{-m-n+k} c^k b^n d^m .e_a \\ e_b.c^k b^n d^m &= c^k b^n d^m .e_b + q^2 \mu [n] q^{-m-n} c^k b^{n-1} d^{m-1} e_a + q^{1-n} \mu [m+n] c^{k+1} b^n d^{m-1} e_a \\ e_c.c^k b^n d^m &= c^k b^n d^m .e_c + q^{k-m} \mu [k] c^{k-1} b^n d^{m+1} e_a \\ e_d.c^k b^n d^m &= q\mu^2 ([k+1][m+n] c^k b^n d^m + q^{-m} [k][n] c^{k-1} b^{n-1} d^m) .e_a + \mu [k] q^n c^{k-1} b^n d^{m+1} e_b \\ &\quad + q^{-k} \mu ([n] c^k b^{n-1} d^{m-1} + q^{-1} (q^{m+n} [m] + [n]) c^{k+1} b^n d^{m-1}) e_c + q^{-k+n+m} c^k b^n d^m .e_d \end{aligned}$$

where terms with negative powers of b, c are omitted.

Proof This is a left-handed version of [1], so we will be brief. Writing $[n] = (q^n - q^{-n})/(q - q^{-1})$ and $u = \begin{pmatrix} a \\ c \end{pmatrix}$, $v = \begin{pmatrix} b \\ d \end{pmatrix}$, we iterate the bimodule relations to obtain

$$\begin{aligned} e_d v^n &= q^n v^n e_d + q^{n-1} [n] \mu u v^{n-1} e_c + q\mu^2 [n] v^n e_a, & e_b v^n &= v^n e_b + q\mu [n] u v^{n-1} e_a \\ e_c v^n &= v^n e_c, & e_a v^n &= q^{-n} v^n e_a \\ e_d u^n &= q^{-n} u^n e_d + q^{1-n} [n] \mu v u^{n-1} e_b, & e_b u^n &= u^n e_b \\ e_c u^n &= u^n e_c + q\mu [n] v u^{n-1} e_a, & e_a u^n &= q^n u^n e_a. \end{aligned}$$

These then give the commutation relations with basis elements as stated. \diamond

From these it is clear how the right handed derivatives in [1] transpose to our left handed ones. We easily obtain

$$\begin{aligned}
d(c^k b^n d^m) &= \mu^{-1}(q^{m+n-k} - 1) c^k b^n d^m e_d + q^{n-k+1} [k]_{q^2} c^{k-1} b^n d^{m+1} e_b \\
&+ q^{-k-n} ([m+n]_{q^2} c^{k+1} b^n d^{m-1} + q[n]_{q^2} c^k b^{n-1} d^{m-1}) e_c \\
&+ \mu q^{-k-m-n+2} ([k+1]_{q^2} [m+n]_{q^2} c^k b^n d^m + q[n]_{q^2} [k]_{q^2} c^{k-1} b^{n-1} d^m) e_a \\
&+ \mu^{-1}(q^{-m-n+k} - 1) c^k b^n d^m e_a
\end{aligned} \tag{4}$$

obeying the Leibniz rule. We write $[n]_{q^2} = (1 - q^{2n})/(1 - q^2)$.

Next, we will need the projection $\tilde{\pi} : \mathcal{A} \rightarrow \Lambda^1$ which characterises the above calculus as a quotient of the universal one, i.e. with $df = f \circ \tilde{\pi}(f_{(2)})$ for all $f \in \mathcal{A}$. Here $\Delta f = f \circ \otimes f_{(2)}$ is the Sweedler notation for the coproduct. Actually $\tilde{\pi}$ can be obtained backwards from d as follows. Let the partial derivatives $\partial^i : \mathcal{A} \rightarrow \mathcal{A}$ be defined by $df = \sum_i (\partial^i f) e_i$. Then

$$\tilde{\pi}(f) = \sum_i e_i \epsilon(\partial^i f). \tag{5}$$

In particular, we obtain

$$\tilde{\pi}(a) = \frac{q}{[2]_q} (q e_a - e_d), \quad \tilde{\pi}(c) = e_b, \quad \tilde{\pi}(b) = e_c, \quad \tilde{\pi}(d) = \frac{1}{[2]_q} (q^2 e_d + (q^2 - q^{-1} - 1) e_a). \tag{6}$$

We use the formula (4) for the exterior derivative. For generic q we compute da separately.

Finally, we need the braided-Lie algebra structure constants

$$\text{ad}_R = (\text{id} \otimes \tilde{\pi}) \Delta_R, \quad \text{ad}_L = (\tilde{\pi} \otimes \text{id}) \Delta_L \tag{7}$$

where $\Delta_L = (S^{-1} \otimes \text{id}) \circ \tau \circ \Delta_R$ is the right coaction converted to a left coaction and τ is the usual vector space flip. We have

$$\begin{aligned}
\text{ad}_R(e_a) &= e_c \otimes e_b - e_b \otimes e_c + \mu(e_a - q^2 e_d) \otimes e_a \\
\text{ad}_R(e_d) &= e_b \otimes e_c - e_c \otimes e_b - \mu(e_a - q^2 e_d) \otimes e_a \\
\text{ad}_R(e_b) &= e_b \otimes (q^2 e_a - e_d) - (q^{-2} e_a - e_d) \otimes e_b \\
\text{ad}_R(e_c) &= e_c \otimes ((q^2 - 1 - q^{-2}) e_a + q^2 e_d) + (e_a - q^2 e_d) \otimes e_c \\
\text{ad}_L(e_a) &= e_c \otimes e_b - e_b \otimes e_c + \mu e_a \otimes (e_a - q^2 e_d) \\
\text{ad}_L(e_d) &= e_b \otimes e_c - e_c \otimes e_b - \mu e_a \otimes (e_a - q^2 e_d) \\
\text{ad}_L(e_c) &= (q^2 e_a - e_d) \otimes e_c - e_c \otimes (q^{-2} e_a - e_d) \\
\text{ad}_L(e_b) &= ((q^2 - 1 - q^{-2}) e_a + q^2 e_d) \otimes e_b + e_b \otimes (e_a - q^2 e_d)
\end{aligned} \tag{8}$$

We define by $\text{ad}(e_i) = \sum_{j,k} \text{ad}(jk|i) e_j \otimes e_k$, where we use the indices i, j, k to run through $\{e_a, e_b, e_c, e_d\}$. They are given as above for all q and are versions of the structure constants of the braided-Lie algebra $\widetilde{sl_{q,2}}$ as explained in [3].

3 Spin connection and Riemannian curvature

Still with q arbitrary, there is a natural Δ_R -covariant Killing metric in [3] of the form

$$\eta = e_c \otimes e_b + q^2 e_b \otimes e_c + \frac{(e_a \otimes e_a - q e_a \otimes e_d - q e_d \otimes e_a + q(q^2 + q - 1)e_d \otimes e_d)}{[2]_q} + \lambda \theta \otimes \theta$$

where $\lambda = q(1 - q - q^2)/(1 + q)$ is the natural choice for the Hodge $*$ operator as explained in [1]. In this case we have more simply

$$\eta = e_c \otimes e_b + q^2 e_b \otimes e_c + \frac{q^4}{[2]_{q^2}}(e_z \otimes e_z - \theta \otimes \theta) \quad (9)$$

which (for real q) is the usual q -Minkowski space metric cf. [9]. This is the local cotangent space of $\mathbb{C}_q[SL_2]$ with θ an intrinsic ‘time’ direction induced by noncommutative geometry. We can add any multiple of $\theta \otimes \theta$ and still retain Δ_R -invariance.

For any such invariant metric, we have symmetry in the sense

$$\wedge(\eta) = 0. \quad (10)$$

Moreover, the equations for a torsion-free and skew-metric-compatible ‘generalised Levi-Civita’ spin connection become independent of η and reduce to the torsion and ‘cotorsion’ equations[3]:

$$de_i + \sum_{j,k} A_j \wedge e_k \operatorname{ad}_L(jk|i) = 0, \quad de_i + \sum_{j,k} e_j \wedge A_k \operatorname{ad}_R(jk|i) = 0, \quad \forall i. \quad (11)$$

In these equations we write $A(e_i) \equiv A_i$, and a generalised spin connection is given by four such 1-forms A_a, A_b, A_c, A_d obeying (11). In principle there is also an optional ‘regularity’ condition as explained in [3] which ensures that the curvature is braided-Lie algebra valued. By the same arguments as above, this regularity condition can be written as

$$\sum_{i,j} A_i \wedge A_j \epsilon(\partial^i \partial^j f) = 0, \quad \forall f \in \ker \tilde{\pi}. \quad (12)$$

Theorem 3.1 *For generic q or for q an odd root of unity there is a unique torsion-free and cotorsion-free spin connection given by*

$$A_a = \frac{q^4}{[4]_{q^2}} e_z = -A_d, \quad A_b = \frac{1}{q^2 + q^{-2}} e_b, \quad A_c = \frac{1}{q^2 + q^{-2}} e_c.$$

The connection is not in general regular.

Proof We define the components of the connection by $A_i = A^j{}_i e_j$ and have to solve for this 4×4 matrix of elements of $\mathbb{C}_q[SL_2]$. Looking first at the torsion equation, we see that the coefficients of A are all to the left and hence its functional dependence is immaterial.

We write out the equations using the form of ad_L and de_a from (8) and match coefficients of a basis of Λ^2 . This is a linear system and we obtain the general solution of the torsion equation

$$A = \begin{pmatrix} x & q^2\mu y & -q^{-2}\mu z & -q^{-2}\delta \\ z & \frac{q^4+x(1-q^8)}{1+q^4} & 0 & q^{-6}z \\ y & 0 & \frac{1+x(q^6-q^{-2})}{1+q^4} & (q^4-1+q^{-2})y \\ -q^2x & -q^4\mu y & \mu z & \delta \end{pmatrix}; \quad \delta = \frac{1+q^6}{[4]_{q^2}} - x(q^4 - q^2 + q^{-2})$$

where x, y, z are arbitrary elements of $\mathbb{C}_q[SL_2]$. If we assume for the moment that A^j_i are numbers (i.e. commute with the 1-forms) then the cotorsion equation is entirely similar, now with ad_R . Inserting the above gives a joint solution of this form iff $y = z = 0$ and $x = q^2/[4]_{q^2}$ which then simplifies to the solution stated in the theorem. We still have to show uniqueness. Denoting our stated solution by A_0 , our above general solution of the torsion equation can be written as $A = A_0 + \tau$, where

$$\begin{aligned} \tau_a &= q^2xe_z + ze_c + ye_b, & \tau_d &= x(q^4 - q^2 + q^{-2})e_z + q^{-6}ze_c + (q^4 - 1 + q^{-2})ye_b \\ \tau_b &= -\mu ze_z - q^{-2}x(1 - q^4)e_b, & \tau_c &= q^4\mu ye_z + x(1 - q^4)e_c. \end{aligned}$$

Here we have relabelled $x - q^2/[4]_{q^2} \rightarrow x$ and x, y, z are arbitrary elements of $\mathbb{C}_q[SL_2]$. We now write out the full cotorsion equation for τ , which refers only to the linear term since A_0 is already a solution. From $\text{ad}_R(e_a)$, $\text{ad}_R(e_c)$ and $\text{ad}_R(e_b)$ we have respectively the equations

$$\begin{aligned} 0 &= -e_c \wedge (\mu ze_z + q^{-2}(1 - q^4)xe_b) - e_b \wedge (q^4\mu ye_z + x(1 - q^4)e_c) + \mu q^2e_z \wedge (q^2xe_z + ze_c + ye_b) \\ 0 &= (q^4 - 1)e_c \wedge (q^2xe_z + q^{-2}\mu ze_c + (q^2 + q^{-2})ye_b) + q^2e_z \wedge (q^4\mu ye_z + x(1 - q^4)e_c) \\ 0 &= (q^4 - 1)e_b \wedge (q^{-2}xe_z + q^{-4}(q^2 + q^{-2})ze_c - \mu ye_b) + e_z \wedge (\mu ze_z + q^{-2}(1 - q^4)xe_b). \end{aligned}$$

and a similar equation from We write $e_z f \equiv \sum_i \Delta_z^i(f)e_i$ for any element $f \in \mathbb{C}_q[SL_2]$, and similarly for Δ_b^i and Δ_c^i . Using this notation we pass all functions to the left and then collect basic two forms in a basis $\{e_{ab}, e_{ac}, e_{bd}, e_{cd}, e_{ad}, e_{bc}\}$, where $e_{ab} \equiv e_a \wedge e_b$ is a shorthand. We use the relations from Section 2.1 to order products of 1-forms. This gives us up to 18 equations for x, y, z . Among these, the coefficient of the e_{bd} in the first displayed equation above and that of e_{bc} in the second, and a similar combination from the first and third equations, gives

$$\Delta_z^d(y) = \frac{1 + 2q^4}{1 - q^2 + q^4} y, \quad \Delta_z^d(z) = \frac{2 + q^4}{1 - q^2 + q^4} z. \quad (13)$$

But from Lemma 2.1 and a similar formula for powers of a , we have $-\Delta_z^d(a^l c^k b^n d^m) = q^{m+n-k-l} a^l c^k b^n d^m$. Here $m + n - k - l$ is the natural \mathbb{Z} -grading on $\mathbb{C}_q[SL_2]$, corresponding to the diagonal \mathbb{C}^* or $U(1)$ action. Taking a homogeneous basis of $\mathbb{C}_q[SL_2]$ (such as in

Lemma 2.1 along with another part with a to the left in place of d on the right for the full non-reduced Hopf algebra), and noting that for typical q the equation $-(1+2q^4) = (1-q^2+q^4)q^N$ has no solution among integers N , we conclude that $y = 0$. Similarly $z = 0$. Also from our 18 equations we have

$$\Delta_z^b(z) = (1+q^{-2})(\Delta_z^d(x) - q^2(x)), \quad \Delta_z^c(y) = (1+q^2)(\Delta_z^d(x) - q^{-2}x) \quad (14)$$

from which we conclude $x = 0$ as well. These arguments apply at least for real positive q and for $|q| = 1$ other than an eighth-root of unity, since then the eigenvalue equations (13) cannot be solved by integer gradings (by looking at the sign or the modulus of the two sides). This covers the cases of particular interest. For all generic q with $|q| \neq 1$ we proceed as follows. (i) If $y = 0$ or $z = 0$ then (14) implies that x is an eigenfunction of the degree operator, which is not possible as $q^P + q^{\pm 2} = 1$ has no solution among integers P , hence $x = 0$. Two more of our 18 equations are

$$\Delta_z^b(x) = \Delta_z^d(y) \frac{(1+q^2)(-1+q^2-2q^4+q^6)}{q^4(1+2q^4)}, \quad \Delta_z^c(x) = -\Delta_z^d(z) \frac{(1+q^2)(-1+2q^2-q^4+q^6)}{q^4(2+q^4)} \quad (15)$$

and these now imply all $x, y, z = 0$. (ii) It remains to consider both $y, z \neq 0$, so that (13) holds with N, M the degrees of y, z . Then (15) implies $x \neq 0$ also. From Lemma 2.1 we see that Δ_z^b raises degree by 2, and Δ_z^c lowers by two. One may then argue from (14) that x is homogeneous of degree $P = M + 2 = N - 2$. Hence $N - M - 4 = 0$ and the ratio of the eigenvalue equations (13) becomes $2 + q^{-4} = 2 + q^4$, which contradicts our assumption that $|q| \neq 1$. Hence $x = y = z = 0$, completing our proof of uniqueness. The odd roots of unity case was already covered in the above analysis. It can also be verified at 3,5,7'th roots of unity by encoding the relations in Lemma 2.1 as a $4r^3 \times 4r^3$ matrix and computing directly. The failure of regularity is also verified at a small root of unity but is expected for all $q \neq 1$.
 \diamond

The covariant derivative $\Omega^1 \rightarrow \Omega^1 \otimes_{\mathcal{A}} \Omega^1$ is computed from[3]

$$\nabla e_i = - \sum_{j,k} A_j \otimes e_k \text{ad}_L(jk|i), \quad \forall i. \quad (16)$$

It obeys the usual derivation-like rule for covariant derivatives, so we need only give it on basic 1-forms. For the above canonical spin connection it comes out (in a similar manner to solving the torsion equation in Theorem 3.1) as

$$\begin{aligned} \nabla e_a &= -\nabla e_d = \frac{1}{q^2 + q^{-2}}(e_b \otimes e_c - e_c \otimes e_b) - \mu \frac{q^6}{[4]_{q^2}} e_z \otimes e_z \\ \nabla e_b &= \frac{q^4(1+q^{-2})}{[4]_{q^2}} e_z \otimes e_b - \frac{q^2}{q^2 + q^{-2}} e_b \otimes e_z \\ \nabla e_c &= -\frac{q^4(1+q^2)}{[4]_{q^2}} e_z \otimes e_c + \frac{1}{q^2 + q^{-2}} e_c \otimes e_z. \end{aligned} \quad (17)$$

Note that $\nabla\theta = 0$, i.e. this form is covariantly constant. It is easy to verify that $\nabla\wedge = d$ on 1-forms (i.e. the torsion indeed vanishes). One may likewise also verify that the cotorsion indeed vanishes, i.e. that skew-metric compatibility holds in the sense[2]

$$(\nabla \wedge \text{id} - \text{id} \wedge \nabla)\eta = 0 \tag{18}$$

for η of the form above. On the other hand one may compute that if we extend ∇ to tensor powers as a derivation (while keeping the left-most output of ∇ to the far left), then

$$\nabla\eta = O(\mu) \tag{19}$$

i.e. for generic q it tends to 0 as $q \rightarrow 1$, but is not zero for $q^2 \neq 1$. This confirms the view in [2] that naive metric compatibility $\nabla\eta = 0$ is too strong for noncommutative geometry and the correct notion is the skew version (18) as expressed in the cotorsion. We see that there is still a unique solution and it obeys $\nabla\eta = 0$ in the classical limit. Indeed, $A_z \equiv q^{-2}A_a - A_d, A_b, A_c$ in Theorem 3.1 tends to the classical spin connection on SU_2 when suitable linear combinations of A_b, A_c are taken. As explained in [2], on the compact real form of a Lie group G associated to a simple Lie algebra \mathfrak{g} it is enough to take the framing group as G itself and $\Lambda^1 = \mathfrak{g}^*$, with a canonical spin connection given by $\frac{1}{2}e$, where e is the Maurer-Cartan form; the invariant Killing form means that $\text{Ad}(G) \subseteq SO(\dim(\mathfrak{g}))$ and ∇ from the connection on this reduced frame bundle is the usual Levi-Civita metric-compatible one. The unique solution in Theorem 3.1 is exactly a q -deformation of this classical construction.

We also note from the theorem that if one wants regularity then one must give up either the torsion or cotorsion conditions, or both. In fact it is the difference between the torsion and cotorsion which is the physically important skew metric-compatibility (18), as explained in [3], i.e. we should at least keep the torsion and cotorsion equal. Among connections with constant components there is a 6-dimensional space of these, of which $A_b \propto e_c$ (other components zero), and a similar one with $A_c \propto e_b$, are the only regular ones. The moduli of all connections with torsion=cotorsion is much bigger (e.g. for $r = 3$ it is 74-dimensional), among which one may expect some interesting but not constant regular ones.

Finally, for any connection, the Riemannian curvature is computed from[3]

$$\text{Riemann} = ((\text{id} \wedge \nabla) - d \otimes \text{id}) \circ \nabla \tag{20}$$

or equally well from the curvature $F = dA + A * A : \ker \epsilon \rightarrow \Omega^2$ of A . When the connection is not regular, the latter Yang-Mills curvature does not descend to a map $\Lambda^1 \rightarrow \Omega^2$ (it is not ‘Lie algebra valued but lives in the enveloping algebra of the braided-Lie algebra). However, this does not directly affect the Riemannian geometry (it merely complicates the geometry ‘upstairs’ on the quantum frame bundle); in the proof of [3, Corol. 3.8] one should simply omit the $\tilde{\pi}$ in the argument of F for the relation to the Riemann curvature.

Proposition 3.2 *The Riemann curvature $\Omega^1 \rightarrow \Omega^2 \otimes_{\mathcal{A}} \Omega^1$ of the canonical spin connection in Theorem 3.1 is*

$$\begin{aligned} \text{Riemann}(e_a) &= -\text{Riemann}(e_d) = \frac{q^6}{(1+q^4)^2} (e_c \wedge e_z \otimes e_b + q^{-2} e_b \wedge e_z \otimes e_c - \mu e_c \wedge e_b \otimes e_z) \\ \text{Riemann}(e_b) &= \frac{q^6}{(1+q^4)^2} (-e_b \wedge e_z \otimes e_z + q^{-2} (1+q^{-2}) e_c \wedge e_b \otimes e_b) \\ \text{Riemann}(e_c) &= \frac{q^6}{(1+q^4)^2} (-e_c \wedge e_z \otimes e_z - (1+q^{-2}) e_c \wedge e_b \otimes e_c) \end{aligned}$$

Proof Direct computation using the relations in the preliminaries and the formula (17) for ∇ . Note that Riemann is a tensor, so that $\text{Riemann}(f e_a) = f \text{Riemann}(e_a)$ for all $f \in \mathcal{A}$, i.e. we need only give it on the basic 1-forms. \diamond

4 Lifting and Ricci curvature

We now want to compute the Ricci tensor of our canonical spin connection. As explained in [3], we need for this a suitable lift $i : \Omega^2 \rightarrow \Omega^1 \otimes_{\mathcal{A}} \Omega^1$. We will look for this as induced by a map $i : \Lambda^2 \rightarrow \Lambda^1 \otimes \Lambda^1$. For quantum groups we explain a natural choice for the map i and we compute it.

First of all, note that the metric η defines a Hodge $*$ operator as in [1]. The formulae can be obtained in our conventions by a conversion process as in Section 2. Using $*$: $\Lambda^2 \rightarrow \Lambda^2$ we then decompose the invariant 2-forms into self-dual and antiself-dual pieces $\Lambda^2 = \Lambda_+^2 \oplus \Lambda_-^2$, where

$$\Lambda_+^2 = \{e_{ac}, e_{bd} - \mu e_{ab}, e_{ad} - q^{-2} e_{cb}\}, \quad \Lambda_-^2 = \{e_{ab}, e_{cd}, e_{ad} + e_{cb}\} \quad (21)$$

and where we use the shorthand $e_{ac} = e_a \wedge e_c$, etc. as in the proof of Theorem 3.1. The Λ_{\pm}^2 each form 3-dimensional comodules under Δ_R . On the other hand the eigenvalues of Ψ in our case are

$$\text{Eigenvalues}(\Psi) = \{1, -q^{-2}, -q^2\}$$

The 1-eigenvalue subspace is the kernel of $\wedge : \Lambda^1 \otimes \Lambda^1 \rightarrow \Lambda^2$. The $-q^{\pm 2}$ -eigenvalue subspaces E_{\pm} are again 3-dimensional and are comodules under Δ_R since Ψ is covariant.

Proposition 4.1 *The map i defined by the eigenspace decomposition of $\Lambda^1 \otimes \Lambda^1$ under Ψ is*

such that $i(\Lambda_{\pm}^2) = E_{\pm}$, with

$$\begin{aligned}
i(e_{ac}) &= [2]_{q^2}^{-1}(q^2 e_a \otimes e_c - e_c \otimes e_a) \\
i(e_{bd} - \mu e_{ab}) &= [2]_{q^2}^{-1} q^2 (e_b \otimes e_d - e_d \otimes e_b - \mu e_a \otimes e_b) \\
i(e_{ad} - q^{-2} e_{cb}) &= [2]_{q^2}^{-1} (q^2 e_a \otimes e_d - e_d \otimes e_a + e_b \otimes e_c - e_c \otimes e_b - \mu e_a \otimes e_a) \\
i(e_{ab}) &= [2]_{q^2}^{-1} (e_a \otimes e_b - q^2 e_b \otimes e_a) \\
i(e_{cd}) &= [2]_{q^2}^{-1} (e_c \otimes e_d - e_d \otimes e_c + \mu e_c \otimes e_a) \\
i(e_{ad} + e_{cb}) &= [2]_{q^2}^{-1} (e_a \otimes e_d - q^2 e_d \otimes e_a + e_c \otimes e_b - e_b \otimes e_c + \mu e_a \otimes e_a)
\end{aligned}$$

Proof The procedure to construct i works for any quantum group bicovariant calculus provided only that we can fully diagonalise $\Psi : \Lambda^1 \otimes \Lambda^1 \rightarrow \Lambda^1 \otimes \Lambda^1$. In this case we let M be the space spanned by all eigenvectors with eigenvalues other than 1. Then the restriction $\wedge|_M : M \rightarrow \Lambda^2$ is a linear isomorphism and we define $i = (\wedge|_M)^{-1}$. We then compute in our example and find that it maps (anti)-self dual 2-forms to the two eigenspaces as stated, thereby giving these $-q^{\pm 2}$ -eigenspaces a geometrical interpretation in terms of the Hodge $*$ operator. \diamond

Let us note that for low odd roots of unity we find that the moduli space of all possible equivariant splittings $i : \Lambda^2 \rightarrow \Lambda^1 \otimes \Lambda^1$ of the wedge product is 2-dimensional. The above construction gives one point in this moduli space but it works also for generic q and is nonsingular as $q \rightarrow 1$.

Next applying i to the first factor of Riemann and for our form of Riemann tensor where there is no functional dependence, we have $i(\text{Riemann}(e_a)) \in \Lambda^1 \otimes \Lambda^1 \otimes \Lambda^1$. We then define the Ricci tensor by a ‘quantum trace’

$$\text{Ricci} = \sum_{i,j,k} \langle f^k, i(\text{Riemann}(e_j)) \eta^{ij} \eta_{ik} \rangle \in \Lambda^1 \otimes \Lambda^1 \quad (22)$$

where $\{f^i\}$ is a dual basis to $\{e_i\}$ and is evaluated against the left most copy. Here $\eta = \eta^{ij} e_i \otimes e_j$ (summation understood) defines the metric components for the metric in Section 3 and η_{ij} is the inverse matrix. Because the matrix is not symmetric the combination $u^j_k \equiv \eta^{ij} \eta_{ik}$ is not the identity matrix as it would be in a usual trace. Instead we use a kind of ‘quantum trace’ in defining the Ricci. This is a small change from the formula in [3] (where we used the usual trace) and has the merit that all constructions remain covariant under the background quantum group (not only under a classical change of basis as discussed there). The physical meaning of this improved Ricci tensor is that we use the metric to raise the input index of Riemann and then contract as usual with another (but transposed) copy of the metric.

Proposition 4.2 *The Ricci tensor for the canonical spin connection in Theorem 3.1 is*

$$\text{Ricci} = -\frac{2q^2}{[4]_{q^2}}(\eta + \frac{q^4}{(1+q^2)}\theta \otimes \theta)$$

i.e. an ‘Einstein space’ up to a shift by $\theta \otimes \theta$.

Proof We first compute

$$\begin{aligned} i(e_b \wedge e_z) &= [2]_{q^2}^{-1} q^2 ((\mu - q^{-2})e_a \otimes e_b + e_b \otimes e_a + e_d \otimes e_b - e_b \otimes e_d) \\ i(e_c \wedge e_z) &= [2]_{q^2}^{-1} ((q^{-2} - \mu)e_c \otimes e_a - e_a \otimes e_c + e_d \otimes e_c - e_c \otimes e_d) \\ i(e_c \wedge e_b) &= [2]_{q^2}^{-2} q^2 2(e_c \otimes e_b - e_b \otimes e_c - \frac{q^2 \mu}{2}(e_a \otimes e_d + e_d \otimes e_a) + \mu e_a \otimes e_a). \end{aligned}$$

We also compute the quantum trace matrix as

$$u = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & q^2 & 0 & 0 \\ 0 & 0 & q^{-2} & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}$$

in the basis $\{e_a, e_c, e_b, e_d\}$. We then contract against the Riemann tensor from Section 3. \diamond

Thus we find the same phenomenon as for the S_3 finite symmetric group example in [3], i.e. the natural ‘Killing metric’ on $\mathbb{C}_q[SL_2]$ has covariantly constant Ricci tensor essentially proportional to the metric up to a shift by $\theta \otimes \theta$. It is interesting that, as for S_3 , this shift is exactly such as to make the shifted metric degenerate. But for all nearby metric we could take the shifted metric as a new metric without changing most of the geometry. Also as in S_3 our contraction conventions (dictated by the noncommutative geometry) are such that negative Ricci curvature corresponds to what in usual geometry would be positive Ricci curvature. In this sense the above result fits our expected picture of $\mathbb{C}_q[SU_2]$ with real q and a suitable $*$ -structure as a noncommutative sphere.

5 Dirac operator

The additional ingredient for a Dirac operator is a choice of spinor representation W of the frame quantum group and equivariant gamma-matrices $\gamma : \Lambda^1 \rightarrow \text{End}(W)$. The spinor bundle in our case is just the tensor product $\mathcal{A} \otimes W$, which is the space of spinors. We take the 2-dimensional representation (i.e. a Weyl spinor) so a spinor has components $\psi^\alpha \in \mathbb{C}_q[SL_2]$ for $\alpha = 1, 2$.

Since Λ^1 for the differential calculus of $\mathbb{C}_q[SL_2]$ was originally given in the endomorphism basis $\{e_\alpha^\beta\}$, the canonical gamma-matrices proposed in [3] are just the identity map in that basis. Or in terms of our above $\{e_i\}$ they provide the conversion according to

$$\gamma(e_i)^\alpha_\beta e_\alpha^\beta = e_i, \quad \gamma(e_a) = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}, \quad \gamma(e_b) = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, \quad \text{etc.} \quad (23)$$

If we take more usual linear combinations e_x, e_y, e_z, θ (where e_x, e_y are linear combinations of e_b, e_c) then the gamma-matrices would have a more usual form of Pauli matrices and the identity, but this is not particularly natural when $q \neq 1$ given that our metric is not symmetric.

Using the $\{e_\alpha^\beta\}$ basis, the canonical Dirac operator in [3] is

$$(\mathcal{D} \psi)^\alpha = \partial^\alpha_\beta \psi^\beta - A(\tilde{\pi} S^{-1} t^\gamma_\beta)^\alpha_\gamma \psi^\beta \quad (24)$$

where

$$A = A^\alpha_\beta e_\alpha^\beta, \quad df = \partial^\alpha_\beta(f) e_\alpha^\beta, \quad \forall A \in \Omega^1, \quad f \in \mathbb{C}_q[SL_2].$$

Proposition 5.1 *The Dirac operator for the canonical spin connection on $\mathbb{C}_q[SL_2]$ in Theorem 3.1 is*

$$\mathcal{D} = \begin{pmatrix} \partial^a + \frac{[3]}{[4]} & \partial^b \\ \partial^c & \partial^d + \frac{[3]}{[4]} \end{pmatrix} = \mathcal{D} + \frac{[3]}{[4]}$$

where $\mathcal{D} = \partial^i \gamma(e_i)$ and $[n] = (q^n - q^{-n})/(q - q^{-1})$.

Proof From (5) for $\tilde{\pi}$, and the specific form of our spin connection from Theorem 3.1, we have

$$\begin{aligned} A(\tilde{\pi} S^{-1} a) &= A(\tilde{\pi} d) = -q^{-1} \alpha e_z, & A(\tilde{\pi} S^{-1} b) &= -q^{-1} A(\tilde{\pi} b) = -q^{-1} \beta e_c \\ A(\tilde{\pi} S^{-1} c) &= -q A(\tilde{\pi} c) = -q \beta e_b, & A(\tilde{\pi} S^{-1} d) &= A(\tilde{\pi} a) = q \alpha e_z. \end{aligned}$$

where $\alpha = q^4/[4]_{q^2}$ and $\beta = 1/(q^2 + q^{-2})$ are the coefficients in Theorem 3.1. We then convert to the spinor basis with $(e_i)^\alpha_\beta = \gamma(e_i)^\alpha_\beta$ so, e.g. $(e_b)^1_2 = 1$ and its other components are zero. We then compute the matrix $\mathcal{A}^\alpha_\beta = A(\tilde{\pi} S^{-1} t^\gamma_\beta)^\alpha_\gamma$ so that $\mathcal{D} = \mathcal{D} - \mathcal{A}$. We have

$$A(\tilde{\pi} S^{-1} t^1_1)^1_1 + A(\tilde{\pi} S^{-1} t^2_1)^1_2 = -\alpha q^{-3} - q\beta, \quad A(\tilde{\pi} S^{-1} t^1_2)^2_1 + A(\tilde{\pi} S^{-1} t^2_2)^2_2 = -q^{-1} \beta - q\alpha$$

and zero for the off-diagonal entries of \mathcal{A} . Remarkably, these two diagonal entries coincide and equal $-q[3]_{q^2}/[4]_{q^2} = -[3]/[4]$ on inserting the values of α, β . \diamond

We see that the Dirac operator is the naive \mathcal{D} that one might write guess without a spin connection, plus an offset of $[3]/[4]$ reflecting the constant background curvature. This is much the same phenomenon as for S_3 in [3] and in keeping with the broad geometrical picture.

Proposition 5.2 *At least for $r = 3, 5, 7$ the above Dirac operator at an r 'th root of unity has r eigenvalues of \mathcal{D} lying equally spaced on a circle in the complex plane. Here*

$$\text{Eigenvalues}(\mathcal{D}) = \left\{ \frac{q^2[m]_q}{[2]_q}, m = 0, 1, \dots, r-1 \right\}$$

The offset value in \mathcal{D} due to the constant curvature background is

$$\frac{\sin(\frac{2\pi 3}{r})}{\sin(\frac{2\pi 4}{r})} = 0, \sqrt{\frac{\sqrt{5}-1}{\sqrt{5}+1}}, -1,$$

for $r = 3, 5, 7$. For $r = 9$ it jumps to approximately 2.53... and thereafter decays asymptotically towards $\frac{3}{4}$ as $r \rightarrow \infty$.

Proof It is easiest (and most natural mathematically) to compute the unnormalised \mathcal{D} where $d = [\theta,]$ without the factor μ^{-1} that was inserted for the classical limit. The eigenvalues in this case are computed using the explicit formulae for d in Section 2 and the corresponding ∂^a without this factor, and one obtains $\{q^m - 1 \mid m = 0, 1, \dots, r-1\}$ i.e. all the conjugate roots of unity, shifted by -1. The normalised \mathcal{D} is then as stated. Note that we have one eigenvalue for each irreducible representation of the reduced quantum group $u_q(sl_2)$. \diamond

We expect the above results for all odd roots of unity and see that the limit as $r \rightarrow \infty$ fits with the real $q \rightarrow 1$ limit. The two real eigenvalues of \mathcal{D} are 0,-1 (for $m = 0, r-2$) and the offset due to the curvature for large r ensures that these appear in \mathcal{D} on either side of zero. For small r it is interesting that the offset vanishes for $r = 3$, while for $r = 5$ it is

$$4 \cos^2\left(\frac{\pi}{5}\right) - 2$$

which is reminiscent of the discrete series $\{4 \cos^2(\frac{\pi}{r})\}$ for the values of the Jones index of a type II_1 subfactor[10]. Likewise the distance between the $m = 0$ and $m = 1$ eigenvalues is $1/(2 \cos(\frac{\pi}{r}))$, while the radius of the circle of eigenvalues is a similar series $|\mu|^{-1} = 1/(2 \sin(\frac{2\pi}{r}))$. Note, however, that these observations, as well as the reality of the offset due to the curvature, depend on the normalisation μ^{-1} in our initial definition of d , which is not the only possible one. We note also that Proposition 5.2 relating the eigenvalues of the Dirac operator to the q -integer dimensions of irreducible representations should be expected to apply also for the full $\mathbb{C}_q[SL_2]$ at generic q , but with m no longer truncated at $r-1$. For generic q a different approach to Dirac operators appeared shortly after [3] in [11], with different results than ours.

Finally, at roots of unity, the Dirac operator is *not* fully diagonalisable, just as the Laplace operator was not in [1]. We demonstrate this for $r = 3$:

Proposition 5.3 For $r = 3$, the 14 zero-modes of $\not{\partial} = \not{D}$ are

$$\begin{aligned} & \begin{pmatrix} 0 \\ 1 \end{pmatrix}, \begin{pmatrix} 0 \\ b^2 d \end{pmatrix}, \begin{pmatrix} 0 \\ b d^2 \end{pmatrix}, \begin{pmatrix} -b \\ q a \end{pmatrix}, \begin{pmatrix} -b^2 d \\ b \end{pmatrix}, \begin{pmatrix} b d^2 \\ d \end{pmatrix}, \begin{pmatrix} c d^2 \\ q c^2 d \end{pmatrix} \\ & \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \begin{pmatrix} c^2 a \\ 0 \end{pmatrix}, \begin{pmatrix} c a^2 \\ 0 \end{pmatrix}, \begin{pmatrix} q^2 d \\ -c \end{pmatrix}, \begin{pmatrix} c \\ -c^2 a \end{pmatrix}, \begin{pmatrix} a \\ c a^2 \end{pmatrix}, \begin{pmatrix} q^2 b^2 a \\ b a^2 \end{pmatrix} \end{aligned}$$

The 12 massive modes with eigenvalue -1 are:

$$\begin{aligned} & \begin{pmatrix} 0 \\ b \end{pmatrix}, \begin{pmatrix} 0 \\ d \end{pmatrix}, \begin{pmatrix} 0 \\ b^2 a \end{pmatrix}, \begin{pmatrix} b \\ q^2 a \end{pmatrix}, \begin{pmatrix} 1 \\ -c d^2 \end{pmatrix}, \begin{pmatrix} c d^2 \\ -q^2 c^2 d \end{pmatrix} \\ & \begin{pmatrix} c \\ 0 \end{pmatrix}, \begin{pmatrix} a \\ 0 \end{pmatrix}, \begin{pmatrix} c^2 d \\ 0 \end{pmatrix}, \begin{pmatrix} q d \\ c \end{pmatrix}, \begin{pmatrix} -b a^2 \\ 1 \end{pmatrix}, \begin{pmatrix} -q b^2 a \\ b a^2 \end{pmatrix} \end{aligned}$$

The 12 massive modes with eigenvalue q^2 are:

$$\begin{aligned} & \begin{pmatrix} 0 \\ b^2 \end{pmatrix}, \begin{pmatrix} 0 \\ d^2 \end{pmatrix}, \begin{pmatrix} 0 \\ b d \end{pmatrix}, \begin{pmatrix} -b^2 \\ q b a \end{pmatrix}, \begin{pmatrix} d^2 \\ -q^2 c d \end{pmatrix}, \begin{pmatrix} -q^2 b d \\ c b - 1 \end{pmatrix} \\ & \begin{pmatrix} c^2 \\ 0 \end{pmatrix}, \begin{pmatrix} a^2 \\ 0 \end{pmatrix}, \begin{pmatrix} c a \\ 0 \end{pmatrix}, \begin{pmatrix} q^2 c d \\ -c^2 \end{pmatrix}, \begin{pmatrix} -q b a \\ a^2 \end{pmatrix}, \begin{pmatrix} c b - 1 \\ -q c a \end{pmatrix} \end{aligned}$$

Proof We first compute the dimensions of the eigenspaces of $\not{\partial}$ as a 54×54 matrix. Due to its size, we then treat it as a numerical matrix to find approximate eigenvectors, which we fit to powers of q to obtain the required linearly independent number of exact eigenvectors verified directly. We are left with 16 non diagonalisable modes. We work in the $\{c^k b^n d^m\}$ basis but simplify final results using the identities $a = (1 + q^2 c b) d^2$ and $a^2 = (1 - q c b + q^2 c^2 b^2) d$. These imply in particular that $c^2 a^2 = c^2 d$, $c^2 d^2 = c^2 a$, $b^2 a^2 = b^2 d$, $b^2 d^2 = b^2 a$. \diamond

We see that the components of the modes of $\not{\partial}$ with real eigenvalues $0, -1$ are given by entries which are the 13 massless spin zero modes $1, a, b, c, d, c^2 d, c d^2, c^2 a, c a^2, b^2 d, b d^2, b^2 a, b a^2$ found in [1]. This is a remnant of the idea that the Dirac operator is the square root of the Laplace one. On the other hand the components of the modes of $\not{\partial}$ of complex mass q^2 have entries from the massive spin zero modes $a^2, b^2, c^2, d^2, c d, b d, c a, b a, c b - 1$ in [1]. Note that at a mathematical level without the μ^{-1} normalisation in $\not{\partial}$, the massive modes would have eigenvalues $q - 1, q^2 - 1$ and we see that these each have an equal number of modes. Their physical make up, however, seems to be quite different as just explained. Finally, we have written out the modes in such a way as to manifest a different symmetry, namely for each mass, they fall into pairs given by swapping the entries of the matrix and replacing $a \leftrightarrow d$, $b \leftrightarrow c$, $q \leftrightarrow q^{-1}$. This appears to be a reasonable charge conjugation operation.

References

- [1] X. Gomez and S. Majid. Noncommutative cohomology and electromagnetism on $\mathbb{C}_q[SL_2]$ at roots of unity. *Lett. Math. Phys.*, in press 2002.
- [2] S. Majid. Quantum and braided group Riemannian geometry. *J. Geom. Phys.*, 30:113–146, 1999.
- [3] S. Majid. Riemannian geometry of quantum groups and finite groups with nonuniversal differentials, *Commun. Math. Phys.*, 225:131–170, 2002.
- [4] I. Heckenberger and K. Schmudgen. Levi-Civita connections on the quantum groups $SL_q(N)$, $O_q(N)$ and $Sp(q)(N)$. *Commun. Math. Phys.*, 185:177–196, 1997.
- [5] S. Major and L. Smolin. Quantum deformation of quantum gravity. *Nucl. Phys. B*, 473:267–290, 1996.
- [6] R. Coquereaux, A.O. Garcia and R. Trinchero. Differential calculus and connections on a quantum plane at a cubic root of unity. *Rev. Math. Phys.* 12:227–285, 2000.
- [7] A. Connes. *Noncommutative Geometry*. Academic Press, 1994.
- [8] S.L. Woronowicz. Differential calculus on compact matrix pseudogroups (quantum groups). *Commun. Math. Phys.*, 122:125–170, 1989.
- [9] S. Majid. *Foundations of Quantum Group Theory*. Cambridge University Press, 1995.
- [10] V.F.R. Jones and V.S. Sunder *Introduction to Subfactors*. Cambridge University Press, 1997.
- [11] I. Heckenberger. Spin geometry on quantum groups via covariant differential calculi. Preprint, math.QA/0006226.