

A NOTE ON JETS OF ENTIRE CURVES IN SEMI-ABELIAN VARIETIES

JUNJIRO NOGUCHI AND JÖRG WINKELMANN

ABSTRACT. We prove a product decomposition of the Zariski closure of the jet lifts of an entire curve $f : \mathbf{C} \rightarrow A$ into a semi-abelian variety A , provided that f is of finite order. On the other hand, by giving an example of f into a three dimensional abelian variety we show that this product decomposition does not hold in general; there was a gap in the proofs of [2], Proposition 1.8 (ii) and of [6], Theorem 2.2.

1. INTRODUCTION

Let A be a semi-abelian variety (i.e. an extension of an abelian variety by a product of copies of the multiplicative group), $f : \mathbf{C} \rightarrow A$ a holomorphic map and $J_k(f) : \mathbf{C} \rightarrow J_k(A)$ its k -th jet lift. Since A has a trivial tangent bundle, all the jet bundles are trivial and $J_k(A)$ decomposes as a direct product $A \times \mathbf{C}^{nk}$ with $n = \dim A$ in a canonical way.

In this note we use Zariski topology in terms of algebraic subsets. A map from a complex space into a complex algebraic variety is said to be Zariski (resp. non)degenerate if its image is (resp. not) Zariski dense.

Now let X_k denote the Zariski closure of the image $J_k(f)(\mathbf{C})$ in $J_k(A)$. Then it was claimed in [2], Proposition 1.8 (ii) and in [6], Theorem 2.2 for the case of an abelian variety A that X_k is a direct product of a translate of a semi-abelian subvariety of A with a subvariety of \mathbf{C}^{nk} . Unfortunately, there was a gap in these proofs, as pointed out by P. Vojta in a private communication.

In this note we prove this assertion for the case where f is of finite order and also for the case where A is a simple abelian variety (see Theorem 2.5). On the other hand, by giving an example we show that the assertion is not true in general (see Proposition 3.1).

2. THE FINITE ORDER CASE

Basically, we follow the arguments of [2]. The key idea is to consider jets of jets. For the better understanding the argument is split in a sequence of lemmas.

Lemma 2.1. *Let X be an irreducible complex algebraic variety and $p \in X \setminus \text{Sing}(X)$ a non-singular point. Then there exists a Zariski nondegenerate holomorphic map f from the unit disk Δ to X with $f(0) = p$.*

Research supported in part by Grant-in-Aid for Scientific Research (A)(1), 13304009.

Proof. Let $m = \dim X$. Let $\alpha_1, \dots, \alpha_m$ be \mathbf{Q} -linearly independent positive real numbers. Let $\Delta_2 = \{z : |z| < 2\}$ and fix an open embedding $\iota : \Delta_2^m \hookrightarrow X$ such that $p \in \iota(\Delta_2^m)$. Recall that there exists an isomorphism $\zeta : \Delta \cong H^+ = \{z : \Im(z) > 0\}$. We define a map $\phi : H^+ \cup \mathbf{R} \rightarrow \Delta_2^m \subset X$ by

$$z \mapsto (e^{\alpha_1 iz}, \dots, e^{\alpha_m iz})$$

Because the real numbers α_i are \mathbf{Q} -linearly independent, $\phi(\mathbf{R})$ is dense in the real torus $(S^1)^m$. Hence $(S^1)^m$ is contained in the closure of $\phi(H^+)$. Next we choose a holomorphic automorphism ψ of Δ_2^m with $\psi(\phi(\zeta(0))) = p$ and define $f = \psi \circ \phi \circ \zeta$.

Because $(S^1)^m$ and therefore also $\psi((S^1)^m)$ is totally real and of real dimension m , it follows that no closed analytic subset of X except X itself can contain $\iota(\psi(\phi(H^+)))$. Thus $f(\Delta)$ must be Zariski dense. \square

Remark 2.2. Regarded as a map from \mathbf{C} to \mathbf{C}^n the map ϕ defined in the above proof is an example for a holomorphic map from \mathbf{C} to \mathbf{C}^n for which the image $\phi(\mathbf{C})$ is not contained in any proper analytic subset of \mathbf{C}^n .

Let A be a semi-abelian variety and \mathbf{C}^m the complex affine space. Let $X \subset A \times \mathbf{C}^m$ be an irreducible algebraic subvariety. We define the stabilizer group by

$$\text{Stab}_A(X) = \{a \in A : (x + a, y) \in X \text{ for } \forall (x, y) \in X \subset A \times \mathbf{C}^m\}^0,$$

where $\{\cdot\}^0$ stands for the identity component. Then $\text{Stab}_A(X)$ is a connected closed algebraic subgroup of A .

For $l \in \mathbf{N}$, let $J_l(X)$ be the l -th jet space of X and let ρ_l denote the natural projection

$$\rho_l : J_l(A \times \mathbf{C}^m) \cong A \times \mathbf{C}^{nl+m(l+1)} \rightarrow J_l(A \times \mathbf{C}^m)/A \cong \mathbf{C}^{nl+m(l+1)}.$$

Lemma 2.3. *Let the notation be as above. Assume that $\text{Stab}_A(X) = \{0\}$. Then for every sufficiently large l the differential of $\rho_l|_{J_l(X)}$ restricted to the jet space $J_l(X)$ has maximal rank at general points of $J_l(X \setminus \text{Sing}(X))$.*

Proof. The proof is similar to that of Lemma (1.2) in [2] (cf. [3], Proof of Lemma (6.3.10), too).

For an arbitrarily fixed point $y_0 \in X \setminus \text{Sing}(X)$ there is a Zariski nondegenerate holomorphic map $f : \Delta \rightarrow X$ with $f(0) = y_0$ by Lemma 2.1. Let $J_l(f) : \Delta \rightarrow J_l(X)$ be the l -th jet lift, and set

$$y_l = J_l(f)(0) \in J_l(X), \quad l = 0, 1, 2, \dots$$

We are going to show that there is a number l_0 satisfying the condition: For every $l \geq l_0$, the differential at y_l

$$d(\rho_l|_{J_l(X)})_{y_l} : \mathbf{T}_{y_l}(J_l(X)) \rightarrow \mathbf{T}_{\rho_l(y_l)}(\mathbf{C}^{nl+m(l+1)}).$$

is injective.

Let $x_0 \in A$ be the image point of y_l by the natural projection, $J_l(X) \rightarrow A$. First we note that

$$\mathbf{T}_{y_l}(J_l(X)) \subset \mathbf{T}_{y_l}(A \times \mathbf{C}^{nl+m(l+1)}) \cong \mathbf{T}_{x_0}(A) \oplus \mathbf{C}^{nl+m(l+1)}.$$

Because of the definition of $\rho_l|_{J_l(X)}$, we have the kernel,

$$\ker d(\rho_l|_{J_l(X)})_{y_l} \subset \mathbf{T}_{x_0}(A).$$

Since

$$\ker d(\rho_l|_{J_l(X)})_{y_l} \supset \ker d(\rho_{l+1}|_{J_{l+1}(X)})_{y_{l+1}},$$

there is a number l_0 such that

$$\ker d(\rho_l|_{J_l(X)})_{y_l} = \ker d(\rho_{l_0}|_{J_{l_0}(X)})_{y_{l_0}} \text{ for all } l \geq l_0.$$

Suppose $\ker d(\rho_{l_0}|_{J_{l_0}(X)})_{y_{l_0}} \neq \{0\}$. Take a vector $v \in \ker d(\rho_{l_0}|_{J_{l_0}(X)})_{y_{l_0}} \setminus \{0\}$. Let $h \in \mathcal{I}_{y_0}(X)$ be a holomorphic function germ in the ideal sheaf of X at y_0 . Then v , considered as a vector field on A , satisfies

$$\left. \frac{d^l}{dz^l} \right|_{z=0} vh(f(z)) = 0, \quad l = 0, 1, 2, \dots$$

Therefore, $vh(f(z)) \equiv 0$ in a neighborhood of 0 (cf. [3], (6.3.12)). Since h was chosen arbitrarily, we obtain $vh(f(z)) \equiv 0$ near 0 for all $h \in \mathcal{I}_{y_0}(X)$. Hence v is tangent to X at every point of $f(\Delta)$. Since $f(\Delta)$ is Zariski dense in X , v is everywhere tangent to X , so that X is invariant by the action of the one parameter subgroup generated by v . This contradicts the assumption. \square

By Lemma 2.3 we immediately have the following.

Lemma 2.4. *Let A be a semi-abelian variety, and $g : \mathbf{C} \rightarrow A \times \mathbf{C}^m$ an entire curve. Let X be the Zariski closure of $g(\mathbf{C})$. Assume that $\text{Stab}_A(X) = \{0\}$. Let l be as in Lemma 2.3. Then for every rational function ϕ on X the induced meromorphic function $\phi \circ g$ on \mathbf{C} is algebraic over the field extension of \mathbf{C} generated by all the components of $\rho_l \circ J_l(g)$.*

Theorem 2.5. *Let A be a semi-abelian variety of dimension n , and $f : \mathbf{C} \rightarrow A$ an entire curve. Let X_k be the Zariski closure of $J_k(f)(\mathbf{C})$ in $J_k(A) \cong A \times \mathbf{C}^{nk}$.*

- (i) *Assume that f is of finite order. Then there exist a semi-abelian subvariety $B \subset A$, $a \in A$, and a subvariety $W_k \subset \mathbf{C}^{nk}$ such that $X_k = (B + a) \times W_k$.*
- (ii) *If A is a simple abelian variety, then there is a subvariety $W_k \subset \mathbf{C}^{nk}$ such that $X_k = A \times W_k$.*

Proof. Let $B = \text{Stab}_A(X_k)$ and apply the preceding Lemma 2.4 to the holomorphic map $F : \mathbf{C} \rightarrow (A/B) \times \mathbf{C}^{nk} \cong J_k(A)/B$ induced by $J_k(f) : \mathbf{C} \rightarrow J_k(A) \cong A \times \mathbf{C}^{nk}$. Let $F_1 : \mathbf{C} \rightarrow A/B$ denote its first component. Then F_1 is the composition of $f : \mathbf{C} \rightarrow A$ with the natural projection from A to A/B .

Assume that f is of finite order. Let $\tilde{f} : \mathbf{C} \rightarrow \mathbf{C}^n$ denote the lift of f to the universal covering of A . Then the lift \tilde{f} is a polynomial map, and hence $\rho_l \circ J_l(F)$ are polynomial map, too (see, e.g., [5]). The preceding Lemma 2.4 now yields that for every rational function ϕ on the semi-abelian variety A/B the induced meromorphic function $\phi \circ F_1$ is algebraic over the rational function field $\mathbf{C}(z)$ in z . It follows that Nevanlinna's order function fulfills $T_{\phi \circ F_1}(r) = O(\log r)$ (cf., e.g., [3], Proposition (5.3.14) and Lemma (6.1.5)). This implies that $\phi \circ F_1$ is a rational function. Since this is true for every rational function ϕ on A/B , it follows that $F_1 : \mathbf{C} \rightarrow A/B$ is an algebraic morphism. But every algebraic morphism from \mathbf{C} to a semi-abelian variety is constant. Thus the projection map from $J_k(A)$ to A/B maps X_k to a point. This implies that $X_k = (B + a) \times W_k$ for some subvariety $W_k \subset J_k(A)/A$ and $a \in A$.

In general, the above argument at least implies $\dim \text{Stab}(X_k) > 0$ (cf. [2], Lemma (1.2)). If A is a simple abelian variety, then $\text{Stab}(X_k) = A$, so that the required assertion follows. \square

Remarks 2.6.

- (i) Using induction, one can deduce from $\dim \text{Stab}_A(X_0) > 0$ that for every such holomorphic map f the Zariski closure of the image $f(\mathbf{C})$ in A is a translate of a semi-abelian subvariety. This fact (“logarithmic Bloch-Ochiai theorem”) was first proved in [1].
- (ii) However, for $k > 0$ and $0 < \dim B < \dim A$ the quotient $J_k(A)/B$ is strictly larger than $J_k(A/B)$. Therefore one cannot use induction to prove that X_k has a product decomposition as in the case where f is of finite order. Basically, this is the gap in the incomplete proofs mentioned in the introduction. This gap cannot be filled: As we will see in the next section there is a counter-example in the case where f is not of finite order.

3. THE COUNTER-EXAMPLE

Proposition 3.1. *There exist a three-dimensional complex abelian variety A and a holomorphic map $f : \mathbf{C} \rightarrow A$ with the following properties:*

- (i) *The image $f(\mathbf{C})$ is Zariski dense in A ;*
- (ii) *Let X denote the Zariski closure of the image of the jet lift $J_1(f) : \mathbf{C} \rightarrow J_1(A) \cong A \times \mathbf{C}^3$. Then X is not a direct product inside $J_1(A)$, i.e., there does not exist a subvariety $W \subset \mathbf{C}^3$ such that $X = A \times W$.*

Proof. Let C be a nonsingular cubic curve in $\mathbf{P}_2(\mathbf{C})$. Then C is an elliptic curve, i.e., a one-dimensional compact complex torus. For a point $x = [x_0 : x_1 : x_2] \in \mathbf{P}_2(\mathbf{C})$ we define $x^\perp = \{[z_0 : z_1 : z_2] : \sum_i z_i x_i = 0\}$. We define a surface $\bar{S} \subset C \times \mathbf{P}_2(\mathbf{C})$ by

$$\bar{S} = \{(x, z) \in \mathbf{P}_2(\mathbf{C}) \times \mathbf{P}_2(\mathbf{C}) : z \in x^\perp \text{ and } x \in C\}$$

Note that the projection $C \times \mathbf{P}_2(\mathbf{C}) \rightarrow C$ induces the structure of a holomorphic fiber bundle on \bar{S} , with fiber $\mathbf{P}_1(\mathbf{C})$ and base C .

Fix $p \in \mathbf{P}_2(\mathbf{C}) \setminus C$. Now for every $x \in C$, the sets p^\perp and x^\perp are two different lines in $\mathbf{P}_2(\mathbf{C})$. In particular $x^\perp \cap p^\perp$ contains exactly one point. Define $S = \bar{S} \setminus (C \times p^\perp)$. Then S is a closed algebraic subvariety of $C \times \mathbf{C}^2 \cong C \times (\mathbf{P}_2(\mathbf{C}) \setminus p^\perp)$. Let $i : S \rightarrow C \times \mathbf{C}^2$ denote this embedding. On the other hand, by the projection on the first factor, S can be realized as a holomorphic fiber bundle with fiber \mathbf{C} and base C . Let $\pi : \tilde{S} \rightarrow S$ be the universal covering. Then $\tilde{S} \cong^{\varphi} \mathbf{C}^2$, because \mathbf{C} is the universal covering of C and every \mathbf{C} -bundle over \mathbf{C} is trivial.

Let $g : \mathbf{C} \rightarrow \mathbf{C}^2$ be a holomorphic map such that no proper analytic subset of \mathbf{C}^2 contains $g(\mathbf{C})$ (cf. Remark 2.2). Let $G = (G_1, G_2) : \mathbf{C} \rightarrow C \times \mathbf{C}^2$ be the holomorphic map given by the compositions

$$\mathbf{C} \xrightarrow{g} \mathbf{C}^2 \xrightarrow{\varphi} \tilde{S} \xrightarrow{\pi} S \xrightarrow{i} C \times \mathbf{C}^2.$$

Note that $i(S)$ is the Zariski closure of $G(\mathbf{C})$ in $C \times \mathbf{C}^2$.

Choose a lattice Λ in \mathbf{C}^2 such that $B = \mathbf{C}^2/\Lambda$ is a simple complex abelian variety. Denote the natural projection by $\tau : \mathbf{C}^2 \rightarrow B$. Let $A = C \times B$. Now we are in a position to define the desired entire curve $f : \mathbf{C} \rightarrow A$: Let $f = (f_1, f_2) : \mathbf{C} \rightarrow C \times B$ with $f_1 = G_1$ and

$$f_2(x) = \tau \left(\int_0^x G_2(t) dt \right).$$

We have to verify that f has the desired properties. By construction $f(\mathbf{C})$ is not contained in (a translate of) one of the two factors of the product $C \times B$. On the other hand, the Zariski closure of $f(\mathbf{C})$ must be (a translate of) a subtorus and B is simple. Therefore $f(\mathbf{C})$ is Zariski dense in A .

Next we consider the first jet lift

$$J_1(f) : \mathbf{C} \rightarrow J_1(A) \cong A \times \mathbf{C}^3 \cong C \times B \times \mathbf{C} \times \mathbf{C}^2.$$

By construction we have

$$J_1(f) = (G_1, f_2, G_1', G_2).$$

Let $q : (x_1, x_2; v_1, v_2) \mapsto (x_1, v_2)$ denote the projection on the first and fourth factor. By our construction the closure of $q(J_1(f)(\mathbf{C}))$ in $C \times \mathbf{C}^2$ coincides with $i(S)$. In particular, it is not a direct product of subvarieties of the two factors. It follows that the Zariski closure of $J_1(f)$ in $J_1(A) = A \times \mathbf{C}^3$ cannot be a direct product of a subvariety of A with a subvariety of \mathbf{C}^3 . \square

Remark 3.2. The projection map p_2 from $i(S) \subset C \times \mathbf{C}^2$ onto the second factor \mathbf{C}^2 maps $i(S)$ surjectively onto \mathbf{C}^2 . As a result, for a holomorphic map $G : \mathbf{C} \rightarrow i(S)$ with Zariski dense image in $i(S)$ the induced map $G_2 = p_2 \circ G : \mathbf{C} \rightarrow \mathbf{C}^2$ has a Zariski dense image. Therefore in

the above construction $G_2 : \mathbf{C} \rightarrow \mathbf{C}^2$ cannot be a polynomial map. As a consequence f_2 and therefore $f : \mathbf{C} \rightarrow A$ is not a holomorphic map of finite order.

Note also that $\text{Stab}_A(X) = \{0\} \times B$.

4. CONSEQUENCES

It should be noted that in the proof of the Main Theorem of [2], Proposition (1.8) (ii) was not used. Hence it is valid without any change.

Similarly, the product decomposition of X_k is not essential in any of the proofs of [5] as in [4], where the product decomposition was not used.

REFERENCES

- [1] Noguchi, J.: Lemma on logarithmic derivatives and holomorphic curves in algebraic varieties. Nagoya Math. J. **83** (1981), 213–233.
- [2] Noguchi, J., On holomorphic curves in semi-Abelian varieties, Math. Z. **228** (1998), 713-721.
- [3] Noguchi, J. and Ochiai, T., Geometric Function Theory in Several Complex Variables, Japanese edition, Iwanami, Tokyo, 1984; English Translation, Transl. Math. Mono. **80**, Amer. Math. Soc., Providence, Rhode Island, 1990.
- [4] Noguchi, J., Winkelmann, J., and Yamanoi, K., The value distribution of holomorphic curves into semi-Abelian varieties, C.R. Acad. Sci. Paris t. **331** (2000), Série I, 235-240.
- [5] Noguchi, J.; Winkelmann, J.; Yamanoi, K.: The second main theorem for holomorphic curves into semi-Abelian varieties, Acta Math. **188** no. 1 (2002), 129-161.
- [6] Siu, Y.-T. and Yeung, S.-K., A generalized Bloch's theorem and the hyperbolicity of the complement of an ample divisor in an Abelian variety. Math. Ann. **306** (1996), 743-758.

NOGUCHI, JUNJIRO, GRADUATE SCHOOL OF MATHEMATICAL SCIENCES, UNIVERSITY OF TOKYO, KOMABA, MEGURO, TOKYO 153-8914, JAPAN

E-mail address: `noguchi@ms.u-tokyo.ac.jp`

WINKELMANN, JÖRG, KOREAN INSTITUTE FOR ADVANCED STUDIES, SCHOOL OF MATHEMATICS, 207-43 CHEONGRYANGRI-DONG, DONGDAEMUN-GU, SEOUL, 130-012 KOREA

E-mail address: `jwinkel@member.ams.org`

Webpage: `http://www.math.unibas.ch/~winkel/`