

**STRICT VOLUME-MINIMIZING PROPERTIES FOR  
LAGRANGIAN SUBMANIFOLDS IN COMPLEX MANIFOLDS  
WITH POSITIVE CANONICAL BUNDLE**

EDWARD GOLDSTEIN

ABSTRACT. It is classically known that closed geodesics on a Riemann surface with a metric of negative curvature strictly minimize length in their free homotopy class. We'd like to generalize this to Lagrangian submanifolds in Kähler manifolds of negative Ricci curvature. The only known result in this direction is a theorem on Y.I. Lee (see [Lee]) for certain Lagrangian submanifolds in a product of two Riemann surfaces of the same negative curvature. We develop an approach to study this problem in higher dimensions. Along the way we prove some weak results (volume-minimization outside of a divisor) and give a counterexample to global volume-minimizing for an immersed minimal Lagrangian submanifold.

1. LOCAL THEORY

In this section we'll present the local volume-minimizing properties for minimal Lagrangian submanifolds in Kähler manifolds with positive canonical bundle. Strictly speaking the rest of the paper does not logically depend on this section and this section serves as a motivation for the global problem.

Let  $(M^{2n}, \omega)$  be a Kähler  $n$ -fold and let  $K(M)$  be the canonical bundle of  $M$ , i.e.  $K(M) = \Lambda^{(n,0)}T^*M$ - the bundle of  $(n,0)$ -forms on  $M$ . The Hermitian metric on  $K(M)$  endowed from  $M$  gives rise to a Hermitian connection on  $K(M)$ . The curvature form of  $K(M)$  is called the Ricci form of  $M$  and denoted by  $Ric$ . It is related to the Ricci tensor of  $M$  (which is a symmetric 2-tensor on  $M$ ) by the following identity:

$$(1) \quad -iRic(X, Y) = Ricci(JX, Y)$$

for any tangent vectors  $X$  and  $Y$  (here  $J$  is the complex structure on  $M$ ).

Let  $L$  be an oriented Lagrangian submanifold of  $M$ . For any point  $l \in L$  there is a unique element  $\kappa_l$  of  $K(M)$  over  $l$  which restricts to the volume form on  $L$  (in the induced metric from  $M$ ). Various  $\kappa_l$  give rise to a section

$$(2) \quad \kappa_L : L \rightarrow K(M)$$

The section  $\kappa_L$  has constant length  $\sqrt{2}^n$  over  $L$  and it defines a connection 1-form  $\xi$  on  $L$  for the connection on  $K(M)$  over  $L$  by the condition  $\xi \otimes \kappa_L = \nabla \kappa_L$ .

Let  $h$  be the mean curvature vector field of  $L$ . So  $h$  is a section of the normal bundle of  $L$  in  $M$  and we have a corresponding 1-form  $\sigma = i_h \omega$  on  $L$ . The following fact was first observed by Y.G. Oh in [Oh1]:

**Lemma 1.**  $\sigma = i\xi$

Thus if  $L$  is *minimal* (i.e.  $h = 0$ ) iff  $\kappa_L$  is parallel over  $L$ . From now on we work with  $L$  which is minimal, embedded and compact. Since  $L$  is a totally-real real-analytic submanifold of  $M$  we easily see that we can extend  $\kappa_L$  to be a holomorphic section of  $K(M)$  in some neighbourhood  $U$  of  $L$  in  $M$ . Consider the function  $f = \ln|\kappa_L|^2$  on  $U$ .  $L$  is a critical submanifold of  $f$  on  $U$ . Moreover since  $K(M)$  is a positive line bundle we see the the Hessian of  $f$  is negative-definite on the normal bundle to  $M$  on  $L$ . Thus  $f$  and hence  $|\kappa_L|$  attains a strict local maximum along  $L$ . Moreover  $\kappa_L$  restricts to the volume form on  $L$ . Thus we get a conclusion:

**Lemma 2.**  *$L$  is calibrated by  $Re\kappa_L$  in some neighbourhood  $U'$  of  $L$  in  $M$  and moreover  $L$  is **strictly** homologically volume-minimizing in  $U'$ .*

We would like to globalize this statement from a neighbourhood of  $L$ . If  $M$  is a Riemann surface then the following result is classical:

**Lemma 3.** *On a Riemann surface with a metric of negative curvature every free homotopy class of loops contains a unique geodesic and this geodesic strictly minimizes length in this class.*

The only progress attained so far in higher dimensions is the result of Y.I. Lee [Lee] in case  $M$  is a product of two Riemann surfaces.

## 2. ALGEBRAIC MINIMAL LAGRANGIAN SUBMANIFOLDS

As a first step we'll consider some "nice" minimal Lagrangian submanifolds, which we'll call Algebraic:

**Definition 1.** *Let  $L$  be a minimal Lagrangian submanifold of  $M$ . We say that  $L$  is algebraic if there is a holomorphic section  $\varphi$  of  $K(M)$  over  $M$  s.t.  $|\varphi|$  has a strict global maximum along  $L$*

Note that if  $L$  is algebraic we can rotate  $\varphi$  by a constant so that  $\varphi$  will restrict to the volume form on  $L$ . Thus we get

**Lemma 4.** *Algebraic minimal Lagrangian submanifolds strictly minimize volume in their homology class.*

Next we'll demonstrate some examples of algebraic minimal Lagrangian submanifolds. Let  $M$  be a complete intersection in  $\mathbb{C}P^n$  given by polynomials  $f_1, \dots, f_m$  with real coefficients. We want  $\Sigma \deg(f_i) - (n+1) = 2k > 0$  for an integer  $k$ . Then  $K(M) \simeq (\gamma^*)^{\otimes 2k}$  where  $\gamma^*$  is the hyperplane bundle of  $\mathbb{C}P^n$ . Let  $L = N \cap \mathbb{R}P^n$ .  $L$  is a minimal Lagrangian submanifold.

Consider the standard Hermitian metric on  $(\gamma^*)^{\otimes 2k}$ . We have a section

$$\varphi = (\Sigma z_i^2)^k$$

of  $(\gamma^*)^{\otimes 2k}$  and  $|\varphi|$  attains strict maximum along  $\mathbb{R}P^n$ . By Yau's resolution of the Calabi conjecture [Yau] we can put a Kähler metric  $\omega_Y$  on  $M$  s.t. the induced metric on  $K(M)$  will be proportional by a constant to the standard metric on  $(\gamma^*)^{\otimes 2k}$ . Thus we'll get:

**Theorem 1.**  *$L$  is an algebraic minimal Lagrangian submanifold of  $(M, \omega_Y)$ .*

## 3. A GLOBAL STRATEGY

We'll now present our strategy for exploring global volume-minimizing properties for Lagrangian submanifolds in Kähler manifolds with positive canonical bundle. We'll explain this approach in the simplest possible case- a fixed point set of an antiholomorphic involution.

So let  $M$  be a complex  $n$ -fold with an antiholomorphic involution  $\sigma$ . Let  $g$  be a Hermitian metric on  $M$  which is  $\sigma$ -invariant. Note we don't assume that  $g$  is Kähler, just Hermitian. The metric  $g$  induces a Hermitian metric  $h$  on the canonical bundle  $K(M)$  of the holomorphic  $(n, 0)$ -forms of  $M$ . Let  $\omega$  be the curvature form of this metric and we assume that  $\omega$  defines a Kähler form on  $M$  (key assumption). If  $g$  were a Kähler metric this would be equivalent to  $g$  having negative Ricci curvature.

Let  $L$  be the fixed point set of  $\sigma$ . We'd like to understand when  $L$  minimizes its  $g$ -volume in its free homotopy class.

Let  $H^0(K(M)^{\otimes m}, M)$  be the space of holomorphic sections of the  $m^{\text{th}}$  power  $K(M)^{\otimes m}$ . Clearly  $\sigma$  acts upon  $H^0(K(M)^{\otimes m}, M)$  by an antiholomorphic involution (the  $\sigma$ -action is by pulling back a section and conjugating it). Let  $H_m^L$  be the fixed point set of  $\sigma$  on  $H^0(K(M)^{\otimes m}, M)$ . Then  $H_m^L$  is a Lagrangian subspace of  $H^0(K(M)^{\otimes m}, M)$ . If we choose an orthonormal basis for  $H_m^L$  then clearly it will give a unitary basis for  $H^0(K(M)^{\otimes m}, M)$ . Moreover the Kodaira embedding  $\phi_m$  that this basis will produce will be intertwining between  $\sigma$  and the conjugation on  $\mathbb{C}P^{N_m}$ . One easily sees that  $L = \phi_m^{-1}(\mathbb{R}P^{N_m})$ .

Let  $\omega_{FS}$  be the Fubini-Study metric on  $\mathbb{C}P^{N_m}$  and let

$$\omega_m = (\phi_m)^* \omega_{FS} / m$$

be the corresponding Bergmann metric on  $M$ . Then by [Tian] and [Ruan],  $\omega_m \rightarrow \omega$  on  $M$ .

The Kodaira embedding  $\phi_m$  also gives an isomorphism between  $K(M)^{\otimes m}$  and the hyperplane bundle  $\gamma^*$  of  $\mathbb{C}P^{N_m}$ . How  $\gamma^*$  has a natural Hermitian metric and so it induces a Hermitian metric on  $K(M)^{\otimes m}$  and hence on  $K(M)$ . Let  $h_m$  be that metric on  $K(M)$ . Then the curvature form of  $h_m$  is  $\omega_m$ . We'd like to conformally scale  $g$  so that the induced metric on  $K(M)$  will equal to  $h_m$  up to a constant multiple. If the scaling is  $g_m = e^{f_m} g$  then  $f_m$  solves:

$$n i \partial \bar{\partial} f_m = \omega_m - \omega$$

Here  $n$  is the dimension of  $M$ . If we choose  $f_m$  to intergate to 0 over  $M$  then we get that  $f_m \rightarrow 0$  in the  $C^\infty$ -topology. Hence  $g_m$  converge to  $g$  in the  $C^\infty$ -topology (actually we just care about  $C^0$  convergence).

Consider now a holomorphic section  $\Sigma z_i^2$  of  $(\gamma^*)^{\otimes 2}$  over  $\mathbb{C}P^{N_m}$ . We pull it back to a holomorphic section  $\varphi_m$  of  $K(M)^{\otimes 2m}$  over  $N$ . Then  $\varphi_m$  is  $\sigma$ -invariant and its length with respect to the Hermitian metric  $h_m$  attains a maximum along  $L$ .

We now would like to use the idea of calibration to study the  $g_m$ -volume minimizing properties of  $L$ . We obviously have the following corollary:

**Corollary 1.** *Suppose we have a 1-parameter family  $L_t$  of homotopies of  $L$  with  $L_0 = L$  and moreover all  $L_t$  avoid the zero set  $D_m = \varphi_m^{-1}(0)$ . Then  $\text{vol}(L_t) \geq \text{vol}(L)$ . Here the volume is with respect to  $g_m$ .*

**Proof:** First will prove this for  $L$  orientable. For non-orientable  $L$  we look on its orientable double cover and copy the proof bellow.

Since  $\varphi_m$  is  $\sigma$ -invariant we have choose its  $2m^{th}$  root  $\xi_m$  along  $L$  such that  $\xi_m$  restricts to a positive real  $n$ -form on  $L$ . We extend  $\xi_m$  to be a  $2m^{th}$  root of  $\varphi_m$  over  $L_t$  along the homotopy. Then  $\xi_m$  has constant length  $C$  (with respect to  $h_m$ ) along  $L$  and the length of  $\xi_m$  is  $\leq C$  on  $L_t$ . We obviously have

$$\int_L \xi_m = \int_{L_t} \xi_m$$

and hence  $vol(L_t) \geq vol(L)$ . Q.E.D.

The last statement about the conservation of the integral of  $\xi_m$  can be better understood using the ramified  $2m$ -cover of  $M$  that  $\varphi_m$  defines. The setup is as follows:

Let  $K(M)$  be the total space of the canonical bundle of  $M^{2n}$  and  $\pi : K(M) \rightarrow M$  be the projection. There is a canonical  $(n, 0)$ -form  $\rho$  on  $K(M)$  defined by  $\rho(a)(v_1, \dots, v_n) = a(\pi_*(v_1), \dots, \pi_*(v_n))$ . Here  $a \in K(M)$  and  $v_1, \dots, v_n$  are tangent vectors to  $K(M)$  at  $a$ . The form  $\varphi = d\rho$  is a holomorphic volume form on  $K(M)$ . If  $z_1, \dots, z_n$  are local coordinates on  $M$  then  $dz_1 \wedge \dots \wedge dz_n$  is a local section of  $K(M)$  over  $M$  which defines a coordinate function  $y$  on  $K(M)$ . The collections of holomorphic functions  $(z_1, \dots, z_n, y)$  are coordinates on  $K(M)$  and

$$(3) \quad \rho = y dz_1 \wedge \dots \wedge dz_n, \quad \varphi = dy \wedge dz_1 \wedge \dots \wedge dz_n$$

Now we have a section  $\varphi_m$  of  $K(M)^{\otimes 2m}$  and it defines a  $2m$ -fold ramified cover  $M_m$  of  $M$  as follows:

$$M_m = (\alpha \in K(M) | \alpha^{\otimes 2m} = \varphi_m(\pi(\alpha)))$$

We can in fact slightly perturb  $\varphi_m$  to keep it holomorphic,  $\sigma$ -invariant and make it transversal to 0. Than way  $M_m$  is a complex  $n$ -fold and the projection  $\pi : M_m \rightarrow M$  is a ramified cover. Moreover the  $(n, 0)$ -form  $\rho$  on  $K(M)$  restricts to a holomorphic form on  $M_m$ .

Now the  $2m^{th}$  root  $\xi_m$  of  $\varphi_m$  along  $L$  can be viewed to a lift  $L^K$  of  $L$  to  $M_m$ . If we have a homotopy  $L_t$  of  $L$  in  $M$  avoiding the zero set  $N_m = \varphi_m^{-1}(0)$  then  $L_t$  can be lifted to a homotopy  $L_t^K$  on  $M_m$ . Obviously we have

$$\int_{L_t^K} \rho = \int_{L_t} \rho$$

Hence the volume comparison property. We have a stronger result:

**Corollary 2.** *Let  $L'$  be a submanifold of  $M_m$  representing the same homology class as  $L^K$  in  $M_m$ . Then  $vol(\pi(L')) \geq vol(L)$ . Here again the volume is  $g_m$ -volume.*

We'd like eventually to understand when  $L$  minimizes  $g$ -volume in its free homotopy class. Note that if  $L$  minimizes  $g$ -volume then it strictly minimizes  $g$ -volume. Indeed suppose that some other  $L'$  homotopic to  $L$  has the same  $g$ -volume. Pick a point  $a$  in  $L$  and not in  $L'$ . Consider a positive  $\sigma$ -invariant function  $f$  supported in a neighbourhood of  $a$  and consider a Hermitian metric  $g_f = e^f g$ . Then the  $g_f$  volume of  $L$  is greater then the  $g_f$ -volume of  $L'$ . We can also choose  $f$  small enough so that the Hermitian metric that  $g_f$  induces on  $K(M)$  still has positive curvature and this will lead us to a contradiction. In particular we'll deduce that if  $\alpha$  is an automorphism of  $M$  then  $L$  and  $\alpha(L)$  are not homotopic if they don't coincide.

Unfortunately the fact that  $L$  minimizes  $g$ -volume in its free homotopy class is not always true as we have an example of a non-orientable  $L$  which doesn't:

#### 4. A COUNTEREXAMPLE IN THE NON-ORIENTABLE CASE

The example is:  $M = (\Sigma z_i^5 = 0 \subset \mathbb{C}P^3)$  endowed with a Kähler-Einstein metric. Let  $L$  be a fixed point set of the conjugation on  $M$ . Then  $L$  is clearly diffeomorphic to  $\mathbb{R}P^2$ . Let  $S^2$  be its orientable double cover. We think of  $L$  as an immersion of  $S^2$  into  $M$ . Clearly  $L$  is a 2-torion in  $H_2(M, \mathbb{Z})$  and since  $M$  is a simply connected 4-manifold,  $H_2(M, \mathbb{Z})$  has no torsion. Hence  $L$  is trivial in  $H_2(M, \mathbb{Z})$  and by the Hurewicz theorem  $L$  is null-homotopic.

From the results of J. Lees [Lees] we in fact can also deduce that this Lagrangian immersion of  $S^2$  can be homotoped through Lagrangian immersions to a Whitney sphere contained in a Darboux chart of some point in  $M$ . We omit the details.

We hope that volume minimization in free homotopy class is true when  $L$  is orientable.

#### REFERENCES

- [Lee] Y.-I. Lee: Lagrangian minimal surfaces in Kähler-Einstein surfaces of negative scalar curvature. *Comm. Anal. Geom.* 2 (1994), no. 4, 579–592.
- [Lees] J. Lees: On the classification of Lagrange immersions. *Duke Math. J.* 43 (1976), no. 2, 217–224
- [Oh1] Y.-G. Oh: Mean curvature vector and symplectic topology of Lagrangian submanifolds in Einstein-Kähler manifolds, *Math. Z.* 216 (1994), 471-482
- [Ruan] W.-D. Ruan: Canonical coordinates and Bergmann [Bergman] metrics. *Comm. Anal. Geom.* 6 (1998), no. 3, 589–631.
- [Tian] G. Tian: On a set of polarized Kähler metrics on algebraic manifolds, *J. Differential Geom.* 32 (1990), no. 1, 99–130.
- [Yau] S. T. Yau : On the Ricci curvature of a compact Kähler manifold and the complex Monge-Ampere equation I, *Comm. Pure Appl. Math.* 31 (1978), no. 3, 339-411

Stanford University  
egold@math.stanford.edu