

Notes on Metrics, Measures, and Dimensions

Stephen Semmes
Department of Mathematics
Rice University

Preface

In this monograph various notions of measure and content for subsets of metric spaces are considered, and both metric and topological notions of dimension. Some familiarity with basic topics in analysis such as Riemann integrals, metric spaces, and continuous functions is assumed, as in [50, 88, 160], for instance, although a number of definitions and results that are needed are reviewed in the text. More sophisticated matters such as Lebesgue integrals are not required, but can be complementary. It is hoped that the present endeavor might be useful in a supplementary role for a course, in conjunction with other material at the discretion of the instructor. It is intended that the presentation be relatively brief and informal, sometimes leaving as an exercise to the reader the task of filling in some details.

Contents

Preface	ii
1 Metric spaces	1
1.1 Real numbers	1
1.2 Some basic notions and results	3
1.2.1 Open and closed subsets and normality of metric spaces	4
1.2.2 Bounded, compact, and totally bounded sets	6
1.2.3 Separable metric spaces	8
1.3 The Baire category theorem	9
1.4 Continuous and uniformly continuous mappings	10
1.4.1 Real-valued functions	12
1.4.2 Spaces of continuous mappings	14
1.5 Connectedness	17
2 Some measure functionals and dimensions	20
2.1 Basic notions	20
2.2 Coverings by open and closed subsets	23
2.3 Subadditivity properties	25
2.4 Dimensions	26
2.5 Snowflake transforms	28
3 More on metric spaces	30
3.1 Lipschitz mappings	30
3.1.1 Diameters, measure functionals, and dimensions	31
3.1.2 Some approximations and extensions	32
3.1.3 Spaces of Lipschitz mappings	34
3.2 The Hausdorff metric	35
3.2.1 The space of closed and bounded subsets of M	36

3.2.2	Subsets, functions, and embeddings	37
3.2.3	Compactness	38
3.2.4	Connectedness	39
4	Examples and special cases	40
4.1	The real line	40
4.2	\mathbf{R}^n for general n	41
	Bibliography	44

Chapter 1

Metric spaces

1.1 Real numbers

Let us begin by reviewing some aspects of the real numbers. As usual, we write \mathbf{R} for the real numbers, and \mathbf{R}^n for the n -tuples of real numbers, where n is a positive integer. We also write \mathbf{Z} for the integers, and \mathbf{Z}_+ for the positive integers.

If a, b are real numbers with $a < b$, then the intervals (a, b) , $[a, b)$, $(a, b]$, and $[a, b]$ can be defined as the sets of real numbers x which satisfy $a < x < b$, $a \leq x < b$, $a < x \leq b$, and $a \leq x \leq b$, respectively. For $[a, b]$ we also allow $a = b$, in which case $[a, b]$ consists of the one point. These intervals are all *bounded*, and their *lengths* are defined to be $b - a$. It can sometimes be convenient to consider unbounded intervals of the form $(-\infty, b)$, $(-\infty, b]$, (a, ∞) , $[a, \infty)$, and $(-\infty, \infty) = \mathbf{R}$, whose lengths are defined to be $+\infty$.

Let A be a subset of \mathbf{R} . A real number b is said to be an *upper bound* if $a \leq b$ for all $a \in A$. A real number c is said to be the *least upper bound*, or *supremum*, if it is an upper bound for A , and if $c \leq b$ for every other upper bound b of A . It is easy to see from the definition that the supremum is unique if it exists. A basic property of the real numbers is completeness for the ordering, which means that every nonempty subset A of \mathbf{R} which has an upper bound has a least upper bound. If A is a nonempty subset of \mathbf{R} which has no upper bound in the real numbers, then we shall make the convention of saying that the supremum is equal to $+\infty$. The supremum of A is denoted $\sup A$.

Similarly, if A is a subset of \mathbf{R} and d is a real number, then d is said to

be a *lower bound* of A if $d \leq a$ for all a in A . A real number h is said to be the *greatest lower bound*, or *infimum*, of A if it is a lower bound for A , and if $d \leq h$ for all other lower bounds of A . Again, it is easy to see from the definition that the infimum is unique if it exists. From the completeness property of the real numbers described in the previous paragraph, one can show that a nonempty subset of \mathbf{R} with a lower bound has an infimum. One way to do this is to show that the infimum of A is equal to the supremum of $-A = \{-a : a \in A\}$, where $-A$ is nonempty and has an upper bound because A is nonempty and has a lower bound. Another approach is to obtain the infimum of A as the supremum of the set of lower bounds of A . This second argument has the nice feature that it applies to any linearly-ordered set. Compare with [160].

If A is a nonempty subset of \mathbf{R} which does not have a lower bound, then we shall make the convention that the infimum is equal to $-\infty$. The infimum of A is denoted $\inf A$ in either case. Often in these notes we shall be concerned with infima and suprema of nonempty sets of nonnegative real numbers, so that the infimum exists as a real number, but the supremum might be $+\infty$.

The real numbers together with $-\infty$, $+\infty$ are called the *extended real numbers*, as in [160]. The usual ordering on the real numbers extends to the extended real numbers, and to some extent the arithmetic operations do too. For instance, if x is a real number, then $x + \infty$ is defined to be $+\infty$, $x - \infty$ is defined to be $-\infty$, while $\infty - \infty$ is normally left undefined. If A is a nonempty set of extended real numbers, the infimum and supremum of A can be defined for A in the same way as above, as extended real numbers.

Let I be a nonempty set, and suppose that to each element i of I is associated a nonnegative real number a_i . Consider the sum

$$(1.1) \quad \sum_{i \in I} a_i.$$

If I is finite, then this sum is defined in the usual way. If I is infinite, then we can define this sum to be the supremum of

$$(1.2) \quad \sum_{i \in F} a_i,$$

where F runs through all finite subsets of I . Thus the sum may be equal to $+\infty$. Sometimes the terms a_i may be allowed to be nonnegative extended

real numbers, and the whole sum is automatically equal to $+\infty$ if any one of the terms is equal to $+\infty$.

If the sum (1.1), then for each $\epsilon > 0$ the number of $i \in I$ such that $a_i \geq \epsilon$ is at most $1/\epsilon$ times the sum (1.1). In particular, the number of such $i \in I$ is finite. As a consequence, if the sum (1.1) is finite, then the set of $i \in I$ such that $a_i > 0$ is at most countable. In these notes, we shall only be concerned with situations in which I is at most countable anyway.

The *absolute value* $|x|$ of a real number x is defined to be equal to x when $x \geq 0$ and to $-x$ when $x \leq 0$. Thus the absolute value of x is always a nonnegative real number, and one can check that $|x + y| \leq |x| + |y|$ for all real numbers x, y . Also, the absolute value of a product of two real numbers is equal to the product of their absolute values.

Let n be a positive integer. If x, y are elements of \mathbf{R}^n and t is a real number, then we can define $x + y$ and tx as elements of \mathbf{R}^n by adding the coordinates of x and y in the first case, and multiplying the coordinates of x by t in the second. For each x in \mathbf{R}^n , we define $|x|$, the standard Euclidean norm of x , by

$$(1.3) \quad |x| = \left(\sum_{j=1}^n x_j^2 \right)^{1/2}.$$

It is easy to see that $|tx| = |t||x|$ for all $x \in \mathbf{R}^n$ and all $t \in \mathbf{R}$, and it is well-known that

$$(1.4) \quad |x + y| \leq |x| + |y|$$

for all $x, y \in \mathbf{R}^n$.

1.2 Some basic notions and results

Let $(M, d(x, y))$ be a metric space. For the record, this means that M is a nonempty set, and that $d(x, y)$ is a nonnegative real-valued function on $M \times M$ such that $d(x, y) = 0$ if and only if $x = y$, $d(x, y) = d(y, x)$ for all $x, y \in M$, and

$$(1.5) \quad d(x, z) \leq d(x, y) + d(y, z)$$

for all $x, y, z \in M$. For each positive integer n , \mathbf{R}^n equipped with the standard Euclidean metric $|x - y|$ defines a metric space. Of course every nonempty subset of M can be viewed as a metric space itself, just by restricting the metric on M to Y .

If x is an element of M and r is a positive real number, then we write $B(x, r)$ for the open ball in M with center x and radius r , defined by

$$(1.6) \quad B(x, r) = \{y \in M : d(x, y) < r\}.$$

Similarly, the closed ball with center x and radius r is denoted $\overline{B}(x, r)$ and defined by

$$(1.7) \quad \overline{B}(x, r) = \{y \in M : d(x, y) \leq r\}.$$

1.2.1 Open and closed subsets and normality of metric spaces

If E is a subset of M and p is an element of M , then p is said to be a limit point of E if for every $\epsilon > 0$ there is a point x in E such that $x \neq p$ and $d(x, p) < \epsilon$. If every limit point of E is also an element of E , then E is said to be a closed subset of M . This is the same as saying that a point p in M lies in E if for every $\epsilon > 0$ there is a point x in E such that $d(x, p) < \epsilon$. A closed ball $\overline{B}(x, r)$ as in (1.7) is always a closed subset of M , and the empty set and M itself are closed subsets of M .

For any subset E , the *closure* is denoted \overline{E} and is defined to be the union of E and the set of limit points of E . Equivalently, a point p lies in \overline{E} if for every $\epsilon > 0$ there is an x in E such that $d(p, x) < \epsilon$. It is not difficult to show that the closure of E is always a closed subset of M . Thus $\overline{E} = E$ if and only if E is closed, and the closure of E can be described as the smallest closed subset of M that contains E . If D is a subset of M , then D is *dense* in M if the closure of D is equal to M .

A subset U of M is said to be *open* if for every x in U there is an $r > 0$ so that $B(x, r)$ is contained in U . An open ball $B(x, r)$ as in (1.6) is always an open subset of M , as are the empty set and M itself. A basic fact is that U is open if and only if $M \setminus U$ is a closed subset of M . For any subset E of M , the *interior* E° of E is the set of points x in E for which there is an $r > 0$ so that $B(x, r)$ is contained in E . One can check that E° is always an open subset of E , $E^\circ = E$ if and only if E is an open set, and E° is the largest open subset of E .

Notice that the interior of E is the same as the complement in M of the closure of the complement of E , i.e.,

$$(1.8) \quad E^\circ = M \setminus \overline{(M \setminus E)}.$$

If E_1, E_2 are subsets of M , then it is not difficult to show that

$$(1.9) \quad \overline{(E_1 \cup E_2)} = \overline{E_1} \cup \overline{E_2}, \quad \overline{(E_1 \cap E_2)} \subseteq \overline{E_1} \cap \overline{E_2}$$

and

$$(1.10) \quad E_1^\circ \cup E_2^\circ \subseteq (E_1 \cup E_2)^\circ, \quad E_1^\circ \cap E_2^\circ = (E_1 \cap E_2)^\circ.$$

The *boundary* of a subset E of M is denoted ∂E and defined by

$$(1.11) \quad \partial E = \overline{E} \setminus E^\circ.$$

This is equivalent to

$$(1.12) \quad \partial E = \overline{E} \cap \overline{(M \setminus E)},$$

and in particular

$$(1.13) \quad \partial E = \partial(M \setminus E).$$

The empty set and M itself have empty boundary in M . Observe that if E_1, E_2 are subsets of M , then

$$(1.14) \quad \partial(E_1 \cup E_2) \subseteq \partial E_1 \cup \partial E_2$$

and

$$(1.15) \quad \partial(E_1 \cap E_2) \subseteq \partial E_1 \cup \partial E_2.$$

As a metric space, M enjoys the Hausdorff property that for every pair of distinct elements x, y in M there are open sets U_1, U_2 such that $x \in U_1, y \in U_2$, and $U_1 \cap U_2 = \emptyset$. For this one can simply take U_1, U_2 to be the balls $B(x, d(x, y)/2), B(y, d(x, y)/2)$. A stronger version of this is called “regularity” and says that if x is an element of M and F is a nonempty closed subset of M which does not contain x as an element, then there are open subsets V_1, V_2 of M such that $x \in V_1, F \subseteq V_2$, and $V_1 \cap V_2 = \emptyset$. To see that this holds, choose $r > 0$ so that $B(x, r) \cap F = \emptyset$, and then choose V_1 to be $B(x, r/2)$, and V_2 to be the set of z in M such that $d(x, z) > r/2$. An even stronger separation property is called normality and states that if F_1 and F_2 are nonempty closed subsets of M such that $F_1 \cap F_2 = \emptyset$, then there are open subsets W_1, W_2 of M such that $F_1 \subseteq W_1, F_2 \subseteq W_2$, and $W_1 \cap W_2 = \emptyset$. To find such a pair W_1, W_2 , choose for each x in F_1 a positive real number $r_1(x)$ so that $B(x, r_1(x)) \cap F_2 = \emptyset$, and choose for each y in F_2 a positive real number $r_2(y)$ so that $B(y, r_2(y)) \cap F_1 = \emptyset$. Define W_1, W_2 by

$$(1.16) \quad W_1 = \bigcup_{x \in F_1} B(x, r_1(x)/2), \quad W_2 = \bigcup_{y \in F_2} B(y, r_2(y)/2).$$

Clearly W_1, W_2 are open subsets of M which contain F_1, F_2 , respectively, as subsets. It is not difficult to show that $W_1 \cap W_2 = \emptyset$. One can strengthen this a bit further by saying that there are open subsets $\widetilde{W}_1, \widetilde{W}_2$ of M such that $F_1 \subseteq \widetilde{W}_1, F_2 \subseteq \widetilde{W}_2$, and the closures of \widetilde{W}_1 and \widetilde{W}_2 are disjoint. In fact, this can be obtained as a consequence of the previous normality property, by first using normality to find disjoint open subsets W_1, W_2 of M such that $F_1 \subseteq W_1, F_2 \subseteq W_2$, and then using normality again to find open subsets $\widetilde{W}_1, \widetilde{W}_2$ of M such that $F_1 \subseteq \widetilde{W}_1, F_2 \subseteq \widetilde{W}_2$ and the closures of $\widetilde{W}_1, \widetilde{W}_2$ are contained in W_1, W_2 , respectively.

1.2.2 Bounded, compact, and totally bounded sets

A subset of M is said to be *bounded* if it is contained in a ball. If A is a bounded subset of M , then the *diameter* $\text{diam } A$ of A is defined by

$$(1.17) \quad \text{diam } A = \sup\{d(x, y) : x, y \in A\}.$$

In this case the diameter of A is a finite real number. If A is an unbounded subset of M , then the diameter of A is defined to be $+\infty$. We make the convention that the diameter of the empty set is equal to 0.

A sequence $\{p_j\}_{j=1}^\infty$ in M is said to converge to a point p in M if for every $\epsilon > 0$ there is a positive integer N such that

$$(1.18) \quad d(p_j, p) < \epsilon \quad \text{for all } j \geq N.$$

In this case we write

$$(1.19) \quad \lim_{j \rightarrow \infty} p_j = p,$$

and one can check that the limit p is unique. A sequence $\{p_j\}_{j=1}^\infty$ in M is said to be a *Cauchy sequence* if for every $\epsilon > 0$ there is a positive integer N so that

$$(1.20) \quad d(p_j, p_k) < \epsilon \quad \text{for all } j, k \geq N.$$

It is easy to see that a convergent sequence is a Cauchy sequence. A metric space in which every Cauchy sequence is a convergent sequence is said to be *complete*. A well-known fact is that the Euclidean spaces \mathbf{R}^n with their standard metrics are complete metric spaces. Also, any metric space can be embedded isometrically into a complete metric space, where the image of the first metric space is dense in the second one, and this “completion” is unique up to isometric equivalence.

A subset E of M is closed if and only if every sequence of points in E which converges in M has its limit in E . For any subset E of M , the closure \overline{E} of E is equal to the set of points in M which occur as limits of sequences of points in E . This includes constant sequences $\{p_j\}_{j=1}^\infty$, in which all the p_j 's are the same, so that the elements of E are limits of sequences of points in E . A subset D of M is dense in M if and only if every element of M occurs as the limit of a sequence of points in D .

If $\{x_j\}_{j=1}^\infty$ is a sequence of points in some set X , and if $\{j_l\}_{l=1}^\infty$ is a strictly increasing sequence of positive integers, so that

$$(1.21) \quad j_1 < j_2 < j_3 < \cdots,$$

then the sequence $\{x_{j_l}\}_{l=1}^\infty$ obtained by restricting the $\{x_j\}_{j=1}^\infty$ to $\{j_l\}_{l=1}^\infty$ is called a *subsequence* of $\{x_j\}_{j=1}^\infty$. In particular, $\{x_j\}_{j=1}^\infty$ is a subsequence of itself, by taking $j_l = l$ for all l . Note that a Cauchy sequence in a metric space converges as soon as it has a convergent subsequence.

In these notes we shall often be concerned with subsets of metric spaces which are *compact*. As usual, compactness can be characterized in a number of equivalent ways. In brief, a subset K of M is compact if (a) every for every covering of K by open subsets of M there is a subcovering of K from this covering with only finitely many elements, or (b) every infinite subset of K has a limit point in K , or (c) every sequence of points in K has a subsequence which converges. If Y is a nonempty subset of M and K is contained in Y , then K is compact as a subset of M if and only if K is compact as a subset of Y , viewed as a metric space itself, with the restriction of the metric from M . In particular, if K is nonempty, we can take $Y = K$, so that K is compact as a subset of M if and only if K is compact as a subset of itself, viewed as a metric space on its own, with the restriction of the metric from M . This is quite different from the properties of being open or closed.

In the case of a Euclidean space \mathbf{R}^n , the compact subsets are exactly the closed and bounded subsets, by the Heine–Borel theorem.

A subset A of M is said to be *totally bounded* if for every $\epsilon > 0$ there is a finite collection of balls in M with radius ϵ whose union contains A as a subset. Compact sets are totally bounded, and subsets of totally bounded sets are totally bounded. As with compactness, a subset A of M is totally bounded as a subset of M if and only if it is totally bounded as a subset of itself, viewed as a metric space on its own, with the restriction of the metric from M .

A well-known theorem states that a metric space is compact if and only if it is totally bounded and complete. This is equivalent to saying that a subset of a complete metric space is compact if and only if it is closed and totally bounded. Another way to look at this is that a subset A of M is totally bounded if and only if every sequence of points in A has a subsequence which is a Cauchy sequence.

1.2.3 Separable metric spaces

Recall that M is said to be *separable* if there is a subset D of M which is dense and at most countable. In this survey we shall be primarily interested in separable metric spaces. For example, the Euclidean spaces \mathbf{R}^n with their standard metrics are separable, because the set of points in \mathbf{R}^n with rational coordinates is countable and dense.

A metric space which is totally bounded is also separable. In fact, a metric space M is separable if and only if for each $\epsilon > 0$ there is a subset D_ϵ of M which is at most countable and such that for each $x \in M$ there is a $y \in D_\epsilon$ that satisfies $d(x, y) < \epsilon$. The property of being totally bounded is the same as saying that one can do this with D_ϵ being finite for each $\epsilon > 0$.

If M is a metric space, a collection \mathcal{B} of open subsets of M is said to be a *basis for the topology* if every open subset of M can be expressed as a union of subsets of M which are elements of \mathcal{B} . This is equivalent to saying that for each x in M and each positive real number r there is an open set U which is an element of \mathcal{B} such that x is an element of U and U is a subset of $B(x, r)$. Separability of M can be characterized by the existence of a basis \mathcal{B} for the topology of M which has at most countably-many elements. For if D is a dense subset of M which is at most countable, then the collection of balls of the form $B(y, 1/n)$, where y runs through the elements of D and n runs through the positive integers, is a basis for the topology of M which is at most countable. Conversely, if \mathcal{B} is a basis for the topology of M which is at most countable, then one can get a subset D of M which is dense and at most countable by choosing a point in each element of \mathcal{B} and putting it into D .

A basic fact about separable metric spaces is that every subset E of a separable metric space M contains a subset which is dense in E and at most countable. In other words, every nonempty subset of M defines a separable metric space itself, with the restriction of the metric from M . This is not difficult to show, but notice that a particular dense subset of M may not

have any elements in any particular subset E of M .

A subset of a metric space is said to be *countably compact* if for every open covering of the set there is a subcovering with at most countably many elements. Clearly a compact set is countably compact. A well-known result states that a metric space M is countably compact if and only if it is separable. Indeed, if M is countably compact, then for every $r > 0$ there is a family of balls of radius r with at most countably many elements which covers M , and this implies that M is separable. Conversely, suppose that M is separable, and that $\{U_\alpha\}_{\alpha \in A}$ is an open covering of M . Let \mathcal{B} be a basis for the topology of M which is at most countable. Define \mathcal{B}_1 to be the subset of \mathcal{B} consisting of those open subsets V of M in \mathcal{B} which are contained in a U_α for some $\alpha \in A$. Since \mathcal{B} is a basis for the topology of M , each U_α is a union of elements of \mathcal{B} . Hence

$$(1.22) \quad \bigcup_{V \in \mathcal{B}_1} V = \bigcup_{\alpha \in A} U_\alpha = M.$$

For each V in \mathcal{B}_1 , choose an $\alpha(V) \in A$ such that $V \subseteq U_{\alpha(V)}$. Then

$$(1.23) \quad \bigcup_{V \in \mathcal{B}_1} U_{\alpha(V)} = M,$$

and $\{U_{\alpha(V)} : V \in \mathcal{B}_1\}$ is a subcovering of M from the covering $\{U_\alpha\}_{\alpha \in A}$ which has at most countably many elements.

1.3 The Baire category theorem

Let $(M, d(x, y))$ be a metric space. If U, V are dense open subsets of M , then the intersection $U \cap V$ is also a dense open subset of M . This is not difficult to verify. More precisely, if U is a dense open subset of M and D is a dense subset of M , then the intersection $U \cap D$ is a dense subset of M . Of course the intersection of two dense sets in general can be dense.

Now suppose that $\{U_j\}_{j=1}^\infty$ is a sequence of dense open subsets of M . In general, the intersection of all of the U_j 's might be the empty set. This occurs when M is the set of rational numbers, equipped with the usual metric, for instance. The Baire category theorem states that if M is a complete metric space, then in fact $\bigcap_{j=1}^\infty U_j$ is a dense subset of M .

Let us briefly review the proof. Let B_0 be any closed ball in M . It suffices to show that B_0 contains an element of $\bigcap_{j=1}^\infty U_j$. Since U_1 is a dense open

subset of M , there is a closed ball B_1 in M such that $B_1 \subseteq B_0$, $B_1 \subseteq U_1$, and the radius of B_1 is less than 1. Similarly, because U_2 is a dense open subset of M , there is a closed ball B_2 in M such that $B_2 \subseteq B_1$, $B_2 \subseteq U_2$, and the radius of B_2 is less than $1/2$. Proceeding in this manner, for each positive integer j there is a closed ball B_j such that $B_j \subseteq B_{j-1}$, $B_j \subseteq U_j$, and the radius of B_j is less than $1/j$. It follows that the sequence of centers of the B_j 's is a Cauchy sequence, and so the assumption that M is complete implies that this sequence converges to a point p in M . One can check that p lies in each B_j , and hence in each U_j , as well as the given ball B_0 .

There is a complementary picture for closed sets with empty interior. In any metric space M , if E and F are closed sets with empty interior, then $E \cup F$ is also a closed set with empty interior. This does not work without the assumption that the sets be closed, since the real line is the union of the sets of rational and irrational numbers, each of which has empty interior. In a complete metric space the union of sequence of closed sets with empty interior also has empty interior, although this union is not a closed set in general.

If F is any nonempty closed subset of M and j is a positive integer, consider the set U_j which is the union of the balls $B(x, 1/j)$ with $x \in F$. This is an open subset of M which contains F as a subset. It is not difficult to check that

$$(1.24) \quad F = \bigcap_{j=1}^{\infty} U_j.$$

Thus every closed subset of a metric space can be expressed as the intersection of a sequence of open sets. By passing to complements one obtains that every open subset of a metric space can be expressed as the union of a sequence of closed subsets.

1.4 Continuous and uniformly continuous mappings

Let $(M, d(x, y))$ and $(N, \rho(u, v))$ be metric spaces, and let E be a nonempty subset of M . A mapping f from E to N is said to be *continuous* if for every x in E and every $\epsilon > 0$ there is a $\delta > 0$ so that

$$(1.25) \quad \rho(f(x), f(y)) < \epsilon \quad \text{for all } y \in E \text{ such that } d(y, x) < \delta.$$

The mapping f is said to be *uniformly continuous* if for every $\epsilon > 0$ there is a $\delta > 0$ such that

$$(1.26) \quad \rho(f(x), f(y)) < \epsilon \quad \text{for all } x, y \in E \text{ such that } d(y, x) < \delta.$$

A basic result states that if E is compact and $f : E \rightarrow N$ is continuous, then f is also uniformly continuous.

Let us assume for simplicity that $E = M$, to which one can reduce anyway. Another basic result states that $f : M \rightarrow N$ is continuous if and only if $f^{-1}(U)$ is an open subset of M for all open subsets U of N . This is equivalent to saying that $f : M \rightarrow N$ is continuous if and only if $f^{-1}(A)$ is a closed subset of M whenever A is a closed subset of N . An alternate characterization states that $f : M \rightarrow N$ is continuous if and only if for every sequence $\{x_j\}_{j=1}^{\infty}$ of points in M which converges to some point x , the sequence $\{f(x_j)\}_{j=1}^{\infty}$ converges in N to $f(x)$.

A mapping $f : M \rightarrow N$ is said to be *bounded* if the image $f(M)$ of f is contained in a bounded subset of N . A basic result states that if $f : M \rightarrow N$ is continuous and K is a compact subset of M , then $f(K)$ is a compact subset of N . In particular, f is bounded in this case. Also, if E is a subset of M which is totally bounded and $f : M \rightarrow N$ is uniformly continuous, then $f(E)$ is totally bounded.

A *homeomorphism* from M to N is a one-to-one mapping f of M onto N such that $f : M \rightarrow N$ and $f^{-1} : N \rightarrow M$ are both continuous. If $f : M \rightarrow N$ is continuous, M is compact, and f is a one-to-one mapping of M onto N , then f is a homeomorphism of M onto N . For instance, f maps closed subsets of M to closed subsets of N in this case, since closed subsets of M are compact.

Suppose that f_1, f_2 are continuous mappings from M to N , and that E is a dense subset of M . If f_1 and f_2 are equal on E , then they are equal on all of M . In other words, a continuous mapping is uniquely determined by its restriction to a dense subset. Conversely, suppose that D is a dense subset of M , and that f is a continuous mapping from D into N . In order for there to be an extension of f to a continuous mapping from M to N , it is sufficient that f be uniformly continuous, and that N be complete. In this case the extension will also be uniformly continuous.

1.4.1 Real-valued functions

Of course constant functions are always continuous. If $f_1(x)$, $f_2(x)$ are real-valued continuous functions on M , then the sum $f_1(x) + f_2(x)$ and the product $f_1(x) \cdot f_2(x)$ are also continuous functions on M . The sum of uniformly continuous real-valued functions is uniformly continuous, but this does not work for the product in general. It does work if at least one of the two functions is bounded. If $f(x)$ is a continuous real-valued function on M and E is a subset of M such that $f(x) \neq 0$ for all $x \in E$, then $1/f(x)$ is a continuous function on E . If we assume that $f(x)$ is uniformly continuous, then it may not be true that $1/f(x)$ is uniformly continuous on E , but this is the case if there is a positive real number c such that $|f(x)| \geq c$ for all x in E .

On \mathbf{R}^n , the n coordinate functions

$$(1.27) \quad x = (x_1, \dots, x_n) \mapsto x_j,$$

$1 \leq j \leq n$, are continuous, as one can easily verify. As a result, polynomials on \mathbf{R}^n are continuous functions, and rational functions, which are quotients of polynomials, are continuous on the set where the denominator does not vanish.

Lemma 1.28 *For each element p of M , $d(x, p)$ is uniformly continuous as a real-valued function of x on M .*

This follows from the triangle inequality. Namely,

$$(1.29) \quad d(x, p) \leq d(y, p) + d(x, y)$$

and

$$(1.30) \quad d(y, p) \leq d(x, p) + d(x, y),$$

so that

$$(1.31) \quad |d(x, p) - d(y, p)| \leq d(x, y).$$

Let A be a nonempty subset of M , and set

$$(1.32) \quad \text{dist}(x, A) = \inf\{d(x, z) : z \in A\}$$

for x in M . Note that

$$(1.33) \quad \text{dist}(x, \overline{A}) = \text{dist}(x, A)$$

for all x in M , and that $\text{dist}(x, A) = 0$ if and only if x lies in the closure of A .

Lemma 1.34 *As a function of x on M , $\text{dist}(x, A)$ is uniformly continuous.*

For this we use the triangle inequality again. If x and y are arbitrary elements of M and z is an arbitrary element of A , then

$$(1.35) \quad \text{dist}(x, A) \leq d(x, z) \leq d(x, y) + d(y, z).$$

By taking the infimum over z it follows that

$$(1.36) \quad \text{dist}(x, A) \leq d(x, y) + \text{dist}(y, A).$$

Similarly,

$$(1.37) \quad \text{dist}(y, A) \leq d(x, y) + \text{dist}(x, A),$$

so that

$$(1.38) \quad |\text{dist}(x, A) - \text{dist}(y, A)| \leq d(x, y).$$

Note that for any nonempty subset A of M ,

$$(1.39) \quad \text{dist}(x, A) = \text{dist}(x, \overline{A})$$

for all x in M , as one can verify from the definitions.

Suppose that F_1, F_2 are disjoint nonempty closed subsets of M . Define a real-valued function $\phi(x)$ on M by

$$(1.40) \quad \phi(x) = \frac{\text{dist}(x, F_1)}{\text{dist}(x, F_1) + \text{dist}(x, F_2)}.$$

Because F_1 and F_2 are disjoint and closed, the denominator in the definition of $\phi(x)$ is never equal to 0. Thus $\phi(x)$ is a continuous function on M which takes values in the interval $[0, 1]$, and it satisfies $\phi(x) = 0$ if and only if $x \in F_1$ and $\phi(x) = 1$ if and only if $x \in F_2$.

In general this function $\phi(x)$ may not be uniformly continuous. For example, if $M = \mathbf{R} \setminus \{0\}$, $F_1 = (-\infty, 0)$, and $F_2 = (0, \infty)$, then F_1, F_2 are disjoint nonempty closed subsets of M , $\phi(x)$ is defined completely by $\phi(x) = 0$ when $x \in F_1$ and $\phi(x) = 1$ when $x \in F_2$, and ϕ is clearly not uniformly continuous.

Let us say that a mapping $f : M \rightarrow N$ is *locally bounded* if for every p in M there is an $r > 0$ so that $f(B(p, r))$ is a bounded subset of N . Thus continuous mappings are automatically locally bounded. Let us say that $f : M \rightarrow N$ is *locally uniformly continuous* if for each $p \in M$ there is an $r > 0$ such that the restriction of f to $B(p, r)$ is uniformly continuous.

Ordinary continuity is already a local condition, and so we do not need to define “local continuity”. For real-valued functions, the sum and product of locally bounded functions is locally bounded, and the sum and product of locally uniformly continuous functions is locally uniformly continuous. If $f(x)$ is a real valued function on M which is locally uniformly continuous and which satisfies $f(x) \neq 0$ for all $x \in M$, then $1/f(x)$ is locally uniformly continuous on M .

In particular, the function $\phi(x)$ defined in (1.40) is always locally uniformly continuous, even if it may not be uniformly continuous. If there is an $\epsilon > 0$ so that $d(y, z) \geq \epsilon$ for all $y \in F_1$ and $z \in F_2$, then $\phi(x)$ is uniformly continuous.

Let us say that a mapping between two metric spaces is *countably uniformly continuous* if the domain can be expressed as the union of an at most countable family of subsets, on each of which the mapping is uniformly continuous. One can also assume that the mapping is bounded on each of these subsets, by making a further decomposition as necessary. As a basic property of such mappings, if the domain can be expressed as the countable union of totally bounded subsets, then so can the image of the mapping.

The sum or product of two real-valued countably uniformly continuous functions is also countably uniformly continuous. This is not difficult to show, using the observation that if $\{E_i\}_{i \in I}$ and $\{F_j\}_{j \in J}$ are families of subsets of M such that I, J are at most countable and

$$(1.41) \quad \bigcup_{i \in I} E_i = \bigcup_{j \in J} F_j = M,$$

then $\{E_i \cap F_j\}_{(i,j) \in I \times J}$ is an at most countable family of subsets of M whose union is also equal to M . Similarly, the reciprocal of a countably uniformly continuous mapping which does not take the value 0 is countably uniformly continuous. A locally uniformly continuous mapping between two metric spaces is countably uniformly continuous if the domain is separable, since that implies countable compactness.

1.4.2 Spaces of continuous mappings

We shall write $C(M, N)$ for the space of continuous mappings from M to N . A mapping $f : M \rightarrow N$ is said to be *bounded* if its image is contained in a bounded subset of N , and we write $C_b(M, N)$ for the space of bounded continuous mappings from M to N . Note that $C(M, N) = C_b(M, N)$ if M

is compact. If f_1, f_2 are two bounded continuous mappings from M to N , then we set

$$(1.42) \quad \theta(f_1, f_2) = \sup\{\rho(f_1(x), f_2(x)) : x \in M\}.$$

It is not difficult to verify that this defines a metric on $C_b(M, N)$. The constant mappings from M to N , which take all of M to a single point in N , are obviously bounded and continuous, and in this way we get a natural embedding of N in $C_b(M, N)$. This is an isometric embedding, which is to say that the given metric $\rho(\cdot, \cdot)$ on N agrees with the metric $\theta(\cdot, \cdot)$ applied to the constant mappings.

If $\{f_j\}_{j=1}^\infty$ is a sequence of mappings from M to N , and if f is another mapping from M to N , then we say that $\{f_j\}_{j=1}^\infty$ *converges pointwise* if

$$(1.43) \quad \lim_{j \rightarrow \infty} f_j(x) = f(x) \quad \text{in } N$$

for all x in M . We say that $\{f_j\}_{j=1}^\infty$ converges to f *uniformly* if for every $\epsilon > 0$ there is an integer L so that

$$(1.44) \quad \rho(f_j(x), f(x)) < \epsilon \quad \text{for all } j \geq L.$$

Thus uniform convergence implies pointwise convergence.

A well-known result states that if $\{f_j\}_{j=1}^\infty$ is a sequence of mappings from M to N which converges uniformly to a mapping f from M to N , and if each f_j is continuous, then f is continuous as well. This does not work in general under the assumption of pointwise convergence instead of uniform convergence. Note that a sequence $\{f_j\}_{j=1}^\infty$ in $C_b(M, N)$ converges to a function f in $C_b(M, N)$ with respect to the metric θ defined above if and only if $\{f_j\}_{j=1}^\infty$ converges uniformly to f . If $\{f_j\}_{j=1}^\infty$ converges uniformly to f and each f_j is uniformly continuous, then f is uniformly continuous as well.

If N is complete as a metric space, then $C_b(M, N)$ is complete as a metric space. In other words, if $\{f_j\}_{j=1}^\infty$ is a Cauchy sequence in $C_b(M, N)$ with respect to the metric θ and N is complete as metric space, then there is a function f in $C_b(M, N)$ to which $\{f_j\}_{j=1}^\infty$ converges uniformly. To show this, one can first use the assumption that $\{f_j\}_{j=1}^\infty$ is a Cauchy sequence in $C_b(M, N)$ to obtain that $\{f_j(x)\}_{j=1}^\infty$ is a Cauchy sequence in N for all x in M . Hence $\{f_j\}_{j=1}^\infty$ converges pointwise to some mapping f from M to N , since N is complete. Using the assumption that $\{f_j\}_{j=1}^\infty$ is a Cauchy sequence with respect to the metric θ , one can check that $\{f_j\}_{j=1}^\infty$ converges to f uniformly. As a consequence, f is continuous, and one can also verify that f is bounded in this case.

Let us write $C(M)$ for the space of continuous real-valued functions on M , and $C_b(M)$ for the space of bounded continuous real-valued functions on M . Thus $C(M)$ and $C_b(M)$ are vector spaces over the real numbers, and on $C_b(M)$ we have the norm

$$(1.45) \quad \|f\|_{sup} = \sup\{|f(x)| : x \in M\}.$$

That is, $\|f\|_{sup}$ is a nonnegative real number which is equal to 0 exactly when f is identically equal to 0, $\|tf\|_{sup} = |t| \|f\|_{sup}$ for all real numbers t and all f in $C_b(M)$, and the triangle inequality for norms holds, namely

$$(1.46) \quad \|f_1 + f_2\|_{sup} \leq \|f_1\|_{sup} + \|f_2\|_{sup}$$

for all f_1, f_2 in $C_b(M)$. These properties are not difficult to verify. The metric $\theta(\cdot, \cdot)$ defined in general above is the same as the metric associated to the norm $\|\cdot\|_{sup}$ on $C_b(M)$, which is to say that

$$(1.47) \quad \theta(f_1, f_2) = \|f_1 - f_2\|_{sup}$$

for all f_1, f_2 in $C_b(M)$. The completeness of $C_b(M)$ as a metric space implies that $C_b(M)$ equipped with the norm $\|\cdot\|_{sup}$ is a *Banach space*. In fact it is a *Banach algebra* with respect to the additional operation of multiplication of functions, which means that

$$(1.48) \quad \|f_1 f_2\|_{sup} \leq \|f_1\|_{sup} \|f_2\|_{sup}$$

for all f_1, f_2 in $C_b(M)$.

There is a natural mapping from M to $C(M)$, in which a point p in M is associated to the continuous function $f_p(x) = d(x, p)$. If M is bounded, then this defines a mapping from M into $C_b(M)$, and one can show that this mapping is an isometric embedding with respect to the metric $\theta(\cdot, \cdot)$ on $C_b(M)$. In other words,

$$(1.49) \quad \sup\{|f_p(x) - f_q(x)| : x \in M\} = d(p, q)$$

for all p, q in M , as one can check. If M is unbounded, then one can adjust this as follows. Fix an element w of M , which will serve as a base point. To each point p in M we can associate the function $f_p(x) - f_w(x)$. It is not hard to see that this is a bounded continuous function on M , and that the distance between $f_p - f_w$ and $f_q - f_w$ is equal to $d(p, q)$, so that we again get an isometric embedding of M into $C_b(M)$.

1.5 Connectedness

Let $(M, d(x, y))$ be a metric space. Two subsets A, B of M are said to be *separated* if

$$(1.50) \quad \overline{A} \cap B = \emptyset, \quad A \cap \overline{B} = \emptyset.$$

A subset E of M is said to be *connected* if it cannot be written as the union of two separated sets.

If E, Y are subsets of M with Y nonempty and $E \subseteq Y$, then E is connected as a subset of M if and only if E is connected as a subset of Y , viewed as a metric itself, with the restriction of the metric from M . This is not difficult to verify, just from the definition. In particular, E is connected as a subset of M if and only if E is connected as a subset of itself, viewed as a metric space, with the restriction of the metric from M .

The empty set and sets with only one element are automatically connected. It is well-known that the connected subsets of the real line are the empty set and the subsets which are intervals, which may be open, closed, or mixed, and which may be bounded or unbounded. This includes the real line itself.

The connectedness of M itself is equivalent to saying that M cannot be written as the disjoint union of two nonempty open subsets, or that M cannot be written as the disjoint union of two nonempty closed subsets.

The closure of a connected set is always connected. The union of two connected sets is connected as long as they have nonempty intersection. More generally, the union of two nonempty connected sets is connected if and only if they are not separated.

Let E be a subset of M , and suppose that for every pair of points p, q in E there is a subset $E_{p,q}$ of E which is connected and which contains p, q . Then E is connected.

A set which contains no connected subset with more than one element is said to be *totally disconnected*. This terminology will be convenient, but it has the feature that the empty set and sets with one element are called connected and totally disconnected.

In general, we can define a relation \sim on a metric space M by saying that $x \sim y$ when x and y are contained in a connected subset of M . This relation is clearly reflexive and symmetric, and it is also transitive, because of the earlier remark about the union of two connected sets being connected when they contain a common element.

Thus \sim defines an equivalence relation on M . The corresponding equivalence classes are called the *connected components* of M . It is not hard to see that the connected components of M are indeed connected subsets of M , and in fact they are the maximal connected subsets of M . The connected components of M are also closed sets, since the closure of a connected set is always connected.

If $(M_1, d_1(x, y))$ and $(M_2, d_2(u, v))$ are two metric spaces and f is a continuous mapping from M_1 to M_2 , and if E is a connected subset of M_1 , then $f(E)$ is a connected subset of M_2 . As a special case of this, suppose that a, b are real numbers with $a \leq b$, and that $p(t)$ is a continuous mapping from $[a, b]$ into M . We call $p(t)$ a *path* in M . The image of a path is always a connected set, since it is the image of a connected set under a continuous mapping.

A subset E of M is said to be *pathwise connected* if for each x, y in E there is a continuous path $p(t)$ defined on an interval $[a, b]$ and taking values in E such that $p(a) = x$ and $p(b) = y$. Because the image of the path is a connected set, it follows that every pair of points in E is contained in a connected subset of E , so that E is itself connected.

Let us define a relation \approx on M by saying that $x \approx y$ if there is a path in M from x to y , i.e., if there is a closed interval $[a, b]$ in the real line and a continuous mapping $p(t)$ from $[a, b]$ into M such that $p(a) = x$, $p(b) = y$. Note that $x \approx y$ implies $x \sim y$. One can check that \approx defines an equivalence relation on M . The equivalence classes associated to the relation \approx are pathwise connected subsets of M , and they are called the *pathwise-connected components* of M . The pathwise-connected components of M are the maximal pathwise-connected subsets of M , and they are contained in the ordinary connected components of M .

Connected subsets of the real line are clearly pathwise connected. Now fix a positive integer n , and suppose that M is an open subset of \mathbf{R}^n , equipped with the standard metric. It is not difficult to show that the pathwise connected components of M are then open subsets of M . In fact the pathwise-connected components are the same as the connected components, and M is pathwise-connected if it is connected. This works more generally under local connectedness conditions for M , which hold in a very simple way for open subsets of \mathbf{R}^n , just using paths along line segments locally.

Fix a positive real number ϵ . By an ϵ -*chain* in M we mean a finite sequence z_1, \dots, z_k of points in M such that $d(z_j, z_{j+1}) < \epsilon$ when $1 \leq j \leq k-1$. This condition is considered to hold automatically when $k = 1$, so that

a single point defines an ϵ -chain. A subset E of M is said to be ϵ -connected if for every pair of points x, y in E there is an ϵ -chain z_1, \dots, z_k in E such that $z_1 = x$ and $z_k = y$.

Let us again assume that $E = M$ for simplicity. Define a relation \simeq_ϵ on M by saying that $x \simeq_\epsilon y$ if there is an ϵ -chain z_1, \dots, z_k in M such that $z_1 = x$ and $z_k = y$. It is easy to see that \simeq_ϵ defines an equivalence relation on M , and that the corresponding equivalence classes are the maximal ϵ -connected subsets of M . By construction, two points in different equivalence classes associated to \simeq_ϵ have distance at least ϵ from each other. Using this, one can show that if M is connected in the ordinary sense, then M is ϵ -connected for all $\epsilon > 0$.

Conversely, if M is compact and ϵ -connected for each $\epsilon > 0$, then M is connected. Note that the set of rational numbers, equipped with the usual metric, is ϵ -connected for each $\epsilon > 0$, but not connected in the usual sense.

Chapter 2

Some measure functionals and dimensions

If M is a set, then by a *measure functional on subsets of M* we mean a function μ which takes subsets of M to nonnegative real numbers or $+\infty$ and which satisfies $\mu(\emptyset) = 0$ and $\mu(A) \leq \mu(B)$ whenever $A, B \subseteq M$ and $A \subseteq B$. In many cases the measure functional satisfies additional nice properties, at least for some reasonably-large class of subsets of M .

2.1 Basic notions

Let $(M, d(x, y))$ be a metric space, and let E be a subset of M . Define $\mathcal{U}_{con}(E)$ to be the collection of finite or countably-infinite families $\{A_i\}_i$ of subsets of M such that E is contained in the union $\bigcup_i A_i$. For each nonnegative real number α , define the α -dimensional Hausdorff content of E , denoted $H_{con}^\alpha(E)$, to be the infimum of the sums

$$(2.1) \quad \sum_i (\text{diam } A_i)^\alpha$$

over all coverings $\{A_i\}_i$ of E in $\mathcal{U}_{con}(E)$. If some A_i is unbounded, so that $\text{diam } A_i = +\infty$, then we interpret $(\text{diam } A_i)^\alpha$ to be $+\infty$ as well, for all $\alpha \geq 0$. If A_i is bounded and $\alpha = 0$, then we interpret $(\text{diam } A_i)^\alpha$ as being 1 when A_i is not empty, and we interpret it as being 0 if A_i is the empty set. In particular, the α -dimensional Hausdorff content of the empty set is taken to be 0. Technically, one might also say that for the empty set

the empty covering is admissible, and that the corresponding empty sum is automatically equal to 0.

A simple estimate is

$$(2.2) \quad H_{con}^\alpha(E) \leq (\text{diam } E)^\alpha.$$

For $\alpha = 0$ we have that $H_{con}^0(E)$ is equal to 0 when E is the empty set, to 1 when E is nonempty and bounded, and to $+\infty$ when E is unbounded.

Now suppose that δ is an extended real number such that $0 < \delta \leq \infty$, and that α is a nonnegative real number. If E is a subset of M , we define $\mathcal{U}_\delta(E)$ to be the collection of finite or countably-infinite families $\{A_i\}_i$ of subsets of M such that E is contained in the union $\bigcup_i A_i$ and $\text{diam } A_i < \delta$ for all i . We then define $H_\delta^\alpha(E)$ to be the infimum of the same sum (2.1) over all $\{A_i\}_i$ in $\mathcal{U}_\delta(E)$ assuming that $\mathcal{U}_\delta(E)$ is nonempty, and otherwise we define $H_\delta^\alpha(E)$ to be $+\infty$. If M is separable, then one can check that $\mathcal{U}_\delta(E)$ is always nonempty. In that situation M can be covered by a family of at most countably many balls of radius r for any fixed $r > 0$, for instance. In fact, $\mathcal{U}_\delta(E)$ is nonempty for all $\delta > 0$ if and only if E contains a countable dense subset. By contrast, $\mathcal{U}_\infty(E)$ is automatically nonempty, because for any fixed p in M , E can be covered by the family of balls $B(p, n)$, where n runs over all positive integers.

It is easy to see that

$$(2.3) \quad H_{\delta_1}^\alpha(E) \geq H_{\delta_2}^\alpha(E)$$

whenever $0 < \delta_1 \leq \delta_2 \leq +\infty$. This is because

$$(2.4) \quad \mathcal{U}_{\delta_1}(E) \subseteq \mathcal{U}_{\delta_2}(E),$$

while the sums involved in the definitions of $H_{\delta_1}^\alpha(E)$, $H_{\delta_2}^\alpha(E)$ are identical. In other words, $H_{\delta_1}^\alpha(E)$ and $H_{\delta_2}^\alpha(E)$ are defined as the infima of the same expressions, but with different collections of coverings, with the coverings for $H_{\delta_1}^\alpha(E)$ being more restricted than the coverings for $H_{\delta_2}^\alpha(E)$. Notice also that

$$(2.5) \quad H_\infty^\alpha(E) = H_{con}^\alpha(E).$$

Again the sums being used are the same, and the coverings of E in $\mathcal{U}_\infty(E)$ are included in $\mathcal{U}_{con}(E)$. The converse does not quite hold, but if $\{A_i\}_i$ lies in $\mathcal{U}_{con}(E)$ and $\sum_i (\text{diam } A_i)^\alpha < \infty$, then $\text{diam } A_i < \infty$ for all i , and $\{A_i\}_i$ lies in $\mathcal{U}_\infty(E)$. It is easy to use this to obtain (2.5).

When $\alpha = 0$ we are reduced to the following. If E is the empty set, then $H_\delta^0(E) = 0$, and otherwise $H_\delta^0(E)$ is equal to the smallest number of sets of diameter less than δ needed to cover E , where this is interpreted as being $+\infty$ if no finite covering of E by sets with diameter less than δ exists.

The α -dimensional Hausdorff measure of E is denoted $H^\alpha(E)$ and defined by

$$(2.6) \quad H^\alpha(E) = \sup\{H_\delta^\alpha(E) : \delta > 0\}.$$

Because of the monotonicity property (2.3), this is the same as the limit as $\delta \rightarrow 0$ of $H_\delta^\alpha(E)$. By definition,

$$(2.7) \quad H_{con}^\alpha(E) \leq H_\delta^\alpha(E) \leq H^\alpha(E)$$

for $0 < \delta \leq \infty$.

If $\alpha = 0$, then one can check that Hausdorff measure reduces to counting measure. That is to say, $H^0(E)$ is equal to the number of elements of E when E is finite, and to $+\infty$ when E is infinite.

Observe that H_{con}^α , H_δ^α , and H^α are all measure functionals on subsets of M in the sense described at the beginning of the chapter. Also, for each $\alpha \geq 0$, $H_{con}^\alpha(E) = 0$ implies that $H_\delta^\alpha(E) = 0$ for all $\delta > 0$, and hence that $H^\alpha(E) = 0$. Thus for a fixed α , the measure functionals H_{con}^α , H_δ^α , and H^α vanish on the same subsets of M .

We can make analogous definitions using finite coverings. Namely, for a subset E of M , let $\mathcal{U}_{con}^f(E)$ denote the collection of finite families $\{A_i\}_i$ of subsets of M such that E is contained in $\bigcup_i A_i$. For each nonnegative real number α , define $HF_{con}^\alpha(E)$ to be the infimum of

$$(2.8) \quad \sum_i (\text{diam } A_i)^\alpha$$

over all finite coverings $\{A_i\}_i$ of E in M . Thus we have the simple inequality

$$(2.9) \quad HF_{con}^\alpha(E) \leq (\text{diam } E)^\alpha.$$

For $0 < \delta \leq \infty$, define $\mathcal{U}_\delta^f(E)$ to be the collection of finite families $\{A_i\}_i$ of subsets of M such that $E \subseteq \bigcup_i A_i$ and $\text{diam } A_i < \delta$ for each i . We define $HF_\delta^\alpha(E)$ to be the infimum of (2.8) over all $\{A_i\}_i$ in $\mathcal{U}_\delta^f(E)$ if $\mathcal{U}_\delta^f(E)$ is not empty, and otherwise to be $+\infty$. Observe that E is totally bounded if and only if $\mathcal{U}_\delta^f(E)$ is finite for all $\delta > 0$, which is also equivalent to saying that $HF_\delta^\alpha(E)$ is finite for all $\delta > 0$. If E is unbounded, then $HF_{con}^\alpha(E)$ and $HF_\delta^\alpha(E)$ are equal to $+\infty$ for all α, δ .

As before, $\mathcal{U}_{\delta_1}^f(E) \subseteq \mathcal{U}_{\delta_2}^f(E)$ when $\delta_1 \leq \delta_2$, and hence

$$(2.10) \quad HF_{\delta_2}^\alpha(E) \leq HF_{\delta_1}^\alpha(E)$$

in this case. Also,

$$(2.11) \quad HF_\infty^\alpha(E) = HF_{con}^\alpha(E),$$

almost by definition. We define $HF^\alpha(E)$ by

$$(2.12) \quad HF^\alpha(E) = \sup\{HF_\delta^\alpha(E) : \delta > 0\},$$

which is the same as $\lim_{\delta \rightarrow 0} HF_\delta^\alpha(E)$, by monotonicity. Thus

$$(2.13) \quad HF_{con}^\alpha(E) \leq HF_\delta^\alpha(E) \leq HF^\alpha(E)$$

for $0 < \delta \leq \infty$.

When $\alpha = 0$, these quantities are the same as their counterparts described before, i.e., $HF_{con}^0(E) = H_{con}^0(E)$, $HF_\delta^0(E) = H_\delta^0(E)$ for $0 < \delta \leq \infty$, and $HF^0(E) = H^0(E)$. For each α and δ , HF_{con}^α , HF_δ^α , and HF^α define measure functionals on subsets of M . If α is fixed, then HF_{con}^α , HF_δ^α , and HF^α are equal to 0 on the same subsets of M .

Because $\mathcal{U}^f(E) \subseteq \mathcal{U}(E)$ and $\mathcal{U}_\delta^f(E) \subseteq \mathcal{U}_\delta(E)$ for all δ , we have that

$$(2.14) \quad H_{con}^\alpha(E) \leq HF_{con}^\alpha(E), \quad H_\delta^\alpha(E) \leq HF_\delta^\alpha(E), \quad H^\alpha(E) \leq HF^\alpha(E).$$

We shall see cases when these inequalities are strict, and cases where equality holds.

There are a number of variations of these definitions, in which one might restrict the sets A_i used in the coverings, or use other measurements of the sizes of the A_i 's, etc. These matters are related to well-known work of Carathéodory, Jordan, and Minkowski, for instance.

Note that if Y is a nonempty subset of M and E is a subset of Y , then $H_{con}^\alpha(E)$, $HF_{con}^\alpha(E)$, $H_\delta^\alpha(E)$, $HF_\delta^\alpha(E)$, $H^\alpha(E)$, and $HF^\alpha(E)$ are the same for E as a subset of M as they are for E as a subset of Y , viewed as metric space itself, with the restriction of the metric $d(x, y)$ from M .

2.2 Coverings by open and closed subsets

Let $(M, d(x, y))$ continue to be some metric space.

Lemma 2.15 *If A is a subset of M , then $\text{diam } \overline{A} = \text{diam } A$.*

This is easy to check. As a result, for the definitions of the various measures of a set E in Section 2.1, one may as well use coverings by families $\{A_i\}_i$ of *closed* subsets of M . In other words, if one starts with a covering $\{A_i\}_i$ of some subset E of M by arbitrary subsets of M , then one can replace this with the family $\{\overline{A_i}\}_i$, and the latter is still be a covering of E . This new covering also satisfies the same side conditions as the initial covering used in Section 2.1, and does not change the value of sums of the form $\sum_i (\text{diam } A_i)^\alpha$.

Lemma 2.16 *If E is a subset of M , then $HF_{\text{con}}^\alpha(E) = HF_{\text{con}}^\alpha(\overline{E})$, $HF_\delta^\alpha(E) = HF_\delta^\alpha(\overline{E})$, and $HF^\alpha(E) = HF^\alpha(\overline{E})$ for all $\alpha \geq 0$ and all $0 < \delta \leq \infty$.*

This follows easily from the remarks preceding the statement of the lemma.

Lemma 2.17 *Let A be a subset of M , and let r be a positive real number. Define A_r by*

$$(2.18) \quad A_r = \bigcup \{B(a, r) : a \in A\}.$$

Then A_r is an open subset of M such that $A \subseteq A_r$ and $\text{diam } A_r \leq \text{diam } A + 2r$.

This is easy to see. As a consequence, if one restricts oneself to coverings by open subsets in Section 2.1, then the resulting values of the various measures of a set E are the same.

Lemma 2.19 *If E is a compact subset of M , then $H_{\text{con}}^\alpha(E) = HF_{\text{con}}^\alpha(E)$, $H_\delta^\alpha(E) = HF_\delta^\alpha(E)$, and $H^\alpha(E) = HF^\alpha(E)$ for all $\alpha \geq 0$ and $0 < \delta \leq \infty$.*

Indeed, one can use coverings of E by open sets, and then reduce to finite subcoverings by compactness.

Lemma 2.20 *Let E be a subset of M . For each $\alpha \geq 0$ there is a subset E_1 of M such that $E \subseteq E_1$, E_1 can be written as the intersection of a sequence of open subsets of M , and $H_{\text{con}}^\alpha(E_1) = H_{\text{con}}^\alpha(E)$. The analogous statements for H_δ^α and H^α also hold for all $\alpha \geq 0$ and $0 < \delta \leq \infty$.*

This is not difficult to show. Fix $\alpha > 0$, and consider $H_{con}^\alpha(E)$. If $H_{con}^\alpha(E) = \infty$, then we can simply take $E_1 = M$. Otherwise, for each positive integer n , let $\{A_i(n)\}_i$ be a covering of E by open subsets of M such that

$$(2.21) \quad \sum_i (\text{diam } A_i(n))^\alpha < H_{con}^\alpha(E) + \frac{1}{n}.$$

Put

$$(2.22) \quad E_1 = \bigcap_{n=1}^{\infty} \bigcup_i A_i(n).$$

Then $E \subseteq E_1$ and E_1 is the intersection of a sequence of open sets by construction, and it is not difficult to see that $H_{con}^\alpha(E_1) = H_{con}^\alpha(E)$. Indeed, $H_{con}^\alpha(E) \leq H_{con}^\alpha(E_1)$ because $E \subseteq E_1$, while $H_{con}^\alpha(E_1) \leq H_{con}^\alpha(E)$ because $\{A_i(n)\}_i$ is a covering of E_1 for all n . The argument for $H_\delta^\alpha(E)$ is quite similar, except that one should also ask that $\text{diam } A_i(n) < \delta$ for all i and n , while for $H^\alpha(E)$ one should ask that $\text{diam } A_i(n) < 1/n$ for all i and n .

2.3 Subadditivity properties

Let M be a set. A measure functional μ on subsets of M is said to be *finitely subadditive* if

$$(2.23) \quad \mu(E_1 \cup E_2) \leq \mu(E_1) + \mu(E_2)$$

for all subsets E_1, E_2 of M . If

$$(2.24) \quad \mu\left(\sum_j E_j\right) \leq \sum_j \mu(E_j)$$

for any at-most-countable family $\{E_j\}_j$ of subsets of M , then μ is said to be *countably subadditive*.

For the rest of this section, we assume that $(M, d(x, y))$ is a metric space.

Lemma 2.25 *For each $\alpha \geq 0$ and $0 < \delta \leq \infty$, HF_{con}^α , HF_δ^α , and HF^α are finitely subadditive, and H_{con}^α , H_δ^α , and H^α are countably subadditive.*

This is an easy consequence of the definitions. In particular, if E is a finite subset of M and $\alpha > 0$, then $HF^\alpha(E) = 0$, and if $E \subseteq M$ is at most countable and $\alpha > 0$ then $H^\alpha(E) = 0$.

If E is a bounded subset of \mathbf{R}^n , then it is not difficult to check that $H^\alpha(E) = HF^\alpha(E) = 0$ for all $\alpha > 0$. Using the countable subadditivity of H^α , it follows that $H^\alpha(\mathbf{R}^n) = 0$ for all $\alpha > n$.

Let us say that two subsets E_1, E_2 are η -separated, where η is a positive real number, if $d(x, y) \geq \eta$ for all x in E_1 and all y in E_2 .

Lemma 2.26 *Let E_1, E_2 be a pair of η -separated subsets of M . For all $\alpha \geq 0$ and $\delta \in (0, \eta]$ we have that*

$$(2.27) \quad HF_\delta^\alpha(E_1 \cup E_2) \geq HF_\delta^\alpha(E_1) + HF_\delta^\alpha(E_2)$$

and

$$(2.28) \quad H_\delta^\alpha(E_1 \cup E_2) \geq H_\delta^\alpha(E_1) + H_\delta^\alpha(E_2).$$

As a result,

$$(2.29) \quad HF^\alpha(E_1 \cup E_2) \geq HF^\alpha(E_1) + HF^\alpha(E_2)$$

and

$$(2.30) \quad H^\alpha(E_1 \cup E_2) \geq H^\alpha(E_1) + H^\alpha(E_2).$$

This is not too difficult to verify.

2.4 Dimensions

Let $(M, d(x, y))$ be a metric space.

Lemma 2.31 *Let α, β be nonnegative real numbers such that $\alpha \leq \beta$, and let E be a subset of M . For $0 < \delta < \infty$ we have that*

$$(2.32) \quad H_\delta^\beta(E) \leq \delta^{\beta-\alpha} H_\delta^\alpha(E)$$

and

$$(2.33) \quad HF_\delta^\beta(E) \leq \delta^{\beta-\alpha} HF_\delta^\alpha(E).$$

In particular, $H_\delta^\beta(E) \leq H_\delta^\alpha(E)$ and $HF_\delta^\beta(E) \leq HF_\delta^\alpha(E)$ when $0 < \delta \leq 1$. As a result, $H^\beta(E) \leq H^\alpha(E)$, $HF^\beta(E) \leq HF^\alpha(E)$, $H_{con}^\alpha(E) = 0$ implies that $H_{con}^\beta(E) = 0$, and $HF_{con}^\alpha(E) = 0$ implies that $HF_{con}^\beta(E) = 0$. Furthermore, if $\alpha < \beta$, then $H^\alpha(E) < \infty$ implies that $H^\beta(E) = 0$, and $HF^\alpha(E) < \infty$ implies that $HF^\beta(E) = 0$.

This is easy to check, using the fact that $r^\beta \leq \delta^{\beta-\alpha} r^\alpha$ when $0 \leq r < \delta$.

The *Hausdorff dimension*, or *H-dimension*, of a subset E of M is denoted $\dim_H(E)$ and defined to be the infimum of the set

$$(2.34) \quad \{\alpha \geq 0 : H^\alpha(E) = 0\}$$

when this set is nonempty, and $+\infty$ otherwise. Similarly, the *HF-dimension* of E is denoted $\dim_{HF}(E)$ and defined to be the infimum of the set

$$(2.35) \quad \{\alpha \geq 0 : HF^\alpha(E) = 0\}$$

when this set is nonempty, and $+\infty$ otherwise. As a variant of the *HF-dimension*, the *HF*-dimension* of E is denoted $\dim_{HF^*}(E)$ and defined to be the supremum of the *HF* dimensions of $E \cap B(p, n)$ over all positive integers n , where p is any fixed point in M . One can easily check that this does not depend on the choice of p . Of course the *HF*-dimension* of E is automatically equal to the *HF-dimension* of E when E is bounded. If E is unbounded, then the *HF-dimension* of E is $+\infty$, while the *HF*-dimension* of E can be finite. Note that the empty set has dimension 0 in each of these senses.

Lemma 2.36 *If E_1, E_2 are subsets of M such that $E_1 \subseteq E_2$, then the Hausdorff dimension of E_1 is less than or equal to the Hausdorff dimension of E_2 , the HF-dimension of E_1 is less than or equal to the HF-dimension of E_2 , and the HF*-dimension of E_1 is less than or equal to the HF*-dimension of E_2 .*

This is an easy consequence of the definitions.

Lemma 2.37 *If E is a compact subset of M , then the Hausdorff and HF-dimensions of E are equal.*

This follows from the fact that the Hausdorff and *HF*-measures are equal for compact sets.

Lemma 2.38 *If E is a closed subset of M such that closed and bounded subsets of E are compact, then the Hausdorff and HF* dimensions of E are equal.*

This is not difficult to verify.

Lemma 2.39 *Let E_1, E_2 be two subsets of M . The HF -dimension of $E_1 \cup E_2$ is equal to maximum of the HF -dimensions of E_1 and E_2 , and the HF^* -dimension of $E_1 \cup E_2$ is equal to the maximum of the HF^* dimensions of E_1 and E_2 .*

Indeed, one equality can be obtained from Lemma 2.36, while the other uses the subadditivity of the HF measures.

Lemma 2.40 *Let $\{E_j\}_{j=1}^\infty$ be a sequence of subsets of M . The Hausdorff dimension of $\bigcup_{j=1}^\infty E_j$ is equal to the supremum of the Hausdorff dimensions of the E_j 's, $j \in \mathbf{Z}_+$.*

Again, one inequality follows from Lemma 2.36, and the other can be derived from countable subadditivity of the Hausdorff measures.

2.5 Snowflake transforms

Lemma 2.41 *Let a be a real number such that $0 < a \leq 1$. If r_1, r_2 are nonnegative real numbers, then $(r_1 + r_2)^a \leq r_1^a + r_2^a$.*

To see this, observe first that

$$(2.42) \quad r_1 + r_2 \leq \max(r_1, r_2)^{1-a} (r_1^a + r_2^a).$$

Also, $\max(r_1, r_2)^a \leq r_1^a + r_2^a$, and therefore

$$(2.43) \quad r_1 + r_2 \leq (r_1^a + r_2^a)^{((1-a)/a)+1} = (r_1^a + r_2^a)^{1/a}.$$

This implies the inequality stated in the lemma.

Let $(M, d(x, y))$ be a metric space, and let a be a real number such that $0 < a \leq 1$. It is not hard to check that $(M, d(x, y)^a)$ is then also a metric space, using the lemma for the triangle inequality. We call $(M, d(x, y)^a)$ the *snowflake transform of order a* of the metric space $(M, d(x, y))$. Observe that the metrics $d(x, y)$ and $d(x, y)^a$ determine the same topology on M , which is the same as saying that the identity mapping on M is a homeomorphism as a map between the metric spaces $(M, d(x, y))$ and $(M, d(x, y)^a)$. Also, a subset of M is bounded with respect to $d(x, y)$ if and only if it is bounded with respect to $d(x, y)^a$.

If A is a subset of M , then the diameter of A with respect to $d(x, y)^a$ is the same as the diameter of A with respect to $d(x, y)$ to the a th power. For each nonnegative real number α , one can use this to check that $H_{con}^\alpha(A)$ and $HF_{con}^\alpha(A)$ with respect to $d(x, y)$ are the same as $H_{con}^{\alpha/a}(A)$ and $HF_{con}^{\alpha/a}(A)$ with respect to $d(x, y)^a$. Similarly, for each $\delta > 0$, $H_\delta^\alpha(A)$ and $HF_\delta^\alpha(A)$ with respect to $d(x, y)$ are the same as $H_{\delta^a}^{\alpha/a}(A)$ and $HF_{\delta^a}^{\alpha/a}(A)$ with respect to $d(x, y)^a$. Taking the supremum over δ , we obtain that $H^\alpha(A)$ and $HF^\alpha(A)$ with respect to $d(x, y)$ is the same as $H^{\alpha/a}(A)$ and $HF^{\alpha/a}(A)$ with respect to $d(x, y)^a$. The Hausdorff, HF , and HF^* -dimensions of A with respect to $d(x, y)$ are equal to the Hausdorff, HF , and HF^* -dimensions of A with respect to $d(x, y)^a$ divided by a .

Chapter 3

More on metric spaces

3.1 Lipschitz mappings

Throughout this section we let $(M, d(x, y))$ and $(N, \rho(u, v))$ be metric spaces. A mapping $f : M \rightarrow N$ is said to be *Lipschitz* if there is a nonnegative real number C such that

$$(3.1) \quad \rho(f(x), f(y)) \leq C d(x, y)$$

for all $x, y \in M$. We also say that f is C -Lipschitz in this case, to mention the constant C explicitly. Lipschitz mappings are clearly continuous, and even uniformly continuous.

The space of Lipschitz mappings from M to N is denoted $\text{Lip}(M, N)$, and the space of C -Lipschitz mappings from M to N is denoted $\text{Lip}_C(M, N)$. We write $\text{Lip}(M)$ for the space of Lipschitz real-valued functions on M , and $\text{Lip}_C(M)$ for the space of C -Lipschitz real-valued functions on M .

In the case of real-valued functions on Euclidean spaces which are continuously differentiable, the Lipschitz condition is equivalent to the norm of the gradient being bounded, and the supremum of the norm of the gradient is equal to the smallest choice of the constant in the Lipschitz condition. As a special case of this, if v is a vector in \mathbf{R}^n , then the linear function on \mathbf{R}^n which maps a point x in \mathbf{R}^n to the inner product of x with v is Lipschitz with constant equal to the norm of v . The orthogonal projection of \mathbf{R}^n onto any linear subspace is Lipschitz.

The sum of two real-valued Lipschitz functions on M is a Lipschitz function, as is the product of a real-valued Lipschitz function and a constant. In general, the product of two real-valued Lipschitz functions is not necessarily

Lipschitz, but this is the case if at least one of the two functions is bounded. If $f(x)$ is a real-valued Lipschitz function on M and there is a positive real number c such that $|f(x)| \geq c$ for all x in M , then $1/f(x)$ is also a Lipschitz function on M .

Lemma 3.2 *If p is an element of M , then $d(x, p)$ is 1-Lipschitz as a real-valued function of x on M . If A is a nonempty subset of M , then $\text{dist}(X, A)$, as defined in (1.32) in Section 1.4, is a 1-Lipschitz function on M .*

This follows from the proofs of Lemmas 1.28 and 1.34 in Section 1.4.

Let us say that a mapping $f : M \rightarrow N$ is *locally Lipschitz* if for every $x \in M$ there is an $r > 0$ so that the restriction of f to $B(x, r)$ is Lipschitz. One can check that the sum and product of real-valued locally Lipschitz functions on M are also locally Lipschitz, and that if $f(x)$ is a real-valued locally Lipschitz function on M such that $f(x) \neq 0$ for all x in M , then $1/f(x)$ is a locally Lipschitz function on M .

A mapping $f : M \rightarrow N$ is said to be *countably Lipschitz* if M can be expressed as the union of at most countably-many subsets on which f is Lipschitz. If $f : M \rightarrow N$ is locally Lipschitz and M is separable, then f is countably Lipschitz. This follows from the countable compactness of M .

If $f : M \rightarrow N$ is countably Lipschitz, then M can be expressed as the union of at most countably-many subsets on which f is both Lipschitz and bounded. This is easy to check, by simply decomposing further the subsets of M on which f is already Lipschitz. Furthermore, the sum and product of real-valued countably Lipschitz functions is countably Lipschitz, as is the reciprocal of a nonzero real-valued countably Lipschitz function.

3.1.1 Diameters, measure functionals, and dimensions

Let us write $\text{diam}_M A$ and $\text{diam}_N B$ for the diameters of subsets A and B of M and N , respectively.

Lemma 3.3 *If A is a subset of M and $f : M \rightarrow N$ is a C -Lipschitz mapping, then $\text{diam}_N f(A) \leq C \text{diam}_M A$.*

This is easy to check from the definitions.

Proposition 3.4 *Let E be a subset of M , α a nonnegative real number, and $f : M \rightarrow N$ a C -Lipschitz mapping. Under these conditions, we have that*

$H_{con}^\alpha(f(E)) \leq C^\alpha H_{con}^\alpha(E)$ and $HF_{con}^\alpha(f(E)) \leq C^\alpha HF_{con}^\alpha(E)$. Similarly, for each δ , $0 < \delta \leq \infty$, $H_{C^\delta}^\alpha(f(E)) \leq C^\alpha H_\delta^\alpha(f(E))$ and $HF_{C^\delta}^\alpha(f(E)) \leq C^\alpha HF_\delta^\alpha(f(E))$. Taking the supremum over δ it follows that $H^\alpha(f(E)) \leq C^\alpha H^\alpha(E)$ and $HF^\alpha(f(E)) \leq C^\alpha HF^\alpha(E)$. In particular, $\dim_H(f(E)) \leq \dim_H(E)$, $\dim_{HF}(f(E)) \leq \dim_{HF}(E)$, and $\dim_{HF^*}(f(E)) \leq \dim_{HF^*}(E)$.

Note that the various measure functionals and dimensions of $f(E)$ are for subsets of N , using the metric $\rho(u, v)$, while the various measure functionals and dimensions for E are for subsets of M , using the metric $d(x, y)$.

The proposition can be shown in a straightforward manner using Lemma 3.3 and the various definitions. When $\alpha = 0$, one can take $C^\alpha = 1$, and the conditions on f can be weakened. For instance, the inequalities $H^0(f(E)) \leq H^0(E)$ and $HF^0(f(E)) \leq HF^0(E)$ just say that the number of elements of $f(E)$ is less than or equal to the number of elements of E , and this is true for any mapping $f : M \rightarrow N$.

Corollary 3.5 *Suppose that $f : M \rightarrow N$ is countably-Lipschitz. If E is a subset of M and α is a nonnegative real number such that $H_{con}^\alpha(E) = 0$, then $H_{con}^\alpha(f(E)) = 0$. As a consequence, $\dim_H(f(E)) \leq \dim_H(E)$.*

3.1.2 Some approximations and extensions

Proposition 3.6 *Let $f(x)$ be a bounded uniformly continuous real-valued function on M . There exists a sequence $\{f_j\}_{j=1}^\infty$ of real-valued Lipschitz functions on M which converges to f uniformly.*

For this one can take

$$(3.7) \quad f_j(x) = \inf\{f(y) + j d(x, y) : y \in M\}.$$

By definition, $f_j(x) \leq f(x)$ for all x in M and $j \geq 1$, and it is not difficult to use the boundedness of f to show that the infimum is always finite. More precisely, if a is a real number such that $f(x) \geq a$ for all x in M , then $f_j(x) \geq a$ for all x in M and $j \geq 1$.

If x, x' , and y are arbitrary elements of M , then

$$(3.8) \quad f_j(x) \leq f(y) + j d(x, y) \leq f(y) + j d(x', y) + j d(x, x').$$

As a result,

$$(3.9) \quad f_j(x) \leq f_j(x') + j d(x, x').$$

From this it follows that f_j is j -Lipschitz. Let us also point out that $f_j(x) = f(x)$ for all x in M when f is j -Lipschitz.

Using the boundedness of f , it is not difficult to show that in fact one only needs to consider $y \in M$ with $d(x, y)$ bounded by a constant times $1/j$ in the infimum in the definition of $f_j(x)$. One can then check that $f_j(x)$ converges to $f(x)$ uniformly when f is uniformly continuous. We leave the details as an exercise.

Proposition 3.10 *Let E be a nonempty subset of M , let C be a nonnegative real number, and let h be a real-valued C -Lipschitz function on E . There exists a C -Lipschitz real-valued function \hat{h} on M such that $\hat{h}(x) = h(x)$ when $x \in E$.*

In this case we set

$$(3.11) \quad \hat{h}(x) = \inf\{h(y) + C d(x, y) : y \in M\}.$$

Thus

$$(3.12) \quad \hat{h}(x) \leq h(z) + C d(x, z)$$

for all x in M and z in E by definition. Next, let us check that

$$(3.13) \quad \hat{h}(x) \geq h(w) - C d(x, w)$$

for all $x \in M$ and $w \in E$, and in particular that the infimum in the definition of $\hat{h}(x)$ is finite. For this it is enough to show that

$$(3.14) \quad h(w) - C d(x, w) \leq h(y) + C d(x, y)$$

for all y in E , which is the same as

$$(3.15) \quad h(w) \leq h(y) + C d(x, w) + C d(x, y).$$

This holds because

$$(3.16) \quad h(w) \leq h(y) + C d(w, y) \leq h(y) + C d(x, w) + C d(x, y),$$

by the hypothesis that h is C -Lipschitz on E and the triangle inequality.

It follows in particular from (3.12) and (3.13) that $\hat{h}(x) = h(x)$ when $x \in E$. If x, x' are arbitrary elements of M and y is any element of E , then

$$(3.17) \quad \hat{h}(x) \leq h(y) + C d(x, y) \leq h(y) + C d(x', y) + C d(x, x'),$$

and hence

$$(3.18) \quad \hat{h}(x) \leq \hat{h}(x') + C d(x, x').$$

This shows that \hat{h} is C -Lipschitz, and the lemma follows.

3.1.3 Spaces of Lipschitz mappings

For the rest of the section we assume that M is a bounded metric space, and that C is a fixed nonnegative real number. Each Lipschitz function on M is then bounded, and we can consider $\text{Lip}_C(M, N)$ as a subset of the space $C_b(M, N)$ of bounded continuous mappings from M into N . As in Section 1.4, we view this space as a metric space equipped with the metric $\theta(f_1, f_2) = \sup\{\rho(f_1(x), f_2(x)) : x \in M\}$. Recall that $C_b(M, N)$ is a complete metric space with respect to this metric when N is complete. Note that $C_b(M, N)$ is bounded if N is bounded.

Lemma 3.19 *As a subset of $C_b(M, N)$, $\text{Lip}_C(M, N)$ is closed.*

This is easy to verify from the definitions.

Lemma 3.20 *Suppose that M is totally bounded and that N is complete. If E is a dense subset of M and $\{f_j\}_{j=1}^\infty$ is a sequence of mappings in $\text{Lip}_C(M, N)$ which converges pointwise on E , which is to say that $\{f_j(x)\}_{j=1}^\infty$ is a convergent sequence in N for each x in E , then $\{f_j\}_{j=1}^\infty$ converges in $C_b(M, N)$ to an element of $\text{Lip}_C(M, N)$.*

Let us write $f(x)$ for the limit of $\{f_j(x)\}_{j=1}^\infty$ in N when $x \in E$. It is not difficult to show that f is C -Lipschitz as a mapping from E into N . As a result, $f(x)$ extends to a C -Lipschitz mapping from M to N . In fact, one can use the C -Lipschitzness of each f_j to show that $\{f_j(x)\}_{j=1}^\infty$ is a Cauchy sequence in N for each x in M , since this is true for x in E by assumption, and one can define $f(x)$ to be the limit of this sequence for each x in M . Using the total boundedness of M one can show that $\{f_j\}_{j=1}^\infty$ converges to f uniformly, and the lemma follows.

Proposition 3.21 *Suppose that M is totally bounded and N is compact. As a subset of $C_b(M, N)$, $\text{Lip}_C(M, N)$ is then compact.*

This is a version of the famous Arzela–Ascoli theorem. Let us mention two approaches to it. For the first approach, since $C_b(M, N)$ is complete, it suffices to show that $\text{Lip}_C(M, N)$ is closed and totally bounded in $C_b(M, N)$. The former works in general, as in Lemma 3.19, and the latter can be verified using the total boundedness of M and N . For the second approach, it is enough to show that every sequence in $\text{Lip}_C(M, N)$ has a convergent subsequence. As in Lemma 3.20, it suffices to find a subsequence which converges

pointwise on a dense subset. Because M is totally bounded, it has a dense subset which is at most countable, and one can use the compactness of N to find subsequences which converge pointwise on any subset of M which is at most countable.

3.2 The Hausdorff metric

Let $(M, d(x, y))$ be a metric space. As in Lemma 2.17, if A is a subset of M and r is a positive real number, then

$$(3.22) \quad A_r = \bigcup \{B(a, r) : a \in A\}.$$

is an open subset of M that contains A and has diameter $\leq \text{diam } A + 2r$.

Let α be a nonnegative real number, and let E be a subset of M such that $H_{\text{con}}^\alpha(E) < \infty$. For every $\epsilon > 0$ there is an $\eta > 0$ so that

$$(3.23) \quad HF_{\text{con}}^\alpha(E_\eta) < HF_{\text{con}}^\alpha(E) + \epsilon.$$

More generally, if $0 < \delta \leq \infty$, $H_\delta^\alpha(E) < \infty$, and $\epsilon > 0$, then there is an $\eta > 0$ so that

$$(3.24) \quad HF_\delta^\alpha(E_\eta) < HF_\delta^\alpha(E) + \epsilon.$$

That is, any covering of E by finitely many subsets of M can be enlarged slightly to get a covering of E_η , and one can use this estimate $HF_{\text{con}}^\alpha(E_\eta)$, $HF_\delta^\alpha(E_\eta)$. This type of argument does not work in general for HF^α itself, as one can see by considering the case where E contains a single point.

If E_1, E_2 are nonempty subsets of M and t is a positive real number, then we say that E_1 and E_2 are t -close if

$$(3.25) \quad \text{for every } x \in E_1 \text{ there is a } y \in E_2 \text{ such that } d(x, y) < t$$

and

$$(3.26) \quad \text{for every } y \in E_2 \text{ there is an } x \in E_1 \text{ such that } d(x, y) < t.$$

This is equivalent to saying that

$$(3.27) \quad E_1 \subseteq (E_2)_t, \quad E_2 \subseteq (E_1)_t.$$

Notice that if E_1, E_2 are bounded nonempty subsets of M which are t -close for some $t > 0$, then

$$(3.28) \quad |\text{diam } E_1 - \text{diam } E_2| \leq 2t.$$

Lemma 3.29 *Suppose that E_1, E_2 , and E_3 are nonempty subsets of M and t, u are positive real numbers such that E_1, E_2 are t -close and E_2, E_3 are u -close. Then E_1, E_3 are $(t + u)$ -close.*

This is easy to verify.

Lemma 3.30 *A subset A of M is totally bounded if and only if for every $t > 0$, A is t -close to a finite subset of M . Moreover, M is separable if and only if for every $t > 0$, M is t -close to a subset of M which is at most countable.*

This is a straightforward consequence of the definitions.

3.2.1 The space of closed and bounded subsets of M

Let us write $\mathcal{S}(M)$ for the collection of all nonempty closed and bounded subsets of M . If E_1, E_2 are two elements of $\mathcal{S}(M)$, define the *Hausdorff distance* $D(E_1, E_2)$ between E_1 and E_2 by

$$(3.31) \quad D(E_1, E_2) = \inf\{t > 0 : E_1, E_2 \text{ are } t\text{-close}\}.$$

The boundedness of E_1 and E_2 ensures that the set of $t > 0$ such that E_1, E_2 are t -close is nonempty. The restriction to closed sets implies that $D(E_1, E_2) = 0$ if and only if $E_1 = E_2$.

Lemma 3.32 *The Hausdorff distance $D(E_1, E_2)$ defines a metric on $\mathcal{S}(M)$.*

This is easy to check, using Lemma 3.29 to show that the Hausdorff distance satisfies the triangle inequality.

Lemma 3.33 *If A is a totally bounded subset of M , then $\mathcal{A} = \{E \in \mathcal{S}(M) : E \subseteq A\}$ is a totally bounded subset of $\mathcal{S}(M)$. Hence if M is totally bounded, then $\mathcal{S}(M)$ is totally bounded, and separable in particular.*

The main point for this lemma is that total boundedness of A implies that A can be approximated by finite sets, and as a result subsets of A can be approximated by finite subsets of a fixed finite set for each fixed degree of approximation.

Lemma 3.34 *Suppose that $\{E_j\}_{j=1}^\infty$ is a sequence of nonempty closed and bounded subsets of M which converges to a nonempty closed and bounded subset E of M in the Hausdorff metric. If each E_j is totally bounded, then so is E .*

This can be derived fairly directly from the definitions.

Lemma 3.35 *Suppose that $\{E_j\}_{j=1}^\infty$ is a sequence of nonempty closed and bounded subsets of M which converges to a nonempty closed and bounded subset E of M . For each x in E , there is a sequence $\{x_j\}_{j=1}^\infty$ of points in M such that $x_j \in E_j$ for all j and $\lim_{j \rightarrow \infty} x_j = x$. Conversely, if $\{j_l\}_{l=1}^\infty$ is a strictly increasing sequence of positive integers, $x_{j_l} \in E_{j_l}$ for each l , and $\{x_{j_l}\}_{l=1}^\infty$ converges in M to a point x , then $x \in E$.*

This characterization of limits of sequences in $\mathcal{S}(M)$ is a straightforward consequence of the definitions.

3.2.2 Subsets, functions, and embeddings

There is a natural embedding of M into $\mathcal{S}(M)$, in which a point p in M is associated to the subset $\{p\}$ of M . This is an isometric embedding, which is to say that

$$(3.36) \quad D(\{p\}, \{q\}) = d(p, q)$$

for all p, q in M .

If E is a nonempty subset of M , let us write $f_E(x)$ for the function $\text{dist}(x, E)$. Thus $f_E(x)$ is a continuous real-valued function on M which is 1-Lipschitz. We may as well restrict our attention to nonempty closed subsets E of M here, since $f_E(x) = f_{\overline{E}}(x)$ for all x in M .

Assume for the moment that M is bounded. Then the mapping $E \mapsto f_E$ defines a mapping from $\mathcal{S}(M)$ into the space $C_b(M)$ of bounded continuous real-valued functions on M . In fact this is an isometric embedding of $\mathcal{S}(M)$ into $C_b(M)$, with respect to the supremum distance on $C_b(M)$. Explicitly, if E_1, E_2 are elements of $\mathcal{S}(M)$, then

$$(3.37) \quad D(E_1, E_2) = \sup\{|f_{E_1}(x) - f_{E_2}(x)| : x \in M\}.$$

This is not too difficult to verify.

If M is not bounded, then we can use a basepoint to get an embedding of $\mathcal{S}(M)$ into $C_b(M)$. That is, we fix an element w of M , and associate to each

nonempty subset E of M the continuous real-valued function $f_E(x) - f_{\{w\}}(x)$ on M . It is easy to see that this is a bounded function on M when E is a bounded nonempty subset of M . One can also check that this leads to an isometric embedding of $\mathcal{S}(M)$ into $C_b(M)$, whether or not M is bounded.

3.2.3 Compactness

Lemma 3.38 *Suppose that $\{K_j\}_{j=1}^\infty$ is a sequence of nonempty compact subsets of M such that $K_{j+1} \subseteq K_j$ for all $j \geq 1$, and put $K = \bigcap_{j=1}^\infty K_j$. Under these conditions, $\{K_j\}_{j=1}^\infty$ converges to K with respect to the Hausdorff metric.*

Note that K is a nonempty compact subset of M in this case, by a well-known result. To prove the lemma, we use essentially the same type of argument as for that result. It suffices to show that for each $\epsilon > 0$ there is a positive integer N such that $K_j \subseteq K(\epsilon)$ for all $j \geq N$, where $K(\epsilon)$ denotes the set of $x \in M$ such that $\text{dist}(x, K) < \epsilon$. This is the same as K_ϵ in the sense of (3.22), but we write $K(\epsilon)$ now to avoid confusion with K_j . Let $\epsilon > 0$ be given. Because $K(\epsilon)$ is an open subset of M , and the union of $K(\epsilon)$ with the complements of the K_j 's is equal to all of M , it follows from the compactness of K_1 that a finite collection of these open sets are enough to cover K_1 . The monotonicity property of the K_j 's implies more precisely that K_1 is contained in $(M \setminus K_N) \cup K(\epsilon)$ for some positive integer N . Hence $K_j \subseteq K(\epsilon)$ when $j \geq N$, as desired.

Lemma 3.39 *If M is compact, then $\mathcal{S}(M)$ is complete as a metric space with respect to the Hausdorff metric.*

In this case $\mathcal{S}(M)$ consists simply of the nonempty compact subsets of M . Let $\{E_j\}_{j=1}^\infty$ be a Cauchy sequence in $\mathcal{S}(M)$, which we would like to show converges in $\mathcal{S}(M)$. By passing to a subsequence we may assume that

$$(3.40) \quad D(E_j, E_{j+1}) < 2^{-j}$$

for all $j \geq 1$. This uses the fact that a Cauchy sequence converges if any of its subsequences converge.

For each j , set

$$(3.41) \quad \tilde{E}_j = \{x \in M : \text{dist}(x, E_j) \leq 2^{-j}\}.$$

Each \tilde{E}_j is a closed subset of M , and hence compact. Also, $D(\tilde{E}_j, E_j) \leq 2^{-j}$, and so if $\{\tilde{E}_j\}_{j=1}^\infty$ converges, then so does $\{E_j\}_{j=1}^\infty$, and they have the same limit. Thus it suffices to show that $\{\tilde{E}_j\}_{j=1}^\infty$ converges.

A key point now is that

$$(3.42) \quad \tilde{E}_{j+1} \subseteq \tilde{E}_j$$

for all j . This follows from (3.40) and the definition of \tilde{E}_j . Therefore $\{\tilde{E}_j\}_{j=1}^\infty$ converges in $\mathcal{S}(M)$ to $\bigcap_{j=1}^\infty \tilde{E}_j$, as in Lemma 3.38.

Proposition 3.43 *If M is compact, then $\mathcal{S}(M)$ is compact, with respect to the Hausdorff metric.*

This follows from Lemmas 3.33 and 3.39, which imply that $\mathcal{S}(M)$ is totally bounded and complete when M is compact.

3.2.4 Connectedness

Lemma 3.44 *Let $\{A_j\}_{j=1}^\infty$ be a sequence of nonempty closed and bounded subsets of M which converges in the Hausdorff metric to a nonempty closed and bounded subset A of M . If M is compact and each A_j is connected, then A is connected.*

Assume for the sake of a contradiction that A is not connected. This implies that A can be expressed as the union of two nonempty disjoint closed subsets A^1, A^2 . Because of compactness, there is an $\epsilon > 0$ so that $d(x, y) \geq \epsilon$ for all $x \in A^1$ and $y \in A^2$. The convergence of $\{A_j\}_{j=1}^\infty$ to A in the Hausdorff metric would then imply that A_j admits a similar decomposition for j large enough, in contradiction to hypothesis. This proves the lemma.

Chapter 4

Examples and special cases

4.1 The real line

If A is any subset of the real line, then there is an interval J contained in the real line such that $A \subseteq J$ and $\text{diam } J = \text{diam } A$. Here one should allow unbounded intervals J to deal with the case where A is unbounded, and it is convenient to consider the empty set as a kind of degenerate interval, to be compatible with some of our conventions. As a result of this simple observation, it follows that for subsets of the real line, if one defines the measure functionals H_{con}^α , H_δ^α , H^α , HF_{con}^α , HF_δ^α , and HF^α in the same manner as in Section 2.1, except for using coverings by families of intervals, then one gets the same answers as for the earlier definitions using coverings by families of general sets.

Now let us specialize to the case of $\alpha = 1$.

Lemma 4.1 *Let E be a subset of the real line. For $0 < \delta \leq \infty$ we have that $H_{con}^1(E) = H_\delta^1(E) = H^1(E)$ and $HF_{con}^1(E) = HF_\delta^1(E) = HF^1(E)$.*

Indeed, if A is any subset of the real line, and $\delta > 0$, then there is a covering of A by intervals with diameter less than δ such that the sum of their lengths is equal to the diameter of A . If A is bounded, then a finite collection of intervals can be used in the covering. This permits one to replace a given covering of E by sets of smaller diameter, while keeping the sums of the diameters of the sets in the covering fixed.

Lemma 4.2 *If $J = [a, b]$ is a closed and bounded interval in the real line, then $H^1(J)$ and $HF^1(J)$ are both equal to the length $|b - a|$ of the interval.*

We already know from Lemma 2.19 that $H^1(J) = HF^1(J)$, since J is compact. It is easy to see that $HF^1(J) \leq |b - a|$, through explicit coverings of J . This follows from (2.9) and Lemma 4.1 as well. The remaining point is that $HF^1(J) \geq |b - a|$, and we leave this as an exercise.

Lemma 4.3 *If E is a bounded subset of \mathbf{R} and $\alpha > 1$, then $H^\alpha(E) = HF^\alpha(E) = 0$. For any subset E of \mathbf{R} and $\alpha > 1$, $H^\alpha(E) = 0$.*

The first statement can be verified using simple choices of coverings of E , while the second follows from countable subadditivity of H^α .

As a consequence of these two lemmas, \mathbf{R} has Hausdorff and HF^* -dimensions equal to 1, and the Hausdorff, HF , and HF^* -dimensions of a closed and bounded interval $J = [a, b]$ with $a < b$ are equal to 1.

Let us now explain how a subset of the real line might be equal to the intersection of a sequence of dense open sets, and also have Hausdorff dimension 0. For each positive integer n , let $\{r_j(n)\}_{j=1}^\infty$ be a sequence of positive real numbers such that $r_j(n) < 1/(2n)$ for all n and j , and

$$(4.4) \quad \sum_{j=1}^\infty (2r_j)^{1/n} < \frac{1}{n}.$$

For instance, one can take $r_j(n) = (j + m_n)^{-2n}$ for sufficiently-large choices of m_n . Next, let $\{x_j\}_{j=1}^\infty$ be an enumeration of the rational numbers. For each n , put

$$(4.5) \quad U_n = \bigcup_{j=1}^\infty (x_j - r_j(n), x_j + r_j(n)).$$

Thus U_n is a dense open subset of \mathbf{R} , and

$$(4.6) \quad H_{1/n}^{1/n}(U_n) \leq \sum_{j=1}^\infty (2r_j)^{1/n} < \frac{1}{n}.$$

If $E = \bigcap_{n=1}^\infty U_n$, then E is the intersection of a sequence of dense open subsets of the real line, and one can check that $H^\alpha(E) = 0$ for all $\alpha > 0$.

4.2 \mathbf{R}^n for general n

Fix a positive integer n . Recall that a subset E of \mathbf{R}^n is said to be *convex* if for every pair of points v, w in E and every real number t with $0 \leq t \leq 1$ we

have that $tv + (1 - t)w$ also lies in E . For example, open and closed balls in \mathbf{R}^n are convex sets.

If l is a positive integer and y_1, \dots, y_l are points in \mathbf{R}^n , then a *convex combination* is a point in \mathbf{R}^n of the form

$$(4.7) \quad \sum_{j=1}^l \lambda_j y_j,$$

where $\lambda_1, \dots, \lambda_l$ are nonnegative real numbers such that

$$(4.8) \quad \sum_{j=1}^l \lambda_j = 1.$$

If A is a subset of \mathbf{R}^n , then the *convex hull* is the set consisting of all points in \mathbf{R}^n which are convex combinations of points in A . The convex hull of A is the smallest convex subset of \mathbf{R}^n which contains A .

The closure of a convex set is also convex, as one can easily verify. The closure of the convex hull of a subset A of \mathbf{R}^n can be described as the smallest closed convex subset of \mathbf{R}^n that contains A . Let us mention the classical fact that the convex hull of a compact subset of \mathbf{R}^n is compact. This can be established by first showing that every element of the convex hull of a subset A of \mathbf{R}^n can be written as a convex combination of less than or equal to $n + 1$ elements of A .

If A is a subset of \mathbf{R}^n , then the diameter of the closure of the convex hull of A is equal to the diameter of A . This simple fact implies that for the measure functionals on \mathbf{R}^n described in Section 2.1, i.e., the various Hausdorff measures and contents and so on, one can restrict to coverings by closed convex sets and get the same values as before. This works as well in any normed vector space. When $n = 1$, this corresponds to the remark in the previous section that one can restrict to coverings by intervals and get the same values for the Hausdorff measures and contents, etc.

Observe that if A is a subset of \mathbf{R}^n , then the diameter of A is equal to the diameter of the image of A under a translation or orthogonal transformation on \mathbf{R}^n . As a result, the measure functionals described in Section 2.1 applied to a subset E of \mathbf{R}^n have the same value as when they are applied to the image of E under a translation or orthogonal transformation. A similar remark works in any metric space, concerning sets which are isometrically-equivalent; Euclidean spaces happen to have a lot of such symmetries.

Now let us specialize to the case where $\alpha = n$ for the measure functionals in Section 2.1.

Lemma 4.9 *For each positive integer n there is a positive real number $C(n)$ such that $HF^n(E) \leq HF_{con}^n(E)$ and $H^n(E) \leq H_{con}^n(E)$ for any subset E of \mathbf{R}^n .*

This follows from the geometric fact that if A is a bounded subset of \mathbf{R}^n and δ is a positive real number such that $\delta < \text{diam } A$, then A can be covered by $\leq C(n) \delta^{-n} (\text{diam } A)^n$ subsets of \mathbf{R}^n with diameter less than δ .

If K is a compact subset of \mathbf{R}^n which is the closure of a region with reasonably-nice boundary, then the classical way to measure the volume of K is by approximating K by a finite union of cubes of the same size. For reasonably-nice compact sets, approximations of K by unions of cubes contained in K and with disjoint interiors and approximations by unions of cubes which contain K as a subset give nearly the same estimate of the volume, and the volume of K is defined to be the limit of these approximations as the mesh size tends to 0.

Using this, it is not too difficult to show that the n -dimensional Hausdorff measure of K is equal to the classical volume of K times the n -dimensional Hausdorff measure of the unit cube $Q_0 = \{x \in \mathbf{R}^n : 0 \leq x_j \leq 1 \text{ for } j = 1, \dots, n\}$. One can also show that $0 < H^n(Q_0) < \infty$.

Bibliography

- [1] M. Adams and V. Guillemin, *Measure Theory and Probability*, Birkhäuser, 1996.
- [2] D. Adams and L. Hedberg, *Function Spaces and Potential Theory*, Springer-Verlag, 1996.
- [3] L. Ambrosio, *Some fine properties of sets of finite perimeter in Ahlfors regular metric spaces*, Advances in Mathematics **159** (2001), 51–67.
- [4] L. Ambrosio and B. Kirchheim, *Currents in metric spaces*, Acta Mathematica **185** (2000), 1–80.
- [5] L. Ambrosio and B. Kirchheim, *Rectifiable sets in metric and Banach spaces*, Mathematische Annalen **318** (2000), 527–555.
- [6] L. Ambrosio and P. Tilli, *Selected Topics on “Analysis on Metric Spaces”*, Scuola Normale Superiore, Pisa, 2000.
- [7] J. Anderson, *Hyperbolic Geometry*, Springer-Verlag, 1999.
- [8] P. Assouad, *Espaces Métriques, Plongements, Facteurs*, Thèse de Doctorat (January, 1977), Université de Paris XI, 91405 Orsay, France.
- [9] P. Assouad, *Étude d’une dimension métrique liée à la possibilité de plongement dans \mathbf{R}^n* , Comptes Rendus de l’Académie des Sciences Paris, Sér. A **288** (1979), 731–734.
- [10] P. Assouad, *Plongements Lipschitziens dans \mathbf{R}^n* , Bulletin de la Société Mathématique de France **111** (1983), 429–448.

- [11] A. Banyaga, H. Movahedi-Lankarani, and R. Wells, editors, *Topics in Low-Dimensional Topology: Proceedings of the Conference in Honor of Steve Armentrout*, World Scientific, 1999.
- [12] D. Bao, S.-S. Chern, and Z. Shen, *An Introduction to Riemann–Finsler Geometry*, Springer-Verlag, 2000.
- [13] M. Barnsley, *Fractals Everywhere*, second edition, Academic Press, 1993.
- [14] M. Barnsley, *Fractal image compression*, Notices of the American Mathematical Society **43** (1996), 657–662.
- [15] S. Bates, W. Johnson, J. Lindenstrauss, D. Preiss, and G. Schechtman, *Affine approximation of Lipschitz functions and nonlinear quotients*, Geometric and Functional Analysis **9** (1999), 1092–1127.
- [16] A. Beardon, *The Geometry of Discrete Groups*, Springer-Verlag, 1983.
- [17] A. Beardon, *Iteration of Rational Functions: Complex Analytic Dynamical Systems*, Springer-Verlag, 1991.
- [18] A. Bellaïche and J.-J. Risler, *Sub-Riemannian Geometry*, Birkhäuser, 1996.
- [19] Y. Benyamini and J. Lindenstrauss, *Geometric Nonlinear Functional Analysis*, Volume 1, Colloquium Publications **48**, American Mathematical Society, 2000.
- [20] R. Bing, *The Geometric Topology of 3-Manifolds*, Colloquium Publications **40**, American Mathematical Society, 1983.
- [21] C. Bishop and T. Steger, *Representation-theoretic rigidity in $\mathrm{PSL}(2, \mathbf{R})$* , Acta Mathematica **170** (1993), 121–149.
- [22] E. Bombieri, editor, *Seminar on Minimal Manifolds*, Annals of Mathematics Studies **103**, Princeton University Press, 1983.
- [23] M. Bonk, J. Heinonen, and P. Koskela, *Uniformizing Gromov Hyperbolic Spaces*, Astérisque **270**, 2001.

- [24] K. Borsuk, *Theory of Retracts*, Monografie Matematyczne **44**, PWN – Polish Scientific Publishers, 1967.
- [25] K. Borsuk, *Theory of Shape*, Monografi Matematyczne **59**, PWN – Polish Scientific Publishers, 1975.
- [26] R. Bott and L. Tu, *Differential Forms in Algebraic Topology*, Springer-Verlag, 1982.
- [27] G. Bredon, *Topology and Geometry*, Springer-Verlag, 1993.
- [28] J. Bryant, S. Ferry, W. Mio, and S. Weinberger, *Topology of homology manifolds*, Bulletin of the American Mathematical Society (N.S.) **28** (1993), 324–328.
- [29] J. Bryant, S. Ferry, W. Mio, and S. Ferry, *Topology of homology manifolds*, Annals of Mathematics (2) **143** (1996), 435–467.
- [30] C. Burgess and J. Cannon, *Embeddings of surfaces in \mathbf{E}^3* , Rocky Mountain Journal of Mathematics **1** (1971), 259–344.
- [31] H. Busemann, *Metric Methods in Finsler Spaces and in the Foundations of Geometry*, Annals of Mathematics Studies **8**, Princeton University Press, 1942.
- [32] J. Cannon, *The recognition problem: What is a topological manifold?*, Bulletin of the American Mathematical Society **84** (1978), 832–866.
- [33] J. Cannon, *The characterization of topological manifolds of dimension $n \geq 5$* , in *Proceedings of the International Congress of Mathematicians, Helsinki, 1978*, 449–454, 1980.
- [34] L. Carleson and T. Gamelin, *Complex Dynamics*, Springer-Verlag, 1993.
- [35] J. Cheeger, *Pinching theorems for a certain class of Riemannian manifolds*, American Journal of Mathematics **91** (1969), 807–834.
- [36] J. Cheeger, *Finiteness theorems for Riemannian manifolds*, American Journal of Mathematics **92** (1970), 61–74.

- [37] J. Cheeger, *Differentiability of Lipschitz functions on metric measure spaces*, Geometric and Functional Analysis **9** (1999), 428–517.
- [38] J. Cheeger and J. Kister, *Counting topological manifolds*, Topology **9** (1970), 149–151.
- [39] R. Coifman and G. Weiss, *Analyse Harmonique Non-Commutative sur certains Espaces Homogènes*, Lecture Notes in Mathematics **242**, Springer-Verlag, 1971.
- [40] R. Coifman and G. Weiss, *Extensions of Hardy spaces and their use in analysis*, Bulletin of the American Mathematical Society **83** (1977), 569–645.
- [41] M. Coornaert, *Mesures de Patterson–Sullivan sur le bord d’un espace hyperbolique au sens de Gromov*, Pacific Journal of Mathematics **159** (1993), 241–270.
- [42] R. Daverman, *Embeddings of $(n - 1)$ -spheres in Euclidean n -space*, Bulletin of the American Mathematical Society **84** (1978), 377–405.
- [43] R. Daverman, *Decompositions of Manifolds*, Academic Press, 1986.
- [44] R. Daverman, *The intimate connections among decomposition theory, embedding theory, and manifold structure theory*, in *Geometric Topology and Shape Theory*, Lecture Notes in Mathematics **1283**, 43–47, 1987.
- [45] R. Daverman and R. Sher, editors, *Handbook of Geometric Topology*, North-Holland, 2002.
- [46] G. David and S. Semmes, *Fractured Fractals and Broken Dreams: Self-Similar Geometry through Metric and Measure*, Oxford University Press, 1997.
- [47] G. David and S. Semmes, *Quasiminimal surfaces of codimension 1 and John domains*, Pacific Journal of Mathematics **183** (1998), 213–277.
- [48] G. David and S. Semmes, *Regular mappings between dimensions*, Publications Mathématiques **44** (2000), 369–417.

- [49] G. David and S. Semmes, *Uniform Rectifiability and Quasiminimizing Sets of Arbitrary Codimension*, Memoirs of the American Mathematical Society **687**, 2000.
- [50] A. Devinatz, *Advanced Calculus*, Holt, Rinehart, and Winston, 1968.
- [51] S. Donaldson and P. Kronheimer, *The Geometry of 4-Manifolds*, Oxford University Press, 1990.
- [52] S. Donaldson and D. Sullivan, *Quasiconformal 4-manifolds*, Acta Mathematica **163** (1989), 181–252.
- [53] J. Doob, *Classical Potential Theory and its Probabilistic Counterpart*, Springer-Verlag, 2001.
- [54] J. Duoandikoetxea, *Fourier Analysis*, translated and revised by D. Cruz-Uribe, SFO, American Mathematical Society, 2001.
- [55] P. Duren, J. Heinonen, B. Osgood, and B. Palka, editors, *Quasiconformal Mappings and Analysis: A Collection of Papers Honoring F.W. Gehring*, Springer-Verlag, 1998.
- [56] R. Edwards, *Dimension theory, I*, in *Geometric Topology: Proceedings of the Topology Conference Held at Park City, Utah, 1974*, 195–211, Lecture Notes in Mathematics **438**, Springer-Verlag, 1975.
- [57] R. Edwards, *The topology of manifolds and cell-like maps*, in *Proceedings of the International Congress of Mathematicians, Helsinki, 1978*, 111–127, 1980.
- [58] L. Evans and R. Gariepy, *Measure Theory and Fine Properties of Functions*, CRC Press, 1992.
- [59] K. Falconer, *The Geometry of Fractal Sets*, Cambridge University Press, 1986.
- [60] K. Falconer, *Fractal Geometry: Mathematical Foundations and Applications*, Wiley, 1990.
- [61] K. Falconer, *Techniques in Fractal Geometry*, Wiley, 1997.
- [62] H. Federer, *Geometric Measure Theory*, Springer-Verlag, 1969.

- [63] J. Fernández and M. Melián, *Rims of Cantor trees*, Illinois Journal of Mathematics **44** (2000), 329–348.
- [64] J. Fernández and M. Melián, *Escaping geodesics of Riemannian surfaces*, Acta Mathematica **187** (2001), 213–236.
- [65] J. Fernández and A. Nicolau, *Boundary behavior of inner functions and holomorphic mappings*, Mathematische Annalen **310** (1998), 423–445.
- [66] S. Ferry, *Topological finiteness theorems for manifolds in Gromov–Hausdorff space*, Duke Mathematical Journal **74** (1994), 95–106.
- [67] S. Ferry, *Notes for Polyhedral and TOP Topology*, preprint.
- [68] S. Ferry, *Controlled Topology and the Characterization of Topological Manifolds*, preprint.
- [69] B. Franchi, R. Serapioni, and F. Serra Cassano, *Sur les ensembles de périmètre fini dans le groupe de Heisenberg*, Comptes Rendus des Séances de l’Académie des Sciences Sér. I Math. **329** (1999), 183–188.
- [70] B. Franchi, R. Serapioni, and F. Serra Cassano, *Rectifiability and perimeter in the Heisenberg group*, Mathematische Annalen **321** (2001), 479–531.
- [71] D. Freed and K. Uhlenbeck, *Instantons and Four-Manifolds*, second edition, Mathematical Sciences Research Institute Publications **1**, Springer-Verlag, 1991.
- [72] M. Freedman, *The topology of four-dimensional manifolds*, Journal of Differential Geometry **17** (1982), 357–453.
- [73] M. Freedman and F. Luo, *Selected Applications of Geometry to Low-Dimensional Topology*, University Lecture Series **1**, American Mathematical Society, 1989.
- [74] M. Freedman and F. Quinn, *Topology of 4-Manifolds*, Princeton University Press, 1990.
- [75] J. Fu, *Curvature of singular spaces via the normal cycle*, in *Differential Geometry: Geometry in Mathematical Physics and Related Topics*, Los Angeles CA, 1990, 211–221, Proceedings of Symposia in Pure Mathematics **54**, Part 2, American Mathematical Society, 1993.

- [76] J. Fu, *Curvature measures and Chern classes of singular varieties*, Journal of Differential Geometry **39** (1994), 251–280.
- [77] J. Fu, *Curvature measures of subanalytic sets*, American Journal of Mathematics **116** (1994), 819–880.
- [78] J. García-Cuerva and J. Rubio de Francia, *Weighted Norm Inequalities and Related Topics*, North-Holland, 1985.
- [79] J. Garnett, *Bounded Analytic Functions*, Academic Press, 1981.
- [80] F. Gehring and J. Väisälä, *Hausdorff dimension and quasiconformal mappings*, Journal of the London Mathematical Society (2) **6** (1973), 504–512.
- [81] F. Gehring, *The Hausdorff measure of sets which link in Euclidean space*, in *Contributions to Analysis: A Collection of Papers Dedicated to Lipman Bers*, 159–167, Academic Press, 1974.
- [82] F. Gehring, *Lower dimensional absolute continuity properties of quasiconformal mappings*, Mathematical Proceedings of the Cambridge Philosophical Society **78** (1975), 81–93.
- [83] F. Gehring, *The L^p -integrability of the partial derivatives of a quasiconformal mapping*, Acta Mathematica **130** (1973), 265–277.
- [84] F. Gehring, *Some metric properties of quasi-conformal mappings*, in *Proceedings of the International Congress of Mathematicians, Vancouver, 1974*, Volume 2, 203–206, 1975.
- [85] M. Giaquinta, G. Modica, and J. Souček, *Cartesian Currents in the Calculus of Variations I: Cartesian Currents*, Springer-Verlag, 1998.
- [86] M. Giaquinta, G. Modica, and J. Souček, *Cartesian Currents in the Calculus of Variations II: Variational Integrals*, Springer-Verlag, 1998.
- [87] E. Giusti, *Minimal surfaces and functions of bounded variation*, Birkhäuser, 1984.
- [88] R. Goldberg, *Methods of Real Analysis*, second edition, Wiley, 1976.

- [89] R. Greene and P. Petersen, *Little topology, big volume*, Duke Mathematical Journal **67** (1992), 273–290.
- [90] M. Gromov et al., *Structures Métriques pour les Variétés Riemanniennes*, Cedric / Fernand Nathan, 1981.
- [91] M. Gromov, *Quantitative homotopy theory*, in *Prospects in Mathematics (Princeton NJ, 1996)*, 45–49, American Mathematical Society, 1999.
- [92] M. Gromov et al., *Metric Structures for Riemannian and Non-Riemannian Spaces*, Birkhäuser, 1999.
- [93] M. Gromov, *Spaces and questions*, in *Visions in Mathematics, Proceedings of the meeting held at Tel Aviv University, 1999*, Geometric and Functional Analysis Special Volume, Part I (2000), 118–161.
- [94] M. Gromov and P. Pansu, *Rigidity of lattices: An introduction*, in *Geometric Topology: Recent Developments (Montecatini Terme, 1990)*, 39–137, Lecture Notes in Mathematics **1504**, Springer-Verlag, 1991.
- [95] M. de Guzmán, *Differentiation of Integrals in \mathbf{R}^n* , with appendices by A. Córdoba, R. Fefferman, and R. Moriyón, Lecture Notes in Mathematics **481**, Springer-Verlag, 1975.
- [96] M. de Guzmán, *Real Variable Methods in Fourier Analysis*, North-Holland, 1981.
- [97] R. Hardt and L. Simon, *Seminar on Geometric Measure Theory*, Birkhäuser, 1986.
- [98] F. Harvey and S. Semmes, *Zero divisors of atomic functions*, Annals of Mathematics (2) **135** (1992), 567–600.
- [99] A. Hatcher, *Algebraic Topology*, Cambridge University Press, 2002.
- [100] J. Heinonen, *Lectures on Analysis on Metric Spaces*, Springer-Verlag, 2001.
- [101] J. Heinonen and S. Rickman, *Quasiregular maps $\mathbf{S}^3 \rightarrow \mathbf{S}^3$ with wild branch sets*, Topology **37** (1998), 1–24.

- [102] J. Heinonen and S. Rickman, *Geometric branched covers between generalized manifolds*, Duke Mathematical Journal **113** (2002), 465–529.
- [103] J. Heinonen and D. Sullivan, *On the locally branched Euclidean metric guage*, Duke Mathematical Journal **114** (2002), 15–41.
- [104] E. Hewitt and K. Stromberg, *Real and Abstract Analysis*, third printing, Springer-Verlag, 1975.
- [105] J. Hocking and G. Young, *Topology*, second edition, Dover, 1988.
- [106] S.-T. Hu, *Theory of Retracts*, Wayne State University Press, 1965.
- [107] W. Hurewicz and H. Wallman, *Dimension Theory*, revised edition, 1948.
- [108] J. Hutchinson, *Fractals and self-similarity*, Indiana University Mathematics Journal **30** (1981), 713–747.
- [109] J. Hutchinson and L. Rüschendorf, *Random fractals and probability metrics*, Advances in Applied Probability **32** (2000), 925–947.
- [110] W. Johnson, J. Lindenstrauss, D. Preiss, and G. Schechtman, *Uniform quotient mappings of the plane*, Michigan Mathematical Journal **47** (2000), 15–31.
- [111] W. Johnson, J. Lindenstrauss, D. Preiss, and G. Schechtman, *Lipschitz quotients from metric trees and from Banach spaces containing ℓ^1* , Journal of Functional Analysis **194** (2002), 332–346.
- [112] F. Jones, *Lebesgue Integration on Euclidean Space*, Jones and Bartlett, 1993.
- [113] J.-L. Journé, *Calderón–Zygmund Operators, Pseudo-Differential Operators, and the Cauchy Integral of Calderón*, Lecture Notes in Mathematics **994**, Springer-Verlag, 1983.
- [114] I. Kaplansky, *Set Theory and Metric Spaces*, second edition, Chelsea, 1977.
- [115] M. Kapovich, *Hyperbolic Geometry and Discrete Groups*, Birkhäuser, 2001.

- [116] S. Katok, *Fuchsian Groups*, University of Chicago Press, 1992.
- [117] Y. Katznelson, *An Introduction to Harmonic Analysis*, second edition, Dover, 1976.
- [118] J. Kigami, *Analysis on Fractals*, Cambridge University Press, 2001.
- [119] R. Kirby, *The Topology of 4-Manifolds*, Lecture Notes in Mathematics **1374**, Springer-Verlag, 1989.
- [120] R. Kirby and L. Siebenmann, *Foundational Essays on Topological Manifolds, Smoothings, and Triangulations*, with notes by J. Milnor and M. Atiyah, Annals of Mathematics Studies **88**, Princeton University Press, 1977.
- [121] B. Kirchheim, *Rectifiable metric spaces: Local structure and regularity of the Hausdorff measure*, Proceedings of the American Mathematical Society **121** (1994), 113–123.
- [122] S. Krantz, *A Panorama of Harmonic Analysis*, Carus Mathematical Monographs **27**, Mathematical Association of America, 1999.
- [123] S. Krantz, *Function Theory of Several Complex Variables*, second edition, AMS Chelsea Publishing, 2001.
- [124] S. Krantz and H. Parks, *The Geometry of Domains in Space*, Birkhäuser, 1999.
- [125] S. Krantz and H. Parks, *A Primer of Real Analytic Functions*, second edition, Birkhäuser, 2002.
- [126] S. Krantz and H. Parks, *The Implicit Function Theorem: History, Theory, and Applications*, Birkhäuser, 2002.
- [127] B. Lawson, *The Theory of Gauge Fields in Four Dimensions*, CBMS Regional Conference Series in Mathematics **58**, American Mathematical Society, 1985.
- [128] M. Levin and Y. Sternfeld, *Monotone basic embeddings of hereditarily indecomposable continua*, Topology and its Applications **68** (1996), 241–249.

- [129] M. Levin and Y. Sternfeld, *Hyperspaces of two-dimensional continua*, Fundamenta Mathematicae **150** (1996), 17–24.
- [130] M. Levin and Y. Sternfeld, *The space of subcontinua of a 2-dimensional continuum is infinite-dimensional*, Proceedings of the American Mathematical Society **125** (1997), 2771–2775.
- [131] J. Luukkainen, *Assouad dimension: Antifractal metrization, porous sets, and homogeneous measures*, Journal of the Korean Mathematical Society **35** (1998), 23–76.
- [132] V. Magnani, *Elements of Geometric Measure Theory on Sub-Riemannian Groups*, thesis, Scuola Normale Superiore, Pisa, 2002.
- [133] R. Mañé, *Ergodic Theory and Differentiable Dynamics*, translated from the Portuguese by S. Levy, Springer-Verlag, 1987.
- [134] O. Martio, S. Rickman, and J. Väisälä, *Topological and metric properties of quasiregular mappings*, Annales Academiæ Scientiarum Fennicæ Ser. A I Math. **488** (1971), 31p.
- [135] W. Massey, *A Basic Course in Algebraic Topology*, Springer-Verlag, 1991.
- [136] P. Mattila, *Geometry of Sets and Measures in Euclidean Spaces*, Cambridge University Press, 1995.
- [137] V. Mazya, *Sobolev Spaces*, Springer-Verlag, 1985.
- [138] V. Milman and G. Schechtman, *Asymptotic Theory of Finite Dimensional Normed Vector Spaces*, with an appendix by M. Gromov, Lecture Notes in Mathematics **1200**, Springer-Verlag, 1986.
- [139] J. Milnor, *Topology from the Differentiable Viewpoint*, Princeton University Press, 1997.
- [140] E. Moise, *Geometric Topology in Dimensions 2 and 3*, Springer-Verlag, 1977.
- [141] R. Montgomery, *A Tour of Subriemannian Geometries, their Geodesics and Applications*, Mathematical Surveys and Monographs **91**, American Mathematical Society, 2002.

- [142] R. Monti and F. Serra Cassano, *Surface measures in Carnot–Carathéodory spaces*, Calculus of Variations and Partial Differential Equations **13** (2001), 339–376.
- [143] F. Morgan, *Riemannian Geometry: A Beginner’s Guide*, second edition, A K Peters, 1998.
- [144] F. Morgan, *Geometric Measure Theory: A Beginner’s Guide*, third edition, Academic Press, 2000.
- [145] J. Nagata, *Modern Dimension Theory*, revised edition, Heldermann Verlag, 1983.
- [146] P. Pansu, *Dimension conforme et sphère à l’infini des variétés à courbure négative*, Annales Academiae Scientiarum Fennicae Ser. A I Math. **14** (1989), 177–212.
- [147] S. Patterson, *The limit set of a Fuchsian group*, Acta Mathematica **136** (1976), 241–273.
- [148] P. Petersen, *A finiteness theorem for metric spaces*, Journal of Differential Geometry **31** (1990), 387–395.
- [149] P. Petersen, *Gromov–Hausdorff convergence of metric spaces*, in *Differential Geometry: Riemannian Geometry, Los Angeles CA, 1990*, 489–504, Proceedings of Symposia in Pure Mathematics **54**, Part 3, American Mathematical Society, 1993.
- [150] P. Petersen, *Riemannian Geometry*, Springer-Verlag, 1998.
- [151] P. Petersen, *Aspects of global Riemannian geometry*, Bulletin of the American Mathematical Society (N.S.) **36** (1999), 297–344.
- [152] G. Pisier, *The Volume of Convex Bodies and Banach Space Geometry*, Cambridge University Press, 1989.
- [153] C. Pommerenke, *Boundary Behavior of Conformal Maps*, Springer-Verlag, 1992.
- [154] A. Pressley, *Elementary Differential Geometry*, Springer-Verlag, 2001.

- [155] F. Quinn, *Local algebraic topology*, Notices of the American Mathematical Society **33** (1986), 895–899.
- [156] Y. Reshetnyak, *Space Mappings with Bounded Distortion*, American Mathematical Society, 1989.
- [157] S. Rickman, *Quasiregular Mappings*, Springer-Verlag, 1993.
- [158] C. Rogers, *Hausdorff Measures*, with a forward by K. Falconer, Cambridge University Press, 1998.
- [159] H. Royden, *Real Analysis*, third edition, Macmillan, 1988.
- [160] W. Rudin, *Principles of Mathematical Analysis*, third edition, McGraw-Hill, 1976.
- [161] W. Rudin, *Function Theory in the Unit Ball of \mathbf{C}^n* , Springer-Verlag, 1980.
- [162] W. Rudin, *Real and Complex Analysis*, third edition, McGraw-Hill, 1987.
- [163] W. Rudin, *Functional Analysis*, second edition, McGraw-Hill, 1991.
- [164] T. Rushing, *Topological Embeddings*, Academic Press, 1973.
- [165] T. Rushing, *Hausdorff dimension of wild fractals*, Transactions of the American Mathematical Society **334** (1992), 597–613.
- [166] C. Sadosky, *Interpolation of Operators and Singular Integral Operators: An Introduction to Harmonic Analysis*, Marcel Dekker, 1979.
- [167] S. Semmes, *Finding structure in sets with little smoothness*, in *Proceedings of the International Congress of Mathematicians, Zürich, 1994*, Volume 2, 875–885, 1995.
- [168] S. Semmes, *Some topics concerning homeomorphic parameterizations*, Publicacions Matemàtiques **45** (2001), 3–67.
- [169] S. Semmes, *Real analysis, quantitative topology, and geometric complexity*, Publicacions Matemàtiques **45** (2001), 265–333.

- [170] Z. Shen, *Lectures on Finsler Geometry*, World Scientific, Singapore, 2001.
- [171] L. Siebenmann, *Topological manifolds*, in *Proceedings of the International Congress of Mathematicians, Nice, 1970*, Volume 2, 133–163, 1971.
- [172] L. Siebenmann and D. Sullivan, *On complexes that are Lipschitz manifolds*, in *Geometric Topology: Proceedings of the Georgia Topology Conference, Athens GA, 1977*, 503–525, Academic Press, 1979.
- [173] G. Simmons, *Introduction to Topology and Modern Analysis*, McGraw-Hill, 1963.
- [174] L. Simon, *Lectures on Geometric Measure Theory*, Australian National University, 1983.
- [175] S. Staples and L. Ward, *Quasisymmetrically thick sets*, *Annales Academiae Scientiarum Fennicae Math.* **23** (1998), 151–168.
- [176] L. Steen and J. Seebach, Jr., *Counterexamples in Topology*, Dover, 1995.
- [177] E. Stein, *Singular Integrals and Differentiability Properties of Functions*, Princeton University Press, 1970.
- [178] E. Stein, *Harmonic Analysis: Real-Variable Methods, Orthogonality, and Oscillatory Integrals*, Princeton University Press, 1993.
- [179] E. Stein and G. Weiss, *Introduction to Fourier Analysis on Euclidean Spaces*, Princeton University Press, 1971.
- [180] Y. Sternfeld, *A short elementary proof of the Dranishnikov–West theorem on stable intersection of compacta in Euclidean spaces*, in *Proceedings of the International Conference on Set-Theoretic Topology and its Applications, Matsuyama, 1994*, *Topology and its Applications* **74** (1996), 177–178.
- [181] Y. Sternfeld, *Extension of mappings of Bing spaces into ANRs*, *Topology and its Applications* **80** (1997), 189–194.

- [182] R. Stong, *Mapping \mathbf{Z}^r into \mathbf{Z}^s with maximal contraction*, Discrete & Computational Geometry **20** (1998), 131–138.
- [183] D. Stroock, *Probability Theory: An Analytic View*, Cambridge University Press, 1993.
- [184] D. Stroock, *A Concise Introduction to the Theory of Integration*, third edition, Birkhäuser, 1999.
- [185] D. Sullivan, *Hyperbolic geometry and homeomorphisms*, in *Geometric Topology: Proceedings of the Georgia Topology Conference, Athens GA, 1977*, 543–555, Academic Press, 1979.
- [186] D. Sullivan, *The density at infinity of a discrete group of hyperbolic motions*, Publications Mathématiques IHES **50** (1979), 171–202.
- [187] A. Thompson, *Minkowski Geometry*, Cambridge University Press, 1996.
- [188] W. Thurston, *Three-Dimensional Geometry and Topology*, edited by Silvio Levy, Princeton University Press, 1997.
- [189] P. Tukia and J. Väisälä, *Quasisymmetric embeddings of metric spaces*, Annales Academiae Scientiarum Fennicae Ser. A I Math. **5** (1980), 97–114.
- [190] J. Väisälä, *Quasisymmetric embeddings in Euclidean spaces*, Transactions of the American Mathematical Society **264** (1981), 191–204.
- [191] J. Väisälä, *Porous sets and quasisymmetric maps*, Transactions of the American Mathematical Society **299** (1987), 525–533.
- [192] J. Väisälä, *Metric duality in Euclidean spaces*, Mathematica Scandinavica **80** (1997), 249–288.
- [193] J. Väisälä, *The wall conjecture for domains in Euclidean spaces*, Manuscripta Mathematica **93** (1997), 515–534.
- [194] S. Weinberger, *The Topological Classification of Stratified Spaces*, University of Chicago Press, 1994.

- [195] G. Whyburn, *Topological Analysis*, second revised edition, Princeton University Press, 1964.
- [196] R. Wilder, *Topology of Manifolds*, Colloquium Publications **32**, American Mathematical Society, 1963.
- [197] R. Wheeden and A. Zygmund, *Measure and Integral: An Introduction to Real Analysis*, Marcel Dekker, 1977.
- [198] H. Whitney, *Geometric Integration Theory*, Princeton University Press, 1957.
- [199] Y. Yomdin, *The set of zeroes of an “almost polynomial” function*, Proceedings of the American Mathematical Society **90** (1984), 538–542.
- [200] Y. Yomdin, *Volume growth and entropy*, Israel Journal of Mathematics **57** (1987), 285–300.
- [201] Y. Yomdin, *C^k -Resolution of semialgebraic mappings: Addendum to “Volume growth and entropy”*, Israel Journal of Mathematics **57** (1987), 310–317.
- [202] W. Ziemer, *Weakly Differentiable Functions: Sobolev Spaces and Functions of Bounded Variation*, Springer-Verlag, 1989.
- [203] A. Zygmund, *Trigonometric Series*, Cambridge University Press, 1979.