# ON THE COMPUTATION OF RATLIFF-RUSH CLOSURE

JUAN ELIAS \*

ABSTRACT. Let R be a Cohen-Macaulay local ring with maximal ideal **m**. In this paper we present a procedure for computing the Ratllif-Rush closure of a **m**-primary ideal  $I \subset R$ .

## INTRODUCTION

Let R be a Cohen-Macaulay Noetherian local ring of dimension  $d \ge 1$  with maximal ideal  $\mathbf{m}$  and residue field  $\mathbf{k}$  that we may assume infinite. Given a  $\mathbf{m}$ -primary ideal  $I \subset R$  in [5] the Ratliff-Rush closure of I is defined by  $\tilde{I} = \bigcup_{k\ge 1} (I^{k+1} : I^k))$ , and it holds that

$$\tilde{I} = \bigcup_{k \ge 1} (I^{k+1} : (x_1^k, \cdots, x_d^k))$$

where  $x_1, \dots, x_d$  is a minimal reduction of *I*.

Although Ratliff-Rush behaves bad under most of the basic operations of commutative algebra it is a basic tool in the study of the Hilbert functions of primary ideals, see for example [6] and its reference list.

Shah defined in [8] a finite chain of ideals between I and its integral closure  $\overline{I}$ 

$$I \subset I_{[d]} \subset \cdots \subset I_{[1]} \subset \overline{I}$$

 $I_{[i]}$  is the *i*-coefficient ideal of I, and  $I_{[d]} = \tilde{I}$  the Ratliff-Rush closure of I. Few results are known about the explicit computation of coefficient ideals. Ciupercă in [4] computed the first coefficient ideal of an ideal  $I \subset R$ , R is an  $(S_2)$  ring, by considering the  $S_2$ -ification of the extended Rees algebra of I.

The aim of this paper is to present an algorithm for the computation of Ratliff-Rush closure. In the first section we prove some results on superficial sequences that enable us to describe, in the section two, an algorithm to compute Ratliff-Rush closure. We end the paper with some explicit computations of the Ratliff-Rush closure of ideals using the algorithm of this paper.

\*Partially supported by DGICYT BFM2001-3584, and

Date: September 23, 2018.

Commutative Algebra program of the MSRI, Berkeley.

<sup>2000</sup> MSC: 13P10, 13H15, 13H10.

#### JUAN ELIAS

We will use freely [2] as a general reference for the algebraic concepts appearing in this paper. The computations of this paper are performed by using CoCoA, [3].

We thank Ciupercă for the useful comments on a previous version of this paper. We also thank M.E. Rossi and W. Vasconcelos for pointing us that [7, Corollary 3.4] holds also for  $\mathbf{m}$ -primary ideals.

### 1. On superficial sequences

Let I be an **m**-primary ideal of R. We denote by  $gr_I(R) = \bigoplus_{k\geq 0} I^k/I^{k+1}$  the associated graded ring of I, and by l(I) the analytic spread of I.

Let  $h_I(n) = length_R(R/I^{n+1})$  be the Hilbert-Samuel function of  $I, n \in \mathbb{N}$ . Hence there exist integers  $e_j(I) \in \mathbb{Z}$  such that

$$p_I(X) = \sum_{j=0}^d (-1)^j e_j(I) \binom{X+d-j}{d-j}$$

is the Hilbert-Samuel polynomial of I, i.e.  $h_I(n) = p_I(n)$  for  $n \gg 0$ . The integer  $e_j(I)$  is the *j*-th Hilbert coefficient of I,  $j = 0, \dots, d$ . Shah proved that coefficient ideals are the largest ideals  $I_{[t]}$  containing I and such that:

(i) 
$$e_i(I) = e_i(I_{[t]})$$
 for  $i = \underline{0}, \cdots, t$ ,

(ii) 
$$I \subset I_{[d]} \subset \cdots \subset I_{[1]} \subset \overline{I}$$

where  $\overline{I}$  is the integral closure of I, [8]. Notice that  $\tilde{I}$  is the largest ideal containing I and such that  $e_i(I) = e_i(\tilde{I})$  for  $i = 0, \dots, d$ .

We say that  $x \in I$  is a superficial element of I if there exists an integer  $k_0$  such that  $(I^{k+1}: x) = I^k$  for  $k \ge k_0$ . Since the residue field is infinite it hold:

- (1) a set elements  $x_1, \dots, x_d \in I$ , such that their cosets  $\overline{x}_1, \dots, \overline{x}_d \in I/\mathbf{m}I$  are generic, form a superficial sequence  $x_1, \dots, x_d$  of I, i.e.  $x_i$  is a superficial element of  $I/(x_1, \dots, x_{i-1})$  for  $i = 1, \dots, d$ ,
- (2) if  $x_1, \dots, x_d \in I$  is a set of elements such that

length 
$$_{R}\left(\frac{R}{(x_{1},\cdots,x_{d})}\right) = e_{0}(I),$$

where  $e_0(I)$  is the multiplicity of I, then  $x_1, \dots, x_d$  a superficial sequence of the ideal I,

(3) if  $x_1, \dots, x_d$  is a superficial sequence of I then  $J = (x_1, \dots, x_d)$  is a minimal reduction of I, [9].

Given a superficial element x of I if we write  $\overline{I} = I/(x)$  then it is well known that

$$p_{\overline{I}}(X) = p_I(X) - p_I(X-1) = \sum_{j=0}^{d-1} (-1)^j e_j(I) \binom{X+d-2-j}{d-1-j}$$

 $\mathbf{2}$ 

in particular  $e_i(\overline{I}) = e_i(I)$  for  $i = 0, \dots, d-1$ . We define the postulation number pn(I) of I as the smallest integer n such that  $h_I(t) = p_I(t)$  for all  $t \ge n$ . Given a superficial sequence  $x_1, \dots, x_d$  of I we denote by  $pn(I; x_1, \dots, x_d)$  the maximum among pn(I) and  $pn(I/(x_i)), i = 1, \dots, d$ .

**Proposition 1.1.** Let I be a m-primary ideal of R and x a superficial element of I. We denote by  $\overline{I} = I/(x)$  the ideal of  $\overline{R} = R/(x)$ . For all  $k \ge pn(I; x) + 1$  it holds  $(I^{k+1}: x) = I^k.$ 

*Proof.* Let us consider the exact sequence

$$0 \longrightarrow \frac{(I^{k+1}:x)}{I^k} \longrightarrow \frac{R}{I^k} \xrightarrow{.x} \frac{R}{I^{k+1}} \longrightarrow \frac{\overline{R}}{\overline{I^{k+1}}} \longrightarrow 0$$

 $\mathbf{SO}$ 

length 
$$_{R}\left(\frac{(I^{k+1}:x)}{I^{k}}\right) = h_{I}(k-1) - h_{I}(k) + h_{\overline{I}}(k)$$

If  $k \ge pn(I; x) + 1$  then we have that  $h_I(k) = p_I(k)$ ,  $h_I(k-1) = p_I(k-1)$  and  $h_{\overline{I}}(k) = p_{\overline{I}}(k)$ , so

length 
$$_{R}\left(\frac{(I^{k+1}:x)}{I^{k}}\right) = p_{I}(k-1) - p_{I}(k) + p_{\overline{I}}(k)$$

On the other hand, since x is a superficial element of I we have that  $p_{\overline{I}}(X) = p_I(X) - p_I(X-1)$  then  $(I^{k+1}: x) = I^k$  for all  $k \ge pn(I; x) + 1$ .

Notice that for the explicit computations of coefficient ideals it is enough to consider the number  $pn(I; x_1, \dots, x_d)$ , Theorem 2.1 (i), but if we look for a explicit formula of the Ratliff-Rush closure avoiding the computation of superficial sequences we have to consider the Castelnuovo-Mumford regularity, Theorem 2.1 (ii).

Given a standard  $A_0$ -algebra  $A = A_0 \oplus A_1 \oplus \cdots$  with  $A_0$  an Artin ring, we denote by reg (A) the Castelnuovo-Mumford regularity of A, i.e. the smallest integer m such that  $H^i_{A_+}(A)_n = 0$  for all  $i = 0, \cdots, d$  and  $n \ge m - i + 1$ , where  $A_+ = A_1 \oplus \cdots$ the irrelevant ideal of A.

We denote by  $f: \mathbb{N}^2 \longrightarrow \mathbb{N}$  the numerical function defined by

$$f(e,d) = \begin{cases} e-1 & \text{if } d=1\\ e^{2(d-1)!-1}(e-1)^{(d-1)!} & \text{if } d \ge 2 \end{cases}$$

Rossi, Trung and Valla prove that f(e, d) is an upper bound of the Castelnuovo-Mumford regularity of the associated graded ring of I, see Proposition 1.2.

Given a minimal reduction J of I we denote by  $r_J(I)$  the reduction number of I with respect to I, i.e. the smallest integer r such that  $I^{r+1} = JI^r$ .

#### JUAN ELIAS

In the next result we relate some of the numerical characters that we already defined in this paper.

**Proposition 1.2.** Let R be a Cohen-Macaulay local ring of dimension  $d \ge 1$ . Let I be a **m**-primary ideal of R and  $J = (x_1, \dots, x_d)$  a minimal reduction of I. Then

- (i)  $r_J(I) \leq reg (gr_I(R)) \leq f(e_0(I), d)$ , and
- (ii)  $pn(I; x_1, \cdots, x_d) \le f(e_0(I), d) + 1.$

*Proof.* (i) The first inequality comes from [10, Proposition 3.2], see also [1, Theorem 18.3.12]. The second inequality is due to Rossi, Trung and Valla, [7, Corollary 3.4]. (ii) Notice that from Serre's formula, [2, Theorem 4.4.3], and the right hand side inequality in (i) we have that  $pn(I) \leq f(e_0(I), d) + 1$  and  $pn(I/(x_i)) \leq f(e_0(I/(x_i)), d-1) + 1$ ,  $i = 1, \dots, d$ . Since  $e_0(I) = e_0(I/(x_i))$  and  $f(e, d-1) \leq f(e, d)$ , we get the claim.

Notice that in [7, Corollary 3.4] the right hand side inequality in (i) of the above result is proved for the maximal ideal  $I = \mathbf{m}$ , but the proof holds also for general  $\mathbf{m}$ -primary ideals.

**Corollary 1.3.** Let x be a superficial element of I. For all  $k \ge f(e_0(I), d) + 2$  we have

$$(I^{k+1}:x) = I^k.$$

*Proof.* It is a consequence of Proposition 1.2 (*ii*) and Proposition 1.1.

## 2. An Algorithm for computing Ratliff-Rush closure

In this section we compute explicitly Ratliff-Rush closure by using Proposition 1.1 and Corollary 1.3. We consider the increasing ideal chain

$$\mathcal{L}_1 \subset \mathcal{L}_2 \subset \cdots \subset \mathcal{L}_k \subset \cdots$$

where

 $\mathcal{L}_k = (I^{k+1} : (x_1^k, \cdots, x_d^k)).$ 

Notice that  $\tilde{I} = \bigcup_{k>1} \mathcal{L}_k$  is the Ratliff-Rush closure of I.

**Theorem 2.1.** Let R be a Cohen-Macaulay local ring of dimension  $d \ge 1$ . Let I be an **m**-primary ideal of R and let  $x_1, \dots, x_d$  be a superficial sequence of I.

(i) For all  $k \ge pn(I; x_1, \cdots, x_d) + 1$  it holds that  $\tilde{I} = (I^{k+1} : (x_1^k, \cdots, x_d^k)).$ 

(ii) For all 
$$k \ge (d+1)(f(e_0(I))+2)$$
 it holds that  
 $\tilde{I} = (I^{k+1} : I^k).$ 

Proof. (i) We have to prove that for all  $k \ge pn(I; x_1, \dots, x_d) + 1$  it holds  $\mathcal{L}_k = \mathcal{L}_{k+1}$ . Notice that for all  $n \ge 1$  we have  $\mathcal{L}_n \subset \mathcal{L}_{n+1}$  so we only need to prove  $\mathcal{L}_{k+1} \subset \mathcal{L}_k$ . Given  $a \in \mathcal{L}_{k+1}$  we have  $ax_i^{k+1} = x_i(ax_i^k) \in I^{k+2}$ , for all  $i = 1, \dots, d$ . Since  $k \ge pn(I; x_1, \dots, x_d) + 1$  from Proposition 1.1 we get  $ax_i^k \in I^{k+1}$  for all  $i = 1, \dots, d$ , so  $a \in \mathcal{L}_k$ .

(*ii*) Notice that  $J = (x_1, \dots, x_d)$  is a minimal reduction of I so for all  $k \ge r_J(I)$  $I^{(d+1)k} = I^{dk}(x_1^k, \dots, x_d^k).$ 

From Proposition 1.2 we have that  $r_J(I) \leq reg(gr_I(R)) \leq f(e_0(I), d)$ . Let  $n \geq f(e_0(I), d) + 2$  be an integer and let  $a \in \tilde{I}$  be an element of the Rattlif-Rush closure of I. Hence from (i) we have  $ax^{[k]} \subset I^{k+1}$  and since  $I^{(d+1)k} = I^{dk}(x_1^k, \cdots, x_d^k)$  we get

$$aI^{(d+1)k} \subset aI^{dk}(x_1^k, \cdots, x_d^k) \subset I^{(d+1)k+1}.$$

In particular we have  $a \in (I^{(d+1)k+1} : I^{(d+1)k})$ , since by definition  $(I^{(d+1)k+1} : I^{(d+1)k}) \subset \tilde{I}$  we get the claim.  $\Box$ 

From the last result we deduce that the problem of computing the Ratliff-Rush closure can be reduced to the computation of the postulation number of I and its quotients  $I/(x_i)$ ,  $i = 1, \dots, d$ . Next we recall how to compute these numbers.

We denote by  $PS_I(X) \in \mathbb{Z}[[X]]$  the Poincaré series of I

$$PS_{I}(X) = \sum_{i \ge 0} length_{R}\left(\frac{I^{i}}{I^{i+1}}\right) X^{i}$$

it is known that there exists a degree s polynomial  $f(X) = \sum_{i=0}^{s} a_i X^i \in \mathbb{Z}[X]$  such that

$$PS_I(X) = \frac{f(X)}{(1-X)^d}.$$

It is easy to prove that  $e_0(I) = \sum_{i=0}^s a_i$  and that pn(I) = s - d.

**Remark 2.2.** It is well known that the computation of the Poincaré series of I and its quotients  $I/(x_i)$  can be reduced to a elimination of variables process, see for example the library primary.lib of CoCoa, [3].

An algorithm for computing the Ratliff-Rush closure.

**Step 1.** Compute the Poincaré series of *I*. Then we know the multiplicity  $e_0(I)$  and the postulation number pn(I) of *I*.

#### JUAN ELIAS

- Step 2. Find d generic elements  $x_1, \dots, x_d$  of the **k**-vector space  $I/\mathbf{m}I$  such that  $length_R(R/(x_1, \dots, x_d)) = e_0(I)$ . Recall that  $x_1, \dots, x_d$  is a superficial sequence of I and generates a minimal reduction of I.
- Step 3. As in Step 1 compute  $PS_{I/(x_i)} = f_i(X)/(1-X)^{d-1}$  for  $i = 1, \dots, d$ . From this and the fact  $pn(I/(x_i)) = deg(f_i) - (d-1)$  we can compute  $pn(I; x_1, \dots, x_d)$ .
- Step 4. For  $k \ge pn(I; x_1, \cdots, x_d) + 1$  we get

$$\tilde{I} = (I^{k+1} : (x_1^k, \cdots, x_d^k)).$$

**Remark 2.3.** Notice that if I is a monomial ideal then **Step 4** can be performed without Gröbner basis computation.

We will show how to compute the Ratliff-Rush closure in some explicit examples of [4] and [6].

**Example 2.4.** Example 1.10 of [6]. Let  $I = (x^{10}, y^5, xy^4, x^8y)$  be an ideal of  $R = \mathbf{k}[x, y]_{(x,y)}$ . The Poincaré series of I is

$$PS_I(X) = \frac{35 + 4X + 4X^2 + 4X^3 - 2X^4}{(1-X)^2},$$

so  $e_0(I) = 45$  and pn(I) = 2. Since the length of  $R/(y^5 + x^{10} + x^8y, xy^4)$  is  $45 = e_0(I)$  we deduce that  $y^5 + x^{10} + x^8y, xy^4$  is a superficial sequence of I. A CoCoA computation shows that

$$PS_{I/(xy^4)}(X) = \frac{35 + 6X + 2X^2 + 2X^3}{1 - X},$$

and

$$PS_{I/(y^5+x^{10}+x^8y)}(X) = \frac{35+6X+4X^2}{1-X},$$

so  $pn(I; y^5 + x^{10} + x^8y, xy^4) = 2$ . Then by Theorem 2.1 (i) we get

$$I \subsetneq \tilde{I} = (I^4 : ((y^5 + x^{10} + x^8 y)^3, (xy^4)^3))) = (x^{10}, y^5, xy^4, x^7 y^2, x^6 y^3, x^8 y).$$

**Remark 2.5.** Let  $x_1 = y^5 + x^{10} + x^8 y$ ,  $x_2 = xy^4$  be the minimal reduction of the ideal I of the last example. Since  $pn(I; x_1, x_2) = 2$  we have that  $\tilde{I} = (I^4 : (x_1^3, x_2^3))$ . On the other hand, Theorem 2.1 (*ii*) gives that  $\tilde{I} = (I^{k+1} : I^k)$  for all  $k \ge 540$ , this is a hard computation.

**Example 2.6.** Example 1.4 of [6]. Let us consider the ideal

$$I = (y^{22}, x^4y^{18}, x^7y^{15}, x^8y^{14}, x^{11}y^{11}, x^{14}y^8, x^{15}y^7, x^{18}y^4, x^{22})$$

of the local ring  $R = \mathbf{k}[x, y]_{(x,y)}$ . A similar computation as we did in the previous example shows that  $I = \tilde{I}$  and

$$I^2 \subsetneq \widetilde{I^2} = I^2 + (x^{24}y^{20}, x^{20}y^{24}).$$

**Example 2.7.** Example 3.3 of [4]. Let us consider the ideal

$$I = (x^8, x^3y^2, x^2y^4, y^8)$$

of the local ring  $R = \mathbf{k}[x, y]_{(x,y)}$ . A similar computation as before shows that  $I = \tilde{I}$ . Ciupercă in [4] computed the first coefficient ideal of I:

$$I = I \subsetneq I_{[1]} = (x^8, x^3y^2, x^2y^4, xy^6, y^8).$$

## References

- M.P. Brodman and R.Y. Sharp. Local Cohomology, volume 60 of Cambridge Studies in Advanced Mathematics. Cambridge University Press, 1998.
- [2] W. Bruns and J. Herzog. Cohen-Macaulay Rings. Cambridge studies in advanced mathematics. Cambridge University Press, 1993.
- [3] A. Capani, G. Niesi, and L. Robbiano. *CoCoA*, a system for doing Computations in Commutative Algebra. Available via anonymous ftp from cocoa.dima.unige.it, 4.0 edition, 2000.
- [4] C. Ciupercă. First coefficient ideals and the S<sub>2</sub>-ification of a Rees algebra. J. of Alg., 242:782– 794, 2001.
- [5] L. Ratliff and D. Rush. Two notes on reductions of ideals. Indiana Univ. Math. J., 27:929–934, 1978.
- [6] M. E. Rossi and I. Swanson. Notes on the behaviour of the Ratliff-Rush filtration. *Preprint*, 2002.
- [7] M.E. Rossi, N.V. Trung, and G. Valla. Castelnuovo-Mumford regularity and extended degree. *Preprint*, 2002.
- [8] K. Shah. Coefficient ideals. Trans. A.M.S., 327(1):373–384, 1991.
- [9] I. Swanson. A note on analytic spread. Comm. in Algebra, 22:407–411, 1994.
- [10] N.V. Trung. Reduction exponent and degree bound for the defining equations of graded rings. Proc. A.M.S., 101-2:229–236, 1987.

DEPARTAMENT D'ÀLGEBRA I GEOMETRIA FACULTAT DE MATEMÀTIQUES UNIVERSITAT DE BARCELONA GRAN VIA 585, 08007 BARCELONA, SPAIN *E-mail address*: elias@mat.ub.es