

**LOW-DIMENSIONAL COHOMOLOGY OF CURRENT LIE
ALGEBRAS AND STRUCTURE FUNCTIONS ASSOCIATED
WITH LOOP MANIFOLDS**

PASHA ZUSMANOVICH

ABSTRACT. We obtain formulas for the first and second cohomology groups of a general current Lie algebra with coefficient in the “current” module, and apply them to computation of structure functions associated with manifolds of loops on compact Hermitian symmetric space.

INTRODUCTION

We deal with the low-dimensional cohomology of current Lie algebras with coefficient in the “current module”. Namely, let L be a Lie algebra, M be a L -module, A be an associative commutative algebra with unit, V be a symmetric unitary A -module. Then the Lie algebra structure on $L \otimes A$ and the $L \otimes A$ -module structure on $M \otimes V$ are defined via obvious formulas:

$$\begin{aligned} [x \otimes a, y \otimes b] &= [x, y] \otimes ab \\ (x \otimes a) \bullet (m \otimes v) &= x \bullet m \otimes a \bullet v \end{aligned}$$

for $x, y \in L$, $m \in M$, $a, b \in A$, $v \in V$.

The aim of this note is threefold. First, we want to establish some facts on the low-dimensional cohomology we will need in our further projects. Second, we want to demonstrate that the problem of describing of such cohomology in terms of the tensor factors L and A probably does not have an adequate general solution, as even the partial answer for the two-dimensional cohomology seems to be overwhelmingly complex. Third, we want to demonstrate an application of this cohomology to some differential geometric question.

In §1 we establish an elementary result from linear algebra which will be useful in the course of subsequent algebraic manipulations. In §2 we get a formula for the first cohomology group. In §3 we compute the second cohomology group in two cases – when L is abelian and when L acts trivially on the whole cohomology group $H^2(L \otimes A, M \otimes V)$. At the end of this section we present a list of 13 types of 2-cocycles (so-called cocycles of rank 1, generated by decomposable elements in the tensor product) in the general case. However, this list is a priori not complete. In §4 a certain spectral sequence is sketched, which may provide a more conceptual framework for computations in preceding sections. However, we do not go into details and other sections are not dependent on that one. The last §5 devoted to application. We show how from previous computations one may derive the results

Date: February 26, 2003; last corrected April 18, 2003.

2000 Mathematics Subject Classification. Primary 17B56, Secondary 17B20, 17B70, 32M15, 53C10, 58H15.

about obstructions to integrability (structure functions) of manifolds of loops on compact Hermitian symmetric space.

One should note that the result about the first cohomology group (particularly, about derivations of the current Lie algebra) can be found in different forms in the literature and is a sort of folklore, and the partial results about the second cohomology were obtained by Cathelineau [C], Haddi [H], Lecomte and Roger [R] and the author [Z]. However, all these results do not provide the whole generality we need - as various restrictions, notably the zero characteristic of the ground field and perfectness of the Lie algebra L were imposed. Moreover, as we see in §5, the case in a sense opposite to the case of perfect L , namely, the case of abelian L , does lead to some interesting application.

The technique used is highly computational in nature and based on applying various symmetrization operators to the cocycle equation.

NOTATIONS

The ground field K assumed to be arbitrary field of characteristic $p \neq 2, 3$ in §1-4, and \mathbb{C} in §5.

$H^n(L, M)$, $C^n(L, M)$, $Z^n(L, M)$, $B^n(L, M)$ stands, respectively, for cohomology, cochains, cocycles and coboundaries of a Lie algebra L with coefficients in a module M .

$S^n(A, V)$ stands for n -linear maps $A \times \cdots \times A \rightarrow V$, symmetric in all arguments.

$\wedge^n(V)$ and $T^n(V)$ stands, respectively, for n -fold skew and tensor products of a module V .

$Har^n(A, V)$ and $Z^n(A, V)$ stands, respectively, for Harrison cohomology and Harrison cycles (for $n = 2$ these are just symmetric Hochschild cocycles) of an associative commutative algebra A with coefficients in a module V .

$Der(A)$ denotes derivation algebra of an algebra A , and, more generally,

$Der(A, V)$ denotes an algebra of derivations of A with values in a A -module V .

All other (nonstandard and unavoidably numerous) notations for different spaces of multilinear mappings and modules are defined as long as they introduced in the text.

The symbol \curvearrowright after an expression refers to the sum of all cyclic permutations (in $S(3)$) of letters and indices occurring in that expression.

1. A LEMMA FROM LINEAR ALGEBRA

If either both L and M or both A and V are finite-dimensional, then each cocycle Φ in $Z^n(L \otimes A, M \otimes V)$ can be represented as an element in $Hom(L^{\otimes n}, M) \otimes Hom(A^{\otimes n}, V)$:

$$(1.1) \quad \Phi = \sum_{i \in I} \varphi_i \otimes \alpha_i$$

where φ_i, α_i are multilinear mappings $L \times \cdots \times L \rightarrow M$ and $A \times \cdots \times A \rightarrow V$ respectively. We restrict our considerations to this case. The minimal possible number $|I|$ such that the cocycle Φ can be written in the form (1.1) will be called the *rank of cocycle*.

In evaluation of the cohomology group $H^n(L \otimes A, M \otimes V)$ in terms of pairs (L, M) and (A, V) , we encounter conditions such as

$$(1.2) \quad \sum_{i \in I} S\varphi_i \otimes T\alpha_i = 0$$

where S and T are some linear operators defined on the spaces of multilinear mappings $L \times \cdots \times L \rightarrow M$ and $A \times \cdots \times A \rightarrow V$ respectively.

For example, the substitution $a_1 = \cdots = a_{n+1} = 1$ in the cocycle equation $d\Phi(x_1 \otimes a_1, \dots, x_{n+1} \otimes a_{n+1}) = 0$, where Φ is as in (1.1), yields

$$\sum_{i \in I} d\varphi_i(x_1, \dots, x_{n+1}) \otimes a_i(1, \dots, 1) = 0.$$

Another example: apply to the cocycle equation the symmetrization operator with respect to letters x_1, \dots, x_{n+1} , which we denote Y for a moment:

$$\sum_{i \in I} \left(Y(x_1 \bullet \varphi_i(x_2, \dots, x_{n+1})) \otimes \sum_{j=1}^{n+1} (-1)^j a_j \bullet \alpha_i(a_1, \dots, \hat{a}_j, \dots, a_{n+1}) \right) = 0.$$

So, suppose that a condition of type (1.2) holds. Since $\text{Ker}(S \otimes T) = \text{Hom}(A^{\otimes n}, V) \otimes \text{Ker} T + \text{Ker} S \otimes \text{Hom}(L^{\otimes n}, M)$, replacing α_i 's and φ_i 's by appropriate linear combinations, one can find a decomposition of the set of indices $I = I_1 \cup I_2$ such that

$$(1.3) \quad S\varphi_i = 0, \quad i \in I_1 \quad \text{and} \quad T\alpha_i = 0, \quad i \in I_2.$$

Suppose that another equality of type (1.2) holds:

$$(1.2)' \quad \sum_{i \in I} S'\varphi_i \otimes T'\alpha_i = 0.$$

Then it determines a new decomposition $I = I'_1 \cup I'_2$ such that $S'\varphi_i = 0$ if $i \in I'_1$ and $T'\alpha_i = 0$ if $i \in I'_2$. It turns out that it is possible to replace φ_i 's and α_i 's by linear combinations in such a way that both decompositions will be hold simultaneously.

Lemma 1.1. *Let U, W be two vector spaces, $S, S' \in \text{Hom}(U, -)$, $T, T' \in \text{Hom}(W, -)$. Then*

$$\begin{aligned} & \text{Ker}(S \otimes T) \cap \text{Ker}(S' \otimes T') \\ &= (\text{Ker} S \cap \text{Ker} S') \otimes W + \text{Ker} S \otimes \text{Ker} T' + \text{Ker} S' \otimes \text{Ker} T + U \otimes (\text{Ker} T \cap \text{Ker} T'). \end{aligned}$$

Proof. Since $\text{Ker}(S \otimes T) = \text{Ker} S \otimes W + U \otimes \text{Ker} T$ and analogously for $\text{Ker}(S' \otimes T')$, the asserted equality is a particular case of

$$(1.4) \quad \begin{aligned} & (U_1 \otimes W + U \otimes W_1) \cap (U_2 \otimes W + U \otimes W_2) \\ &= (U_1 \cap U_2) \otimes W + U_1 \otimes W_2 + U_2 \otimes W_1 + U \otimes (W_1 \cap W_2) \end{aligned}$$

provided U_1, U_2 and W_1, W_2 are subspaces of U and W respectively.

Assume for the moment that $U_1 \cap U_2 = W_1 \cap W_2 = 0$. Then writing $U = U_1 \oplus U_2 \oplus U'$ and $W = W_1 \oplus W_2 \oplus W'$ for some subspaces U', W' and substituting this in the left side of (1.4), we get:

$$\begin{aligned} & (U_1 \otimes W \oplus U_2 \otimes W_1 \oplus U' \otimes W_1) \cap (U_1 \otimes W_2 \oplus U_2 \otimes W \oplus U' \otimes W_2) \\ &= U_1 \otimes W_2 \oplus U_2 \otimes W_1. \end{aligned}$$

To prove (1.4) in the general case, pass to quotient by $(U_1 \cap U_2) \otimes W + U \otimes (W_1 \cap W_2)$ and receive by the just proved $U_1 \otimes W_2 + U_2 \otimes W_1$. ■

Below, in numerous applications of Lemma 1.1, we will use, by abuse of language, phrases like “by (1.2) and (1.2)', one gets a decomposition $I = I_1 \cup I_2 \cup I_3 \cup I_4$ such that $S\varphi_i = S'\varphi_i = 0$ for $i \in I_1$, $S\varphi_i = T'\alpha_i = 0$ for $i \in I_2$, $S'\varphi_i = T\alpha_i = 0$ for $i \in I_3$ and $T\alpha = T'\alpha_i = 0$ for $i \in I_4$ ”. This means that one can find a new expression $\Phi = \sum_{i \in I} \varphi_i \otimes \alpha_i$ with indicated properties (where the new φ_i 's and α_i 's are linear combinations of the old ones).

Unfortunately, for “triple intersection” $\text{Ker}(S \otimes T) \cap \text{Ker}(S' \otimes T') \cap \text{Ker}(S'' \otimes T'')$ the analogous decomposition is no longer true. That is why dealing with the second cohomology group in §3, we are unable to obtain a general result and restrict our considerations with cocycles of rank 1 or with some special cases. In the case $n = 1$, however, Lemma 1.1 suffices to consider a general case, but at the end of the proof it turns out that it is possible to choose a basis consisting of cocycles of rank 1.

2. THE FIRST COHOMOLOGY GROUP

From now on (in this and subsequent sections), either both L and M or both A and V are finite-dimensional.

In this section we prove

Theorem 2.1.

$$(2.1) \quad H^1(L \otimes A, M \otimes V) \simeq H^1(L, M) \otimes V$$

$$\oplus \text{Hom}_L(L, M) \otimes \text{Der}(A, V) \oplus \text{Hom}(L/[L, L], M^L) \otimes \frac{\text{Hom}(A, V)}{V + \text{Der}(A, V)}.$$

Each cocycle in $Z^1(L \otimes A, M \otimes V)$ is a linear combination of cocycles of the three following types (which correspond to the summands in (2.1)):

- (i) $x \otimes a \mapsto \varphi(x) \otimes a \bullet v$, $\varphi \in Z^1(L, M), v \in V$
- (ii) $x \otimes a \mapsto \varphi(x) \otimes \alpha(a)$, $\varphi \in \text{Hom}_L(L, M), \alpha \in \text{Der}(A, V)$
- (iii) as in (ii) with $\varphi(L) \subseteq M^L, \varphi([L, L]) = 0, \alpha \in \text{Hom}(A, V)$.

Remark. Theorem 2.1 was obtained earlier by Santharoubane [Sa] in particular case when $M = L^*, V = A^*$ and L is 1-generated as $U(L)^+$ -module, and by Haddi [H] (in homological form) in the case of zero characteristic and perfect L .

Proof. Let $\Phi = \sum_{i \in I} \varphi_i \otimes \alpha_i$ be a cocycle in $Z^1(L \otimes A, M \otimes V) \subset \text{Hom}(L, M) \otimes \text{Hom}(A, V)$. The cocycle equation $d\Phi = 0$ reads

$$(2.2) \quad \sum_{i \in I} (x \bullet \varphi_i(y) \otimes a \bullet \alpha_i(b) - y \bullet \varphi_i(x) \otimes b \bullet \alpha_i(a) - \varphi_i([x, y]) \otimes \alpha_i(ab)) = 0.$$

Symmetrizing this equation with respect to x, y , we get:

$$\sum_{i \in I} (x \bullet \varphi_i(y) + y \bullet \varphi_i(x)) \otimes (a \bullet \alpha_i(b) - b \bullet \alpha_i(a)) = 0.$$

Substitute in (2.2) $a = b = 1$:

$$\sum_{i \in I} d\varphi_i(x, y) \otimes \alpha_i(1) = 0.$$

Applying Lemma 1.1 to the last two equalities, we get a decomposition $I = I_1 \cup I_2 \cup I_3 \cup I_4$ such that

$$\begin{aligned} d\varphi_i &= 0, x \bullet \varphi_i(y) + y \bullet \varphi_i(x) = 0, & i \in I_1 \\ d\varphi_i &= 0, a \bullet \alpha_i(b) = b \bullet \alpha_i(a), & i \in I_2 \\ x \bullet \varphi_i(y) + y \bullet \varphi_i(x) &= 0, \alpha_i(1) = 0, & i \in I_3 \\ a \bullet \alpha_i(b) &= b \bullet \alpha_i(a), \alpha_i(1) = 0, & i \in I_4. \end{aligned}$$

It is easy to see that for each $i \in I_2$, $\alpha_i(a) = a \bullet \alpha_i(1)$ and the mappings $x \otimes a \mapsto \varphi_i(x) \otimes \alpha_i(a)$ are cocycles of type (i) from the statement of the theorem, and for each $i \in I_4$, $\alpha_i = 0$.

Substitute $b = 1$ in the cocycle equation (2.2):

$$\sum_{i \in I_1 \cup I_3} (x \bullet \varphi_i(y) - \varphi_i([x, y])) \otimes (\alpha_i(a) - a \bullet \alpha_i(1)) = 0.$$

Now apply Lemma 1.1 again. For elements $\varphi_i, i \in I_1$, the vanishing of $x \bullet \varphi_i(y) - \varphi_i([x, y])$ implies $\varphi_i = 0$, and the vanishing of $\alpha_i(a) - a \bullet \alpha_i(1)$ gives cocycles of type (i), an already considered case. We have $x \bullet \varphi_i(y) = \varphi_i([x, y])$ for all (remaining) $i \in I_3$.

Hence (2.2) can be rewritten as

$$\sum_{i \in I_3} \varphi_i([x, y]) \otimes (a \bullet \alpha_i(b) + b \bullet \alpha_i(a) - \alpha_i(ab)) = 0.$$

The vanishing of the first and second tensor factors gives rise to cocycles of type (iii) and (ii) respectively.

Hence we have

$$\begin{aligned} Z^1(L \otimes A, M \otimes V) &= Z^1(L, M) \otimes V \\ &\quad + Hom_L(L, M) \otimes Der(A, V) + Hom(L/[L, L], M^L) \otimes Hom(A, V) \end{aligned}$$

what can be rewritten as

$$\begin{aligned} Z^1(L \otimes A, M \otimes V) &= Z^1(L, M) \otimes V \\ &\quad \oplus Hom_L(L, M) \otimes Der(A, V) \oplus Hom(L/[L, L], M^L) \otimes \frac{Hom(A, V)}{V + Der(A, V)}. \end{aligned}$$

From the considerations above we easily deduce:

$$B^1(L \otimes A, M \otimes V) = B^1(L, M) \otimes V$$

and (2.1) now follows. ■

Corollary 2.2. *The derivation algebra of a current Lie algebra $L \otimes A$ is isomorphic to*

$$Der(L) \otimes A \oplus Hom_L(L, L) \otimes Der(A) \oplus Hom(L/[L, L], Z(L)) \otimes \frac{End(A)}{A + Der(A)}.$$

This overlaps with [B], Theorem 7.1.

3. THE SECOND COHOMOLOGY GROUP

In this section we shall obtain some particular results on the second cohomology group $H^2(L \otimes A, M \otimes V)$. The computations go by the same scheme as for H^1 but are more complicated.

As we want to express H^2 through the tensor products of modules depending on (L, M) and (A, V) , it is natural to do so for underlying modules of the Chevalley–Eilenberg complex. We have (under the same finiteness assumptions as previous):

$$(3.1) \quad \begin{aligned} C^1(L \otimes A, M \otimes V) &\simeq C^1(L, M) \otimes C^1(A, V) \\ C^2(L \otimes A, M \otimes V) &\simeq C^2(L, M) \otimes S^2(A, V) \oplus S^2(L, M) \otimes C^2(A, V). \end{aligned}$$

To obtain a similar decomposition in the third degree, let us denote (by abuse of language) the Young symmetrizer corresponding to tableau λ by the same symbol λ . We have decompositions of a unit element in the group algebra KS_3 :

$$e = \frac{1}{6} \begin{array}{|c|} \hline 1 \\ \hline 2 \\ \hline 3 \\ \hline \end{array} + \frac{1}{3} \left(\begin{array}{|c|c|} \hline 1 & 3 \\ \hline 2 & \\ \hline \end{array} + \begin{array}{|c|c|} \hline 1 & 2 \\ \hline 3 & \\ \hline \end{array} \right) + \frac{1}{6} \begin{array}{|c|c|c|} \hline 1 & 2 & 3 \\ \hline \end{array}.$$

Then, using the natural isomorphism $i : T^3(L \otimes A) \simeq T^3(L) \otimes T^3(A)$ and a projection $p : T^3(L \otimes A) \rightarrow \wedge^3(L \otimes A)$, one can decompose the third exterior power of a tensor product as follows:

$$\begin{aligned} \wedge^3(L \otimes A) &= p \circ (e \times e) \circ i(T^3(L \otimes A)) \\ &\simeq \wedge^3(L) \otimes S^3(A) \oplus \begin{array}{|c|c|} \hline 1 & 2 \\ \hline 3 & \\ \hline \end{array} (L) \otimes \begin{array}{|c|c|} \hline 1 & 3 \\ \hline 2 & \\ \hline \end{array} (A) + \begin{array}{|c|c|} \hline 1 & 3 \\ \hline 2 & \\ \hline \end{array} (L) \otimes \begin{array}{|c|c|} \hline 1 & 2 \\ \hline 3 & \\ \hline \end{array} (A) \oplus S^3(L) \otimes \wedge^3(A) \end{aligned}$$

(all other components appearing in $T^3(L) \otimes T^3(A)$ vanish under projection). One directly verifies that

$$\begin{array}{|c|c|} \hline 1 & 2 \\ \hline 3 & \\ \hline \end{array} \times \begin{array}{|c|c|} \hline 1 & 3 \\ \hline 2 & \\ \hline \end{array} (u) = 0 \quad \text{iff} \quad \begin{array}{|c|c|} \hline 1 & 3 \\ \hline 2 & \\ \hline \end{array} \times \begin{array}{|c|c|} \hline 1 & 2 \\ \hline 3 & \\ \hline \end{array} (u) = 0 \quad \text{for each } u \in \wedge^3(L \otimes A).$$

Hence we get a (noncanonical) isomorphism:

$$\wedge^3(L \otimes A) \simeq \wedge^3(L) \otimes S^3(A) \oplus \begin{array}{|c|c|} \hline 1 & 3 \\ \hline 2 & \\ \hline \end{array} (L) \otimes \begin{array}{|c|c|} \hline 1 & 2 \\ \hline 3 & \\ \hline \end{array} (A) \oplus S^3(L) \otimes \wedge^3(A).$$

Passing to $Hom(-, A \otimes V) \simeq Hom(-, M) \otimes Hom(-, V)$, one gets:

$$(3.2) \quad \begin{aligned} C^3(L \otimes A, M \otimes V) \\ \simeq C^3(L, M) \otimes S^3(A, V) \oplus Y^3(L, M) \otimes \tilde{Y}^3(A, V) \oplus S^3(L, M) \otimes C^3(A, V) \end{aligned}$$

where $Y^3(L, M) = Hom(\begin{array}{|c|c|} \hline 1 & 3 \\ \hline 2 & \\ \hline \end{array} (L), M)$, $\tilde{Y}^3(A, V) = Hom(\begin{array}{|c|c|} \hline 1 & 2 \\ \hline 3 & \\ \hline \end{array} (A), V)$.

According to (3.1)–(3.2) one can decompose H^2 as

$$(3.3) \quad H^2(L \otimes A, M \otimes V) = (H^2)' \oplus (H^2)''$$

where $(H^2)'$ are the classes of cocycles lying in $C^2(L, M) \otimes S^2(A, V)$ and $(H^2)''$ are the classes of cocycles of the form $\Phi + \Psi$, where $\Phi \in S^2(L, M) \otimes C^2(A, V)$, $\Psi \in C^2(L, M) \otimes S^2(A, V)$, $\Phi \neq 0$. We will compute $(H^2)'$ and obtain some particular

results on $(H^2)''$ (actually $(H^2)'$ and $(H^2)''$ are limit terms of a certain spectral sequence; see §4).

The differentials of the low degree in the piece of the standard Chevalley-Eilenberg complex $C^1(L \otimes A, M \otimes V) \xrightarrow{d^1} C^2(L \otimes A, M \otimes V) \xrightarrow{d^2} C^3(L \otimes A, M \otimes V)$ can be decomposed as follows:

$$d^1 = d_1 + d_2$$

$$d^2 = \sum_{\substack{1 \leq i \leq 2 \\ 1 \leq j \leq 3}} d_{ij}$$

where

$$\begin{aligned} d_1 &: C^1(L, M) \otimes C^1(A, V) \rightarrow C^2(L, M) \otimes S^2(A, V) \\ d_2 &: C^1(L, M) \otimes C^1(A, V) \rightarrow S^2(L, M) \otimes C^2(A, V) \\ d_{11} &: C^2(L, M) \otimes S^2(A, V) \rightarrow C^3(L, M) \otimes S^3(A, V) \\ d_{12} &: C^2(L, M) \otimes S^2(A, V) \rightarrow Y^3(L, M) \otimes \tilde{Y}^3(A, V) \\ d_{13} &: C^2(L, M) \otimes S^2(A, V) \rightarrow S^3(L, M) \otimes C^3(A, V) \\ d_{21} &: S^2(L, M) \otimes C^2(A, V) \rightarrow C^3(L, M) \otimes S^3(A, V) \\ d_{22} &: S^2(L, M) \otimes C^2(A, V) \rightarrow Y^3(L, M) \otimes \tilde{Y}^3(A, V) \\ d_{23} &: S^2(L, M) \otimes C^2(A, V) \rightarrow S^3(L, M) \otimes C^3(A, V). \end{aligned}$$

Direct computations show:

$$\begin{aligned} d_1(\varphi \otimes \alpha)(x_1 \otimes a_1, x_2 \otimes a_2) &= \\ \frac{1}{2}(x_2 \bullet \varphi(x_1) - x_1 \bullet \varphi(x_2)) \otimes (a_1 \bullet \alpha(a_2) + a_2 \bullet \alpha(a_1)) - \varphi([x_1, x_2]) \otimes \alpha(a_1 a_2); \end{aligned}$$

$$\begin{aligned} d_2(\varphi \otimes \alpha)(x_1 \otimes a_1, x_2 \otimes a_2) &= \\ \frac{1}{2}(x_1 \bullet \varphi(x_2) + x_2 \bullet \varphi(x_1)) \otimes (a_2 \bullet \alpha(a_1) - a_1 \bullet \alpha(a_2)); \end{aligned}$$

$$\begin{aligned} d_{11}(\varphi \otimes \alpha)(x_1 \otimes a_1, x_2 \otimes a_2, x_3 \otimes a_3) &= \\ \frac{1}{3}(\varphi([x_1, x_2], x_3) + \curvearrowright) \otimes (\alpha(a_1 a_2, a_3) + \curvearrowright) \\ - \frac{1}{3}(x_1 \bullet \varphi(x_2, x_3) + \curvearrowright) \otimes (a_1 \bullet \alpha(a_2, a_3) + \curvearrowright); \end{aligned}$$

$$\begin{aligned} d_{12}(\varphi \otimes \alpha)(x_1 \otimes a_1, x_2 \otimes a_2, x_3 \otimes a_3) &= \\ (2\varphi([x_1, x_2], x_3) + \varphi([x_1, x_3], x_2) - \varphi([x_2, x_3], x_1)) \otimes (\alpha(a_1 a_2, a_3) - \alpha(a_2 a_3, a_1)) \\ + (-x_1 \bullet \varphi(x_2, x_3) + x_2 \bullet \varphi(x_1, x_3) + 2x_3 \bullet \varphi(x_1, x_2)) \otimes (a_1 \bullet \alpha(a_2, a_3) - a_3 \bullet \alpha(a_1, a_2)); \end{aligned}$$

$$d_{13}(\varphi \otimes \alpha)(x_1 \otimes a_1, x_2 \otimes a_2, x_3 \otimes a_3) = 0;$$

$$\begin{aligned} d_{22}(\varphi \otimes \alpha)(x_1 \otimes a_1, x_2 \otimes a_2, x_3 \otimes a_3) &= \\ (2\varphi([x_1, x_2], x_3) + \varphi([x_1, x_3], x_2) - \varphi([x_2, x_3], x_1)) \otimes (\alpha(a_1 a_2, a_3) - \alpha(a_2 a_3, a_1)) \\ + (-x_1 \bullet \varphi(x_2, x_3) + x_2 \bullet \varphi(x_1, x_3)) \otimes (a_1 \bullet \alpha(a_2, a_3) + a_3 \bullet \alpha(a_1, a_2) + 2a_2 \bullet \alpha(a_1, a_3)); \end{aligned}$$

$$d_{23}(\varphi \otimes \alpha)(x_1 \otimes a_1, x_2 \otimes a_2, x_3 \otimes a_3) = \frac{1}{3}(x_1 \bullet \varphi(x_2, x_3) + \curvearrowright) \otimes (a_1 \bullet \alpha(a_2, a_3) + \curvearrowright)$$

(the absence of d_{21} in this list is merely a technical matter: on the relevant stage of computations, it will be convenient to use the entire differential d rather than d_{21}).

Now the reader should prepare for a bunch of tedious and cumbersome definitions. We apologize for this, but our excuse is that all this stuff provides building blocks for $H^2(L \otimes A, M \otimes V)$ and one can hardly imagine that it may be defined in a simpler way. Taking a glance at the expressions below, one can believe that the general formula for $H^n(L \otimes A, M \otimes V)$ hardly exists – if it does, one should give correct n -dimensional generalizations of definitions below (in a few cases this is evident – like Harrison or cyclic cohomology, but in most it is not).

Definitions.

- (i) Define $d^{[1]}, d^\bullet : Hom(L^{\otimes 2}, M) \rightarrow Hom(L^{\otimes 3}, M)$ as follows:

$$\begin{aligned} d^{[1]}\varphi(x, y, z) &= \varphi([x, y], z) + \curvearrowright \\ d^\bullet\varphi(x, y, z) &= x \bullet \varphi(y, z) + \curvearrowright. \end{aligned}$$

- (ii) Define $\wp, D : Hom(A^{\otimes 2}, V) \rightarrow Hom(A^{\otimes 3}, V)$ as follows:

$$\begin{aligned} \wp\alpha(a, b, c) &= \alpha(ab, c) + \curvearrowright \\ D\alpha(a, b, c) &= a \bullet \alpha(b, c) + \curvearrowright. \end{aligned}$$

- (iii) $\mathcal{B}(L, M) = \{\varphi \in C^2(L, M) \mid \varphi([x, y], z) + z \bullet \varphi(x, y) = 0; d^{[1]}\varphi(x, y, z) = 0\}$.

- (iv) $Q^2(L, M) = \{d\psi \mid \psi \in Hom(L, M); x \bullet \psi(y) = y \bullet \psi(x)\};$
 $H_M^2(L) = (Z^2(L, M^L) + Q^2(L, M))/Q^2(L, M)$.

- (v) $\mathcal{K}(L, M) = \{\varphi \in C^2(L, M) \mid d^{[1]}\varphi(x, y, z) = 2x \bullet \varphi(y, z)\};$
 $\mathcal{J}(L, M) = \{\varphi \in C^2(L, M) \mid \varphi(x, y) = \psi([x, y]) - \frac{1}{2}x \bullet \psi(y) + \frac{1}{2}y \bullet \psi(x) \text{ for } \psi \in Hom(L, M)\};$
 $\mathcal{H}(L, M) = (\mathcal{K}(L, M) + \mathcal{J}(L, M))/\mathcal{J}(L, M)$.

- (vi) $\mathcal{X}(L, M) = \{\varphi \in C^2(L, M) \mid 2\varphi([x, y], z) = z \bullet \varphi(x, y);$
 $\varphi([x, y], z) = \varphi([z, x], y)\}$.

- (vii) $\mathcal{J}(L, M) = \{\varphi \in C^2(L, M) \mid 3\varphi([x, y], z) = 2z \bullet \varphi(x, y); \varphi([x, y], z) = \varphi([z, x], y)\}$.

- (viii) $Poor_-(L, M) = \{\varphi \in C^2(L, M^L) \mid \varphi([L, L], L) = 0\};$
 $Poor_+(L, M) = \{\varphi \in S^2(L, M^L) \mid \varphi([L, L], L) = 0\}$.

- (ix) $Sym^2(L, M) = \{\varphi \in S^2(L, M) \mid x \bullet \varphi(y, z) = y \bullet \varphi(x, z)\};$
 $SB^2(L, M) = \{\varphi \in S^2(L, M) \mid \varphi(x, y) = x \bullet \psi(y) + y \bullet \psi(x) \text{ for } \psi \in Hom(L, M)\};$
 $SH^2(L, M) = (Sym^2(L, M) + SB^2(L, M))/SB^2(L, M)$.

- (x) Define an action of L on $Hom(L^{\otimes 2}, M)$ via

$$z \circ \varphi(x, y) = z \bullet \varphi(x, y) + \varphi([x, z], y) + \varphi(x, [y, z]).$$

$$S^2(L, M) = \{\varphi \in S^2(L, M)^L \mid \varphi([x, y], z) + \curvearrowright = 0\}.$$

- (xi) $D(A, V) = \{\beta \in Hom(A, V) \mid \beta(abc) = a \bullet \beta(bc) - bc \bullet \beta(a) + \curvearrowright\}$.

- (xii) $HC^1(A, V) = \{\alpha \in C^2(A, V) \mid \wp\alpha = 0\}$.

- (xiii) $\mathcal{C}^2(A, V) = \{\alpha \in C^2(A, V) \mid \alpha(ac, b) - \alpha(bc, a) + a \bullet \alpha(b, c) - b \bullet \alpha(a, c) + 2c \bullet \alpha(a, b) = 0\}$.
- (xiv) $\mathcal{P}_-(A, V) = \{\alpha \in C^2(A, V) \mid \alpha(ab, c) = a \bullet \alpha(b, c) + b \bullet \alpha(a, c)\}$;
 $\mathcal{P}_+(A, V) = \{\alpha \in S^2(A, V) \mid \alpha(ab, c) = a \bullet \alpha(b, c) + b \bullet \alpha(a, c)\}$.
- (xv) $\mathcal{A}(A, V) = \{\alpha \in S^2(A, V) \mid 2D\alpha = \wp\alpha\}$.

Spaces defined in (xi), (xv) are relevant to evaluation of $Ker d_{11}$ (Lemma 3.2), spaces defined in (iii)–(viii), (xiv) are relevant to evaluation of $Ker d_{11} \cap Ker d_{12}$ (see (3.6)), spaces defined in (ix) are relevant to computation for particular case when L is abelian (Proposition 3.5), and spaces defined in (x), (xii)–(xiii) are relevant to computation of the relative cohomology group $H^2(L \otimes A; L, M \otimes V)$ (Proposition 3.8).

Remarks.

- (i) d (the Chevalley-Eilenberg differential) $= d^{[1]} + d^\bullet$.
- (ii) As $B^2(L, M^L) \subseteq Q^2(L, M)$, there is a surjection

$$H^2(L) \otimes M^L \rightarrow H_M^2(L).$$

- (iii) If $V = K$, $HC^1(A, V)$ is just the first-order cyclic cohomology $HC^1(A)$.
- (iv) The following relations hold:

$$\begin{aligned} Poor_-(L, M) &\subseteq \mathcal{B}(L, M) \subseteq Z^2(L, M) \\ \mathcal{B}(L, M) \cap Z^2(L, M^L) &= Poor_-(L, M) \\ S^2(L, M) \cap S^2(L, M^L)^L &= Poor_+(L, M) \\ \mathcal{C}^2(A, V) \cap HC^1(A, V) &= \mathcal{P}_-(A, V) \\ \mathcal{Z}^2(A, V) \cap \mathcal{A}(A, V) &= \mathcal{P}_+(A, V) \\ Der(A, V) &\subseteq D(A, V). \end{aligned}$$

Proposition 3.1.

$$\begin{aligned} (H^2)' &\simeq H^2(L, M) \otimes V \oplus H_M^2(L) \otimes \frac{Hom(A, V)}{V \oplus Der(A, V)} \oplus \mathcal{H}(L, M) \otimes Der(A, V) \\ &\oplus \mathcal{B}(L, M) \otimes \frac{Har^2(A, V)}{\mathcal{P}_+(A, V)} \oplus C^2(L, M)^L \otimes \mathcal{P}_+(A, V) \\ &\oplus \mathcal{X}(L, M) \otimes \frac{\mathcal{A}(A, V)}{\mathcal{P}_+(A, V)} \oplus \mathcal{J}(L, M) \otimes \frac{D(A, V)}{Der(A, V)} \\ &\oplus Poor_-(L, M) \otimes \frac{S^2(A, V)}{Hom(A, V) + D(A, V) + Har^2(A, V) + \mathcal{A}(A, V)}. \end{aligned}$$

Each cocycle which lies in $C^2(L, M) \otimes S^2(A, V)$ is a linear combination of cocycles of the eight following types (which correspond respectively to the direct summands in the isomorphism):

- (i) $x \otimes a \wedge y \otimes b \mapsto \varphi(x, y) \otimes ab \bullet v$, $\varphi \in Z^2(L, M), v \in V$
- (ii) $x \otimes a \wedge y \otimes b \mapsto \varphi(x, y) \otimes \beta(ab)$, $\varphi \in Z^2(L, M^L), \beta \in Hom(A, V)$
- (iii) as in (ii) with $\varphi \in \mathcal{X}(L, M), \beta \in Der(A, V)$
- (iv) $x \otimes a \wedge y \otimes b \mapsto \varphi(x, y) \otimes \alpha(a, b)$, $\varphi \in \mathcal{B}(L, M), \alpha \in \mathcal{Z}^2(A, V)$
- (v) as in (iv) with $\varphi \in C^2(L, M)^L, \alpha \in \mathcal{P}_+(A, V)$
- (vi) as in (iv) with $\varphi \in \mathcal{X}(L, M), \alpha \in \mathcal{A}(A, V)$

- (vii) $x \otimes a \wedge y \otimes b \mapsto \varphi(x, y) \otimes (3a \bullet \beta(b) + 3b \bullet \beta(a) - 2\beta(ab)), \varphi \in \mathcal{H}(L, M), \beta \in D(A, V)$
- (viii) *as in (iv) with $\varphi \in \text{Poor}_-(L, M), \alpha \in S^2(A, V)$.*

Proof. We have

$$(3.4) \quad (H^2)' = \frac{\text{Ker } d_{11} \cap \text{Ker } d_{12}}{\text{Im } d_1}.$$

We compute the relevant spaces in the subsequent series of lemmas.

Lemma 3.2.

$$\begin{aligned} \text{Ker } d_{11} &= Z^2(L, M) \otimes V \\ &+ \{\varphi \in C^2(L, M) \mid 2d^{\llbracket} \varphi + d^{\bullet} \varphi = 0\} \otimes \mathcal{A}(A, V) \\ &+ \{\varphi \in C^2(L, M) \mid 3d^{\llbracket} \varphi + 2d^{\bullet} \varphi = 0\} \otimes D(A, V) \\ &+ \{\varphi \in C^2(L, M) \mid d^{\llbracket} \varphi = d^{\bullet} \varphi = 0\} \otimes S^2(A, V). \end{aligned}$$

Proof. Substituting $a_1 = a_2 = a_3 = 1$ into the equation $d_{11}\Phi = 0$ (as usual, $\Phi = \sum_{i \in I} \varphi_i \otimes \alpha_i$), one derives the equality

$$(3.5) \quad \sum_{i \in I} d\varphi_i(x_1, x_2, x_3) \otimes \alpha_i(1, 1) = 0$$

and a decomposition $I = I_1 \cup I_2$ with $d\varphi_i = 0$ for $i \in I_1$ and $\alpha_i(1, 1) = 0$ for $i \in I_2$.

Substituting then $a_2 = a_3 = 1$ into the same equation, one gets

$$\sum_{i \in I} (3d^{\llbracket} \varphi_i + 2d^{\bullet} \varphi_i) \otimes (\alpha_i(1, a_1) - a_1 \bullet \alpha_i(1, 1)) = 0$$

and by Lemma 1.1 there is a decomposition $I = I_{11} \cup I_{12} \cup I_{21} \cup I_{22}$ with

$$\begin{aligned} d\varphi_i &= 0; 3d^{\llbracket} \varphi_i + 2d^{\bullet} \varphi_i = 0, & i \in I_{11} \\ d\varphi_i &= 0; \alpha_i(1, a) = \alpha_i(1, 1), & i \in I_{12} \\ 3d^{\llbracket} \varphi_i + 2d^{\bullet} \varphi_i &= 0; \alpha_i(1, 1) = 0, & i \in I_{21} \\ \alpha_i(1, 1) &= 0; \alpha_i(1, a) = a \bullet \alpha_i(1, 1), & i \in I_{22}. \end{aligned}$$

Obviously $d\varphi_i = d^{\bullet} \varphi_i = 0$ for $i \in I_{11}$, and these components lie in $\text{Ker } d_{11}$, and $\alpha_i(1, a) = 0$ for $i \in I_{22}$.

Further, substituting $a_3 = 1$ in our equation, we get

$$\begin{aligned} \sum_{i \in I} (2d^{\llbracket} \varphi_i + d^{\bullet} \varphi_i) \otimes (\alpha_i(a_1, a_2) - 3a_1 \bullet \alpha_i(1, a_2) - 3a_2 \bullet \alpha_i(1, a_1) \\ + 2\alpha_i(1, a_1 a_2) + 3a_1 a_2 \bullet \alpha_i(1, 1)) = 0. \end{aligned}$$

In order to apply Lemma 1.1 again, we join the sets I_{12} and I_{22} (with the common defining condition $\alpha_i(1, a) = a \bullet \alpha_i(1, 1)$) and obtain a decomposition $I = I'_1 \cup I'_2 \cup I'_3 \cup I'_4$ such that

$$\begin{aligned} 2d^{\llbracket} \varphi_i + d^{\bullet} \varphi_i &= 0; \alpha_i(1, a) = a \bullet \alpha_i(1, 1), & i \in I'_1 \\ \alpha_i(a, b) &= ab \bullet \alpha_i(1, 1), & i \in I'_2 \\ d^{\llbracket} \varphi_i &= d^{\bullet} \varphi_i = 0; \alpha_i(1, 1) = 0, & i \in I'_3 \\ 3d^{\llbracket} \varphi_i + 2d^{\bullet} \varphi_i &= 0; \alpha_i(a, b) = 3a \bullet \alpha_i(1, b) + 3b \bullet \alpha_i(1, a) - 2\alpha_i(1, ab), & i \in I'_4. \end{aligned}$$

Note that components $\varphi_i \otimes \alpha_i$ with $i \in I'_3$ included in those with $i \in I_{11}$ (and lie in $\text{Ker } d_{11}$).

Now, since the contribution of terms with $i \in I'_4$ to the left side of (3.5) vanish, applying Lemma 1.1 again, we obtain a decomposition $I'_1 \cup I'_2 = I'_{11} \cup I'_{12} \cup I'_{21} \cup I'_{22}$ such that

$$\begin{aligned} 2d^{[1]}\varphi_i + d^\bullet\varphi_i &= 0; \alpha_i(1, a) = 0, & i \in I'_{12} \\ d\varphi_i &= 0; \alpha_i(a, b) = ab \bullet \alpha_i(1, 1), & i \in I'_{21} \end{aligned}$$

and the two remaining types of component do not contribute to the whole picture: those with indices from I'_{11} satisfy $d^{[1]}\varphi_i = d^\bullet\varphi_i = 0$ which covered by previous cases, and those with indices from I'_{22} vanish, as $\alpha_i(a, b) = ab \bullet \alpha_i(1, 1) = 0$. Moreover, components with indices from I'_{21} lie in $\text{Ker } d_{11}$.

The remaining part of the equation $d_{11}\Phi = 0$ now reads:

$$\begin{aligned} \sum_{i \in I'_{12} \cup I'_4} d^{[1]}\varphi_i(x_1, x_2, x_3) \otimes (\wp\alpha(a_1, a_2, a_3) - 2D\alpha_i(a_2, a_3) \\ + 3a_1a_2 \bullet \alpha_i(1, a_3) - a_3 \bullet \alpha_i(1, a_1a_2) + \curvearrowright) = 0. \end{aligned}$$

Applying Lemma 1.1 again, and noting that the vanishing of the first tensor factor in the last equality yields the already considered case $d^{[1]}\varphi_i = d^\bullet\varphi_i = 0$, we obtain that the second tensor factor vanishes for all $i \in I'_{12} \cup I'_4$.

Consequently, we receive two types of components $\varphi_i \otimes \alpha_i$ lying in $\text{Ker } d_{11}$:

$$2d^{[1]}\varphi_i + d^\bullet\varphi_i = 0; \wp\alpha_i = 2D\alpha_i$$

and

$$3d^{[1]}\varphi_i + 2d^\bullet\varphi_i = 0; \wp\alpha_i = \frac{3}{2}D\alpha_i; \alpha_i \text{ satisfies the defining condition for } i \in I'_4.$$

The last two conditions imposed on α_i imply $\alpha_i(1, \cdot) \in D(A, V)$.

Summarizing all this, we receive the statement of the Lemma. ■

Lemma 3.3.

$$\begin{aligned} \text{Ker } d_{12} &= C^2(L, M) \otimes V + \{\varphi \in C^2(L, M) \mid x \bullet \varphi(y, z) = z \bullet \varphi(x, y)\} \otimes \text{Hom}(A, V) \\ &+ \{\varphi \in C^2(L, M) \mid \varphi([x, y], z) - \varphi([y, z], x) - x \bullet \varphi(y, z) + z \bullet \varphi(x, y) = 0\} \otimes \mathcal{Z}^2(A, V) \\ &+ \{\varphi \in C^2(L, M) \mid x \bullet \varphi(y, z) = z \bullet \varphi(x, y); \varphi([x, y], z) = \varphi([y, z], x)\} \otimes S^2(A, V). \end{aligned}$$

Proof. Let $\Phi = \sum_{i \in I} \varphi_i \otimes \alpha_i \in \text{Ker } d_{12}$. Substituting $a_2 = 1$ in the equation $d_{12}\Phi = 0$, one gets:

$$\begin{aligned} \sum_{i \in I} (-x_1 \bullet \varphi_i(x_2, x_3) + x_2 \bullet \varphi_i(x_1, x_3) + 2x_3 \bullet \varphi_i(x_1, x_2)) \\ \otimes (a_1 \bullet \alpha_i(1, a_3) - a_3 \bullet \alpha_i(1, a_1)) = 0. \end{aligned}$$

Hence we have a decomposition $I = I_1 \cup I_2$ such that for $i \in I_1$ the first tensor factor in the last expression vanishes, and for $i \in I_2$ the second one vanishes. Elementary transformations show that

$$\begin{aligned} x \bullet \varphi_i(y, z) &= z \bullet \varphi_i(x, y), & i \in I_1 \\ \alpha_i(1, a) &= a \bullet \alpha_i(1, 1), & i \in I_2. \end{aligned}$$

Then substituting $a_3 = 1$ in the same initial equation $d_{12}\Phi = 0$, one gets

$$\begin{aligned} & \sum_{i \in I} (2\varphi_i([x_1, x_2], x_3) + \varphi_i([x_1, x_3], x_2) - \varphi_i([x_2, x_3], x_1) \\ & \quad - x_1 \bullet \varphi_i(x_2, x_3) + x_2 \bullet \varphi_i(x_1, x_3) + 2x_3 \bullet \varphi_i(x_1, x_2)) \\ & \quad \otimes (\alpha_i(1, a_1 a_2) - \alpha_i(a_1, a_2)) = 0. \end{aligned}$$

Applying Lemma 1.1 and the fact that the vanishing of the first tensor factor here is equivalent to the condition $\varphi_i(x, y), z) - \varphi_i([y, z], x) - x \bullet \varphi_i(y, z) + z \bullet \varphi_i(x, y) = 0$, we get a decomposition $I = I_{11} \cup I_{12} \cup I_{21} \cup I_{22}$ such that

$$\begin{aligned} x \bullet \varphi_i(y, z) &= z \bullet \varphi_i(x, y); \varphi_i([x, y], z) = \varphi_i([y, z], x), & i \in I_{11} \\ x \bullet \varphi_i(y, z) &= z \bullet \varphi_i(x, y); \alpha_i(a, b) = \alpha_i(1, ab), & i \in I_{12} \\ \varphi_i([x, y], z) - \varphi_i([y, z], x) - x \bullet \varphi_i(y, z) + z \bullet \varphi_i(x, y) &= 0; \quad \alpha_i(1, a) = a \bullet \alpha_i(1, 1), \\ & & i \in I_{21} \\ \alpha_i(a, b) &= ab \bullet \alpha_i(1, 1), & i \in I_{22}. \end{aligned}$$

It is easy to see that the components $\varphi_i \otimes \alpha_i$ with indices belonging to I_{11} , I_{12} and I_{22} , already lie in $\text{Ker } d_{12}$.

The remaining part of the equation $d_{12}\Phi = 0$ becomes:

$$\sum_{i \in I_{21}} (2\varphi_i([x_1, x_2], x_3) + \varphi_i([x_1, x_3], x_2) - \varphi_i([x_2, x_3], x_1)) \otimes \delta \alpha_i(a_1, a_2, a_3) = 0$$

where δ is Harrison(=Hochschild) differential. Thus there is a decomposition $I_{21} = I'_1 \cup I'_2$, where φ_i for $i \in I'_1$ satisfy the same relations as for $i \in I_{11}$, and $\alpha_i \in \mathcal{Z}^2(A, V)$, $i \in I'_2$.

Putting all these computations together yields the desired formula (the four summands there correspond to the defining conditions for I_{22} , I_{12} , I'_2 and I_{11} respectively; the sum, in general, is not direct). ■

Elementary but tedious transformations of expressions entering in defining conditions of summands of $\text{Ker } d_{11}$ and $\text{Ker } d_{12}$, allow to write their intersection as the following direct sum:

$$\begin{aligned} (3.6) \quad \text{Ker } d_{11} \cap \text{Ker } d_{12} &\simeq Z^2(L, M) \otimes V \oplus Z^2(L, M^L) \otimes \frac{\text{Hom}(A, V)}{V \oplus \text{Der}(A, V)} \\ &\oplus \mathcal{K}(L, M) \otimes \text{Der}(A, V) \oplus \mathcal{B}(L, M) \otimes \frac{\mathcal{Z}^2(A, V)}{\mathcal{P}_+(A, V)} \\ &\oplus C^2(L, M)^L \otimes \mathcal{P}_+(A, V) \oplus \mathcal{X}(L, M) \otimes \frac{\mathcal{A}(A, V)}{\mathcal{P}_+(A, V)} \\ &\oplus \mathcal{T}(L, M) \otimes \frac{D(A, V)}{\text{Der}(A, V)} \\ &\oplus \text{Poor}_-(L, M) \otimes \frac{S^2(A, V)}{\text{Hom}(A, V) + D(A, V) + \mathcal{Z}^2(A, V) + \mathcal{A}(A, V)}. \end{aligned}$$

According to (3.4), to compute $(H^2)'$ we must consider the equation $\Phi = d_1\Psi$ for $\Phi \in \text{Ker } d_{11} \cap \text{Ker } d_{12}$ and $\Psi \in \text{Hom}(L \otimes A, M \otimes V)$, what is equivalent to elucidation of all possible cohomological dependencies between obtained classes of cocycles.

Lemma 3.4. *Let $\{\varphi_i\}$ be cohomologically independent cocycles in $Z^2(L, M)$, $\{\theta_i\}$ be cocycles in $Z^2(L, M^L)$ independent modulo $Q^2(L, M)$, $\{\kappa_i\}$ be elements of $\mathcal{K}(L, M)$ independent modulo $\mathcal{T}(L, M)$, $\{\varepsilon_i\}$ be linearly independent cocycles in $\mathcal{B}(L, M)$, $\{\rho_i\}$ be linearly independent elements in $C^2(L, M)^L$, $\{\chi_i\}$ be linearly independent elements in $\mathcal{X}(L, M)$, $\{\tau_i\}$ be linearly independent elements in $\mathcal{T}(L, M)$, $\{\xi_i\}$ be linearly independent cocycles in $\text{Poor}_-(L, M)$, $\{v_j\}$ be linearly independent elements in V , $\{\delta_j\}$ be linearly independent derivations in $\text{Der}(A, V)$, $\{\beta_j\}$ be mappings in $D(A, V)$ independent modulo $\text{Der}(A, V)$, $\{\gamma_j\}$ be mappings in $\text{Hom}(A, V)$ independent both modulo $\text{Der}(A, V)$ and modulo mappings $a \mapsto a \bullet v$ for all $v \in V$, $\{F_j\}$ be cocycles in $\mathcal{Z}^2(A, V)$ independent both cohomologically and modulo $\mathcal{P}_+(A, V)$, $\{P_j\}$ be linearly independent elements in $\mathcal{P}_+(A, V)$, $\{A_j\}$ be elements in $\mathcal{A}(A, V)$ independent modulo $\mathcal{P}_+(A, V)$, $\{G_j\}$ be mappings in $S^2(A, V)$ independent modulo mappings $a \wedge b \mapsto \gamma(ab)$ for all $\gamma \in \text{Hom}(A, V)$, modulo mappings $a \wedge b \mapsto 3a \bullet \beta(b) + 3b \bullet \beta(a) - 2\beta(ab)$ for all $\beta \in D(A, V)$, and modulo $\mathcal{Z}^2(A, V) + \mathcal{A}(A, V)$.*

Then elements of $\text{Ker } d_{11} \cap \text{Ker } d_{12}$: $\varphi_i \otimes (R_{v_j} \circ m)$, $\theta_i \otimes (\gamma_j \circ m)$, $\kappa_i \otimes (\delta_j \circ m)$, $\varepsilon_i \otimes F_j$, $\rho_i \otimes P_j$, $\chi_i \otimes A_j$, $\tau_i \otimes (3\delta\beta_j - \beta_j \circ m)$, $\xi_i \otimes G_j$ (m stands for multiplication in algebra A and R_v is an element in $\text{Hom}(A, V)$ defined by $a \mapsto a \bullet v$), are independent modulo $\text{Im } d_1$.

Proof. We must to prove that if

$$\begin{aligned}
& \sum \varphi_i(x, y) \otimes ab \bullet v_j + \sum \theta_i(x, y) \otimes \gamma_j(ab) + \sum \kappa_i(x, y) \otimes \delta_j(ab) \\
& + \sum \varepsilon_i(x, y) \otimes F_j(a, b) + \sum \rho_i(x, y) \otimes P_j(a, b) + \sum \chi_i(x, y) \otimes A_j(a, b) \\
& + \sum \tau_i(x, y) \otimes (3a \bullet \beta_j(b) + 3b \bullet \beta_j(a) - 2\beta_j(ab)) + \sum \xi_i(x, y) \otimes G_j(a, b) \\
(3.7) \quad & = \sum_{i \in I} \psi_i([x, y]) \otimes \alpha_i(ab) + \frac{1}{2}(-x \bullet \psi_i(y) + y \bullet \psi_i(x)) \otimes (a \bullet \alpha_i(b) + b \bullet \alpha_i(a))
\end{aligned}$$

for some $\sum_{i \in I} \psi_i \otimes \alpha_i \in \text{Hom}(L, M) \otimes \text{Hom}(A, V)$ (the right side here is the generic element in $\text{Im } d_1$), then all left side terms vanish.

One has $\delta_j(1) = \beta_j(1) = P_j(1, a) = A_j(1, a) = 0$ and one may assume that $\gamma_j(1) = F_j(1, a) = G_j(1, a) = 0$. Substitute in (3.7) $a = b = 1$:

$$\sum \varphi_i(x, y) \otimes v_j = \sum_{i \in I} d\psi_i(x, y) \otimes \alpha_i(1).$$

As φ_i 's are cohomologically independent and v_j 's are linearly independent, the last equality implies that all summands $\varphi_i(x, y) \otimes v_j$ vanish and there is a decomposition $I = I_1 \cup I_2$ with $d\psi_i = 0$ for $i \in I_1$ and $\alpha_i(1) = 0$ for $i \in I_2$.

Now substitute in (3.7) $b = 1$:

$$\begin{aligned}
(3.8) \quad & \sum \theta_i(x, y) \otimes \gamma_j(a) + \sum \kappa_i(x, y) \otimes \delta_j(a) + \sum \tau_i(x, y) \otimes \beta_j(a) \\
& = \sum_{i \in I} (\psi_i([x, y]) + \frac{1}{2}(-x \bullet \psi_i(y) + y \bullet \psi_i(x))) \otimes (\alpha_i(a) - a \bullet \alpha_i(1)).
\end{aligned}$$

Substituting (3.8) in (3.7), one gets:

$$\begin{aligned} & \sum \varepsilon_i(x, y) \otimes F_j(a, b) + \sum \rho_i(x, y) \otimes P_j(a, b) + \sum \chi_i(x, y) \otimes A_j(a, b) \\ & + 3 \sum \tau_i(x, y) \otimes \delta\beta_j(a, b) + \sum \xi_i(x, y) \otimes G_j(a, b) \\ & = \frac{1}{2} \sum (-x \bullet \psi_i(y) + y \bullet \psi_i(x)) \otimes (\delta\alpha_i(a, b) - ab \bullet \alpha_i(1)). \end{aligned}$$

The independency conditions of Lemma imply that all summands in the left side vanish, and due to Lemma 1.1, for element $\sum_{i \in I} \psi_i \otimes \beta_i$ there exists a decomposition $I = I_{11} \cup I_{12} \cup I_{21} \cup I_{22}$ with

$$\begin{aligned} d\psi_i &= 0; \quad x \bullet \psi_i(y) = y \bullet \psi_i(x), \quad i \in I_{11} \\ d\psi_i &= 0; \quad \delta\alpha_i(a, b) = ab \bullet \alpha_i(1), \quad i \in I_{12} \\ x \bullet \psi_i(y) &= y \bullet \psi_i(x); \quad \alpha_i(1) = 0, \quad i \in I_{21} \\ \alpha_i(1) &= 0; \quad \delta\alpha_i(a, b) = ab \bullet \alpha_i(1), \quad i \in I_{22}. \end{aligned}$$

Denoting $\alpha'_i(a) = \alpha_i(a) - a \bullet \alpha_i(1)$ for $i \in I_{12}$, we get $\alpha'_i \in \text{Der}(A, V)$.

Substituting all this information back into (3.8), one finally obtains

$$\begin{aligned} & \sum \theta_i(x, y) \otimes \gamma_j(a) + \sum \kappa_i(x, y) \otimes \delta_j(a) \\ & = \frac{1}{2} \sum_{i \in I_{12}} \psi_i([x, y]) \otimes (\alpha_i(a) - a \bullet \alpha_i(1)) + \sum_{i \in I_{21}} \psi_i([x, y]) \otimes \alpha_i(a) \\ & \quad + \sum_{i \in I_{22}} (\psi_i([x, y]) + \frac{1}{2}(-x \bullet \psi_i(y) + y \bullet \psi_i(x))) \otimes \alpha_i(a). \end{aligned}$$

The independency conditions of Lemma imply that all terms appearing in the last equality vanish, and the desired assertion follows. ■

Conclusion of the proof of proposition 3.1.

Lemma 3.3 implies that

$$\begin{aligned} \text{Im } d_1 &\simeq B^2(L, M) \otimes V \oplus (Q^2(L, M) \cap Z^2(L, M^L)) \otimes \frac{\text{Hom}(A, V)}{V \oplus \text{Der}(A, V)} \\ &\quad \oplus (\mathcal{T}(L, M) \cap \mathcal{K}(L, M)) \otimes \text{Der}(A, V) \oplus \mathcal{B}(L, M) \otimes \text{Der}(A, V) \end{aligned}$$

which together with (3.6) entails the asserted isomorphism. ■

Now we turn to computation of the second summand in (3.3), $(H^2)''$.

We are unable to compute it in general (and are doubtful about the existence of a closed general formula for $(H^2)''$) and restrict ourselves with consideration of two particular cases (in both of them it turns out that $(H^2)''$ coincides with the classes of cocycles lying in $S^2(L, M) \otimes C^2(A, V)$).

Proposition 3.5. *Suppose L is abelian. Then*

$$\begin{aligned} (H^2)'' &\simeq S^2(L, M^L) \otimes \frac{C^2(A, V)}{\{a \bullet \beta(b) - b \bullet \beta(a) \mid \beta \in \text{Hom}(A, V)\}} \\ &\quad \oplus SH^2(L, M) \otimes \{a \bullet \beta(b) - b \bullet \beta(a) \mid \beta \in \text{Hom}(A, V)\}. \end{aligned}$$

First we establish a lemma valid in the general situation (when L is not necessarily abelian).

Lemma 3.6.

- (i) $Ker d_{23} = Ker d^\bullet \otimes C^2(A, V)$
 $+ S^2(L, M) \otimes \{\alpha \in C^2(A, V) \mid \alpha(a, b) = a \bullet \beta(b) - b \bullet \beta(a)\}$;
- (ii) $Im d_2 = \{\varphi \in S^2(L, M) \mid \varphi(x, y) = x \bullet \psi(y) + y \bullet \psi(x)\}$
 $\otimes \{\alpha \in C^2(A, V) \mid \alpha(a, b) = a \bullet \beta(b) - b \bullet \beta(a)\}$.

Proof. The only thing which perhaps needs a proof here is the equality

$$Ker D = \{\alpha \in C^2(A, V) \mid \alpha(a, b) = a \bullet \beta(b) - b \bullet \beta(a)\}.$$

This is achieved by appropriate substitution of 1's. ■

Proof of proposition 3.5. Let $\Phi = \sum_{i \in I} \varphi_i \otimes \alpha_i \in Ker d_{23}$, with a decomposition on the set of indices $I = I_1 \cup I_2$ such that

$$\begin{aligned} d^\bullet \varphi_i(x_1, x_2, x_3) &= 0, \quad i \in I_1 \\ \alpha_i(a, b) &= -a \bullet \beta_i(b) + b \bullet \beta_i(a), \quad i \in I_2. \end{aligned}$$

By Lemma 3.6(i), we may assume also that elements $\alpha_i, i \in I_1$ are independent modulo $\{a \bullet \beta(b) - b \bullet \beta(a)\}$ and hence $\alpha_i(1, a) = 0$ for each $i \in I_1$.

Suppose there is $\Psi = \sum_{i \in I'} \varphi'_i \otimes \alpha'_i \in C^2(L, M) \otimes S^2(A, V)$ such that the class of $\Phi - \Psi$ belongs to $(H^2)''$. This particularly means that $d_{22}\Phi = d_{12}\Psi$:

$$\begin{aligned} &\sum_{i \in I} (-x_1 \bullet \varphi_i(x_2, x_3) + x_2 \bullet \varphi_i(x_1, x_3)) \\ &\quad \otimes (a_1 \bullet \alpha_i(a_2, a_3) + a_3 \bullet \alpha_i(a_1, a_2) + 2a_2 \bullet \alpha_i(a_1, a_3)) \\ &= \sum_{i \in I'} (-x_1 \bullet \varphi'_i(x_2, x_3) + x_2 \bullet \varphi'_i(x_1, x_3) + 2x_3 \bullet \varphi'_i(x_1, x_2)) \\ &\quad \otimes (a_1 \bullet \alpha'_i(a_2, a_3) - a_3 \bullet \alpha'_i(a_1, a_2)). \end{aligned}$$

Substituting here $a_2 = 1$, one gets

$$\begin{aligned} &2 \sum_{i \in I_1} (-x_1 \bullet \varphi_i(x_2, x_3) + x_2 \bullet \varphi_i(x_1, x_3)) \otimes \alpha_i(a_1, a_3) \\ &\quad + 3 \sum_{i \in I_1} (-x_1 \bullet \varphi_i(x_2, x_3) + x_2 \bullet \varphi_i(x_1, x_3)) \otimes \alpha_i(a_1, a_3) \\ &= \sum_{i \in I'} (-x_1 \bullet \varphi'_i(x_2, x_3) + x_2 \bullet \varphi'_i(x_1, x_3) + 2x_3 \bullet \varphi'_i(x_1, x_2)) \\ &\quad \otimes (a_1 \bullet \alpha'_i(1, a_3) - a_3 \bullet \alpha'_i(1, a_1)). \end{aligned}$$

Hence, due to the independency condition imposed on $\alpha_i, i \in I_1$,

$$(3.9) \quad -x_1 \bullet \varphi_i(x_2, x_3) + x_2 \bullet \varphi_i(x_1, x_3) = 0, \quad i \in I_1.$$

This, together with condition $\varphi_i \in Ker d^\bullet$, evidently implies $\varphi_i(L, L) \subseteq M^L$ for each $i \in I_1$. Note that the terms from $S^2(L, M^L) \otimes C^2(A, V)$ lie in $Z^2(L \otimes A, M \otimes V)$.

Now write the cocycle equation for elements from $S^2(L, M) \otimes \{a \bullet \beta(b) - b \bullet \beta(a)\}$:

$$\begin{aligned} \sum_{i \in I_2} (x_1 \bullet \varphi_i(x_2, x_3) \otimes (a_1 a_2 \bullet \beta_i(a_3) - a_1 a_3 \bullet \beta_i(a_2)) \\ + x_2 \bullet \varphi_i(x_1, x_3) \otimes (-a_1 a_2 \bullet \beta_i(a_3) + a_2 a_3 \bullet \beta_i(a_1)) \\ + x_3 \bullet \varphi_i(x_1, x_2) \otimes (a_1 a_3 \bullet \beta_i(a_2) - a_2 a_3 \bullet \beta_i(a_1))) = 0. \end{aligned}$$

Substituting $a_2 = a_3 = 1$, we get

$$\sum_{i \in I_2} (x_2 \bullet \varphi_i(x_1, x_3) - x_3 \bullet \varphi_i(x_1, x_2)) \otimes (\beta_i(a_1) - a_1 \bullet \beta_i(1)) = 0.$$

As the vanishing of the second tensor factor here leads to the vanishing of the whole α_i , we see that the condition (3.9) holds also in this case, i.e. for all $i \in I_2$. Conversely, if (3.9) holds, then the cocycle equation is satisfied. Thus the space of cocycles in $Z^2(L \otimes A, M \otimes V)$ whose cohomology classes lie in $(H^2)''$, coincides with

$$\begin{aligned} S^2(L, M^L) \otimes \frac{C^2(A, V)}{\{a \bullet \beta(b) - b \bullet \beta(a) \mid \beta \in \text{Hom}(A, V)\}} \\ \oplus \frac{\text{Sym}^2(L, M) + SB^2(L, M)}{SB^2(L, M)} \otimes \{a \bullet \beta(b) - b \bullet \beta(a) \mid \beta \in \text{Hom}(A, V)\} \end{aligned}$$

(note that we can always take $\Psi = 0$).

To conclude the proof, one can observe that all these cocycles are cohomologically independent. This is proved in a pretty standard way, as in Lemma 3.4. ■

Summarizing Proposition 3.1 (for the case when L is abelian) and Proposition 3.5, we obtain

Theorem 3.7. *Let L be an abelian Lie algebra. Then*

$$\begin{aligned} H^2(L \otimes A, M \otimes V) \simeq H^2(L, M) \otimes V \oplus \mathcal{H}(L, M) \otimes \text{Der}(A, V) \\ \oplus C^2(L, M^L) \otimes \frac{S^2(A, V)}{V \oplus \text{Der}(A, V)} \\ \oplus S^2(L, M^L) \otimes \frac{C^2(A, V)}{\{a \bullet \beta(b) - b \bullet \beta(a) \mid \beta \in \text{Hom}(A, V)\}} \\ \oplus SH^2(L, M) \otimes \{a \bullet \beta(b) - b \bullet \beta(a) \mid \beta \in \text{Hom}(A, V)\}. \end{aligned}$$

Each cocycle in $Z^2(L \otimes A, M \otimes V)$ is a linear combination of cocycles of the four following types (which correspond respectively to the first, the sum of the second and the third, the fourth and the fifth summands in the isomorphism):

- (i) $x \otimes a \wedge y \otimes b \mapsto \varphi(x, y) \otimes ab \bullet v$, $\varphi \in Z^2(L, M), v \in V$
- (ii) $x \otimes a \wedge y \otimes b \mapsto \varphi(x, y) \otimes \alpha(a, b)$, $\varphi \in C^2(L, M^L), \alpha \in S^2(A, V)$
- (iii) as in (ii) with $\varphi \in S^2(L, M^L)$, $\alpha \in C^2(A, V)$
- (iv) $x \otimes a \wedge y \otimes b \mapsto \varphi(x, y) \otimes (a \bullet \beta(b) - b \bullet \beta(a))$, $\varphi \in \text{Sym}^2(L, M), \beta \in \text{Hom}(A, V)$.

Remark. It is easy to see that if L is abelian, then there is an inclusion $\mathcal{H}(L, M) \subseteq H^2(L, M)$ ($\mathcal{H}(L, M)$ consists of classes of cocycles taking values in M^L). Consequently, singling out the appropriate terms from the first three direct summands in the isomorphism above, we obtain a direct summand $\mathcal{H}(L, M) \otimes S^2(A, V)$.

Now we want to perform another particular computations of the second cohomology group, namely, to compute the relative cohomology $H^2(L \otimes A; L, M \otimes V)$.

One may easily see that all constructions can be restricted to the relative complex $Hom(\wedge^*(L \otimes A/K1), M \otimes V)$ with a single (but simplifying the matters greatly) difference that all mappings from $C^3(A, V)$, $Y^3(A, V)$ and $S^3(A, V)$ vanish whenever one of its arguments is 1.

We will use notations $(H_L^2)'$ and $(H_L^2)''$ to denote the corresponding components of $H^2(L \otimes A; L, M \otimes V)$.

Proposition 3.8.

$$\begin{aligned} (H_L^2)'' &\simeq S^2(L, M^L)^L \otimes HC^1(A, V) \oplus S^2(L, M) \otimes \frac{\mathcal{C}^2(L, M)}{\mathcal{P}_-(A, V)} \\ &\oplus \frac{S^2(L, M)^L}{S^2(L, M^L)^L} \otimes \mathcal{P}_-(A, V) \oplus Poor_+(L, M) \otimes \frac{C^2(A, V)}{HC^1(A, V) + \mathcal{C}^2(A, V)} \end{aligned}$$

Proof. The proof goes by the same scheme as of Proposition 3.1. By Lemma 3.6, $Ker d_{23} = Ker d^\bullet \otimes C^2(A, V)$ (as the second tensor factor in the second component there vanishes in this case).

The condition $d_{22}\Phi = d_{12}\Psi$ for $\Phi = \sum \varphi_i \otimes \alpha_i \in Ker d_{23}$ and $\Psi = \sum \varphi'_i \otimes \alpha'_i \in C^2(L, M) \otimes S^2(L, M)$ reads:

$$\begin{aligned} &\sum_{i \in I} (2\varphi_i([x_1, x_2], x_3) + \varphi_i([x_1, x_3], x_2) - \varphi_i([x_2, x_3], x_1)) \\ &\quad \otimes (\alpha_i(a_1 a_2, a_3) - \alpha_i(a_2 a_3, a_1)) \\ &+ (-x_1 \bullet \varphi_i(x_2, x_3) + x_2 \bullet \varphi_i(x_1, x_3)) \otimes (a_1 \bullet \alpha_i(a_2, a_3) + a_3 \bullet \alpha_i(a_1, a_2) + 2a_2 \bullet \alpha_i(a_1, a_3)) \\ &= \sum_{i \in I'} (2\varphi'_i([x_1, x_2], x_3) + \varphi'_i([x_1, x_3], x_2) - \varphi'_i([x_2, x_3], x_1)) \\ &\quad \otimes (\alpha'_i(a_1 a_2, a_3) - \alpha'_i(a_2 a_3, a_1)) \\ &+ (-x_1 \bullet \varphi'_i(x_2, x_3) + x_2 \bullet \varphi'_i(x_1, x_3) + 2x_3 \bullet \varphi'_i(x_1, x_2)) \\ &\quad \otimes (a_1 \bullet \alpha'_i(a_2, a_3) - a_2 \bullet \alpha'_i(a_1, a_2)). \end{aligned}$$

Substituting here $a_2 = 1$, we receive (remember about vanishing of all α 's if one of arguments is 1):

$$\begin{aligned} &\sum_{i \in I} (2\varphi_i([x_1, x_2], x_3) + \varphi_i([x_1, x_3], x_2) - \varphi_i([x_2, x_3], x_1)) \\ &\quad - x_1 \bullet \varphi_i(x_2, x_3) + x_2 \bullet \varphi_i(x_1, x_3)) \otimes \alpha_i(a_1, a_3) = 0. \end{aligned}$$

This entails

$$(3.10) \quad \begin{aligned} &2\varphi_i([x_1, x_2], x_3) + \varphi_i([x_1, x_3], x_2) - \varphi_i([x_2, x_3], x_1) \\ &\quad - x_1 \bullet \varphi_i(x_2, x_3) + x_2 \bullet \varphi_i(x_1, x_3) = 0, \quad i \in I. \end{aligned}$$

Since $\varphi \in Ker d^\bullet$,

$$(3.11) \quad x_1 \bullet \varphi_i(x_2, x_3) + x_3 \bullet \varphi_i(x_1, x_3) + x_3 \bullet \varphi_i(x_1, x_2) = 0, \quad i \in I.$$

With the help of elementary transformations, (3.10) and (3.11) yield

$$\varphi_i([x_1, x_3], x_2) + \varphi_i([x_2, x_3], x_1) + x_3 \bullet \varphi_i(x_1, x_2) = 0$$

or, in other words, $\varphi_i \in S^2(L, M)^L$ for each $i \in I$.

Now writing the cocycle equation for $\sum_{i \in I} \varphi_i \otimes \alpha_i \in S^2(L, M)^L \otimes C^2(A, V)$, one gets

$$\begin{aligned} & \sum_{i \in I} \varphi_i([x_1, x_2], x_3) \otimes (\alpha_i(a_1 a_2, a_3) - a_1 \bullet \alpha_i(a_2, a_3) - a_2 \bullet \alpha_i(a_1, a_3)) \\ & + \varphi_i([x_1, x_3], x_2) \otimes (-\alpha_i(a_1 a_3, a_2) - a_1 \bullet \alpha_i(a_2, a_3) + a_3 \bullet \alpha_i(a_1, a_2)) \\ & + \varphi_i([x_2, x_3], x_1) \otimes (\alpha_i(a_2 a_3, a_1) + a_2 \bullet \alpha_i(a_1, a_3) + a_3 \bullet \alpha_i(a_1, a_2)) = 0. \end{aligned}$$

Antisymmetrize this expression with respect to a_1, a_2 :

$$\begin{aligned} & \sum_{i \in I} (\varphi_i([x_1, x_3], x_2) + \varphi_i([x_2, x_3], x_1)) \otimes (-\alpha_i(a_1 a_3, a_2) + \alpha_i(a_2 a_3, a_1) \\ & - a_1 \bullet \alpha_i(a_2, a_3) + a_2 \bullet \alpha_i(a_1, a_3) + 2a_3 \bullet \alpha_i(a_1, a_2)) = 0. \end{aligned}$$

Consequently, we have a decomposition $I = I_1 \cup I_2$ with

$$(3.12) \quad \varphi_i([x_1, x_3], x_2) + \varphi_i([x_2, x_3], x_1) = 0, \quad i \in I_1$$

$$(3.13) \quad \alpha_i \in \mathcal{C}^2(A, V), \quad i \in I_2.$$

Note that (3.12) together with condition $\varphi_i \in S^2(L, M)^L$ imply $\varphi_i(L, L) \subseteq M^L$ for $i \in I_1$. Applying to the condition (3.13) the symmetrizer $e - (13) + (123)$, we get

$$(3.14) \quad \begin{aligned} & a_1 \bullet \alpha_i(a_2, a_3) + a_2 \bullet \alpha_i(a_1, a_3) \\ & = \frac{1}{3}(2\alpha_i(a_1 a_2, a_3) - \alpha_i(a_2 a_3, a_1) - \alpha_i(a_1 a_3, a_2)), \quad i \in I_2. \end{aligned}$$

Taking into account (3.12)–(3.14), the cocycle equation can be rewritten as

$$\begin{aligned} & \sum_{i \in I} (\varphi_i([x_1, x_2], x_3) - \varphi_i([x_1, x_3], x_2) + \varphi_i([x_2, x_3], x_1)) \\ & \otimes (\alpha_i(a_1 a_2, a_3) + \alpha_i(a_1 a_3, a_2) + \alpha_i(a_2 a_3, a_1)) = 0. \end{aligned}$$

By Lemma 1.1, there is a decomposition $I = I_{11} \cup I_{12} \cup I_{21} \cup I_{22}$ such that

$$\varphi_i([x, y], z) = \varphi_i(x, [y, z]); \varphi_i([x, y], z) + \curvearrowright = 0, \quad i \in I_{11}$$

$$\varphi_i([x, y], z) = \varphi_i(x, [y, z]); \alpha_i \in HC^1(A, V), \quad i \in I_{12}$$

$$\varphi_i([x, y], z) + \curvearrowright = 0; \alpha_i \in \mathcal{C}^2(A, V), \quad i \in I_{21}$$

$$\alpha_i \in \mathcal{C}^2(A, V) \cap HC^1(A, V), \quad i \in I_{22}.$$

Evidently $\varphi_i([L, L], L) = 0$ for $i \in I_{11}$ and $\alpha_i \in \mathcal{P}_-(A, V)$ for $i \in I_{22}$. All these four types of components are cocycles in $Z^2(L \otimes A, M \otimes V)$.

Therefore, the space of cocycles whose cohomology classes lie in $(H_L^2)''$, looks as follows:

$$\begin{aligned} {}^L Z^{02} \simeq & S^2(L, M^L)^L \otimes HC^1(A, V) + S^2(L, M) \otimes \mathcal{C}^2(A, V) + S^2(L, M)^L \otimes \mathcal{P}_-(A, V) \\ & + Poor_+(L, M) \otimes C^2(A, V). \end{aligned}$$

(the four summands here correspond to the components indexed by I_{12}, I_{21}, I_{22} and I_{11} respectively; note that we may let $\Psi = 0$ again).

Rewriting this as a direct sum, we get:

$$\begin{aligned} S^2(L, M^L)^L \otimes HC^1(A, V) \oplus \mathfrak{S}^2(L, M) \otimes \frac{\mathcal{C}^2(A, V)}{\mathcal{P}_-(A, V)} \\ \oplus \frac{S^2(L, M)^L}{S^2(L, M^L)^L} \otimes \mathcal{P}_-(A, V) \oplus Poor_+(L, M) \otimes \frac{C^2(A, V)}{HC^1(A, V) + \mathcal{C}^2(A, V)}. \end{aligned}$$

And finally, one may show in the same fashion as previously, that all these cocycles are cohomologically independent, and the assertion of the Lemma follows. ■

Summarizing Propositions 3.1 and 3.8, we obtain:

$$H^2(L \otimes A; L, M \otimes V) \simeq (H_L^2)' \oplus (H_L^2)''$$

where

$$\begin{aligned} (H_L^2)' \simeq \mathfrak{B}(L, M) \otimes \frac{Har^2(A, V)}{\mathcal{P}_+(A, V)} \oplus C^2(L, M)^L \otimes \mathcal{P}_+(A, V) \oplus \mathfrak{X}(L, M) \otimes \frac{\mathcal{A}(A, V)}{\mathcal{P}_+(A, V)} \\ \oplus Poor_-(L, M) \otimes \frac{S^2(A, V)}{Hom(A, V) + D(A, V) + Har^2(A, V) + \mathcal{A}(A, V)} \end{aligned}$$

and $(H_L^2)''$ is described by Proposition 3.8.

We conclude this section with enumeration (for the case of general L) of all possible cocycles of rank 1, i.e. those which can be written in a form $\varphi \otimes \alpha \in Hom(L^{\otimes 2}, M) \otimes Hom(A^{\otimes 2}, V)$.

In view of (3.3), Propositions 3.1 and 3.8, one suffices to consider cocycles of rank 1 whose cohomology classes lie in $(H^2)''$ and which are independent modulo $(H_L^2)''$. Let us denote this space of cocycles as Z'' .

Proposition 3.9. *Each cocycle in Z'' is cohomologous to a sum of cocycles of the following two types:*

- (i) $x \otimes a \wedge y \otimes b \mapsto \varphi(x, y) \otimes (a \bullet \beta(b) - b \bullet \beta(a)), \varphi \in Sym^2(L, M), \varphi([L, L], L) = 0, \beta \in Hom(A, V)$
- (ii) *as in (i) with $\varphi \in S^2(L, M), 2\varphi([x, y], z) = x \bullet \varphi(y, z) - y \bullet \varphi(x, z), \beta \in Der(A, V)$.*

Proof. Mainly repetition of previous arguments. ■

Therefore, there are, in general, 13 types of cohomologically independent cocycles of rank 1 (7 coming from Proposition 3.1 + 4 coming from Proposition 3.8 + 2 coming from Proposition 3.9). Of course, in particular cases some of these types of cocycles may vanish.

We see that for $H^2(L \otimes A; L, M \otimes V)$ and for $H^2(L \otimes A, M \otimes V)$, L abelian, one possible to choose a basis consisting of rank 1 cocycles. In general this is, however, not true. The case of $H^2(W_1(n) \otimes A, W_1(n) \otimes A)$, where $W_1(n)$ is a Zassenhaus algebra of positive characteristic, treated in [Z], shows that there are cocycles of rank 2 not cohomologous to (a sum of) cocycles of rank 1.

4. A SKETCH OF A SPECTRAL SEQUENCE

The computations performed in preceding sections can be described (and generalized) in terms of a certain spectral sequence. Let us indicate briefly the main idea (the full treatment with further applications hopefully will appear elsewhere).

One has a Cauchy formula

$$\wedge^n(L \otimes A) \simeq \bigoplus_{\lambda \vdash n} Y_\lambda(L) \otimes Y_{\lambda^\sim}^C(A)$$

where Y_λ is a Schur functor associated with the Young diagram λ , and λ^\sim is a Young diagram obtained from λ by interchanging its rows and columns.

Applying to it the functor $Hom(-, M \otimes V) \simeq Hom(-, M) \otimes Hom(-, V)$, one gets a decomposition of the underlying modules in the Chevalley-Eilenberg complex:


$$(4.1) \quad C^n(L \otimes A, M \otimes V) \simeq \sum_{\lambda \vdash n} C_\lambda(L, M) \otimes C_{\lambda^\sim}(A, V)$$

where $C_\lambda(U, W) = Hom(Y_\lambda(U), W)$. The two extreme terms here are $C^n(L, M) \otimes S^n(A, V)$ and $S^n(L, M) \otimes C^n(A, V)$.

So each differential $d : C^n(L \otimes A, M \otimes V) \rightarrow C^{n+1}(L \otimes A, M \otimes V)$ in the Chevalley-Eilenberg complex decomposes according (4.1) into components


$$d_\lambda^{\lambda'} : C_{\lambda'}(L, M) \otimes C_{\lambda'^\sim}(A, V) \rightarrow C_\lambda(L, M) \otimes C_{\lambda^\sim}(A, V)$$

for each pair $\lambda' \vdash n, \lambda \vdash (n + 1)$. Therefore the following graph of all Young diagrams



may be interpreted in the following way: each Young diagram λ of size n designate a module $C_\lambda(L, M) \otimes C_{\lambda^\sim}(A, V)$ and an arrow from λ' to λ represents $d_\lambda^{\lambda'}$.

One can prove that nonzero arrows $d_\lambda^{\lambda'}$ are exactly the following: all arrows going “from right to left” and those going “from left to right” for which either λ' is a vertical diagram of size n and λ is of size $n + 1$ and of the following form:



or λ' is included in λ .

Using this, we can define a decreasing nonnegative filtration $F^k C^*$ on the complex $(C^*(L \otimes A, M \otimes V), d)$ as the sum of all $C_\lambda(L, M) \otimes C_{\lambda^\sim}(A, V)$ with λ belonging to a “closure” under nonzero arrows of a single vertical diagram of size $k + 1$.

Now we may consider a (first quadrant) spectral sequence $\{E_r^{**}, d_r\}$ associated with this filtration. Since the filtration is finite in each degree, the spectral sequence converges to the desired cohomology group $H^*(L \otimes A, M \otimes V)$.

Then $E_\infty^{20} = 0$ and $(H^2)'$ and $(H^2)''$ from §3 are nothing else than E_∞^{11} and E_∞^{02} respectively.

5. STRUCTURE FUNCTIONS

In this section we show how result from §3 may be applied to a geometric problem of calculation of structure functions on manifolds of loops on compact Hermitian symmetric spaces.

Let us recall briefly necessary notions and results. Let M be a (complex) manifold endowed with a G -structure (so G is a complex Lie group). *Structure functions* (SF for short) are sections of certain vector bundles over M with fiber over a point $m \in M$. Their importance stems from the fact that they constitute the complete set of obstructions to integrability (= possibility of local flattening) of a given G -structure. A remarkable fact is that SF admit a purely algebraic description. Starting with $\mathfrak{g}_{-1} = T_m(M)$, the tangent space at a point m , and $\mathfrak{g}_0 = Lie G$, one may construct, via apparatus of Cartan prolongations, a graded Lie algebra $\mathfrak{g} = \bigoplus_{i \geq -1} \mathfrak{g}_i$. Namely,

$$(5.1) \quad \mathfrak{g}_i = \{X \in Hom(\mathfrak{g}_{-1}, \mathfrak{g}_{i-1}) \mid [X(v), w] = [X(w), v] \quad \forall v, w \in \mathfrak{g}_{-1}\}$$

for $i > 0$.

For a such graded Lie algebra one may define the *Spencer cohomology groups* $H_{\mathfrak{g}_0}^{pq}(\mathfrak{g}_{-1})$. Then the space of SF of order k , i.e. obstructions to identification of the k th infinitesimal neighborhood of a point $m \in M$ with that of a point of the flat manifold, is isomorphic to the group $H_{\mathfrak{g}_0}^{k2}(\mathfrak{g}_{-1})$. Note that since $H^2(\mathfrak{g}_{-1}, \mathfrak{g}) = \bigoplus_{k \geq 1} H_{\mathfrak{g}_0}^{k2}(\mathfrak{g}_{-1})$, to compute SF for a given G -structure on a manifold, one merely need to evaluate the usual Chevalley-Eilenberg cohomology group $H^2(\mathfrak{g}_{-1}, \mathfrak{g})$ of an abelian Lie algebra \mathfrak{g}_{-1} with coefficients in the whole \mathfrak{g} and to identify SF of order k with the graded component $\{\bar{\varphi} \in H^2(\mathfrak{g}_{-1}, \mathfrak{g}) \mid Im \varphi \subseteq \mathfrak{g}_{k-2}\}$, $k \geq 1$. We refer for the classical text [St], Chapter VII, for details.

One of the nice examples of manifolds endowed with a G -structure are (irreducible) compact Hermitian symmetric spaces (CHSS). There are naturally distinguishable two cases: $\text{rank} M > 1$ and $\text{rank} M = 1$.

If $\text{rank} M > 1$, \mathfrak{g} turns out to be a classical simple Lie algebra with a grading of length 1: $\mathfrak{g} = \mathfrak{g}_{-1} \oplus \mathfrak{g}_0 \oplus \mathfrak{g}_1$ (particularly, Cartan prolongations of order > 1 vanish, so we have SF of orders 1, 2 and 3 only). We will refer for this case as a *general case*.

The case of $\text{rank} M = 1$ is exceptional in a sense. The Cartan prolong of $(\mathbb{C}e_{-1}, \mathbb{C}e_0)$ (with $[e_{-1}, e_0] = e_{-1}$) is the infinite-dimensional algebra

$$W_1 = \langle e_i \mid i \geq -1; [e_i, e_j] = (j - i)e_{i+j} \rangle,$$

a version of the famous Witt algebra. $\mathbb{C}e_{-1} \oplus \mathbb{C}e_0 \oplus \mathbb{C}e_1 \simeq sl(2)$ is a subalgebra of W_1 , and we will refer for this case as a *$sl(2)$ case*.

Corresponding SF (in both cases) were determined by Goncharov [G]. During the last decade, there was a big amount of activity by Grozman, Leites, Poletaeva and Serganova in determining SF of various classes of (super)manifolds and G -structures on them (see e.g. [GL] and [LPS] with a transitive closure of references therein).

What we are going to do here, is to describe SF of manifolds M^{S^1} of loops with values in a (finite-dimensional) CHSS. The group G here is formally no longer a Lie group, but its infinite-dimensional analogue, the group of loops, and a corresponding Lie algebra is a current Lie algebra $\mathfrak{g} \otimes \mathbb{C}[t, t^{-1}]$ with a grading inherited from \mathfrak{g} :

$$\begin{aligned} \mathfrak{g} \otimes \mathbb{C}[t, t^{-1}] &= \mathfrak{g}_{-1} \otimes \mathbb{C}[t, t^{-1}] \oplus \mathfrak{g}_0 \otimes \mathbb{C}[t, t^{-1}] \oplus \mathfrak{g}_1 \otimes \mathbb{C}[t, t^{-1}] \\ W_1 \otimes \mathbb{C}[t, t^{-1}] &= \bigoplus_{i \geq -1} e_i \otimes \mathbb{C}[t, t^{-1}] \end{aligned}$$

for the general and $sl(2)$ cases respectively.

The last statement follows from the next simple but handy observation:

Proposition 5.1. *Let $\bigoplus_{i \geq -1} \mathfrak{g}_i$ be a Cartan prolongation of a pair $(\mathfrak{g}_{-1}, \mathfrak{g}_0)$, $\mathfrak{g}_{-1}, \mathfrak{g}_0$ are finite-dimensional. Then $\bigoplus_{i \geq -1} (\mathfrak{g}_i \otimes A)$ is a Cartan prolongation of a pair $(\mathfrak{g}_{-1} \otimes A, \mathfrak{g}_0 \otimes A)$.*

Proof. Induction on i . As all \mathfrak{g}_i are finite-dimensional, an element $X \in \text{Hom}(\mathfrak{g}_{-1} \otimes A, \mathfrak{g}_{i-1} \otimes A)$ in inductive definition (5.1) of Cartan prolongation may be written in the form $\sum_{i \in I} \varphi_i \otimes \alpha_i$ for $\varphi_i \in \text{Hom}(\mathfrak{g}_{-1}, \mathfrak{g}_{i-1})$, $\alpha_i \in \text{End}(A)$. Then it goes as in proof of Theorem 2.1. ■

Thus, we shall obtain, so to speak, a “loopization” of Goncharov’s result.

In November 1993, the author was shown by Dmitry Leites handwritten notes by Elena Poletaeva on computation of SF of manifolds of loops corresponding to the $sl(2)$ case and the special case $\mathfrak{g} = sl(4)$ with graded components $\mathfrak{g}_{-1} = V \otimes V^*$, $\mathfrak{g}_0 = gl(2) \oplus gl(2) \oplus \mathbb{C}z$, $\mathfrak{g}_1 = V^* \otimes V$ where V is a standard 2-dimensional $gl(2)$ -module.

However, this never was published, the notes seem be lost, and 10 years later nobody from the involved parties cannot recollect the details.

So for the sake of completeness we reproduce the Poletaeva’s result for the $sl(2)$ case obtained with our method.

Theorem 5.2 (Poletaeva). *For a manifold M^{S^1} of loops with values in CHSS M of rank 1 (i.e. M is a complex projective space), the space of SF of order 1 is isomorphic to $e_{-1} \otimes C^2(\mathbb{C}[t, t^{-1}], \mathbb{C}[t, t^{-1}])$ and the space of SF of order $k > 1$ is isomorphic to $e_{k-2} \otimes (End(\mathbb{C}[t, t^{-1}])/\mathbb{C}[t, t^{-1}])$.*

The corresponding basic cocycles are given by

$$\begin{aligned} e_{-1} \otimes a \wedge e_{-1} \otimes b &\mapsto e_{-1} \otimes \alpha(a, b), \quad \alpha \in C^2(\mathbb{C}[t, t^{-1}], \mathbb{C}[t, t^{-1}]) \\ e_{-1} \otimes a \wedge e_{-1} \otimes b &\mapsto e_{k-2} \otimes (a\beta(b) - b\beta(a)), \quad \beta \in End(\mathbb{C}[t, t^{-1}]), k > 1. \end{aligned}$$

Proof. Our task is to compute $H^2(e_{-1} \otimes \mathbb{C}[t, t^{-1}], W_1 \otimes \mathbb{C}[t, t^{-1}])$. Substitute our specific data to the formula of Theorem 3.7. As $L = \mathbb{C}e_{-1}$ is one-dimensional, the first three terms of that formula, being skew-symmetric in L , vanish, and we get:

$$\begin{aligned} &H^2(e_{-1} \otimes \mathbb{C}[t, t^{-1}], W_1 \otimes \mathbb{C}[t, t^{-1}]) \\ &\simeq S^2(\mathbb{C}e_{-1}, W_1^{e_{-1}}) \otimes \frac{C^2(\mathbb{C}[t, t^{-1}], \mathbb{C}[t, t^{-1}])}{\{a\beta(b) - b\beta(a) \mid \beta \in End(\mathbb{C}[t, t^{-1}])\}} \\ &\quad \oplus SH^2(\mathbb{C}e_{-1}, W_1) \otimes \{a\beta(b) - b\beta(a) \mid \beta \in End(\mathbb{C}[t, t^{-1}])\}. \end{aligned}$$

Now, evidently $W_1^{e_{-1}} = \mathbb{C}e_{-1}$, $S^2(\mathbb{C}e_{-1}, \mathbb{C}e_{-1}) = \langle e_{-1} \vee e_{-1} \mapsto e_{-1} \rangle \simeq \mathbb{C}e_{-1}$, and $SH^2(\mathbb{C}e_{-1}, W_1) = S^2(\mathbb{C}e_{-1}, W_1) = \{e_{-1} \vee e_{-1} \mapsto u \mid u \in W_1\} \simeq W_1$.

Regrouping direct summands in an obvious way, splitting Chevalley-Eilenberg cohomology into graded Spencer cohomology components, and noting that the kernel of a mapping

$$\begin{aligned} T : End(\mathbb{C}[t, t^{-1}]) &\rightarrow C^2(\mathbb{C}[t, t^{-1}], \mathbb{C}[t, t^{-1}]) \\ \beta &\mapsto (T\beta)(a, b) = a\beta(b) - b\beta(a), \end{aligned}$$

coincides with multiplications by an element of $\mathbb{C}[t, t^{-1}]$, we get the statement of the Theorem. ■

Now turn to consideration of the general case.

Definitions.

- (i) SF (identified with the elements of the second cohomology group) generated by cocycles of the kind

$$x \otimes a \wedge y \otimes b \mapsto \varphi(x, y) \otimes abu$$

where φ is SF of CHSS and $u \in \mathbb{C}[t, t^{-1}]$, will be called *trivial*.

- (ii) SF generated by cocycles of the kind

$$x \otimes a \wedge y \otimes b \mapsto \varphi(x, y) \otimes \alpha(a, b)$$

where $\varphi \in C^2(\mathfrak{g}_{-1}, \mathfrak{g}_{-1})$, $\alpha \in S^2(\mathbb{C}[t, t^{-1}], \mathbb{C}[t, t^{-1}])$, will be called *almost trivial*.

- (iii) Define a symmetric analogue of $H_{\mathfrak{g}_0}^{1,2}(\mathfrak{g}_{-1})$, denoted as $SH_{\mathfrak{g}_0}^{1,2}(\mathfrak{g}_{-1})$, to be a factorspace

$$\frac{S^2(\mathfrak{g}_{-1}, \mathfrak{g}_{-1})}{\{\varphi \in S^2(\mathfrak{g}_{-1}, \mathfrak{g}_{-1}) \mid \varphi(x, y) = [x, \psi(y)] + [y, \psi(x)] \text{ for } \psi \in Hom(\mathfrak{g}_{-1}, \mathfrak{g}_0)\}}.$$

For convenience, we gather here the necessary properties of the grading $\mathfrak{g} = \mathfrak{g}_{-1} \oplus \mathfrak{g}_0 \oplus \mathfrak{g}_1$ of a simple classical Lie \mathbb{C} -algebra (cf., for example, [D] and references therein).

First,

$$(5.2) \quad \{x \in \mathfrak{g}_i \mid [x, \mathfrak{g}_{-1}] = 0\} = 0, \quad i = 0, 1.$$

Particularly, \mathfrak{g}_{-1} is a faithful \mathfrak{g}_0 -module.

Next, all such gradings may be obtained in the following way. Let R be a root system of \mathfrak{g} corresponding to a Cartan subalgebra \mathfrak{h} , B a basis of R , $\{h_\beta, e_\alpha \mid \beta \in B, \alpha \in R\}$ a Chevalley basis of \mathfrak{g} . Fix a root $\beta \in B$ such that β enters in decomposition of each root only with coefficients $-1, 0, 1$ (the existence of such root implies that R is not of type G_2, F_4 or E_8). Denote by $R_i, i = -1, 0, 1$, the set of roots in which β enters with coefficient i . Then

$$\mathfrak{g}_{-1} = \bigoplus_{\alpha \in R_{-1}} \mathbb{C}e_\alpha, \quad \mathfrak{g}_0 = \mathfrak{h} \oplus \bigoplus_{\alpha \in R_0} \mathbb{C}e_\alpha, \quad \mathfrak{g}_1 = \bigoplus_{\alpha \in R_1} \mathbb{C}e_\alpha.$$

Theorem 5.3. *For a manifold M^{S^1} of loops with values in CHSS M of rank > 1 (i.e. M is not a complex projective space), all SF of orders 2 and 3 are trivial. The space of SF of order 1 modulo almost trivial SF is isomorphic to*

$$\begin{aligned} B_{\mathfrak{g}_0}^{1,2}(\mathfrak{g}_{-1}) \otimes \frac{S^2(\mathbb{C}[t, t^{-1}], \mathbb{C}[t, t^{-1}])}{(\mathbb{C}1 \oplus \mathbb{C}\frac{d}{dt}) \otimes \mathbb{C}[t, t^{-1}]} \oplus S^2(\mathfrak{g}_{-1}, \mathfrak{g}_{-1}) \otimes \frac{C^2(\mathbb{C}[t, t^{-1}], \mathbb{C}[t, t^{-1}])}{\text{End}(\mathbb{C}[t, t^{-1}])} \\ \oplus SH_{\mathfrak{g}_0}^{1,2}(\mathfrak{g}_{-1}) \otimes \frac{\text{End}(\mathbb{C}[t, t^{-1}])}{\mathbb{C}[t, t^{-1}]} \end{aligned}$$

Remarks.

(i) $B_{\mathfrak{g}_0}^{1,2}(\mathfrak{g}_{-1})$ is the space of corresponding Spencer coboundaries, i.e. the space of mappings $\varphi \in C^2(\mathfrak{g}_{-1}, \mathfrak{g}_{-1})$ of the form $\varphi(x, y) = [x, \psi(y)] - [y, \psi(x)]$ for some $\psi \in \text{Hom}(\mathfrak{g}_{-1}, \mathfrak{g}_0)$.

(ii) Theorem 3.7 suggests the way in which denominator is imbedded into numerator in the factorspaces involving $\mathbb{C}[t, t^{-1}]$. In the first factorspace, an element $(\lambda 1 + \mu \frac{d}{dt})t^n$ corresponds to a mapping

$$\alpha(t^i, t^j) = \lambda t^{i+j+n} + \mu(i+j)t^{i+j+n-1}.$$

In the second one, a mapping $\beta(t^i) = \sum_n \lambda_{in} t^n$ corresponds to a mapping

$$\alpha(t^i, t^j) = \sum_n (\lambda_{j, n-i} - \lambda_{i, n-j}) t^n.$$

And in the third one, an element t^n corresponds to multiplication by t^n :

$$\beta(t^i) = t^{i+n}.$$

Proof. To compute $H^2(\mathfrak{g}_{-1} \otimes \mathbb{C}[t, t^{-1}], \mathfrak{g} \otimes \mathbb{C}[t, t^{-1}])$, we start as in the proof of Theorem 5.2. As it turns out that the concrete structure of the Laurent polynomial ring $\mathbb{C}[t, t^{-1}]$ is not important in our approach, for notational convenience we replace it by an arbitrary (associative commutative unital) algebra A .

Substitute our specific data to the formula of Theorem 3.7:

$$\begin{aligned}
 (5.3) \quad H^2(\mathfrak{g}_{-1} \otimes A, \mathfrak{g} \otimes A) &\simeq H^2(\mathfrak{g}_{-1}, \mathfrak{g}) \otimes A \oplus \mathcal{H}(\mathfrak{g}_{-1}, \mathfrak{g}) \otimes \text{Der}(A) \\
 &\oplus C^2(\mathfrak{g}_{-1}, \mathfrak{g}^{\mathfrak{g}_{-1}}) \otimes \frac{S^2(A, A)}{A \oplus \text{Der}(A)} \\
 &\oplus S^2(\mathfrak{g}_{-1}, \mathfrak{g}^{\mathfrak{g}_{-1}}) \otimes \frac{C^2(A, A)}{\{a\beta(b) - b\beta(a) \mid \beta \in \text{End}(A)\}} \\
 &\oplus SH^2(\mathfrak{g}_{-1}, \mathfrak{g}) \otimes \{a\beta(b) - b\beta(a) \mid \beta \in \text{End}(A)\}.
 \end{aligned}$$

The next technical lemma is devoted to determination of components appearing in this isomorphism.

Lemma 5.4.

- (i) $\mathfrak{g}^{\mathfrak{g}_{-1}} = \mathfrak{g}_{-1}$
- (ii) $\mathcal{H}(\mathfrak{g}_{-1}, \mathfrak{g}) = H_{\mathfrak{g}_0}^{1,2}(\mathfrak{g}_{-1})$
- (iii) $SH^2(\mathfrak{g}_{-1}, \mathfrak{g}) = SH_{\mathfrak{g}_0}^{1,2}(\mathfrak{g}_{-1})$.

Proof. (i) Evident in view of (5.2).

(ii) Follows from definitions of appropriate spaces, (5.2) and part (i).

(iii) According to the definition,

$$Sym^2(\mathfrak{g}_{-1}, \mathfrak{g}) = \{\varphi \in S^2(\mathfrak{g}_{-1}, \mathfrak{g}) \mid [x, \varphi(y, z)] = [y, \varphi(x, z)] \ \forall x, y, z \in \mathfrak{g}_{-1}\}.$$

We may decompose φ into the sum of three components $\varphi_{-1} + \varphi_0 + \varphi_1$, $\varphi_i \in Sym^2(\mathfrak{g}_{-1}, \mathfrak{g}_i) = Sym^2(\mathfrak{g}_{-1}, \mathfrak{g}) \cap S^2(\mathfrak{g}_{-1}, \mathfrak{g}_i)$.

One immediately seen that $Sym^2(\mathfrak{g}_{-1}, \mathfrak{g}_{-1}) = S^2(\mathfrak{g}_{-1}, \mathfrak{g}_{-1})$, and for each fixed $z \in \mathfrak{g}_{-1}$, $\varphi_i(\cdot, z)$ belongs to the $(i+1)$ st Cartan prolongation of the pair $(\mathfrak{g}_{-1}, \mathfrak{g}_0)$, $i = 0, 1$. Hence $Sym^2(\mathfrak{g}_{-1}, \mathfrak{g}_1) = 0$ and there is a bilinear mapping

$$\begin{aligned}
 F : Sym^2(\mathfrak{g}_{-1}, \mathfrak{g}_0) \times \mathfrak{g}_{-1} &\rightarrow \mathfrak{g}_1 \\
 \varphi_0(x, z) &= [x, F(\varphi_0, z)] \quad \forall x \in \mathfrak{g}_{-1}.
 \end{aligned}$$

But the symmetry of φ_0 implies that an element $F(\varphi_0, \cdot) \in Hom(\mathfrak{g}_{-1}, \mathfrak{g}_1)$ lies in the second Cartan prolongation of $(\mathfrak{g}_{-1}, \mathfrak{g}_0)$ and hence vanish. Thus $Sym^2(\mathfrak{g}_{-1}, \mathfrak{g}_0) = 0$.

Therefore, $Sym^2(\mathfrak{g}_{-1}, \mathfrak{g}) = S^2(\mathfrak{g}_{-1}, \mathfrak{g}_{-1})$ and the asserted equality follows. ■

Continuation of the proof of Theorem 5.3. Substituting the results of Lemma 5.4 to (5.3), and rearranging the summands as indicated in the Remark after Theorem 3.7, we obtain:

$$\begin{aligned}
 H^2(\mathfrak{g}_{-1} \otimes A, \mathfrak{g} \otimes A) &\simeq H_{\mathfrak{g}_0}^{1,2} \otimes S^2(A, A) \oplus (H_{\mathfrak{g}_0}^{2,2}(\mathfrak{g}_{-1}) \oplus H_{\mathfrak{g}_0}^{3,2}(\mathfrak{g}_{-1})) \otimes A \\
 &\oplus B_{\mathfrak{g}_0}^{1,2}(\mathfrak{g}_{-1}) \otimes \frac{S^2(A, A)}{A \oplus \text{Der}(A)} \\
 &\oplus S^2(\mathfrak{g}_{-1}, \mathfrak{g}_{-1}) \otimes \frac{C^2(A, A)}{\{a\beta(b) - b\beta(a) \mid \beta \in \text{End}(A)\}} \\
 &\oplus SH_{\mathfrak{g}_0}^{1,2}(\mathfrak{g}_{-1}) \otimes \{a\beta(b) - b\beta(a) \mid \beta \in \text{End}(A)\}.
 \end{aligned}$$

The first tensor product here consists of almost trivial SF of order 1, the second one consists of trivial SF of order 2 and 3, and the Theorem follows (well, after the final substitution $A = \mathbb{C}[t, t^{-1}]$). ■

Theorem 5.3 tells how to describe structure functions of a manifolds of loops with values in CHSS in terms of SF of underlying CHSS (Spencer cohomology groups), and the space $SH_{\mathfrak{g}_0}^{1,2}(\mathfrak{g}_{-1})$, which is a sort of a symmetric analogue of the Spencer cohomology group.

The thorough treatment of the latter symmetric analogue, including its calculation for all \mathfrak{g} 's, as well as related construction of a symmetric analogue of Cartan prolongation and some questions pertained to Jordan algebras, hopefully will appear elsewhere. Here we only briefly outline how $SH_{\mathfrak{g}_0}^{1,2}(\mathfrak{g}_{-1})$ can be determined in terms of a corresponding root system R .

Indeed, consider a mapping

$$T : Hom(\mathfrak{g}_{-1}, \mathfrak{g}_0) \rightarrow S^2(\mathfrak{g}_{-1}, \mathfrak{g}_{-1})$$

$$\psi(x) \mapsto (T\psi)(x, y) = [x, \psi(y)] + [y, \psi(x)].$$

The question of determining $SH_{\mathfrak{g}_0}^{1,2}(\mathfrak{g}_{-1})$ evidently reduces to evaluation of $Ker T$. Writing

$$\psi(e_r) = \sum_{\alpha \in B} \lambda_\alpha^r h_\alpha + \sum_{\alpha \in R_0} \mu_\alpha^r e_\alpha$$

for $r \in R_{-1}$ and certain parameters $\lambda_\alpha^r, \mu_\alpha^r \in \mathbb{C}$, one seen that the equality $[x, \psi(y)] + [y, \psi(x)] = 0$ is equivalent to the following three conditions:

$$\sum_{\alpha \in B} \lambda_\alpha^s r(h_\alpha) = \mu_{r-s}^r N_{s, r-s} \quad \forall r, s \in R_{-1} : r - s \in R_0$$

$$\sum_{\alpha \in B} \lambda_\alpha^s r(h_\alpha) = 0 \quad \forall r, s \in R_{-1} : r - s \notin R$$

$$\mu_\alpha^r N_{s, \alpha} = 0 \quad \forall r, s \in R_{-1}, \alpha \in R_0 : r - s \neq \alpha$$

which serve as (linear) defining relations for the space $Ker T$ and may be computed in each concrete case.

ACKNOWLEDGEMENTS

This note was basically written at the beginning of 1990s while being a Ph.D. student at Bar-Ilan University under the guidance of Steve Shnider.

Back to those days, my thanks due to Steve Shnider for support, attention and conceptual improvements of a preliminary version of manuscript, and to Dmitry Leites for numerous interesting conversations and sharing his tremendous archive of mathematical texts. Particularly, Leites drew my attention to the problem of computation of structure functions in the first place. I am also grateful to Alexander Feldman for facilitating access to some otherwise hardly available mathematical literature and to Elena Poletaeva for sending her recent reprints.

REFERENCES

- [B] R. E. Block, *Determination of the differentiably simple rings with a minimal ideal*, Ann. Math. **90** (1969), 433–459.
- [C] J. L. Cathelineau, *Homologie de degré trois d'algèbres de Lie simple déployées étendues à une algèbre commutative*, Enseign. Math. **33** (1987), 159–173.
- [D] M. Demazure, *Classification des algèbres de Lie filtrées*, Sémin. Bourbaki, Exposé 326, 1967.
- [G] A. B. Goncharov, *Infinitesimal structures related to Hermitian symmetric spaces*, Funct. Anal. Appl. **15** (1981), 221–223.

- [GL] P. Grozman and D. Leites, *From supergravity to ball bearings. Cohomology for physicists and engineers*, Supersymmetries and quantum symmetries (Dubna, 1997), Lecture Notes Phys. **524** (1999), 58–67.
- [H] A. Haddi, *Homologie de degré trois des algèbres de Lie étendues par une algèbre commutative*, Comptes Rendus Acad. Sci. Paris **321** (1995), 965–968.
- [LPS] D. Leites, E. Poletaeva and V. Serganova, *On Einstein equations on manifolds and supermanifolds*, J. Nonlin. Math. Phys. **9** (2002), 394–425.
- [R] C. Roger, *Cohomology of current Lie algebras*, Deformation Theory of Algebras and Structures and Applications (M. Hazewinkel and M. Gerstenhaber, eds.), Kluwer, 1988, 357–374.
- [Sa] L. J. Santharoubane, *The second cohomology group for Kac-Moody Lie algebras and Kähler differentials*, J. Algebra **125** (1989), 13–26.
- [St] S. Sternberg, *Lectures on Differential Geometry*, 2nd edition, AMS Chelsea Publ., 1983.
- [Z] P. Zusmanovich, *Deformations of $W_1(n) \otimes A$ and modular semisimple Lie algebras with a solvable maximal subalgebra*, arXiv: math.RA/0204004, accepted by J. Algebra.

SLAAKSTRAAT 3 - II, 1078BC AMSTERDAM, THE NETHERLANDS

E-mail address: pasha@member.ams.org

URL: <http://www.justpasha.org/math/>