

# NONSTANDARD LAGRANGIAN SUBMANIFOLDS IN $\mathbb{CP}^n$

RIVER CHIANG

**ABSTRACT.** We use Hamiltonian actions to construct nonstandard (as opposed to  $\mathbb{RP}^n$  and  $T^n$ ) Lagrangian submanifolds of  $\mathbb{CP}^n$ . First of all, a quotient of  $\mathbb{RP}^3$  by the dihedral group  $D_3$  is a Lagrangian submanifold of  $\mathbb{CP}^3$ . Secondly,  $SU(n)/\mathbb{Z}_n$  are Lagrangian submanifolds of  $\mathbb{CP}^{n^2-1}$ .

## 1. INTRODUCTION

Let  $(M, \omega)$  be a symplectic manifold. A submanifold  $L \subset M$  is **Lagrangian** if  $\dim L = \frac{1}{2} \dim M$  and  $\omega$  vanishes on  $T(L)$ .

Using Floer homology, Paul Biran has obtained information on the topology of Lagrangian submanifolds of some closed manifolds. One of Paul Biran's theorems concerns the Lagrangian submanifolds of  $\mathbb{CP}^n$  [B-1, B-2]:

**Theorem 1.1.** *Let  $L$  be a Lagrangian submanifold of  $\mathbb{CP}^n$  such that  $H_1(L; \mathbb{Z})$  is 2-torsion (that is,  $2H_1(L; \mathbb{Z}) = 0$ ). Then*

- (1)  $H^*(L; \mathbb{Z}_2) \cong H^*(\mathbb{RP}^n; \mathbb{Z}_2)$  as graded vector spaces.
- (2) If  $n$  is even, the isomorphism in (1) is an isomorphism of graded algebras.

Seidel [Se] also has a theorem with regard to the same subject:

**Theorem 1.2.** *Let  $L$  be a Lagrangian submanifold of  $\mathbb{CP}^n$ . Then*

- (1)  $H^1(L; \mathbb{Z}_{2n+2}) \neq 0$ .
- (2)  $H^1(L; \mathbb{Z}_{2n+2}) \not\cong (\mathbb{Z}_2)^g$  for any  $g \geq 2$ .
- (3) If  $H^1(L; \mathbb{Z}_{2n+2}) \cong \mathbb{Z}_2$ , then  $H^i(L; \mathbb{Z}_2) \cong \mathbb{Z}_2$  for all  $i = 0, \dots, n$ .

To test the limits of these topological restriction on Lagrangian submanifolds of  $\mathbb{CP}^n$ , we construct new examples using Hamiltonian actions. The constructions in this paper relies on the following well-known fact:

**Proposition 1.3.** *Let a compact Lie group  $G$  act on a symplectic manifold  $(M, \omega)$  in a Hamiltonian fashion with a moment map  $\Phi: M \rightarrow \mathfrak{g}^*$ . Assume  $\alpha \in \mathfrak{g}^*$  is a fixed point of the coadjoint action. Then for any  $m \in \Phi^{-1}(\alpha)$ , the orbit  $G \cdot m$  is an isotropic submanifold of  $M$ .*

*Proof.* An orbit is smooth. We only need to show that it is isotropic. Let  $\xi$  and  $\eta$  be any vector fields in the Lie algebra  $\mathfrak{g}$  of  $G$  and let  $\xi_M$  and  $\eta_M$  be their induced vector fields on  $M$ .

$$\begin{aligned} \omega(m)(\xi_M(m), \eta_M(m)) &= d\langle \Phi(m), \xi \rangle(\eta_M(m)) \\ &= \langle d\Phi(\eta_M), \xi \rangle \\ &= \langle \eta_{\mathfrak{g}^*}(\Phi(m)), \xi \rangle \\ &= \left\langle \frac{d}{dt} \Big|_0 \exp t\eta \cdot \alpha, \xi \right\rangle \\ &= \langle 0, \xi \rangle \end{aligned}$$

Therefore  $\omega(m) = 0$  for all  $m \in \Phi^{-1}(\alpha)$ . In particular, the orbit  $G \cdot m$  is a smooth isotropic submanifold of  $M$ .  $\square$

In particular, any orbit of an abelian action is isotropic as well as any orbit in the zero level set of a nonabelian action.

**Example 1.4.** Consider the action of the torus  $T^n$  on  $\mathbb{CP}^n$  given by

$$(\lambda_1, \dots, \lambda_n) \cdot [z_0, z_1, \dots, z_n] = [z_0, \lambda_1^{-1} z_1, \dots, \lambda_n^{-1} z_n].$$

This action is Hamiltonian and its moment map is

$$\Phi([z_0, z_1, \dots, z_n]) = \frac{1}{2} \left( \frac{|z_1|^2}{\|z\|^2}, \dots, \frac{|z_n|^2}{\|z\|^2} \right),$$

where  $\|z\|^2 = \sum_{i=0}^n |z_i|^2$ . The free orbit  $T^n$  is a Lagrangian submanifold of  $\mathbb{CP}^n$ .

Our first example is a Lagrangian submanifold of  $\mathbb{CP}^3$ . Its topology has an interesting deviation from Theorems 1.1 and 1.2.

**Theorem 1.5.** *There exists a Lagrangian submanifold of  $\mathbb{CP}^3$ , which is a quotient of  $\mathbb{RP}^3$  by the dihedral group  $D_3$ .*

We discuss the topology of this submanifold in Section 3.

We use different actions to construct a second family of examples. It provides a generalization of the standard Lagrangian submanifold  $\mathbb{RP}^3$  of  $\mathbb{CP}^3$ .

**Theorem 1.6.**  *$\mathrm{SU}(n)/\mathbb{Z}_n$  is a Lagrangian submanifold of  $\mathbb{CP}^{n^2-1}$ .*

## 2. FIRST EXAMPLE

View  $\mathbb{C}^4$  as the space of homogeneous polynomials of degree 3 in two variables. We can define the irreducible  $\mathrm{SU}(2)$  representation  $\rho: \mathrm{SU}(2) \rightarrow \mathrm{GL}(\mathbb{C}^4)$  by

$$\rho(A)(p)(z) = p(zA)$$

where  $p \in \mathbb{C}^4$ ,  $A \in \mathrm{SU}(2)$  and  $z = (x, y) \in \mathbb{C}^2$ . The moment map for this representation as a map from  $\mathbb{C}^4$  into  $\mathbb{R} \times \mathbb{C} \cong \mathfrak{su}(2)^*$  is given by

$$(u_0, u_1, u_2, u_3) \mapsto \left( \frac{3}{2}|u_0|^2 + \frac{1}{2}|u_1|^2 - \frac{1}{2}|u_2|^2 - \frac{3}{2}|u_3|^2, u_0\overline{u_1} + u_1\overline{u_2} + u_2\overline{u_3} \right).$$

So the zero level set of the moment map must satisfy

$$\begin{cases} 3|u_0|^2 + |u_1|^2 = |u_2|^2 + 3|u_3|^2, \\ u_0\overline{u_1} + u_1\overline{u_2} + u_2\overline{u_3} = 0. \end{cases}$$

This  $\mathrm{SU}(2)$  action commutes with the diagonal  $S^1$  action on  $\mathbb{C}^4$ , and therefore descends to an action on  $\mathbb{CP}^3$ .

Clearly the point  $[1, 0, 0, 1]$  is in the zero level set. We know that

$$(x, y) \begin{pmatrix} \alpha & \beta \\ -\overline{\beta} & \overline{\alpha} \end{pmatrix} = (\alpha x - \overline{\beta} y, \beta x + \overline{\alpha} y),$$

where  $|\alpha|^2 + |\beta|^2 = 1$ . To compute the stabilizer, we set  $\lambda(x^3 + y^3) = (\alpha x - \overline{\beta} y)^3 + (\beta x + \overline{\alpha} y)^3$  for any possible  $\lambda$  in  $S^1$ . Direct computation shows that  $\lambda = \pm 1$  when  $\beta = 0$  and  $\lambda = \pm i$  when  $\alpha = 0$ . So the stabilizer is generated by

$$(2.1) \quad \begin{pmatrix} 0 & i \\ i & 0 \end{pmatrix} \quad \text{and} \quad \left\{ \begin{pmatrix} \alpha & 0 \\ 0 & \overline{\alpha} \end{pmatrix}, \alpha^6 = 1 \right\}$$

Note this  $\mathrm{SU}(2)$  action is not effective on  $\mathbb{CP}^3$ . It induces an effective  $\mathrm{SO}(3)$  action and the stabilizer for  $[1, 0, 0, 1]$  is the semidirect product  $\mathbb{Z}_2 \ltimes \mathbb{Z}_3$ , namely, the dihedral group  $D_3$ .

Therefore, we have a Lagrangian  $L = \mathrm{SO}(3)/D_3 \cong \mathbb{RP}^3/D_3$  in  $\mathbb{CP}^3$ .

In fact, this is the only orbit in the zero level set. According to the Guillemin-Marle-Sternberg local normal form theorem, a neighborhood of the orbit of  $[1, 0, 0, 1]$  is equivariantly symplectomorphic to a neighborhood of the zero section in  $Y = \mathrm{SO}(3) \times_{D_3} \mathbb{R}^3$ . It is easy to check that there exists one single orbit in the zero level set in  $Y$ . Since the zero level set is connected and since the orbit is closed, it follows that the zero level set in  $\mathbb{CP}^3$  consists of one orbit only.

### 3. TOPOLOGY FOR THE FIRST EXAMPLE

Let  $\Gamma$  denote the group generated by the elements in (2.1). Then the fundamental group of  $L$  is  $\pi_1(L) = \Gamma$  since  $L = \mathbb{RP}^3/D_3 = S^3/\Gamma$ .

Let  $b$  denote  $\begin{pmatrix} 0 & i \\ i & 0 \end{pmatrix}$  and  $a$  denote  $\begin{pmatrix} \alpha & 0 \\ 0 & \bar{\alpha} \end{pmatrix}$ ,  $\alpha^6 = 1$ . Then  $b^2 = -1$  and  $ba = a^5b$ . The commutator subgroup  $A = \{x^{-1}y^{-1}xy \mid x, y \in \pi_1(L)\}$  is  $\{1, a^2, a^4\}$ . By Hurewicz Theorem, the first homology group  $H_1(L; \mathbb{Z})$  is  $\{1, b, b^2, b^3\} \cong \mathbb{Z}_4$ .

Using the Universal Coefficient Theorem and Poincaré duality, we can calculate that

$$\begin{aligned} H_0(L; \mathbb{Z}) &= \mathbb{Z}, & H^0(L; \mathbb{Z}) &= \mathbb{Z}, & H^0(L; \mathbb{Z}_2) &= \mathbb{Z}_2, \\ H_1(L; \mathbb{Z}) &= \mathbb{Z}_4, & H^1(L; \mathbb{Z}) &= 0, & H^1(L; \mathbb{Z}_2) &= \mathbb{Z}_2, \\ H_2(L; \mathbb{Z}) &= 0, & H^2(L; \mathbb{Z}) &= \mathbb{Z}_4, & H^2(L; \mathbb{Z}_2) &= \mathbb{Z}_2, \\ H_3(L; \mathbb{Z}) &= \mathbb{Z}, & H^3(L; \mathbb{Z}) &= \mathbb{Z}, & H^3(L; \mathbb{Z}_2) &= \mathbb{Z}_2. \end{aligned}$$

Also,  $H^1(L; \mathbb{Z}_8) = \mathbb{Z}_4$ .

The Lagrangian  $L$  we found does not satisfy the assumption of 2-torsion as in Theorem 1.1 and Theorem 1.2(3), but  $H^*(L; \mathbb{Z}_2)$  is isomorphic to  $H^*(\mathbb{RP}^3; \mathbb{Z}_2)$  as graded vector spaces.

### 4. SECOND FAMILY OF EXAMPLES<sup>1</sup>

Consider  $\mathbb{C}^{n^2}$  as the set of  $n$  by  $n$  matrices.  $\mathrm{SU}(n)$  acts naturally by left multiplication. This action is Hamiltonian and its moment map is given by

$$\Phi(Z) = \frac{i}{2} ZZ^* + \frac{1}{2i} I$$

where  $I$  is the identity matrix.

This  $\mathrm{SU}(n)$  action commutes with the circle action of multiplication by  $\lambda I$ . Therefore, it descends to an action on  $\mathbb{CP}^{n^2-1}$  where the center  $\mathbb{Z}_n$  of  $\mathrm{SU}(n)$  acts trivially. Therefore, a free orbit in the zero level set of  $\mathbb{C}^{n^2}$  descends to an orbit in the zero level set of  $\mathbb{CP}^{n^2-1}$  with stabilizer  $\mathbb{Z}_n$ . Since the dimension of  $\mathrm{SU}(n)$  is  $n^2 - 1$ , by Proposition 1.3,  $\mathrm{SU}(n)/\mathbb{Z}_n$  is a Lagrangian submanifold of  $\mathbb{CP}^{n^2-1}$ .

For  $n = 2$ , the above-mentioned orbit  $\mathrm{SU}(2)/\mathbb{Z}_2$  is the standard Lagrangian  $\mathbb{RP}^3$  in  $\mathbb{CP}^3$ . Indeed it is the projective space of the real 4 dimensional space  $V$  spanned

---

<sup>1</sup>I would like to thank Michèle Audin for showing me a beautiful alternative construction of these examples in terms of real forms of Lie groups [Au].

by the following matrices:

$$\begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \begin{pmatrix} i & 0 \\ 0 & i \end{pmatrix}, \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \begin{pmatrix} 0 & i \\ -i & 0 \end{pmatrix}.$$

#### REFERENCES

- [Au] M. Audin, Unpublished notes.
- [A-L-P] M. Audin, F. Lalonde and L. Polterovich, *Symplectic rigidity: Lagrangian submanifolds*. In Holomorphic curves in symplectic geometry. Edited by M. Audin and J. Lafontaine. Progress in Mathematics, **117**. Birkhäuser Verlag, Basel, 1994.
- [B-1] P. Biran, *Homological uniqueness of Lagrangian submanifolds*. In preparation.
- [B-2] P. Biran, *Geometry of symplectic intersections*. Proceedings of the International Congress of Mathematicians (Beijing 2002), Vol. II, 241–255.
- [Se] P. Seidel, *Graded Lagrangian submanifolds*. Bull. Soc. Math. France **128** (2000), no. 1, 103–149.