ESTIMATES FOR THE MINIMAL CROSSING NUMBER

HERMANN GRUBER

ABSTRACT. First, I give an elementary proof for the fact that the minimal crossing number is additive under composition of torus links. This result is generalized to the composition of homogeneous braids with alternating fibered links. Then there follow estimates for the crossing number of satellite knots. In the last chapter, I discuss a conjecture concerning the HOMFLY and the Kauffman polynomial.

1. INTRODUCTION.

A link is called composite iff there is a 2-sphere in S^3 which meets the link in exactly two points and decomposes it into two sublinks K and L, neither of which is an unknotted arc. Write K#L if the link is the composition of K and L. Whether the minimal crossing number of a composite link is simply the sum of the minimal crossing numbers of the factor links, is a very natural and very old question already posed by Tait some hundred years ago. It is trivial to see that it cannot be greater than the sum, but the question if equality holds in general remains unsolved up to now. However, the question is answered if the factor links are all from a certain family of knots, called adequate knots[10], problem 1.65. Here, another such family is exposed, which contains the torus links and the alternating fibered links. Some of the arguments exposed here are closely following the ideas of Murasugi's paper [12], and some results are re-expressed and proved more shortly here. It turns out to be astonishingly easy to show that the crossing number of torus links is additive under composition. But no one has written up these things up to now, and it is one purpose of this paper to do this.

Another aim is to turn around the classical points of view, gaining beautiful formulas estimating the minimal crossing number via the geometric concepts of braid index and (canonical) genus. For example, you will see that, in this application here, the canonical genus can be a considerably more powerful concept than the Seifert genus.

2. Preliminaries.

Throughout this paper, it is immaterial if the knots are chiral or not. For this cause, don't regard chiral pairs as distinct.

For a (nontrivial and non-split) link K and a regular diagram D(K) (or simply D), let c(D) be the number of crossings in D. Write c(K) for the minimal crossing number, and b(K) for the braid index of K.

Give the knot K an arbitrary orientation. By cutting out each crossing, respecting the orientation, convert the diagram D into a number of oriented closed curves in the (extended) plane, called Seifert circles of D. Write s(D) for the number of Seifert circles of D. From these circles, construct a spanning surface for K. The genus of this surface depends of the diagram: $g(D) = \frac{1}{2}(c(D) - s(D) - |K| + 2)$, where $|\mathbf{K}|$ is the number of components of K. This can easily be seen by calculating the Euler characteristics of the constructed surface.

The (Seifert) genus of a link K is defined as the minimal genus among all orientable surfaces bounded by K, the canonical (or weak) genus as

$$\widetilde{g}(K) = \min_{D = D(K)} \{g(D)\}$$

and the free genus $g_f(K)$ is the minimum genus of all Seifert surfaces for K whose complement in S^3 is a handlebody. We mention $g(K) \leq g_f(K) \leq \tilde{g}(K)$.

3. TORUS LINKS, ALTERNATING FIBERED LINKS AND HOMOGENEOUS BRAIDS.

I shall note first the following self-evident lemma, which plays a central role in our discussion:

Lemma 3.1. For any link K,

$$c(K) = \min_{D = D(K)} \left(2g(D) + s(D) \right) + |K| - 2$$

By minimizing both the number of Seifert circles and the genus independently, we get the following estimate:

Lemma 3.2. For every link K,

$$c(K)\geq 2\widetilde{g}(K)+b(K)+|K|-2\geq 2g(K)+b(K)+|K|-2$$

Proof. What remains to show is

$$\min_{D=D(K)} \{s(D)\} = b(K)$$

See Yamada's proof [24].

Both of the above inequalities are sharp for torus links, so we can give a short and elementary proof of a result of Murasugi:

Proposition 3.3. [12]Let K be a (p,q) torus link with $p \ge q \ge 2$. Then c(K) = 2g(K) + b(K) + |K| - 2 = pq - p

Proof. Consider the standard representation D of K as a closed braid with q strands. Here, c(D) = pq - p and 2g(D) = pq - p - q + |K| - 2. g(D) is minimal over all Seifert surfaces, see [4]. As the diagram D shows, the braid index can be at most q, and since it cannot be lower than the bridge number, it must be equal to q. \Box

This result motivates the following definition:

Definition 3.4. Say that a link K is in the family \mathcal{F} (or, shortly, \mathcal{F} -link) if c(K) = 2g(K) + b(K) + |K| - 2.

Next, see that the weaker of the lower bounds is additive under composition:

Proposition 3.5. Let K be a composite link with the factor links $K_1, ..., K_n$. Then

$$c(K) \ge \sum_{i=1}^{n} [2g(K_i) + b(K_i) + |K_i| - 2]$$

Proof. Use the facts that the genus is additive under composition (see e. g. [1]), and for the braid index it holds $b(K_1 \# K_2) = b(K_1) + b(K_2) - 1[2]$.

Remark. The free genus is also additive under knot composition, see [18]. Thus for knots, we may also write here $g_f(K)$ instead of g(K), sometimes leading to better results.

Corollary 3.6. If K is a composite link with factor links $K_1, ..., K_n$, and all factor links are in \mathcal{F} , then

$$c(K) = \sum_{i=1}^{n} c(K_i)$$

At this stage, we already see that the crossing number is additive for torus links; a problem repeatedly posed by Adams (see [1], chapter 5.1); we see that the answer is almost as simple as the question in this special case.

With two other results by Murasugi, we have that \mathcal{F} also contains other classes of links: We repeat here almost exactly corollary 2 in [12]:

Proposition 3.7. [12] If K is an alternating fibered link, then K is in \mathcal{F} .

Proof. Murasugi's statement is $c(K) = deg\Delta_K + b(K) - 1$, and with $deg\Delta_K - |K| + 1 \le 2g(K)$, the rest follows.

The following proposition is very closely related to proposition 7.4 in [12]. However, I redo it here to point out a different aspect: to fit the definition of the family \mathcal{F} .

Proposition 3.8. Let γ be a homogeneous n-braid, and L be the closure of γ . If b(L) = n, then L is in \mathcal{F} .

Proof. Draw a diagram D by simply closing γ . You have for the degree of the reduced Alexander polynomial $deg\Delta_L = c(D) - s(D) + 1$, since γ is homogeneous. With $deg\Delta_L + 1 - |K| \leq 2g(L)$, s(D) = b(L), this leads to $2g(L) \geq c(D) - b(L) - |K| + 2$, and because of $g(D) \geq g(L)$, c(D) = 2g(L) + b(L) + |K| - 2. By Lemma 3.2, you see that c(D) is minimal.

4. EXAMPLES.

In the previous chapter, we have seen some families of \mathcal{F} -links. You may have observed that all links presented there were homogeneous (for a definition see[4]), and fibered. However, it is neither true that every fibered homogeneous link is an \mathcal{F} -link nor that every \mathcal{F} -link is fibered. We take a look at an example:

Example 4.1. The Perko knot (denoted by 10_{161} in Rolfsen's table [19]) is fibered and homogeneous, and its (canonical) genus is 3, as well as its braid index. But its crossing number is 10. Therefore, the Perko knot is not in \mathcal{F} .

I proceed with a discussion of alternating \mathcal{F} -links.

Definition 4.2. A number n(D) associated with an alternating link diagram D is called an alternating link invariant if $n(D_1) = n(D_2)$ for any two reduced alternating diagrams D_1, D_2 of the same link.

Consider again Seifert circles:

Proposition 4.3. Let D(L) be an alternating link diagram. Let $s_a(D)$ be the number of Seifert circles in D. Then $s_a(D) =: s_a(L)$ is an alternating link invariant.

Proof. Let D be a reduced alternating link diagram of L. Then every other reduced alternating diagram of L can be reached from D by a finite number of flyping moves [11]. The figure shows the flyping move.



Hence, it is enough to show that s(D) is not altered by the flyping move. When considering only the underlying Seifert circles in the plane, then flyping does not affect the connectedness of the circles inside the disk. If the strands on the right-hand side of the above figure have the same orientation, the flyping move does not change the number of Seifert circles. For the remaining case, we use the idemposition algebra for arrangements of (unoriented) circles in the plane (see[21]). In this case, the effect of flyping outside the disk looks like:



When counting Seifert circles, we must consider several cases for the above disk diagram on the left: The outgoing arcs of the diagram can be connected outside the diagram in two ways, and there are two possibilities how the ingoing arcs into the disk can be connected inside the disk:

- (1) the upper left arc is connected to the lower left arc outside the diagram (called numerator closure)
 - (a) the upper left arc is connected inside the disk to the lower left arc
 - (b) the upper left arc is connected inside the disk to the right ingoing arc
- (2) the upper left arc is connected to the upper right arc outside the diagram (called denominator closure)
 - (a) the upper left arc is connected inside the disk to the lower left arc inside the disk
 - (b) the upper left arc is connected to the right ingoing arc inside the disk.

Now the important thing is to see that the left arcs are in all four cases connected to each other. We further need two idemposition types, see the following picture:



To imitate flyping, do the following: Apply first a type 1 idemposition Φ on the left side. In all cases, theorem 14 in [21] gives $s(\Phi(D)) = s(D) + 1$. The disc is now disconnected from the rest of the link. Turn it in the plane by π . Next, apply a type 2 idemposition Θ on the right side. Since the arcs on the right side of the

image cannot be connected before applying Θ , again by theorem 14, we have $s(\Theta(\Phi(D))) = (s(D) + 1) - 1$, and the proof is complete.

We obtain immediately the following condition for alternating \mathcal{F} -links:

Corollary 4.4. An alternating link is in \mathcal{F} iff $s_a(L) = b(L)$.

Proof. Assume first $s_a(L) = b(L)$. It is a long known fact that the Seifert surface from an alternating diagram D is minimal, i. e. g(D) = g(L). Since $s_a(L) = s(D) = b(L)$, and a reduced alternating diagram has minimal number of crossings [1], c(L) = 2g(L) + b(L) - 1. On the other hand, if $b(L) < s_a(L)$, for a reduced alternating diagram we have c(D) > 2g(D) + b(L) - 1. \Box

We proceed with a statement on planar Murasugi sums. For the definition of the terms used here, see [12].

Corollary 4.5. Let L be an alternating link, and D(L) be the planar Murasugi sum of special alternating link diagrams $D(L_1), ..., D(L_n)$. then

$$s_a(L) - 1 = \sum_{i=1}^n (s_a(L_i) - 1)$$

Proof. Follows immediately by the definition of planar Murasugi sum.

It is now natural to ask whether a similar equality holds for the braid index. This is conjectured in [12]:

Conjecture 4.6. Let L be the planar Murasugi sum of special alternating links $L_1, ..., L_n$. then

$$b(L) - 1 = \sum_{i=1}^{n} (b(L_i) - 1)$$

I note that the special alternating \mathcal{F} -links are exactly those satisfying theorem 8.1 in [12], called there "special alternating links of the nonmultiple type". This is an important point for the following discussion.

Example 4.7. The pretzel links $\mathcal{P}(a_1, a_2, ..., a_{2n})$ with all $a_i \geq 2$ are special \mathcal{F} -links (which are not fibered).

Proposition 4.8. Let D(L) be the planar Murasugi sum of special alternating link diagrams $D(L_1), ..., D(L_n)$. Assume at least one of $L_1, ..., L_n$ is a non- \mathcal{F} -link. Then L is a non- \mathcal{F} -link.

Proof. In the proof of theorem 8.1 in [12], a local diagrammatic move upon a special alternating diagram of a non- \mathcal{F} link reduces the number of Seifert circles. Use this move in D(L) to obtain a diagram D' with fewer Seifert circles. But now, $b(L) \leq s(D') < s_a(L)$. Recall 4.4 to conclude that L is not in \mathcal{F} .

Corollary 4.9. Let L be a \mathcal{F} -link. Then conjecture 4.6 holds.

Proof. Since L is in \mathcal{F} , $b(L) = s_a(L) = \sum_{i=1}^n (s_a(L_i) - 1)$. On the other hand, for all L_i , $s_a(L_i) = b(L_i)$ because they must also be in \mathcal{F} (which follows from the above proposition).

I believe furthermore the alternating links constructed from special alternating \mathcal{F} -links are exactly the alternating \mathcal{F} -links. This would follow immediately from conjecture 4.6 and complete the classification of alternating \mathcal{F} -links.

5. Application to satellite knots.

We can apply Lemma 3.2 to give a "good" estimate for the crossing number of (p,q)-cable knots about a knot in \mathcal{F} . (Here, p means the linking number of the knot with a meridian of the essential knotted torus in which L lies.)

Corollary 5.1. Let p,q be positive integers with gcd(p,q) = 1. Let K be the (p,q)-cable knot about a \mathcal{F} -knot C. Then $c(K) \ge q(p-1) + p \cdot c(C)$.

Proof. Schubert showed in ([20], pp. 247 seq.) that

$$2g(K) = (p-1)(q-1) + 2p \cdot g(C)$$

Furthermore, the braid index of K is $p \cdot b(C)$ [3, 23]. Since gcd(p,q) = 1, |K| = 1. Use 3.2 to obtain $c(K) \ge q(p-1) + p \cdot c(C)$.

Remark. However, this bound is far from being sharp, as we conjecture that the crossing number is $q(p-1) + p^2 \cdot c(C)$. This may illustrate the difficulties in proving statements about the crossing number of satellite knots.

Similar estimates can be easily established in the same spirit, which cover other special cases, as you may figure out yourself. Nevertheless, we state here another example.

(The definitions for pattern types and the weights w_i used in the following are the same as in [15, 3].)

Corollary 5.2. Let K be a satellite knot with a \mathcal{F} -knot companion C and a closed alternating braid B as type 0 pattern. Then, $c(K) \ge c(B) + b(B) \cdot c(C)$.

Proof. Another result of Schubert ([20],p.192) states that in our case

$$2g(K) \ge 2b(B) \cdot g(C) + 2g(B)$$

Since B is an \mathcal{F} -knot, 2g(B) = c(B) - b(B) + 1; the same holds for C. Here again $b(K) = b(B) \cdot b(C)$ holds, for B is a type 0 pattern [3]. Now, 3.2 together with both B and C being \mathcal{F} -knots:

$$c(K) \ge c(B) + b(B) \cdot c(C)$$

We can also show an unconditional estimate for satellite knots, as follows:

Proposition 5.3. Let K be a satellite knot with companion C. Let B be the knot from the pattern. Then

$$c(K) \ge \begin{cases} w_0 \cdot (2g(C) + b(C) - 1) + g(B) + w_0 \text{ for pattern type } 0\\ w_0 \cdot (2g(C) + b(C) - 1) + g(B) + w_0 + w_1 \text{ for pattern type } 1\\ 2\alpha(C) - 2 \text{ else} \end{cases}$$

There, $\alpha(K)$ is the arc index of K as defined in [5].

Proof. First consider type 0 and type 1 patterns: Schubert's theorem on the genus of satellite knots reads for the general case as follows: $g(K) \ge n \cdot g(C) + g(B)$, where B is the knot from the pattern n is the linking number of a meridian of the essential torus with B. Here, $n = w_0$. The braid index of K is $w_0 \cdot b(C)$ for a type 0 pattern and $w_0 \cdot b(C) + w_1$ for a type 1 pattern respectively [3]. Using Lemma 3.2, we get $c(K) \ge w_0 \cdot (2g(C) + b(C) - 1) + g(B) + w_0$ for a type 0 pattern, resp. $c(K) \ge w_0 \cdot (2g(C) + b(C) - 1) + g(B) + w_0$ for a type 1 pattern.

For the other case, we use a different estimate given by Ohyama [17]: $c(K) \ge 2b(K) - 2$. Since we have a type k pattern, $b(K) \ge \alpha(C)$ ([15], prop.3.3.5). We can conclude $c(K) \ge 2\alpha(C) - 2$ in this case.

We can furthermore solve here the problem whether the crossing number of a satellite link is greater than that of the companion ([10], problem 1.67) if the latter is an alternating fibered knot:

Corollary 5.4. Let K be a satellite knot and let its companion C be an alternating fibered knot. Then $c(K) \ge 2c(C) + 2$.

Proof. Cromwell showed that $\alpha(K) \ge c(K) + 2$ if K is an alternating knot [5]. Since $K \in \mathcal{F}$, and for a proper satellite knot, the weight w_0 is greater than 1, the rest follows from the above proposition.

The result can be generalized to more companion types and sharpened at the same time if we restrict the pattern type:

Corollary 5.5. Let K be a satellite knot and let its companion C be a \mathcal{F} -knot. Furthermore, assume K has a pattern of type 0 or 1. Then $c(K) > w_0 \cdot c(C)$.

Proof. Obvious from 5.3.

The results worked out here are all dealing with the Seifert genus. But the canonical genus gives us more power; we deal with this in the next section. However, I don't believe that any satellite link is in \mathcal{F} .

6. CROSSING NUMBER, CANONICAL GENUS AND KNOT POLYNOMIALS

Of course, the question of determining the minimal crossing number remains open in general. But we can state another beautiful lower bound in terms of the HOMFLY polynomial, which is again additive under composition. Define the twovariable Laurent polynomial in the variables v, z P[v, z] = P(K) as in [7], and write shortly e (resp. m) and E (resp. M) for the minimal and maximal non-zero exponent in v (resp. z). Corollary 6.1. For every link K,

$$c(K) \ge M + \frac{1}{2}(E - e)$$

Proof. Use Lemma 3.2 together with $M \leq \min_{D=D(K)} c(D) - s(D) + 1$ [13] and the MWF- inequality [13, 6]: $b(K) \geq \frac{1}{2}(E-e) + 1$ Furthermore, |K| = m. \Box

Notice that the right-hand side is again additive under composition, since for a composite link L # M the polynomial is $P(L) \cdot P(M)$.

Remark. Sometimes, this bound can lead to better results than the inequality using the genus. For instance, M = 6 for the untwisted double of the trefoil, whereas the free genus is 2, and the genus is only 1. On the other hand, Stoimenow showed an example where 2g > M [22]; and there are also some examples for which the MWF- inequality is not sharp.

We continue with showing up a connection between the crossing number of a knot and the canonical genus of its double. (I noticed this when reading in [13] that the canonical genus of the Whitehead double of the trefoil is 3, which is also the minimal crossing number)

Proposition 6.2. Let K be a knot and W_K be a Whitehead double of K. Then

$$c(K) \ge \widetilde{g}(W_K)$$

Proof. First, assume that W_K is untwisted. The proof is constructive: Draw a diagram with minimal crossing number of K. Then choose a parallel to run along the curve in the inverse direction of the original curve (diagrams of this kind are often called blackboard diagrams). Then, in a region which is not near a crossing, replace the antiparallel by a (say) positive clasp to obtain a diagram D of W_K with 4c(K) + 2 crossings.

Cut out the crossings to obtain a bunch of Seifert circles with a pattern as shown in the figure.



Near each underlying crossing of K, you have four emerging arcs. Since the underlying diagram of K is minimal (and hence reduced), every one of these four arcs belongs to a distinct Seifert circle. In the region with the clasp from the Whitehead doubling, you see a closed Seifert circle and two lines above and below it, belonging to the same Seifert circle. The following picture should illustrate this.



Since every arc closes up to a Seifert circle with another arc, we count s(D) = 2c(K) + 1 Seifert circles in the diagram, and clearly, this leads for the genus of this diagram to $g(D) = c(K) \ge \tilde{g}(W_K)$.

Next, consider the case W_K is twisted. Let D_n (resp. D_{n+1}) be a diagram with n (resp. n+1) half-twists. The following picture illustrates that one more half-twist produces splits a Seifert circle in D_n into two, and at the same time we have more crossing in D_{n+1} , thus every pair of diagrams D_n , D_{n+1} has the same genus. See the figure.



Complete the proof by induction on n.

Remark. I was informed that this proposition was found independently by Kidwell and Stoimenow [9].

Together with Morton's canonical genus inequality, we can estimate the minimal crossing number from the polynomial of a satellite about it. A similar technique has already successfully been used in [14] for the braid index in special cases.

Problem 6.3. Does an infinite family of knots exist with $\tilde{g}(W_K) = c(K)$?

Very recently, Stoimenow and Kidwell asked a question (see [16], Problem 1.15) which turns out to be closely related to this problem:

Conjecture 6.4. Let K be a knot and W_K be its Whitehead double. Let F(K) be the Kauffman polynomial of K in the variables a and z. Then

 $2(\max \deg_z F(K) + 1) = \max \deg_z P(W_K)$

The truth of the conjecture would imply that there exists such an infinite family, namely the prime alternating knots. With the inequality found here, I can give further "evidence" for the truth of the above equality (both sides are bounded above by 2c(K)), and attack a part of it:

Proposition 6.5. Let K be a prime alternating knot and Wh(K) a Whitehead double of K. Then

$$2(\max \deg_z F(K) + 1) \ge \max \deg_z P(W_K)$$

Proof. Kidwell showed that $c(K) \ge \max \deg_z F(K) + 1$ with equality if K is a prime alternating knot [8]. Morton's inequality and proposition 6.2 give us $2c(K) \ge \max \deg_z P(W_K)$, and the proof is complete.

Since we have also found now

$$2c(K) \geq \max \deg_z P(W_K)$$

and

$2c(K) \ge 2(\max \deg_z F(K) + 1)$

I feel free to mention the following, which is quite similar:

Let G(K) be the Kauffman polynomial with coefficients reduced modulo 2. Then $b(W_K) \ge \alpha(K)$ (see [15], prop.3.3.5) together with $\alpha(K) \ge \max \deg_a G(K) - \min \deg_a G(K) + 2$ (see [15], prop.4.4.1) imply:

$$2b(W_K) - 2 \ge 2(\max \deg_a G(K) - \min \deg_a G(K) + 1)$$

whereas

$$2b(W_K) - 2 \ge \max \deg_v P(W_K) - \min \deg_v P(W_K)$$

is the MWF-inequality. (Note that the degrees in v of the HOMFLY polynomial depend heavily on how W_K is twisted, whereas G(K) is of course not affected.)

References

- [1] Adams, Colin C. : "The knot book" (german), Berlin 1995
- [2] Birman, J.S.: "Studying links via closed braids IV: composite links and split links." Invent. Math. 102 (1990), 115-139.
- Birman, J.S.; Menasco, W.: "Special Positions for essential tori in link complements", Topology 33, No. 3 (1994) 525-556
- [4] Cromwell, P. R.: "Homogeneous Links", J. London Math. Soc. (series 2) 39 (1989), 535-552
- [5] Cromwell, P. : "Arc presentations of knots and links", in: Knot Theory (Proc. Conference, Warsaw 1995), eds. V. F. R. Jones et al, Banach Center Publications 42, Warsaw, 1998, 57–64
- [6] Franks, J.; Williams, R. F.: "Braids and the Jones-Conway polynomial" Trans. Amer. Math. Soc. 303 (1987), 97-108.
- [7] Freyd, P.; Hoste, J.; Lickorish, W. B. R.; Millet, K.; Ocneanu, A.; Yetter, D.: "A new polynomial invariant for knots and links", Bull. Amer. Math. Soc. 12 (1985), 239-246.
- [8] Kidwell, M. :"On the degree of the Brandt-Lickorish-Millet-Ho polynomial of a link", Proc. Amer. Math. Soc. 100 (1987), 755-761.
- [9] Kidwell, M.; Stoimenow, A.: "Examples Relating to the Crossing Number, Writhe, and Maximal Bridge Length of Knot Diagrams", to appear in Mich. Math. J.
- [10] Kirby, R. :"Problems in low-dimensional topology", on the web.
- [11] Menasco, W. ; Thistlethwaite, M. : "The classification of alternating links", Ann. Math. 138 (1993), 113-171 AMS Bulletin, Vol. 25, No. 2, October 1991, 403-412.
- [12] Murasugi, K. : "On the braid index of alternating links", Trans. Amer. Math. Soc. 326, No. 1 (1991), 237-260.
- [13] Morton, H. : "Seifert circles and knot polynomials", Proc. Cambridge Phil. Soc. 99 (1986), 107-109.
- [14] Morton, H. : "The 2-variable polynomial of cable knots", Proc. Cambridge Phil. Soc. 101 (1987), 267-278.
- [15] Nutt, I. : "Braid index of satellite links", PhD thesis, University of Liverpool (1995)
- [16] Ohtsuki, T. (ed.): "Problems on invariants of knots and 3-manifolds", Geometry and Topology Monographs, Vol. 4
- [17] Ohyama, Y. :"On the minimal crossing number and the braid index of links", Can. J. Math. 45, No. 1 (1993), 117-131.
- [18] Ozawa, M.: "Additivity of free genus of knots", Topology 40 (2001), no. 4, 659-665.
- [19] Rolfsen, D. :"Knots and links", Berkeley 1976.
- [20] Schubert, H. :"Knoten und Vollringe", Acta Mathematica 90 (1953), 131-286
- [21] Spencer-Brown, G. "Laws of form", international edition, Lübeck 1997, "Appendix 5: Two proofs of the four-color map theorem"
- [22] Stoimenow, A.: "On the crossing number of positive knots and braids and braid index criteria of Jones and Morton-Williams-Franks", Trans. Amer. Math. Soc. 354 (2002), no. 10, 3927–3954
- [23] R. F. Williams:"The braid index of generalized cables", Pacific J. Math. 155 (1992), no. 2, 369–375.
- [24] Yamada, S. : "The minimal number of Seifert circles equals the braid index" Invent. Math. 88 (1987), 347-356.