Indices of vector fields or 1-forms and characteristic numbers

W.Ebeling and S.M.Gusein-Zade *

Abstract

We define an index of a collection of 1-forms on a complex isolated complete intersection singularity corresponding to a Chern number and, in the case when the 1-forms are complex analytic, express it as the dimension of a certain algebra.

Introduction

An isolated singular point (zero) p of a vector field X on a (real or complex analytic) manifold M has an index: the degree of the map X/||X|| from a small sphere around the point p to the unit sphere. If the manifold M is closed (compact without boundary) and the vector field X has only isolated singular points, the sum of their indices is equal to the Euler characteristic $\chi(M)$ of the manifold M. If M is a complex analytic manifold of dimension n, then its Euler characteristic $\chi(M)$ is the characteristic number $\langle c_n(TM), [M] \rangle$, where TM is the tangent bundle of the manifold M and c_n is the corresponding Chern class.

If the vector field X is complex analytic, the index of its (isolated) singular point p can be computed as the dimension of a certain algebra. Namely, if, in local coordinates centred at the point p, $X = \sum_{i=1}^{n} X_i(\partial/\partial x_i)$, then $\operatorname{ind}_p X = \dim \mathcal{O}_{\mathbb{C}^n,0}/(X_1,\ldots,X_n)$, where $\mathcal{O}_{\mathbb{C}^n,0}$ is the ring of germs of holomorphic functions of n variables, (X_1,\ldots,X_n) is the ideal generated by the

^{*}Partially supported by the DFG-programme "Global methods in complex geometry", grants RFBR-01-01-00739, INTAS-00-259, NWO-RFBR-047.008.005. Keywords: characteristic numbers, 1-forms, vector fields, singular points, indices. AMS Math. Subject Classification: 32S50, 57R20, 57R25.

components of the field X. The Eisenbud-Levine-Khimshiashvili formula [EL, Kh] expresses the index of an algebraically isolated singular point of a (smooth) vector field on a real smooth manifold as the signature of a certain quadratic form defined on the real analogue of the algebra mentioned above. Similar statements hold for 1-forms on manifolds with the only difference that the sum of indices of singular points of a 1-form on a complex analytic manifold M of dimension n is equal to the characteristic number $\langle c_n(T^*M), [M] \rangle = (-1)^n \chi(M)$.

In the framework of the described properties there is almost no difference between vector fields and 1-forms (except the fact that non-zero analytic vector fields and 1-forms exist, generally speaking, on different complex manifolds). The difference becomes crucial if one considers complex-analytic vector fields or 1-forms on singular varieties [EG1, EG2]. The main difference is related with the fact that a complex analytic 1-form can be restricted to a submanifold or to a subvariety (staying complex analytic, at least, at the non-singular points of the subvariety), while there is no invariant notion of a restriction of a vector field to a submanifold.

Here we start to discuss generalizations of these notions and statements for other characteristic numbers (different from $\langle c_n, [M] \rangle$). For smooth varieties, it is not very essential whether we discuss vector fields, or 1-forms, or simply sections of a vector bundle. For singular varieties (namely for isolated complete intersection singularities: ICIS) the properties of 1-forms and vector fields with respect to the discussed problems are quite different. Here we restrict ourselves almost exclusively to the complex analytic case.

1 An index for collections of sections of a vector bundle

We consider a general setting. Let $\pi : E \to M$ be a complex analytic vector bundle of rank m over a complex analytic manifold M of dimension n. It is known that the Poincaré dual to the characteristic class $c_k(E)$ (k = 1, ..., m)is represented by the 2(n - k)-dimensional cycle which consists of points of the manifold M where m - k + 1 sections of the vector bundle E are linearly dependent (cf., e.g., [GH, p. 413]).

For natural numbers p and q with $p \ge q$, let $\mathcal{M}(p,q)$ be the space of $p \times q$ matrices with complex entries and let $D_{p,q}$ be the subspace of $\mathcal{M}(p,q)$

consisting of matrices of rank less than q. The subset $D_{p,q}$ is an irreducible subvariety of $\mathcal{M}(p,q)$ of codimension p-q+1 (see below). The complement $W_{p,q} = \mathcal{M}(p,q) \setminus D_{p,q}$ is the Stiefel manifold of q-frames (collections of q linearly independent vectors) in \mathbb{C}^p . It is known that $W_{p,q}$ is (2p-2q)-connected and $H_{2p-2q+1}(W_{p,q}) \cong \mathbb{Z}$ (see, e.g., [H]). The latter fact also proves that $D_{p,q}$ is irreducible. Since $W_{p,q}$ is the complement of an irreducible complex analytic subvariety of codimension p-q+1 in $\mathcal{M}(p,q)$, there is a natural choice of a generator of the homology group $H_{2p-2q+1}(W_{p,q}) \cong \mathbb{Z}$. Namely, the ("positive") generator is the boundary of a small ball in a smooth complex analytic slice transversal to $W_{p,q}$ at a non-singular point.

Let $\mathbf{k} = (k_1, \ldots, k_s)$ be a sequence of positive integers with $\sum_{i=1}^{s} k_i = k$. Consider the space $\mathcal{M}_{m,\mathbf{k}} = \prod_{i=1}^{s} \mathcal{M}(m, m-k_i+1)$ and the subvariety $D_{m,\mathbf{k}} = \prod_{i=1}^{s} D_{m,m-k_i+1}$ in it. The variety $D_{m,\mathbf{k}}$ consists of sets $\{A_i\}$ of $m \times (m-k_i+1)$ matrices such that $\operatorname{rk} A_i < m - k_i + 1$ for each $i = 1, \ldots, s$. Since $D_{m,\mathbf{k}}$ is irreducible of codimension k, its complement $W_{m,\mathbf{k}} = \mathcal{M}_{m,\mathbf{k}} \setminus D_{m,\mathbf{k}}$ is (2k-2)-connected, $H_{2k-1}(W_{m,\mathbf{k}}) \cong \mathbb{Z}$, and there is a natural choice of a generator of the latter group. This choice defines a degree (an integer) of a map from an oriented manifold of dimension 2k - 1 to the manifold $W_{m,\mathbf{k}}$.

Let $\{\omega_j^{(i)}\}\ (i=1,\ldots,s; j=1,\ldots,m-k_i+1; \sum_{i=1}^s k_i=n)$ be a collection of sections of the vector bundle $\pi: E \to M$ such that there are only isolated points where, for each $i=1,\ldots,s$, the sections $\omega_j^{(i)}, j=1,\ldots,m-k_i+1$, are linearly dependent. Let us choose a trivialization of the vector bundle $\pi: E \to M$ in a neighbourhood of a point p, let $(\omega_1^{(i)}(x),\ldots,\omega_{m-k_i+1}^{(i)}(x))$ be the $m \times (m-k_i+1)$ -matrix the columns of which consist of the components of the sections $\omega_j^{(i)}(x), j=1,\ldots,m-k_i+1, x \in M$, with respect to this trivialization. Let Ψ_p be the mapping from a neighbourhood of the point p to $\mathcal{M}_{m,\mathbf{k}}$ which sends a point x to the collection of matrices $\{(\omega_1^{(i)}(x),\ldots,\omega_{m-k_i+1}^{(i)}(x))\}, i=1,\ldots,s$. Its restriction ψ_p to a small sphere $S_{p,\varepsilon}^{2n-1}$ around the point p maps this sphere to the subset $W_{m,\mathbf{k}}$.

Definition The degree of the mapping $\psi_p : S_{p,\varepsilon}^{2n-1} \to W_{m,\mathbf{k}}$ is called the *index* of the collection of sections $\{\omega_i^{(i)}\}$ at the point p and is denoted by $\operatorname{ind}_p\{\omega_i^{(i)}\}$.

In other words, the index $\operatorname{ind}_p\{\omega_j^{(i)}\}\$ is equal to the intersection number of the germ of the image of the (complex analytic) map Ψ_p with the variety $D_{m,\mathbf{k}}$.

The point p is non-singular for the collection $\{\omega_i^{(i)}\}$ if at least for some i

the values $\omega_j^{(i)}(p)$ of the sections $\omega_j^{(i)}$ at the point p are linearly independent. The index $\operatorname{ind}_p\{\omega_j^{(i)}\}$ of a non-singular point is equal to zero. The description of the Poincaré dual cycles to the Chern classes $c_{k_i}(E)$ of the vector bundle E yields the following statement.

Theorem 1 Let $\sum_{i=1}^{s} k_i = n$ and suppose that the collection $\{\omega_j^{(i)}\}$ $(i = 1, \ldots, s; j = 1, \ldots, m - k_i + 1)$ of sections of the vector bundle $\pi : E \to M$ over a closed complex manifold M has only isolated singular points. Then the sum of the indices of these points is equal to the characteristic number $\langle \prod_{i=1}^{s} c_{k_i}(E), [M] \rangle$ of the vector bundle E.

2 An algebraic formula for the index

Suppose that, in local coordinates centred at the point p, the components of the section $\omega_j^{(i)}$ are complex analytic functions on M (e.g., this happens if the fibre bundle $\pi : E \to M$ is complex analytic and the sections $\omega_j^{(i)}$ are complex analytic in a neighbourhood of the point p = 0). Let $I_{\{\omega_j^{(i)}\}}$ be the ideal in the ring $\mathcal{O}_{\mathbb{C}^n,0}$ of germs of analytic functions of n variables generated by the $(m - k_i + 1) \times (m - k_i + 1)$ -minors of the matrices $(\omega_1^{(i)}, \ldots, \omega_{m-k_i+1}^{(i)})$ for all $i = 1, \ldots, s$.

Theorem 2

$$\operatorname{ind}_{p}\{\omega_{j}^{(i)}\} = \dim_{\mathbb{C}} \mathcal{O}_{\mathbb{C}^{n},0}/I_{\{\omega_{j}^{(i)}\}}.$$

Proof. The proof follows the same lines as the proof of [EG2, Theorem 1]. Let $\widetilde{\omega}_j^{(i)}$ be suitable analytic perturbations of the sections $\omega_j^{(i)}$ and let \widetilde{D} be the corresponding degeneracy locus. The essential point in the proof is the fact that \widetilde{D} is Cohen-Macaulay which follows from [BR, Corollary (2.7)]. \Box

3 Indices on ICIS

Let $f = (f_1, \ldots, f_\ell) : (\mathbb{C}^n, 0) \to (\mathbb{C}^\ell, 0)$ be an analytic map which defines an $(n-\ell)$ -dimensional isolated complete intersection singularity (ICIS) $V = f^{-1}(0) \subset (\mathbb{C}^n, 0) \ (f_r : (\mathbb{C}^n, 0) \to (\mathbb{C}, 0))$. Let $\{\omega_j^{(i)}\}$ be a collection of 1forms on a neighbourhood of the origin in $(\mathbb{C}^n, 0)$ with $i = 1, \ldots, s, j = 1, \ldots, n-\ell - k_i + 1, \sum k_i = n-\ell$, the restriction of which to V has no singular points on V outside of the origin in a neighbourhood of it. (Here it is not necessary to demand that the 1-forms $\omega_j^{(i)}$ are complex analytic. It is sufficient to suppose that $\omega_j^{(i)}$ are continuous complex linear functions on the tangent bundle $T\mathbb{C}^n$.) Let U be a neighbourhood of the origin in \mathbb{C}^n where all the functions f_r $(r = 1, \ldots, \ell)$ and the 1-forms $\omega_j^{(i)}$ are defined and such that the restriction of the collection $\{\omega_j^{(i)}\}$ of 1-forms to $(V \cap U) \setminus \{0\}$ has no singular points. Let $S_\delta \subset U$ be a sufficiently small sphere around the origin which intersects V transversally and denote by $K = V \cap S_\delta$ the link of the ICIS (V, 0). Let $\hat{\mathbf{k}} = (k_1 + \ell, \ldots, k_s + \ell)$ and let Ψ_V be the mapping from $V \cap U$ to $\mathcal{M}_{n,\hat{\mathbf{k}}}$ which sends a point $x \in V \cap U$ to the collection of $n \times (n - k_i + 1)$ -matrices

$$\{(df_1(x),\ldots,df_\ell(x),\omega_1^{(i)}(x),\ldots,\omega_{n-\ell-k_i+1}^{(i)}(x))\}, \quad i=1,\ldots,s.$$

Its restriction ψ_V to the link K maps K to the subset $W_{n\hat{k}}$.

Definition The *index* $\operatorname{ind}_{V,0}\{\omega_j^{(i)}\}$ of the collection of 1-forms $\{\omega_j^{(i)}\}$ on the ICIS V is the degree of the mapping $\psi_V : K \to W_{n,\hat{\mathbf{k}}}$.

As above, the index $\operatorname{ind}_{V,0}\{\omega_j^{(i)}\}\$ is equal to the intersection number of the germ of the image of the complex analytic mapping Ψ_V with the variety $D_{n,\hat{\mathbf{k}}}$.

For s = 1, $k_1 = n - \ell$, this definition coincides with the one of the index of a 1-form from [EG1].

Let $V \subset \mathbb{CP}^n$ be a complete intersection with isolated singular points. Let L be a complex line bundle on V and let $\{\omega_j^{(i)}\}$ be a collection of continuous 1-forms on V with values in L. Here this means that the forms $\omega_j^{(i)}$ are continuous sections of the fibre bundle $T^*V \otimes L$ outside of the singular points of V. Since, in a neighbourhood of each point p, the vector bundle L is trivial, one can define the index $\operatorname{ind}_p\{\omega_j^{(i)}\}$ of the collection of 1-forms $\{\omega_j^{(i)}\}$ at the point p just in the same way as in the local setting above. Moreover, one can consider L as a line bundle on a smoothing \widetilde{V} of the complete intersection V as well (e.g., using the pull back along a projection of \widetilde{V} to V).

Theorem 3 One has

$$\sum_{p \in V} \operatorname{ind}_p \{ \omega_j^{(i)} \} = \langle \prod_{i=1}^s c_{k_i}(T^* \widetilde{V} \otimes L), [\widetilde{V}] \rangle,$$

where \widetilde{V} is a smoothing of the complete intersection V.

Now, as above, let $(V, 0) \subset (\mathbb{C}^n, 0)$ be the ICIS defined by the equations $f_1 = \cdots = f_{\ell} = 0$ and let $\{\omega_j^{(i)}\}$ $(i = 1, \ldots, s; j = 1, \ldots, n - \ell - k_i + 1)$ be a collection of complex analytic 1-forms on a neighbourhood of the origin in \mathbb{C}^n the restriction of which has no singular points on V outside of the origin in a neighbourhood of it. Let $I_{V,\{\omega_j^{(i)}\}}$ be the ideal in the ring $\mathcal{O}_{\mathbb{C}^n,0}$ generated by the functions f_1, \ldots, f_ℓ and by the $(n - k_i + 1) \times (n - k_i + 1)$ minors of all the matrices

$$(df_1(x), \dots, df_{\ell}(x), \omega_1^{(i)}(x), \dots, \omega_{n-\ell-k_i+1}^{(i)}(x))$$

for all $i = 1, \ldots, s$.

Theorem 4

$$\operatorname{ind}_{V,0}\{\omega_j^{(i)}\} = \dim_{\mathbb{C}} \mathcal{O}_{\mathbb{C}^n,0}/I_{V,\{\omega_j^{(i)}\}}$$

See the "proof" of Theorem 2.

Remark. Let $\{X_j^{(i)}\}$ be a collection of vector fields on a neighbourhood of the origin in $(\mathbb{C}^n, 0)$ $(i = 1, ..., s; j = 1, ..., n - \ell - k_i + 1; \sum k_i = n - \ell)$ which are tangent to the ICIS $(V, 0) = \{f_1 = \cdots = f_\ell = 0\} \subset (\mathbb{C}^n, 0)$ at nonsingular points of V. One can define the index $\operatorname{ind}_{V,0}\{X_j^{(i)}\}$ as the degree of the mapping $K \to W_{n,\hat{\mathbf{k}}}$ which sends a point $x \in K$ to the collection of $n \times (n - k_i + 1)$ matrices

{(grad
$$f_1(x), \ldots, \text{grad } f_\ell(x), X_1^{(i)}(x), \ldots, X_{n-\ell-k_i+1}^{(i)}(x))$$
}, $i = 1, \ldots, s$.

Here

grad
$$f_r(x) = \left(\frac{\overline{\partial f_r}}{\partial x_1}(x), \dots, \frac{\overline{\partial f_r}}{\partial x_n}(x)\right),$$

where \overline{z} is the complex conjugate of the complex number z. For s = 1, $k_1 = n - \ell$, this definition coincides with the definition of the index of a vector field on an ICIS from [GSV, SS]. For vector fields the analogue of Theorem 3 holds with the only difference that the sum of the indices is equal to the characteristic number $\langle \prod_{i=1}^{s} c_{k_i}(T\widetilde{V} \otimes L), [\widetilde{V}] \rangle$. However, a formula similar to that of Theorem 4 does not exist. A reason for that is that in this case the index is the intersection number with $D_{n,\hat{\mathbf{k}}}$ of the image of the ICIS (V, 0) under a map which is not complex analytic. Moreover, in some cases this index can be negative.

References

- [BR] Buchsbaum, D. A., Rim, D. S.: A generalized Koszul complex. II. Depth and multiplicity. Trans. Am. Math. Soc. **111**, 197–224 (1964).
- [EG1] Ebeling, W., Gusein-Zade, S. M.: On the index of a holomorphic 1form on an isolated complete intersection singularity. Russian Math. Doklady, 380, 458–461 (2001).
- [EG2] Ebeling, W., Gusein-Zade, S. M.: Indices of 1-forms on an isolated complete intersection singularity. Moscow Math. J. (to appear), Preprint math.AG/0105242.
- [EL] Eisenbud, D., Levine, H.: An algebraic formula for the degree of a C^{∞} map germ. Ann. Math. **106**, 19–38 (1977).
- [GSV] Gómez-Mont, X., Seade, J., Verjovsky, A.: The index of a holomorphic flow with an isolated singularity. Math. Ann. 291, 737–751 (1991).
- [GH] Griffiths, Ph., Harris, J.: Principles of Algebraic Geometry. John Wiley & Sons, New York etc., 1978.
- [H] Husemoller, D.: Fibre bundles. Second Edition. Graduate Texts in Math. 20, Springer-Verlag, New York Heidelberg Berlin 1975.
- [Kh] Khimshiashvili, G. N.: On the local degree of a smooth map. Comm. Acad. Sci. Georgian SSR. **85**, no. 2, 309–311 (1977) (in Russian).
- [SS] Seade, J. A., Suwa, T.: A residue formula for the index of a holomorphic flow. Math. Ann. **304**, 621–634 (1996).

Universität Hannover, Institut für Mathematik Postfach 6009, D-30060 Hannover, Germany E-mail: ebeling@math.uni-hannover.de

Moscow State University, Faculty of Mechanics and Mathematics Moscow, 119992, Russia E-mail: sabir@mccme.ru