

THE MODULAR GROUP ACTION ON REAL $SL(2)$ -CHARACTERS OF A ONE-HOLED TORUS

WILLIAM M. GOLDMAN

ABSTRACT. The group Γ of automorphisms of the polynomial

$$\kappa(x, y, z) = x^2 + y^2 + z^2 - xyz - 2$$

is isomorphic to

$$\mathrm{PGL}(2, \mathbb{Z}) \ltimes (\mathbb{Z}/2 \oplus \mathbb{Z}/2).$$

We study the dynamics of the Γ -action on $\kappa^{-1}(t) \cap \mathbb{R}^3$, for $t \in \mathbb{R}$. The action of Γ preserves a Poisson structure defining a Γ -invariant area form on each $\kappa^{-1}(t) \cap \mathbb{R}^3$. For $t < 2$, the action of Γ is properly discontinuous on the four contractible components of $\kappa^{-1}(t) \cap \mathbb{R}^3$ and ergodic on the compact component (which is empty if $t < -2$). The contractible components correspond to Teichmüller spaces of (possibly singular) hyperbolic structures on a torus. For $t = 2$, the level set $\kappa^{-1}(t) \cap \mathbb{R}^3$ consists of characters of reducible representations and comprises two ergodic components corresponding to actions of $\mathrm{GL}(2, \mathbb{Z})$ on $(\mathbb{R}/\mathbb{Z})^2$ and \mathbb{R}^2 respectively. For $2 < t \leq 18$, the action of Γ on $\kappa^{-1}(t) \cap \mathbb{R}^3$ is ergodic. Corresponding to the Fricke space of a three-holed sphere is a Γ -invariant open subset $\Omega \subset \mathbb{R}^3$ whose components are permuted freely by Γ . The level set $\kappa^{-1}(t) \cap \mathbb{R}^3$ intersects Ω if and only if $t > 18$, in which case the Γ -action on the complement $(\kappa^{-1}(t) \cap \mathbb{R}^3) - \Omega$ is ergodic.

Date: February 8, 2020.

1991 *Mathematics Subject Classification.* 57M05 (Low-dimensional topology), 20H10 (Fuchsian groups and their generalizations), 30F60 (Teichmüller theory).

Key words and phrases. surface, fundamental group, character variety, representation variety, mapping class group, ergodic action, proper action, hyperbolic structure with cone singularity, Fricke space, Teichmüller space.

The author gratefully acknowledges partial support from National Science Foundation grant DMS-9803518, DMS-0103889 and a Semester Research Award from the General Research Board of the University of Maryland in Fall 1998.

CONTENTS

| | |
|---|----|
| Introduction | 2 |
| 1. The modular group and the moduli space | 4 |
| 2. Structure of the character variety | 6 |
| 3. Hyperbolic structures on tori ($t < 2$) | 15 |
| 4. Reducible characters ($t = 2$) | 22 |
| 5. Three-holed spheres and ergodicity ($t > 2$) | 24 |
| Appendix: Elements of the modular group | 32 |
| References | 37 |

INTRODUCTION

Let M be a compact oriented surface of genus one with one boundary component, a *one-holed torus*. Its fundamental group π is free of rank two. Its mapping class group $\pi_0(\text{Homeo}(M))$ is isomorphic to the outer automorphism group $\text{Out}(\pi)$ of π and acts on the space of equivalence classes of representations $\pi \rightarrow \text{SL}(2)$. We investigate the dynamics of this action on the set of real points on this moduli space.

Corresponding to the boundary of M is an element $K \in \pi$ which is the commutator of free generators $X, Y \in \pi$. By a theorem of Fricke [8, 9], the moduli space of $\text{SL}(2, \mathbb{C})$ -representations naturally identifies with affine 3-space \mathbb{C}^3 , via the quotient map

$$\text{Hom}(\pi, \text{SL}(2, \mathbb{C})) \rightarrow \mathbb{C}^3$$

$$\rho \mapsto \begin{bmatrix} x \\ y \\ z \end{bmatrix} := \begin{bmatrix} \text{tr}(\rho(X)) \\ \text{tr}(\rho(Y)) \\ \text{tr}(\rho(XY)) \end{bmatrix}.$$

In terms of these coordinates, the trace $\text{tr} \rho(K)$ equals:

$$\kappa(x, y, z) := x^2 + y^2 + z^2 - xyz - 2$$

which is preserved under the action of $\text{Out}(\pi)$. The action of $\text{Out}(\pi)$ on \mathbb{C}^3 is commensurable with the action of the group Γ of polynomial automorphisms of \mathbb{C}^3 which preserve κ (Horowitz [18]).

Theorem. *Let $\kappa(x, y, z) = x^2 + y^2 + z^2 - xyz - 2$ and let $t \in \mathbb{R}$.*

- *For $t < -2$, the group Γ acts properly on $\kappa^{-1}(t) \cap \mathbb{R}^3$;*
- *For $-2 \leq t < 2$, there is a compact connected component C_t of $\kappa^{-1}(t) \cap \mathbb{R}^3$ and Γ acts properly on the complement $\kappa^{-1}(t) \cap \mathbb{R}^3 - C_t$;*
- *For $t = 2$, the action of Γ is ergodic on the compact subset $\kappa^{-1}(2) \cap [-2, 2]^3$ and the action is ergodic on the complement $\kappa^{-1}(2) - [-2, 2]^3$;*
- *For $18 \geq t > 2$, the group Γ acts ergodically on $\kappa^{-1}(t) \cap \mathbb{R}^3$;*
- *For $t > 18$, the group Γ acts properly and freely on an open subset $\Omega_t \subset \kappa^{-1}(t) \cap \mathbb{R}^3$, permuting its components. The Γ -action on the complement of Ω_t is ergodic.*

The paper begins with background on the punctured torus and its mapping class group (§1.1). The automorphism group Γ of the cubic polynomial κ relates to the mapping class group (§1.3), the linear automorphisms being generated by sign-changes and permutations (§1.3.2). Characters of reducible representations comprise $\kappa^{-1}(2)$ (§1.5). An invariant Poisson structure (§1.6) determines the Γ -invariant measure on the set $\kappa^{-1}(t) \cap \mathbb{R}^3$ of \mathbb{R} -points of $\kappa^{-1}(t)$, which correspond to representations in real forms of $\mathrm{SL}(2, \mathbb{C})$ (§1.7).

§2 describes examples of automorphisms of π , their representation by polynomial automorphisms of (\mathbb{C}^3, κ) and their representation in the modular group. For $t \leq -2$, the action of Γ on $\kappa^{-1}(t) - \{(0, 0, 0)\}$ is proper, since (§3) $\kappa^{-1}(t) - \{(0, 0, 0)\}$ is the Fricke space of complete hyperbolic structures on a 1-holed torus. Similar ideas are used (§4) to prove that Γ acts properly on the noncompact components of $\kappa^{-1}(t) \cap \mathbb{R}^3$ where $-2 < t < 2$. However, most of the representations now are not discrete. Nonetheless these representations may be interpreted as moduli spaces of singular hyperbolic structures on a torus, with a single cone point singularity. Ergodicity on $\kappa^{-1}(2)$ (the reducible characters) follows (§5) from Moore's Ergodicity Theorem [28].

For $t > 2$, the dynamics of Γ acting on $\kappa^{-1}(t) \cap \mathbb{R}^3$ splits into two cases (§6), depending on whether $t \leq 18$ or $t > 18$. When $t > 18$, the moduli space contains characters of representations which are discrete embeddings, but for which the quotient hyperbolic surface is not a punctured torus, but rather a three-holed sphere (§6.1). The main technical tool (§6.2) is the trace-reduction algorithm of Kern-Isberner and Rosenberger [19] (see also Gilman-Maskit [10]) to produce either simple loops with elliptic representation or a hyperbolic structure on a pants. Finally arguments from Goldman [14] are used to deduce

ergodicity of the Γ -action from generators of π with represented by elliptic elements.

Notation and terminology. We work in Poincaré’s model of the hyperbolic plane \mathbf{H}^2 as the upper half-plane. We denote the identity map (identity matrix) by \mathbb{I} . Commutators are denoted:

$$[A, B] = ABA^{-1}B^{-1}$$

and inner automorphisms are denoted:

$$\iota_g : x \longmapsto gxg^{-1}.$$

For any group π we denote the group of all automorphisms by $\text{Aut}(\pi)$ and the normal subgroup of inner automorphisms by $\text{Inn}(\pi)$. We denote the quotient group $\text{Aut}(\pi)/\text{Inn}(\pi)$ by $\text{Out}(\pi)$. A compact surface S with n boundary components will be called *n-holed*. Thus a *one-holed torus* is the complement of an open disc inside a torus and a *three-holed sphere* (sometimes called a “pair-of-pants”) is the complement of three disjoint discs inside a sphere.

Acknowledgements. I am grateful to Matt Bainbridge, Robert Benedetto, Lawrence Ein, Cathy Jones, Misha Kapovich, Bernhard Leeb, John Millson, Walter Neumann, John Parker, Dan Rudolph, Peter Shalen, Richard Schwartz, Adam Sikora, George Stantchev and Scott Wolpert for helpful conversations during the course of this work.

1. THE MODULAR GROUP AND THE MODULI SPACE

In this section we define the modular group and the moduli space for the punctured torus. The modular group is isomorphic to

$$\pi_0(\text{Homeo}(M)) \cong \text{Out}(\pi) \cong \text{GL}(2, \mathbb{Z})$$

and the moduli space is affine space \mathbb{C}^3 . We explain how the invariant function κ originates. We describe various elements of the modular group and how they act on the moduli space.

1.1. The mapping class group. Let M denote a compact connected orientable surface of genus one with one boundary component. Since attaching a disc to M yields a torus, we refer to M as a *punctured torus*. The *mapping class group* of M is the group $\pi_0(\text{Homeo}(M, \partial M))$ of isotopy classes of homeomorphisms of M . We investigate the action of $\pi_0(\text{Homeo}(M, \partial M))$ on the moduli space of flat $\text{SL}(2)$ -connections on M .

1.1.1. *Relation to $\pi_1(M)$.* Choose a basepoint $x_0 \in M$ and let $\pi := \pi_1(M; x_0)$. Any homeomorphism of M is isotopic to one which fixes x_0 and hence defines an automorphism of π . Two such isotopic homeomorphisms determine automorphisms of π which differ by an inner automorphism, so there is a well-defined homomorphism

$$(1.1) \quad N : \pi_0(\text{Homeo}(M)) \longrightarrow \text{Out}(\pi) \cong \text{Aut}(\pi)/\text{Inn}(\pi).$$

If M is a closed surface, then Dehn (unpublished) and Nielsen [31] proved that N is an isomorphism. (See Stillwell [36] for a proof of the Dehn-Nielsen theorem.) When $\partial M \neq \emptyset$, then each component $\partial_i M$ determines a conjugacy class C_i of elements of $\pi_1(M)$ and the image of N consists of elements of $\text{Out}(\pi)$ represented by automorphisms which preserve each C_i .

Let M be a one-holed torus. Its fundamental group admits a geometric redundant presentation

$$(1.2) \quad \pi = \langle X, Y, K \mid [X, Y] = K \rangle$$

where K corresponds to the generator of $\pi_1(\partial M)$. Of course, π is freely generated by X, Y .

1.1.2. *Nielsen's theorem.* The following remarkable property of M is due to Jakob Nielsen [30] and does not generalize to other hyperbolic surfaces with boundary. For a proof see Magnus-Karrass-Solitar [24], Theorem 3.9 or Lyndon-Schupp [22], Proposition 5.1.

Proposition 1.1. *Any automorphism of the rank two group*

$$\pi = \langle X, Y, K \mid K = [X, Y] \rangle$$

takes K to a conjugate of either K itself or its inverse K^{-1} .

An equivalent geometric formulation is:

Proposition 1.2. *Every homotopy-equivalence $M \rightarrow M$ is homotopic to a homeomorphism of M .*

Thus the homomorphism (1.1) defines an isomorphism

$$N : \pi_0(\text{Homeo}(M)) \longrightarrow \text{Out}(\pi).$$

1.2. **The structure of the modular group.** We say that an automorphism of π which takes K to either K or K^{-1} is *normalized*. The normalized automorphisms form a subgroup $\text{Aut}(\pi, K)$ of $\text{Aut}(\pi)$.

Let ϕ be an automorphism of π . Proposition 1.1 implies that $\phi(K)$ is conjugate to $K^{\pm 1}$. Thus there exists an inner automorphism ι_g such that

$$\phi(K) = gK^\epsilon g^{-1},$$

that is

$$\iota_g^{-1} \circ \phi \in \text{Aut}(\pi, K)$$

where $\epsilon = \pm 1$. (In fact, $\epsilon = \det(h([\phi]))$, where h is the homomorphism defined by (1.3) below.) Since the centralizer of K in π equals the cyclic group $\langle K \rangle$, the automorphism ϕ determines the coset of g modulo $\langle K \rangle$ uniquely.

We obtain a short exact sequence

$$1 \longrightarrow \langle \iota_K \rangle \hookrightarrow \text{Aut}(\pi, K) \longrightarrow \text{Out}(\pi) \longrightarrow 1$$

The action on the homology $H_1(M; \mathbb{Z}) \cong \mathbb{Z}^2$ defines a homomorphism

$$(1.3) \quad h : \text{Out}(\pi) \longrightarrow \text{GL}(2, \mathbb{Z}).$$

By Nielsen [30], h is an isomorphism. (Surjectivity follows by realizing an element of $\text{GL}(2, \mathbb{Z})$ as a *linear* homeomorphism of the torus $\mathbb{R}^2/\mathbb{Z}^2$. See Lyndon-Schupp [22], Prop. 4.5 or Magnus-Karrass-Solitar [24], §3.5, Corollary N4.) We obtain an isomorphism

$$\mu : \text{GL}(2, \mathbb{Z}) \longrightarrow \text{Aut}(\pi, K)/\langle \iota_K \rangle.$$

Restriction of the composition $\mu \circ h$ to $\text{Aut}(\pi, K)$ equals the quotient homomorphism

$$\text{Aut}(\pi, K) \longrightarrow \text{Aut}(\pi, K)/\langle \iota_K \rangle.$$

2. STRUCTURE OF THE CHARACTER VARIETY

2.1. Trace functions. The relevant moduli space is the *character variety*, the algebro-geometric quotient of $\text{Hom}(\pi, G)$ by the G -action by inner automorphisms, where $G = \text{SL}(2, \mathbb{C})$. Since π is freely generated by two elements X and Y , the set $\text{Hom}(\pi, G)$ of homomorphisms $\pi \longrightarrow G$ identifies with the set of pairs $(\xi, \eta) \in G \times G$, via the mapping

$$\begin{aligned} \text{Hom}(\pi, G) &\longrightarrow G \times G \\ \rho &\longmapsto (\rho(X), \rho(Y)). \end{aligned}$$

This mapping is equivariant respecting the action of G on $\text{Hom}(\pi, G)$ by

$$g : \rho \longmapsto \iota_g \circ \rho$$

and the action of G on $G \times G$ by

$$g : (\xi, \eta) \longmapsto (g\xi g^{-1}, g\eta g^{-1}).$$

The moduli space $\text{Hom}(\pi, G)//G$ consists of equivalence classes of elements of $\text{Hom}(\pi, G) \cong G \times G$ where the equivalence class of a homomorphism ρ is defined as the closure of the G -orbit $G\rho$. Then $\text{Hom}(\pi, G)//G$ is the algebro-geometric quotient in the sense that its

coordinate ring identifies with the ring of G -invariant regular functions on $\text{Hom}(\pi, G)$. For a single element $g \in G$, the conjugacy class $\text{Inn}(G)(g)$ is determined by the trace $t = \text{tr}(g)$ if $t \neq \pm 2$. That is,

$$(2.1) \quad \text{Inn}(G)(g) = \text{tr}^{-1}(t).$$

For $t = \pm 2$, then

$$(2.2) \quad \text{tr}^{-1}(\pm 2) = \{\pm \mathbb{I}\} \cup \text{Inn}(G) \cdot \left(\pm \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix} \right).$$

By Fricke [8] and Fricke-Klein [9], the traces of the generators X, Y, XY parametrize $\text{Hom}(\pi, G)//G$ as the affine space \mathbb{C}^3 . As X and Y freely generate π , we may identify:

$$\begin{aligned} \text{Hom}(\pi, G) &\longleftrightarrow G \times G \\ \rho &\longleftrightarrow (\rho(X), \rho(Y)) \end{aligned}$$

The character mapping

$$\begin{aligned} \chi : \text{Hom}(\pi, G)//G &\longrightarrow \mathbb{C}^3 \\ [\rho] &\longmapsto \begin{bmatrix} x(\rho) \\ y(\rho) \\ z(\rho) \end{bmatrix} = \begin{bmatrix} \text{tr}(\rho(X)) \\ \text{tr}(\rho(Y)) \\ \text{tr}(\rho(XY)) \end{bmatrix} \end{aligned}$$

is an isomorphism. (Compare the discussion in Goldman [13], 4.1, [14], §§4–5 and [15].)

For example, given $(x, y, z) \in \mathbb{C}^3$, the representation ρ defined by

$$(2.3) \quad \rho(X) = \begin{bmatrix} x & -1 \\ 1 & 0 \end{bmatrix}, \quad \rho(Y) = \begin{bmatrix} 0 & \zeta^{-1} \\ -\zeta & y \end{bmatrix}$$

satisfies $\chi(\rho) = (x, y, z)$ where $\zeta \in \mathbb{C}$ is chosen so that

$$\zeta + \zeta^{-1} = z.$$

Conversely, if $(x, y, z) \in \mathbb{C}^3$ then $\chi^{-1}(x, y, z)$ consists of a single G -orbit if and only if $\kappa(x, y, z) \neq 2$ where κ is defined below in (2.4). (This is the condition that ρ is an *irreducible representation* of π . Compare Lubotzky-Magid [21], Brumfiel-Hilden [4], Culler-Shalen [6].)

For any word $w(X, Y)$, the function

$$\begin{aligned} \text{Hom}(\pi, G) &\longrightarrow \mathbb{C} \\ [\rho] &\longmapsto \text{tr}(\rho(w(X, Y))) \end{aligned}$$

is G -invariant. Hence there exists a polynomial $f_w(x, y, z) \in \mathbb{C}[x, y, z]$ such that

$$\text{tr}(\rho(w(X, Y))) = f_w(x(\rho), y(\rho), z(\rho)).$$

A particularly important example occurs for $w(X, Y) = [X, Y] = K$, in which case we denote $f_w(x, y, z)$ by $\kappa(x, y, z)$. By an elementary calculation (see, for example [15]),

$$(2.4) \quad \text{tr}(\rho(K)) = \kappa(x, y, z) := x^2 + y^2 + z^2 - xyz - 2.$$

The level set $\kappa^{-1}(t)$ consists of equivalence classes of representations $\rho : \pi \rightarrow G$ where $\rho(K)$ is constrained to lie in the conjugacy class $\text{tr}^{-1}(t)$. (Compare (2.1) and (2.2) above.)

2.2. Automorphisms. Let $G = \text{SL}(2, \mathbb{C})$. The group $\text{Aut}(\pi)$ acts on the character variety $\text{Hom}(\pi, G) // G$ by:

$$\phi_*([\rho]) = [\rho \circ \phi^{-1}]$$

and since

$$\rho \circ \iota_\gamma = \iota_{\rho(\gamma)} \circ \rho$$

the subgroup $\text{Inn}(\pi)$ acts trivially. Thus $\text{Out}(\pi)$ acts on

$$\text{Hom}(\pi, G) // G \cong \mathbb{C}^3$$

and since an automorphism ϕ of π is determined by

$$(\phi(X), \phi(Y)) = (w_1(X, Y), w_2(X, Y)),$$

the action of ϕ on \mathbb{C}^3 is given by the three polynomials

$$f_{w_1}(x, y, z), f_{w_2}(x, y, z), f_{w_1 w_2}(x, y, z).$$

Hence $\text{Out}(\pi)$ acts on \mathbb{C}^3 by *polynomial automorphisms*. Nielsen's theorem (Proposition 1.1) implies that any such automorphism preserves $\kappa : \mathbb{C}^3 \rightarrow \mathbb{C}$, that is

$$\kappa(f_{w_1}(x, y, z), f_{w_2}(x, y, z), f_{w_1 w_2}(x, y, z)) = \kappa(x, y, z).$$

2.2.1. Sign-change automorphisms. Some automorphisms of the character variety are not induced by automorphisms of π . Namely, the homomorphisms of π into the center $\{\pm \mathbb{I}\} \subset G$ form a group acting on $\text{Hom}(\pi, G)$ by pointwise multiplication. Let $\zeta \in \text{Hom}(\pi, \{\pm \mathbb{I}\})$ and $\rho \in \text{Hom}(\pi, G)$. Then

$$\zeta \cdot \rho : \gamma \mapsto \zeta(\gamma)\rho(\gamma)$$

is a homomorphism. This defines an action

$$\text{Hom}(\pi, \{\pm \mathbb{I}\}) \times \text{Hom}(\pi, G) \rightarrow \text{Hom}(\pi, G).$$

Furthermore, since Z is central in G and $K \in \pi$ is a commutator

$$(2.5) \quad (\zeta \cdot \rho)(K) = \rho(K).$$

Since π is free of rank two,

$$\text{Hom}(\pi, Z) \cong \mathbb{Z}/2 \times \mathbb{Z}/2.$$

The three nontrivial elements $(0, 1), (1, 0), (1, 1)$ act on representations by $\sigma_1, \sigma_2, \sigma_3$ respectively:

$$\sigma_1 \cdot \rho : \begin{cases} X & \mapsto \rho(X) \\ Y & \mapsto -\rho(Y) \end{cases}$$

$$\sigma_2 \cdot \rho : \begin{cases} X & \mapsto -\rho(X) \\ Y & \mapsto \rho(Y) \end{cases}$$

$$\sigma_3 \cdot \rho : \begin{cases} X & \mapsto -\rho(X) \\ Y & \mapsto -\rho(Y) \end{cases}$$

The corresponding action on characters is:

$$\begin{aligned} (\sigma_1)_* : \begin{bmatrix} x \\ y \\ z \end{bmatrix} &\mapsto \begin{bmatrix} x \\ -y \\ -z \end{bmatrix} \\ (\sigma_2)_* : \begin{bmatrix} x \\ y \\ z \end{bmatrix} &\mapsto \begin{bmatrix} -x \\ y \\ -z \end{bmatrix} \\ (\sigma_3)_* : \begin{bmatrix} x \\ y \\ z \end{bmatrix} &\mapsto \begin{bmatrix} -x \\ -y \\ z \end{bmatrix}. \end{aligned}$$

We call this group the *group of sign-changes* and denote it by Σ .

Evidently Σ preserves $\kappa(x, y, z)$. (Alternatively apply (2.5).)

2.2.2. Permutations. Since $\kappa(x, y, z)$ is symmetric in x, y, z the full symmetric group \mathfrak{S}_3 also acts on \mathbb{C}^3 preserving κ . Unlike Σ , elements of \mathfrak{S}_3 are induced by automorphisms of π (compare §A-5). The group of all *linear* automorphisms of (\mathbb{C}^3, κ) is generated by Σ and \mathfrak{S}_3 and forms a semidirect product $\Sigma \rtimes \mathfrak{S}_3$.

\mathfrak{S}_3 is actually a quotient of Γ as follows. The action of $\mathrm{GL}(2, \mathbb{Z})$ on $(\mathbb{Z}/2)^2$ defines a homomorphism

$$\mathrm{GL}(2, \mathbb{Z}) \longrightarrow \mathrm{GL}(2, \mathbb{Z}/2) \cong \mathfrak{S}_3$$

whose kernel $\mathrm{GL}(2, \mathbb{Z})_{(2)}$ is generated by the involutions

$$(2.6) \quad \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}, \begin{bmatrix} 1 & -2 \\ 0 & -1 \end{bmatrix}, \begin{bmatrix} 1 & 0 \\ 2 & -1 \end{bmatrix}$$

and $-\mathbb{I} \in \mathrm{GL}(2, \mathbb{Z})_{(2)}$. The sequence

$$\mathrm{PGL}(2, \mathbb{Z})_{(2)} := \mathrm{GL}(2, \mathbb{Z})_{(2)} / \{\pm \mathbb{I}\} \longrightarrow \mathrm{PGL}(2, \mathbb{Z}) \longrightarrow \mathfrak{S}_3$$

is exact. Let $\Gamma_{(2)}$ denote the kernel of the composite

$$\Gamma \longrightarrow \mathrm{PGL}(2, \mathbb{Z}) \longrightarrow \mathfrak{S}_3,$$

the semidirect product

$$\mathrm{PGL}(2, \mathbb{Z})_{(2)} \ltimes (\mathbb{Z}/2 \oplus \mathbb{Z}/2).$$

2.3. Reducible characters. Any representation ρ having character in $\kappa^{-1}(2)$ is reducible (the invariant bilinear form is degenerate). In that case ρ is conjugate to an upper-triangular representation:

$$(2.7) \quad \begin{aligned} \rho(X) &= \begin{bmatrix} \xi & * \\ 0 & \xi^{-1} \end{bmatrix} \\ \rho(Y) &= \begin{bmatrix} \eta & * \\ 0 & \eta^{-1} \end{bmatrix} \\ \rho(XY) &= \begin{bmatrix} \zeta & * \\ 0 & \zeta^{-1} \end{bmatrix}. \end{aligned}$$

where $\xi\eta = \zeta$. Thus

$$(2.8) \quad \begin{aligned} x &= \xi + \xi^{-1}, \\ y &= \eta + \eta^{-1}, \\ z &= \zeta + \zeta^{-1}. \end{aligned}$$

(2.8) corresponds to the following factorization (2.9) of

$$\kappa(x, y, z) - 2 = x^2 + y^2 + z^2 - xyz - 4.$$

Under the embedding

$$\mathbb{C}[x, y, z] \longrightarrow \mathbb{C}[\xi, \xi^{-1}, \eta, \eta^{-1}, \zeta, \zeta^{-1}]$$

defined by (2.8), the polynomial $\kappa(x, y, z) - 2$ factors:

$$(2.9) \quad \begin{aligned} \kappa(\xi + \xi^{-1}, \eta + \eta^{-1}, \zeta + \zeta^{-1}) - 2 = \\ \zeta^{-2}(1 - \xi\eta\zeta)(1 - \xi^{-1}\eta\zeta)(1 - \xi\eta^{-1}\zeta)(1 - \xi\eta\zeta^{-1}). \end{aligned}$$

Given (x, y, z) with $\kappa(x, y, z) = 2$, the triple (ξ, η, ζ) is only defined up to $\mathbb{Z}/2 \oplus \mathbb{Z}/2$. Namely the automorphisms

$$\begin{cases} X & \longmapsto X^{-1} \\ Y & \longmapsto Y \end{cases}$$

$$\begin{cases} X & \longmapsto X \\ Y & \longmapsto Y^{-1} \end{cases}$$

$$\begin{cases} X & \longmapsto X^{-1} \\ Y & \longmapsto Y^{-1} \end{cases}$$

act on (ξ, η, ζ) by:

$$\begin{cases} \xi & \longmapsto \xi^{-1} \\ \eta & \longmapsto \eta \\ \zeta & \longmapsto \zeta^{-1} \end{cases}$$

$$\begin{cases} \xi & \longmapsto \xi \\ \eta & \longmapsto \eta^{-1} \\ \zeta & \longmapsto \zeta^{-1} \end{cases}$$

$$\begin{cases} \xi & \longmapsto \xi^{-1} \\ \eta & \longmapsto \eta^{-1} \\ \zeta & \longmapsto \zeta \end{cases}$$

respectively.

2.4. The Poisson structure. As in Goldman [14], §5.3, the automorphisms preserving κ are unimodular and therefore preserve the exterior bivector field as well

$$\begin{aligned}\Xi &= \frac{1}{2\pi^2} d\kappa \cdot \frac{\partial}{\partial x} \wedge \frac{\partial}{\partial y} \wedge \frac{\partial}{\partial z} \\ &= \frac{1}{2\pi^2} \left((2x - yz) \frac{\partial}{\partial y} \wedge \frac{\partial}{\partial z} \right. \\ &\quad \left. + (2y - zx) \frac{\partial}{\partial z} \wedge \frac{\partial}{\partial x} \right. \\ &\quad \left. + (2z - xy) \frac{\partial}{\partial x} \wedge \frac{\partial}{\partial y} \right)\end{aligned}$$

which restricts to (the dual of) an area form on each level set $\kappa^{-1}(t)$ which is invariant under $\text{Out}(\pi)$. We shall always consider this invariant measure on $\kappa^{-1}(t) \cap \mathbb{R}^3$. In the case when $\kappa^{-1}(t)$ has a rational parametrization by an affine plane, that is, when $t = 2$, the above bivector field is the image of the constant bivector field under the parametrization:

$$\begin{aligned}\mathbb{C} \times \mathbb{C} &\longrightarrow \kappa^{-1}(2) \\ (\xi, \eta) &\longmapsto \begin{bmatrix} e^\xi + e^{-\xi} \\ e^\eta + e^{-\eta} \\ e^{\xi+\eta} + e^{-(\xi+\eta)} \end{bmatrix} \\ \frac{\partial}{\partial \xi} \wedge \frac{\partial}{\partial \eta} &\longmapsto 2\pi^2 \Xi\end{aligned}$$

The bivector field Ξ defines a *Poisson structure* on the moduli space \mathbb{C}^3 for which κ defines a *Casimir function*.

(The coefficient $1/(2\pi^2)$ occurs to normalize the area of the compact component C_K of reducible $\text{SU}(2)$ -characters to 1.) Here we study the action of $\text{Out}(\pi)$ on the set $\kappa^{-1}(t) \cap \mathbb{R}^3$ of \mathbb{R} -points of $\kappa^{-1}(t)$ where $t \in \mathbb{R}$, with respect to this invariant measure.

2.5. The orthogonal representation. To describe the structure of the character variety more completely, we use a 3-dimensional orthogonal representation associated to a character $(x, y, z) \in \mathbb{C}^3$. (Compare §4.2 of Goldman [13].) Let $V = \mathbb{C}^3$ with the standard basis $\{e_1, e_2, e_3\}$ and bilinear form defined by the symmetric matrix

$$\mathbb{B} = \begin{bmatrix} 2 & z & y \\ z & 2 & x \\ y & x & 2 \end{bmatrix}.$$

Since

$$\det(\mathbb{B}) = -2(\kappa(x, y, z) - 2)$$

(where $\kappa(x, y, z)$ is defined in (2.4)), the symmetric bilinear form is nondegenerate if and only if $\kappa(x, y, z) \neq 2$. For any vector $v \in V$ such that $\mathbb{B}(v, v) \neq 0$, the reflection

$$R_v : x \longmapsto x - 2 \frac{\mathbb{B}(v, x)}{\mathbb{B}(v, v)} v$$

is a \mathbb{B} -orthogonal involution. We define an orthogonal representation $\rho_{\mathbb{B}}$ of π on (V, \mathbb{B}) by

$$\rho_{\mathbb{B}} : \begin{cases} X & \longmapsto R_{e_2} R_{e_3} \\ Y & \longmapsto R_{e_3} R_{e_1} \end{cases}$$

The adjoint representation Ad of $\text{SL}(2, \mathbb{C})$ is a representation on a three-dimensional vector space V over \mathbb{C} . The group π embeds as the torsionfree index-two subgroup in the group generated by orthogonal reflections in the coordinate axes in the vector space V . When $\kappa(x, y, z) \neq 2$, this form is nondegenerate and the original representation ρ is the lift of a representation in the orthogonal group $\text{SO}(3, \mathbb{C})$ under the double covering

$$\text{SL}(2, \mathbb{C}) \longrightarrow \text{SO}(3, \mathbb{C}).$$

2.6. The set of \mathbb{R} -points. A real character is the character of a representation into a *real form* of G . There are two conjugacy classes of real forms of G , compact or noncompact. Every compact real form of G is conjugate to $\text{SU}(2)$ and every noncompact real form of G is conjugate to $\text{SL}(2, \mathbb{R})$.

Thus every real character $(x, y, z) \in \mathbb{R}^3$ is the character of a representation into either $\text{SU}(2)$ or $\text{SL}(2, \mathbb{R})$. (Proposition III.1.1 of Morgan-Shalen [27]).

Given $(x, y, z) \in \mathbb{R}$, the formula (2.3) gives an explicit representation in $\text{SL}(2, \mathbb{R})$ when $|z| \geq 2$. Consider the orthogonal representation described in §2.5 to construct an invariant symmetric bilinear form \mathbf{B} on a three-dimensional vector space V over \mathbb{C} . The group π embeds as the torsionfree index-two subgroup in the group generated by orthogonal reflections in the coordinate axes in the vector space V . When $\kappa(x, y, z) \neq 2$, this form is nondegenerate and the original representation ρ is the lift of a representation in the orthogonal group $\text{SO}(3, \mathbb{C})$ under the double covering

$$\text{SL}(2, \mathbb{C}) \longrightarrow \text{SO}(3, \mathbb{C}).$$

When $x, y, z \in \mathbb{R}$, the restriction of \mathbf{B} to \mathbb{R}^3 is definite if and only if $-2 \leq x, y, z \leq 2$ and $\kappa(x, y, z) < 2$.

The coordinates of characters of representations $\pi_1(M) \longrightarrow \mathrm{SU}(2)$ satisfy:

$$-2 \leq x, y, z \leq 2, \quad x^2 + y^2 + z^2 - xyz - 2 \leq 2$$

and, for fixed $t \in [-2, 2]$, comprise a component of $\kappa^{-1}(t) \cap \mathbb{R}^3$. The other four connected components of $\kappa^{-1}(t) \cap \mathbb{R}^3$ consist of characters of $\mathrm{SL}(2, \mathbb{R})$ -representations. These four components are freely permuted by Σ . For $t < -2$, all four components of the relative character variety $\kappa^{-1}(t) \cap \mathbb{R}^3$ consist of character of $\mathrm{SL}(2, \mathbb{R})$ -representations and are freely permuted by Σ . For $t > 2$, the relative character variety $\kappa^{-1}(t) \cap \mathbb{R}^3$ is connected and consists of characters of $\mathrm{SL}(2, \mathbb{R})$ -representations.

The two critical values ± 2 of $\mathrm{tr} : G \longrightarrow \mathbb{C}$ deserve special attention. When $t = -2$, a representation $\rho \in \mathrm{Hom}(\pi, G)$ with $\mathrm{tr} \rho(K) = -2$ is a regular point of the mapping

$$\begin{aligned} E_K : \mathrm{Hom}(\pi, G) &\longrightarrow G \\ \rho &\longmapsto \rho(K). \end{aligned}$$

Such a representation is a regular point of the composition

$$\mathrm{tr} \circ E_K : \mathrm{Hom}(\pi, G) \longrightarrow \mathbb{C}$$

unless $\rho(K) = -I$. In the latter case (when $[\rho(X), \rho(Y)] = -I$), the representation is conjugate to the *quaternion representation* in $\mathrm{SU}(2)$

$$(2.10) \quad \rho(X) = \begin{bmatrix} i & 0 \\ 0 & -i \end{bmatrix}, \quad \rho(Y) = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}.$$

In both cases, G acts locally freely on the subset $(\mathrm{tr} \circ E_K)^{-1}(-2)$ of $\mathrm{Hom}(\pi, G)$ with quotient $\kappa^{-1}(-2)$.

The \mathbb{R} -points correspond to characters of representations which lie in the Cartan subgroups (maximal tori) of the real forms $\mathrm{SU}(2)$ and $\mathrm{SL}(2, \mathbb{R})$. Every Cartan subgroup of $\mathrm{SU}(2)$ is conjugate to $\mathrm{U}(1)$, and every Cartan subgroup of $\mathrm{SL}(2, \mathbb{R})$ is conjugate to either $\mathrm{SO}(2)$ or $\mathrm{SO}(1, 1)$. Characters of reducible $\mathrm{SU}(2)$ -representations form the compact set

$$C_K = \kappa^{-1}(2) \cap [-2, 2]^3$$

which identifies with the quotient of the 2-torus $\mathrm{U}(1) \times \mathrm{U}(1)$ by $\{\pm \mathbb{I}\}$ under the extension (2.8).

Characters of reducible $\mathrm{SL}(2, \mathbb{R})$ -representations comprise four components

$$\begin{aligned} C_0 &= \kappa^{-1}(2) \cap \left([2, \infty) \times [2, \infty) \times [2, \infty) \right), \\ C_1 &= \kappa^{-1}(2) \cap \left([2, \infty) \times (-\infty, -2] \times (-\infty, -2] \right), \\ C_2 &= \kappa^{-1}(2) \cap \left((-\infty, -2] \times [2, \infty) \times (-\infty, -2] \right), \\ C_3 &= \kappa^{-1}(2) \cap \left((-\infty, -2] \times (-\infty, -2] \times [2, \infty) \right), \end{aligned}$$

each of which identifies with the quotient of $\mathbb{R}_+ \times \mathbb{R}_+$ by $\{\pm \mathbb{I}\}$ under (2.8). These four components are freely permuted by Σ . There is a compact component of equivalence classes of $\mathrm{SL}(2, \mathbb{R})$ -representations which are irreducible but not *absolutely irreducible* — that is, although \mathbb{R}^2 is an irreducible π -module, its complexification is reducible. In that case the representation is conjugate to a representation in $\mathrm{SO}(2)$. This component agrees with the component C_K consisting of characters of reducible $\mathrm{SU}(2)$ -representations.

3. HYPERBOLIC STRUCTURES ON TORI ($t < 2$)

The topology and dynamics changes dramatically as t changes from $t < -2$ to $t > 2$. The level sets $t < -2$ correspond to Fricke spaces of one-holed tori with geodesic boundary. The level set for $t = -2$ corresponds to the Teichmüller space of the punctured torus. The level sets for $-2 < t < 2$ correspond to Teichmüller spaces of singular hyperbolic structures with a single point, as well as a component of unitary representations. Except for the component of unitary representations, there are four components, freely permuted by Σ . Except for the component of unitary representations, the Γ -action is proper.

3.1. Complete hyperbolic structures on punctured tori ($t \leq -2$). The theory of deformations of geometric structures implies that Γ acts properly on certain components of $\kappa^{-1}(t)$. When $t < 2$, the moduli space $\kappa^{-1}(t)$ contains 4 contractible noncompact components, freely permuted by Σ . (When $-2 \leq t < 2$, an additional compact component corresponds to $\mathrm{SU}(2)$ -representations.) These contractible components correspond to $\mathrm{SL}(2, \mathbb{R})$ -representations, and each one identifies with the Teichmüller space of M , with certain boundary conditions. Specifically, if $t < -2$, then these components correspond to hyperbolic structures on $\mathrm{int}(M)$ with geodesic boundary of length $2 \cosh^{-1}(-t/2)$. For $t =$

-2 , these components correspond to complete hyperbolic structures on $\text{int}(M)$ and identify with the usual Teichmüller space \mathfrak{T}_M . For $-2 < t < 2$, these components correspond to singular hyperbolic structures on a torus whose singularity is an isolated point with cone angle

$$\theta = 2 \cos^{-1}(-t/2).$$

3.2. Complete structures and proper actions. We begin with the case $t < -2$. Then $\kappa^{-1}(t) \cap \mathbb{R}^3$ parametrizes complete hyperbolic structures on $\text{int}(M)$ with a closed geodesic parallel to ∂M having length $2 \cosh^{-1}(-t/2)$. Thus the union

$$\bigcup_{t \leq -2} \kappa^{-1}(t)$$

is the Fricke space of M . This space, denoted $\mathfrak{F}(M)$, contains equivalence classes of marked complete hyperbolic structures on $\text{int}(M)$, not necessarily of finite area. The properness of the action of the Teichmüller modular group $\pi_0(\text{Homeo}(M))$ on $\mathfrak{F}(M)$ implies properness of the Γ -action of $\kappa^{-1}(t)$. (For proof, see §2.2 of Abikoff [1], Bers-Gardiner [3] 6.5.6 (p.156) of Buser [5], 6.3 of Iwayoshi-Tanigawa [20], 2.4.1 of Harvey [17], or §2.7 of Nag [29].)

When $t = -2$, then $\kappa^{-1}(t)$ is the union of the single point $(0, 0, 0)$ and four copies of the Teichmüller space \mathfrak{T}_M of M , comprising equivalence classes of marked complete finite-area hyperbolic structures on M . The mapping class group $\pi_0(\text{Homeo}(M))$ acts properly on $\kappa^{-1}(t)$.

3.3. Singular hyperbolic structures on tori ($-2 < t < 2$). We next consider the case $2 > t > -2$. This case is similar to the previous case, except that the ends are replaced by cone points. Using the properness of the mapping class group action on Teichmüller space, the action on these components of the relative character variety remains proper. However, none of the corresponding representations in $\text{Hom}(\pi, G)$ are discrete embeddings; typically the representations will be isomorphisms onto dense subgroups of $\text{SL}(2, \mathbb{R})$.

We first show that every representation in these components are holonomy representations of singular hyperbolic structures. §3.5 proves that Γ acts properly on the level sets $\kappa^{-1}(t)$ where $-2 < t < 2$.

3.4. Construction of hyperbolic structures on T^2 with one cone point. Let $\theta > 0$ and let \mathfrak{C}_θ denote the space of hyperbolic structures on T^2 with a conical singularity of cone angle θ . By results of McOwen [26] and Troyanov [37], \mathfrak{C}_θ identifies with the Teichmüller space $\mathfrak{T}(\theta)$, the deformation space of conformal structures on T^2 singular at one point with a cone angle θ there.

Theorem 3.1. *Suppose $\xi, \eta \in \mathrm{SL}(2, \mathbb{R})$ and $\gamma := [\xi, \eta]$ is elliptic of rotation angle θ . Let M denote a punctured torus whose fundamental group has the presentation (1.2). Then there exists a singular hyperbolic structure on \bar{M} with a singularity of cone angle θ having holonomy representation ρ defined by*

$$\rho(X) = \xi, \quad \rho(Y) = \eta, \quad \rho(K) = \gamma.$$

The proof will be based on the following.

Lemma 3.2. *Let $p \in \mathbf{H}^2$ be a point fixed by γ and consider the points*

$$\begin{aligned} p_4 &= p, \\ p_3 &= \eta^{-1}p, \\ p_2 &= \xi^{-1}\eta^{-1}p, \\ p_1 &= \eta\xi^{-1}\eta^{-1}p. \end{aligned}$$

The four points p_1, p_2, p_3, p_4 are the vertices of an embedded quadrilateral \mathbf{Q} . In other words the four segments

$$\begin{aligned} l_1 &= \overline{p_1 p_2} \\ l_2 &= \overline{p_2 p_3} \\ l_3 &= \overline{p_3 p_4} \\ l_4 &= \overline{p_4 p_1} \end{aligned}$$

are disjoint and bound a quadrilateral.

Proof of Theorem 3.1 assuming Lemma 3.2.

ξ maps p_1, p_2 respectively to p_4, p_3 respectively and η maps p_2, p_3 respectively to p_1, p_4 respectively. Therefore ξ maps the directed edge l_1 to l_3 with the opposite orientation and η maps l_2 to l_4 with the opposite orientation.

From the embedding of \mathbf{Q} and the identifications of its opposite sides by ξ and η , we construct a developing map for a (singular) hyperbolic structure on M as follows. Let \mathcal{Q} denote an *abstract quadrilateral*, that is, a cell complex with a single 2-cell, four 1-cells (the *edges*), and four 0-cells (the *vertices*). Denote the oriented edges \mathcal{L}_i numbered in cyclic order and the vertices p_i ($i = 1, 2, 3, 4$) where

$$\begin{aligned} \partial\mathcal{L}_1 &= \{p_1, p_2\}, & \partial\mathcal{L}_2 &= \{p_2, p_3\} \\ \partial\mathcal{L}_3 &= \{p_3, p_4\}, & \partial\mathcal{L}_4 &= \{p_4, p_1\}. \end{aligned}$$

Let $\bar{\mathcal{L}}_i$ denote \mathcal{L}_i with the opposite orientation. Write

$$\ddot{\mathcal{Q}} = \mathcal{Q} - \{p_1, p_2, p_3, p_4\}.$$

Choose homeomorphisms

$$\tilde{\xi} : \mathcal{L}_1 \longrightarrow \bar{\mathcal{L}}_3, \quad \tilde{\eta} : \mathcal{L}_2 \longrightarrow \bar{\mathcal{L}}_4.$$

Let $\Pi = \langle X, Y \rangle$ denote the free group generated by X, Y . A model for the universal covering \tilde{M} of M is the quotient space of the Cartesian product

$$\ddot{\mathcal{Q}} \times \Pi$$

by the equivalence relation generated by the identifications

$$(x, \eta) \longleftarrow (\tilde{\xi}(x), X\eta)$$

for $x \in \mathcal{L}_1$, and

$$(x, \eta) \longleftarrow (\tilde{\eta}(x), Y\eta)$$

for $x \in \mathcal{L}_2$. The action of Π on $\mathcal{Q} \times \Pi$ defined by the trivial action on \mathcal{Q} and right-multiplication on Π descends to a free proper action of Π on \tilde{M} . This action corresponds to the action of deck transformations. Let $\rho : \Pi \longrightarrow G$ be the homomorphism defined by:

$$\rho(X) = \xi$$

$$\rho(Y) = \eta$$

The embedding $\ddot{\mathcal{Q}} \longrightarrow \mathbf{Q}$ together with the identifications of the sides of \mathbf{Q} extends uniquely to a ρ -equivariant local homeomorphism $\tilde{M} \longrightarrow \mathbf{H}^2$. The holonomy around the puncture

$$\rho([X, Y]) = [\xi, \eta] = \gamma$$

is elliptic of rotation angle θ fixing p . The resulting hyperbolic structure extends to a singular hyperbolic structure on \bar{M} with a singularity of cone angle θ . Thus Theorem 3.1 reduces to Lemma 3.2. \square

The proof of Lemma 3.2 is based on the following.

Lemma 3.3. *If the sequence of points p_1, p_2, p_3, p_4 are not the vertices of an embedded quadrilateral, then either l_1 and l_3 intersect, or l_2 and l_4 intersect.*

Proof. If $x, y \in \mathbf{H}^2$ are distinct points, let \overleftrightarrow{xy} denote the line passing through x and y . If $z \notin \overleftrightarrow{xy}$, let $H_{x,y}(z)$ denote the component (a half-space) of $\mathbf{H}^2 - \overleftrightarrow{xy}$ containing z , and $\hat{H}_{x,y}(z)$ denote the complementary component:

$$\mathbf{H}^2 = H_{x,y}(z) \cup \overleftrightarrow{xy} \cup \hat{H}_{x,y}(z)$$

Assume, to start, that no three of the points p_1, p_2, p_3, p_4 are collinear. There are four cases, depending on where p_4 lies in relation to the decompositions determined by the lines $\overleftrightarrow{p_1p_2}$ and $\overleftrightarrow{p_2p_3}$:

- (1) $p_4 \in H_{p_1p_2}(p_3) \cap H_{p_2p_3}(p_1)$;

- (2) $p_4 \in \hat{H}_{p_1p_2}(p_3) \cap H_{p_2p_3}(p_1)$;
- (3) $p_4 \in H_{p_1p_2}(p_3) \cap \hat{H}_{p_2p_3}(p_1)$;
- (4) $p_4 \in \hat{H}_{p_1p_2}(p_3) \cap \hat{H}_{p_2p_3}(p_1)$.

In the first case p_1, p_2, p_3, p_4 are the vertices of a convex quadrilateral. (Compare Figure 9.) In the second case, $\overline{p_3p_4}$ meets $\overline{p_1p_2}$. (Compare Figure 10.) In the third case, $\overline{p_2p_3}$ meets $\overline{p_4p_1}$. (Compare Figure 11.) In the fourth case p_1, p_2, p_3, p_4 are the vertices of an embedded (nonconvex) quadrilateral. The case when p_1, p_2, p_3, p_4 are collinear is similar. (Compare Figure 12.) This completes the proof of Lemma 3.3. \square

Returning to the proof of Lemma 3.2, we show that l_1 and l_3 cannot intersect; an identical proof implies l_2 and l_4 cannot intersect.

Claim. *Suppose that l_1 and l_3 intersect and let $q = l_1 \cap l_3$. The triangles $\triangle(qp_1p_4)$ and $\triangle(qp_3p_2)$ are congruent.*

Proof of Claim. Since $\xi(l_1) = l_3$ and $\eta(l_2) = l_4$, the lengths of opposite sides are equal:

$$\begin{aligned} d(p_1, p_2) &= \ell(l_1) = \ell(l_3) = d(p_4, p_3), \\ d(p_2, p_3) &= \ell(l_2) = \ell(l_4) = d(p_1, p_4) \end{aligned}$$

Since the lengths of the corresponding sides are equal, the triangle $\triangle(p_2p_1p_4)$ is congruent to $\triangle(p_4p_3p_2)$. In particular

$$\angle(p_2p_1p_4) = \angle(p_4p_3p_2).$$

Similarly, $\angle(p_3p_4p_1) = \angle(p_1p_2p_3)$. Now

$$\angle(qp_1p_4) = \angle(p_2p_1p_4) = \angle(p_4p_3p_2) = \angle(qp_3p_2)$$

and similarly, $\angle(qp_4p_1) = \angle(qp_2p_3)$. Since $l_4 = \overline{p_1p_4}$ is congruent to $l_2 = \overline{p_3p_2}$, triangles $\triangle(qp_1p_4)$ and $\triangle(qp_3p_2)$ are congruent as claimed. \square

Now let l be the line through q bisecting the angle $\angle(p_1qp_3)$ such that reflection R in l interchanges $\triangle(qp_1p_4)$ and $\triangle(qp_3p_2)$. Since

$$\begin{aligned} R : \begin{cases} p_1 & \mapsto p_3 \\ p_2 & \mapsto p_4 \\ p_3 & \mapsto p_1 \\ p_4 & \mapsto p_2, \end{cases} \\ \xi : \begin{cases} p_1 & \mapsto p_4 \\ p_2 & \mapsto p_3 \end{cases} \\ \eta : \begin{cases} p_2 & \mapsto p_1 \\ p_3 & \mapsto p_4, \end{cases} \end{aligned}$$

$\xi \circ R$ interchanges p_3 and p_4 , and $R \circ \eta$ interchanges p_2 and p_3 . Since an orientation-reversing isometry of \mathbf{H}^2 which interchanges two points must be reflection in a line, $\xi \circ R$ and $R \circ \eta$ have order two. It follows that R conjugates ξ to ξ^{-1} and η to η^{-1} .

These conditions imply either one of two possibilities:

- At least one of ξ and η is elliptic or parabolic;
- Both ξ and η are hyperbolic, and their invariant axes are each orthogonal to l .

Neither possibility occurs, due to the following:

Lemma 3.4. *Let $\xi, \eta \in \mathrm{SL}(2, \mathbb{R})$. The following conditions are equivalent:*

- $\mathrm{tr}[\xi, \eta] < 2$;
- ξ, η are hyperbolic elements and their invariant axes cross.

Proof. Assuming $\mathrm{tr}[\xi, \eta] < 2$, we first show that ξ and η must be hyperbolic. We first show that ξ is not elliptic. If ξ is elliptic (or $\pm\mathbb{I}$), we may assume that $\xi \in \mathrm{SO}(2)$, that is, we represent ξ, η by matrices

$$\xi = \begin{bmatrix} \cos(\theta) & -\sin(\theta) \\ \sin(\theta) & \cos(\theta) \end{bmatrix}, \quad \eta = \begin{bmatrix} a & b \\ c & d \end{bmatrix},$$

where $ad - bc = 1$, whence

$$\mathrm{tr}[\xi, \eta] = 2 + \sin^2(\theta)(a^2 + b^2 + c^2 + d^2 - 2) \geq 2.$$

Similarly if ξ is parabolic, represent ξ, η by matrices

$$\xi = \pm \begin{bmatrix} 1 & s \\ 0 & 1 \end{bmatrix}, \quad \eta = \begin{bmatrix} a & b \\ c & d \end{bmatrix},$$

where $s \neq 0$ $ad - bc = 1$, whence

$$\mathrm{tr}[\xi, \eta] = 2 + s^2 c^2 \geq 2.$$

Thus ξ is hyperbolic. Since $[\eta, \xi] = [\xi, \eta]^{-1}$, an identical argument shows that η is hyperbolic. Denote their invariant axes by l_ξ, l_η respectively.

It remains to show that $\mathrm{tr}[\xi, \eta] < 2$ if and only if $l_\xi \cap l_\eta \neq \emptyset$.

By conjugation, we may assume that the fixed points of ξ are ± 1 and that the fixed points of η are r, ∞ . Thus $l_\xi \cap l_\eta \neq \emptyset$ if and only if $-1 < r < 1$. Represent ξ, η by matrices

$$\xi = \begin{bmatrix} \cosh(\theta) & \sinh(\theta) \\ \sinh(\theta) & \cosh(\theta) \end{bmatrix}, \quad \eta = \begin{bmatrix} e^\phi & -2r \sinh(\phi) \\ 0 & e^{-\phi} \end{bmatrix},$$

where $\theta, \phi \neq 0$ and

$$\mathrm{tr}[\xi, \eta] = 2 + 4(r^2 - 1) \sinh^2(\theta) \sinh^2(\phi).$$

Then $\text{tr}[\xi, \eta] < 2$ if and only if $-1 < r < 1$ and $\text{tr}[\xi, \eta] = 2$ if and only if $r = \pm 1$, as desired. \square

This completes the proof of Lemma 3.2 (and also Theorem 3.1). \square

3.5. Properness. That Γ acts properly on $\kappa^{-1}(t) \cap \mathbb{R}^3$ now follows easily. The construction in Theorem 3.1 gives a map

$$(3.1) \quad \kappa^{-1}(t) \cap \mathbb{R}^3 \longrightarrow \mathfrak{T}(\theta)$$

which is evidently $\text{Out}(\pi)$ -equivariant. Every hyperbolic structure with conical singularities has an underlying *singular conformal structure*, where the singularities are again conical singularities, that is, they are defined by local coordinate charts to a model space, which in this case is a cone. However, there is an important difference. Conical singularities in conformal structures are *removable*, while conical singularities in Riemannian metrics are not.

Here is why conformal conical singularities are removable: Let $D^2 \subset \mathbb{C}$ be the unit disk and let

$$S_\theta := \{z \in D^2 \mid 0 \leq \arg(z) \leq \theta\}$$

be a sector of angle θ . Then

$$\begin{aligned} \Pi_\theta : S_\theta - \{0\} &\longrightarrow D^2 - \{0\} \\ z &\longmapsto z^{2\pi/\theta} \end{aligned}$$

is conformal. The model coordinate patch for a cone point of angle θ is the *cone* C_θ of angle θ , defined as identification space C_θ of S_θ by the equivalence relation defined by

$$z \longleftarrow e^{\pm i\theta} z$$

for $z \in \partial S_\theta$. That is, a cone point p has a coordinate patch neighborhood U and a coordinate chart $\psi : U \longrightarrow C_\theta$ in the atlas defining the singular geometric structure. The power map Π_θ defines a conformal isomorphism between the C_θ punctured at the cone point and a punctured disk (a “cone” of angle 2π). Replacing the coordinate chart $\psi : U \longrightarrow \mathbb{C}$ at a cone point p of angle θ by the composition $\Pi_\theta \circ \psi$ gives a coordinate atlas for a conformal structure which is nonsingular at p and isomorphic to the original structure on the complement of p .

The resulting map $\mathfrak{T}(\theta) \longrightarrow \mathfrak{T}_M$ is evidently $\text{Out}(\pi)$ -equivariant. Since $\text{Out}(\pi)$ acts properly on \mathfrak{T}_M , $\text{Out}(\pi)$ acts properly on $\mathfrak{T}(\theta)$, and hence on $\kappa^{-1}(t) \cap \mathbb{R}^3$ as well.

With more work one can show that (3.1) is an isomorphism. For any hyperbolic structure on T^2 with a cone point p of angle $0 < \theta < 2\pi$, the Arzelá-Ascoli theorem (as in Buser [5] §1.5) applies to represent ξ, η by

geodesic loops based at p . Furthermore these geodesics intersect only at p . By developing this singular geometric structure one obtains a polygon \mathbf{Q} as above.

4. REDUCIBLE CHARACTERS ($t = 2$)

The level set $\kappa^{-1}(2)$ consists of characters of reducible representations. Over \mathbb{C} such a representation is upper-triangular (2.7) with character defined by (2.8). By (2.9), every $(x, y, z) \in \kappa^{-1}(2)$ lies in the image of the map

$$(4.1) \quad \Phi : \mathbb{C}^* \times \mathbb{C}^* \longrightarrow \kappa^{-1}(2)$$

$$(\xi, \eta) \longmapsto \begin{bmatrix} \xi + \xi^{-1} \\ \eta + \eta^{-1} \\ \xi\eta + \xi^{-1}\eta^{-1} \end{bmatrix}.$$

The set of \mathbb{R} -points $\kappa^{-1}(2) \cap \mathbb{R}^3$ is a singular algebraic hypersurface in \mathbb{R}^3 , with singular set

$$S_0 = \left\{ \begin{bmatrix} 2 \\ 2 \\ 2 \end{bmatrix}, \begin{bmatrix} 2 \\ -2 \\ -2 \end{bmatrix}, \begin{bmatrix} -2 \\ 2 \\ -2 \end{bmatrix}, \begin{bmatrix} -2 \\ -2 \\ 2 \end{bmatrix} \right\}.$$

Characters in S_0 correspond to unipotent representations twisted by central characters. A *central character* is a homomorphism taking values in the center $\{\pm\mathbb{I}\}$ of G and a *unipotent representation* is a representation in $\pm U$, where U is a unipotent subgroup of G . A reductive representation with character in S_0 is itself a central character. The most general representation with character in S_0 is one taking values in $\pm U$, where U is a unipotent subgroup of G . The character $(2, 2, 2)$ is the character of any unipotent representation, for example the trivial representation. The other three points are images of $(2, 2, 2)$ by the three nontrivial elements of Σ .

The smooth stratum of $\kappa^{-1}(2)$ is the complement

$$\kappa^{-1}(2) \cap \mathbb{R}^3 - S_0$$

and has five components, denoted by C_K and C_i , where $i = 0, 1, 2, 3$. Here C_0 denotes the component $\kappa^{-1}(2) \cap (\mathbb{R}_+)^3$ and $C_i = (\sigma_i)_* C_0$ where $(\sigma_i)_* \in \Sigma$ is the sign-change automorphism which fixes the i -th coordinate. The component C_K corresponds to reducible $\mathrm{SU}(2)$ -representations which are *non-central*, that is, their image does not lie in the center $\{\pm\mathbb{I}\}$ of $\mathrm{SU}(2)$. The closure of C_K is the union of C_K with

S_0 . The map

$$\begin{aligned} \mathrm{U}(1) \times \mathrm{U}(1) &\longrightarrow \bar{C}_K \\ (\xi, \eta) &\longmapsto (\xi + \xi^{-1}, \eta + \eta^{-1}, \xi\eta + \xi^{-1}\eta^{-1}) \end{aligned}$$

is a double branched covering space, with deck transformation

$$(\xi, \eta) \longmapsto (\xi^{-1}, \eta^{-1})$$

and four branch points

$$(\xi, \eta) = (\pm 1, \pm 1)$$

which map to S_0 .

Similarly, each of the other components identifies with the quotient $(\mathbb{R}_+)^2/\{\pm \mathbb{I}\}$ with the action of $\mathrm{GL}(2, \mathbb{Z})$. For example, \bar{C}_0 is the image of the double branched covering

$$\begin{aligned} \mathbb{R}_+ \times \mathbb{R}_+ &\longrightarrow \bar{C}_0 \\ (\xi, \eta) &\longmapsto (\xi + \xi^{-1}, \eta + \eta^{-1}, \xi\eta + \xi^{-1}\eta^{-1}), \end{aligned}$$

with deck transformation

$$(\xi, \eta) \longmapsto (\xi^{-1}, \eta^{-1}).$$

For $i = 1, 2, 3$, the component C_i is the image of C_0 under the sign change $(\sigma_i)_*$. Composing with the $\mathrm{GL}(2, \mathbb{Z})$ -equivariant diffeomorphisms

$$\begin{aligned} \mathbb{R}^2 &\longrightarrow \mathbb{R}_+ \times \mathbb{R}_+ \\ (\tilde{\xi}, \tilde{\eta}) &\longmapsto (\exp(\tilde{\xi}), \exp(\tilde{\eta})) \end{aligned}$$

and

$$\begin{aligned} (\mathbb{R}/\mathbb{Z})^2 &\longrightarrow \mathrm{U}(1) \times \mathrm{U}(1) \\ (\tilde{\xi}, \tilde{\eta}) &\longmapsto (\exp(2\pi i \tilde{\xi}), \exp(2\pi i \tilde{\eta})) \end{aligned}$$

respectively, yields $\mathrm{GL}(2, \mathbb{Z})$ -equivariant double branched coverings

$$\mathbb{R}^2 \longrightarrow C_i, \text{ and } (\mathbb{R}/\mathbb{Z})^2 \longrightarrow C_K$$

respectively. Furthermore these mappings pull back the invariant area form on $\kappa^{-1}(2) \cap \mathbb{R}^3$ to Lebesgue measure on \mathbb{R}^2 .

Since $\mathrm{SL}(2, \mathbb{Z})$ is a lattice in $\mathrm{SL}(2, \mathbb{R})$ and $\mathrm{SL}(2, \mathbb{R})$ acts transitively on \mathbb{R}^2 with noncompact isotropy group, Moore's ergodicity theorem (Moore [28]; see also Feres [7], Zimmer [38] or Margulis [25]) implies that $\mathrm{SL}(2, \mathbb{Z})$ acts ergodically on \mathbb{R}^2 . It follows easily that $\mathrm{GL}(2, \mathbb{Z})$ acts ergodically on $\mathbb{R}^2/\{\pm \mathbb{I}\}$ and $(\mathbb{R}/\mathbb{Z})^2/\{\pm \mathbb{I}\}$, and hence on each of the components

$$C_0, C_1, C_2, C_3, C_K \subset \kappa^{-1}(2) \cap \mathbb{R}^3.$$

Since Σ permutes C_0, C_1, C_2, C_3, C_K , the Γ -action on their union

$$C_0 \cup C_1 \cup C_2 \cup C_3 = \kappa^{-1}(2) \cap \mathbb{R}^3 - C_K \cup S_0 \cong (\mathbb{R}^* \times \mathbb{R}^*) / \{\pm \mathbb{I}\}$$

is ergodic. (In the case of C_K , any hyperbolic element in $\mathrm{GL}(2, \mathbb{Z})$ acts ergodically on $\mathrm{U}(1) \times \mathrm{U}(1)$ and hence on \bar{C}_0 , a much stronger result.)

5. THREE-HOLED SPHERES AND ERGODICITY ($t > 2$)

Next we consider the level sets where $t > 2$. There is an important difference between the cases when $t > 18$ and $2 < t \leq 18$. When $t \leq 18$, the Γ -action is ergodic, but when $t > 18$, there are wandering domains arising from the Fricke spaces of a three-holed sphere. The three-holed sphere is the only other orientable surface homotopy-equivalent to a one-holed torus, and homotopy equivalences to hyperbolic manifolds homeomorphic to a three-holed sphere define points in these level sets when $t > 18$. However, the Γ -action on the complement is ergodic.

5.1. The Fricke space of a three-holed sphere. When $t > 18$, the octant

$$\Omega_0 = (-\infty, -2) \times (-\infty, -2) \times (-\infty, -2)$$

intersects $\kappa^{-1}(t)$ in a wandering domain and the images of $\Omega_0 \cap \kappa^{-1}(t)$ are freely permuted by Γ . Let

$$\Omega = \Gamma \cdot \Omega_0.$$

Characters in Ω correspond to discrete embeddings $\rho : \pi \rightarrow \mathrm{SL}(2, \mathbb{R})$ where the complete hyperbolic surface $\mathbf{H}^2 / \rho(\pi)$ is diffeomorphic to a three-holed sphere. We call such a discrete embedding a *discrete P -embedding*.

The fundamental group $\pi_1(P)$ is free of rank two. A pair of boundary components ∂_1, ∂_2 , an orientation on P , and a choice of arcs α_1, α_2 from the basepoint to ∂_1, ∂_2 determines a pair of free generators of $\pi_1(P)$:

$$X := (\alpha_1)^{-1} \star \partial_1 \star \alpha_1$$

$$Y := (\alpha_2)^{-1} \star \partial_2 \star \alpha_2.$$

A third generator $Z := (XY)^{-1}$ corresponds to the third boundary component, obtaining a presentation of $\pi_1(P)$ as

$$\langle X, Y, Z \mid XYZ = 1 \rangle.$$

Elements of Ω_0 are discrete P -characters such that the generators $X, Y, Z := (XY)^{-1}$ of π correspond to the boundary components of the quotient hyperbolic surface $\mathbf{H}^2 / \rho(\pi)$.

Lemma 5.1. *A representation $\rho \in \text{Hom}(\pi, \text{SL}(2, \mathbb{C}))$ has character $[\rho] = (x, y, z) \in \Sigma \cdot \Omega_0$ if and only if ρ is a discrete P -embedding such that X, Y, Z correspond to the components of $\partial \mathbf{H}^2 / \rho(\pi)$.*

Proof. The condition that $[\rho] \in \Omega_0$ is equivalent to $x, y, z < -2$, which implies that the generators $\rho(X)$, $\rho(Y)$ and $\rho(XY)$ are hyperbolic and their invariant axes bound an open hexagon H with three ideal boundary curves (Goldman [11], Gilman-Maskit [10], I-7, p. 15). This hexagon is the intersection of an ultra-ideal triangle in the Klein model (compare Fig. 13)) with the hyperbolic plane, and can be truncated along the geodesics polar to the ultra-ideal vertices (the dotted lines in Fig. 13) to a right-angled hexagon. The union of H with its reflected image in the invariant axis for $\rho(XY)$ is a fundamental domain for $\rho(\pi)$ acting on \mathbf{H}^2 . (The identifications corresponding to the generators X, Y are depicted in Fig. 13.) The quotient is necessarily homeomorphic to a three-holed sphere P and the holonomy around components of ∂P are the three generators $\rho(X), \rho(Y), \rho(XY)$. \square

Recall (§2.2.2) that Γ decomposes as the semidirect product

$$\Gamma \cong \Gamma_{(2)} \ltimes \mathfrak{S}_3.$$

The mapping class group of a three-holed sphere P is isomorphic to $\mathbb{Z}/2 \times \mathfrak{S}_3$. The \mathfrak{S}_3 -factor corresponds to the group of permutations of $\pi_0(\partial P)$. The $\mathbb{Z}/2$ -factor is generated by the elliptic involution (5.3), which acts by an orientation-reversing homeomorphism of P , whose fixed-point set is the union of three disjoint arcs joining the boundary components. Accordingly, \mathfrak{S}_3 preserves the Teichmüller space of P . The following result shows that these are the only automorphisms of the character variety which preserve the discrete characters representing three-holed spheres.

Proposition 5.2. *Ω equals the disjoint union $\coprod_{\gamma \in \Gamma_{(2)}} \gamma \Omega_0$.*

Proof. We show that if for some $\gamma \in \Gamma_{(2)}$, the intersection $\Omega_0 \cap \gamma \Omega_0$ is nonempty, then $\gamma = 1$. Suppose that $[\rho] \in \Omega_0 \cap \gamma \Omega_0$. By Lemma 5.1, both ρ and $\rho \circ \gamma$ are discrete P -embeddings such that $X, Y, (XY)^{-1}$ and $\gamma(X), \gamma(Y), \gamma((XY)^{-1})$ correspond to $\partial \mathbf{H}^2 / \rho(\pi)$.

The automorphism γ of the character space \mathbb{C}^3 corresponds to an automorphism $\tilde{\gamma}$ of π such that

$$[\rho \circ \tilde{\gamma}] = \sigma_* \circ \gamma([\rho])$$

for a sign-change automorphism $\sigma_* \in \Sigma$. Thus $\rho \circ \tilde{\gamma}$ is also a discrete embedding with quotient a three-holed sphere bounded by curves corresponding to $\tilde{\gamma}(X), \tilde{\gamma}(Y)$ and $\tilde{\gamma}(XY)$. Then $\tilde{\gamma}(X)$ (respectively

$\tilde{\gamma}(Y), \tilde{\gamma}(XY))$ is conjugate to $X^{\pm 1}$ (respectively $Y^{\pm 1}, (XY)^{\pm 1}$). Such an automorphism is induced by a diffeomorphism of the three-holed sphere P whose mapping class group is generated by the elliptic involution. Thus $\tilde{\gamma}$ must be an inner automorphism composed with the elliptic involution of π , and hence must act trivially on characters. Thus $\gamma = 1$ as desired. \square

For $t \leq 18$, the domain Ω does not meet $\kappa^{-1}(t)$. (The closure $\bar{\Omega}$ intersects $\kappa^{-1}(18)$ in the Γ -orbit of the character $(-2, -2, -2)$ of the holonomy representation of a complete finite-area hyperbolic structure on P . For $t > 18$, observe that $\kappa^{-1}(t) - \Omega$ contains the open subset

$$\left((-2, 2) \times \mathbb{R} \times \mathbb{R} \right) \cap \kappa^{-1}(t).$$

Thus $\kappa^{-1}(t) - \Omega$ has positive measure in $\kappa^{-1}(t)$.

Proposition 5.3. *For any $t > 2$, the action of Γ on $\kappa^{-1}(t) - \Omega$ is ergodic.*

The proof uses an iterative procedure (Theorem 5.4) due to Kern-Isberner and Rosenberger [19], although their proof seems to contain a slight gap near the end. A more geometric treatment may be found in Gilman and Maskit [10]. We present a self-contained (algebraic) proof, which we defer to the end of the paper.

5.2. The equivalence relation defined by Γ . Write $u \sim v$ if there exists $\gamma \in \Gamma$ such that $\gamma u = v$. Since κ is Γ -invariant, $u \sim v$ implies that $\kappa(u) = \kappa(v)$.

Theorem 5.4. *Suppose that $u \in \mathbb{R}^3$ satisfies $\kappa(u) > 2$. Then there exists $(\tilde{x}, \tilde{y}, \tilde{z}) = \tilde{u} \sim u$ such that either*

- $\tilde{u} \in (-\infty, -2]^3$, in which case \tilde{u} is the character of a Fuchsian representation whose quotient is a hyperbolic structure on a three-holed sphere P , with boundary mapping to either cusps or closed geodesics;
- $\tilde{x} \in [-2, 2]$, in which case \tilde{u} is the character of a representation mapping X to a non-hyperbolic element.

Recall that a measurable equivalence relation is *ergodic* if the only invariant measurable sets which are unions of equivalence classes are either null or conull. Equivalently an equivalence relation is ergodic if and only if every function constant on equivalence classes is constant almost everywhere. A group action defines an equivalence relation. However, equivalence relations are more flexible since every subset of a space with an equivalence relation inherits an equivalence relation

(whether it is invariant or not). If every point in X is equivalent to a point of S , then ergodicity of X is equivalent to ergodicity of S .

Proof of Proposition 5.3. Assuming Theorem 5.4, we show that Γ acts ergodically on the complement of Ω . Let

$$\mathfrak{E}^{(1)} := \left([-2, 2] \times \mathbb{R} \times \mathbb{R} \right) \cup \left(\mathbb{R} \times [-2, 2] \times \mathbb{R} \right) \cup \left(\mathbb{R} \times \mathbb{R} \times [-2, 2] \right),$$

$$\begin{aligned} \mathfrak{E}^{(2)} := & \left([-2, 2] \times [-2, 2] \times \mathbb{R} \right) \cup \left([-2, 2] \times \mathbb{R} \times [-2, 2] \right), \\ & \cup \left(\mathbb{R} \times [-2, 2] \times [-2, 2] \right) \end{aligned}$$

$$\mathfrak{E}^{(3)} := \left([-2, 2] \times [-2, 2] \times [-2, 2] \right), \text{ and}$$

$$\mathfrak{E}_t^{(i)} := \kappa^{-1}(t) \cap \mathfrak{E}^{(i)} \text{ for } i = 1, 2, 3.$$

Lemma 5.5. (1) If $t > 18$, then $\mathfrak{E}_t^{(i)} = \emptyset$.

(2) If $t > 2$ and $u \in \mathfrak{E}_t - \mathfrak{I}$, then

$$\Gamma u \cap \mathfrak{E}_t^{(1)} \neq \emptyset.$$

If $2 < t < 18$, then $\mathfrak{I} = \emptyset$.

(3) If $2 < t$ and $u \in \mathfrak{E}_t^{(1)}$, then

$$\Gamma u \cap \mathfrak{E}_t^{(2)} \neq \emptyset.$$

(4) If $2 < t$, then the equivalence relation on $\mathfrak{E}_t^{(2)}$ induced by Γ is ergodic.

Ergodicity of the Γ -action on $\kappa^{-1}(t) - \Omega$ follows from ergodicity of the equivalence relation on

$$(\kappa^{-1}(t) - \Omega) \cap \mathfrak{E}$$

induced by the Γ -action on $\kappa^{-1}(t)$. That is, we show that a measurable function

$$f : \mathfrak{E} \cap \kappa^{-1}(t) \longrightarrow \mathbb{R}$$

constant on equivalence classes $\Gamma u \cap \mathfrak{E}$ is constant almost everywhere. By applying permutations, we further reduce to showing the induced equivalence relation on $(\kappa^{-1}(t) - \Omega) \cap \mathfrak{E}^{(1)}$ is ergodic.

Theorem 5.4 implies that every $u \in \kappa^{-1}(t)$ either lies in Ω or is equivalent to a point in \mathfrak{E}_1 . Thus it suffices to show that the equivalence relation on \mathfrak{E}_1 induced by the Γ -action on $\kappa^{-1}(t)$ is ergodic, that is every measurable function

$$f : \mathfrak{E} \cap \kappa^{-1}(t) \longrightarrow \mathbb{R}$$

constant on equivalence classes $\Gamma u \cap \mathfrak{E}$ is constant almost everywhere.

Assuming Lemma 5.5, ergodicity follows as in §5.2 of [14].

Applying a permutation, we may assume that $-2 < x < 2$. For almost every $x_0 \in [-2, 2]$, the action is by a rotation of infinite order of the level set $x^{-1}(x_0) \cap \kappa^{-1}(t)$, which is ergodic. By disintegration of the measure on $\kappa^{-1}(t)$, the invariant function f factors through the coordinate projection $x : \kappa^{-1}(t) \rightarrow \mathbb{R}$.

In a similar way, f factors through the coordinate projections y and z and therefore is constant almost everywhere.

Apply a permutation to assume $-2 < x < 2$. The level set

$$\kappa^{-1}(t) \cap (\{x\} \times \mathbb{R}^2)$$

is an ellipse upon which (for almost every $u \in \kappa^{-1}(t) \cap \mathfrak{E}$) acts by an irrational rotation and is thus ergodic. As in §5.2 of [14], this implies that f factors through the other two coordinates (y, z) ; that is, there is a measurable function g defined on a measurable subset of \mathbb{R}^2 such that

$$f(x, y, z) = g(x)$$

for almost every $(x, y, z) \in \kappa^{-1}(t) \cap \mathfrak{E}$.

To eliminate the dependence on the variables y and z , apply permutation automorphisms to arrange $-2 < y < 2$ (respectively $-2 < z < 2$) and repeat the argument using Dehn twists about Y and XY . \square

5.3. The trace-reduction algorithm. We now prove Theorem 5.4. The proof is based on the following:

Lemma 5.6. *Let $2 < x \leq y \leq z$. Suppose that $\kappa(x, y, z) > 2$. Let $z' = xy - z$. Then $z' < y$ and $z - z' \geq 2\sqrt{\kappa(x, y, z) - 2}$.*

The following expression for $\kappa(x, y, z)$ will be useful:

$$\begin{aligned} (5.1) \quad \kappa(x, y, z) - 2 &= \frac{1}{4} \{ (2z - xy)^2 - (x^2 - 4)(y^2 - 4) \} \\ &= \frac{1}{4} \{ (z - z')^2 - (x^2 - 4)(y^2 - 4) \}. \end{aligned}$$

For $x, y > 2$, write $\kappa_{x,y} : \mathbb{R} \rightarrow \mathbb{R}$ for the quadratic function

$$\kappa_{x,y} : z \mapsto \kappa(x, y, z) - 2 = z^2 - xyz + (x^2 + y^2 - 4).$$

Then

$$\kappa_{x,y}^{-1}((-\infty, 0]) = [\zeta_-(x, y), \zeta_+(x, y)]$$

where

$$\zeta_{\pm}(x, y) = \frac{1}{2} \left(xy \pm \sqrt{(x^2 - 4)(y^2 - 4)} \right).$$

Furthermore reflection

$$z \mapsto z' = xy - z$$

interchanges the two intervals

$$\begin{aligned} J_- &= (-\infty, \zeta_-(x, y)) \\ J_+ &= (\zeta_+(x, y), \infty) \end{aligned}$$

comprising $\kappa_{x,y}^{-1}((0, \infty))$.

Lemma 5.7. *Suppose that $2 < x \leq y$. Then $\zeta_-(x, y) < y < \zeta_+(x, y)$*

Proof. The conclusion is equivalent to $\kappa(x, y, y) < 2$, which is what we prove. First observe that $y > 2$ implies $y - 1/2 > 3/2$ so that

$$\begin{aligned} y - y^2 &= -\left(y - \frac{1}{2}\right)^2 + \frac{1}{4} \\ &< -\left(\frac{3}{2}\right)^2 + \frac{1}{4} = -2 \end{aligned}$$

and $x < y$ implies that $x + 2 - y^2 < y + 2 - y^2 < 0$. Therefore

$$\begin{aligned} \kappa(x, y, y) - 2 &= x^2 + 2y^2 - xy^2 - 4 \\ &= (x^2 - 4) + (2y^2 - xy^2) \\ &= (x - 2)(x + 2 - y^2) < 0. \end{aligned}$$

Since $x > 2$, the proof of Lemma 5.7 is complete. \square

Conclusion of Proof of Lemma 5.6. By Lemma 5.7, the quadratic function $\kappa_{x,y}$ is negative at y . By hypothesis $\kappa_{x,y}$ is positive at $z \geq y$. Therefore

$$z > \zeta_+(x, y) > y.$$

Thus $z \in J_+$. Reflection $z \mapsto z'$ interchanges the intervals J_+ and J_- , so $z' \in J_-$, that is $z' < \zeta_-(x, y) < y$ (Lemma 5.7). By (5.1),

$$\begin{aligned} (z - z')^2 &= 4\left(\kappa(x, y, z) - 2\right) + (x^2 - 4)(y^2 - 4) \\ &> 4(\kappa(x, y, z) - 2) \end{aligned}$$

whence (because $z \geq y > z'$)

$$z - z' > 2\sqrt{\kappa(x, y, z) - 2}$$

completing the proof of Lemma 5.6. \square

Now we conclude the proof of Theorem 5.4. Fix

$$u = (x, y, z) \in \mathbb{R}^3$$

with $\kappa(u) > 2$. We seek $\tilde{u} \sim u$ such that one of the following possibilities occurs:

- one of the coordinates $\tilde{x}, \tilde{y}, \tilde{z}$ lies in the interval $[-2, 2]$;
- $\tilde{x}, \tilde{y}, \tilde{z} < -2$.

By applying linear automorphisms, it suffices to find $\tilde{u} \sim u$ which lies in $(-\infty, 2]^3$. Define a sequence

$$u = u_0 \sim u_1 \sim u_2 \sim \cdots \sim u_n \sim \cdots$$

by the following iterative procedure. Suppose $u_n \in \mathbb{R}^3$ with

$$\kappa(u_n) = k > 2.$$

Then, applying a linear automorphism if necessary, $u_n \sim \tilde{u}_n$ where either:

- $\tilde{u}_n \in (-\infty, 2]^3$; or
- $2 < \tilde{x}_n \leq \tilde{y}_n \leq \tilde{z}_n$.

In the first case, the procedure terminates at \tilde{u}_n . In the second case, reflect \tilde{u}_n in the z -coordinate

$$u_n = \begin{bmatrix} x_n \\ y_n \\ z_n \end{bmatrix} \mapsto u_{n+1} = \begin{bmatrix} x_n \\ y_n \\ z'_n \end{bmatrix}$$

where $z'_n = x_n y_n - z_n$. If $z'_n \leq 2$, the procedure terminates at u_{n+1} . Otherwise, apply a linear automorphism to u_{n+1} to replace u_{n+1} by an equivalent vector \tilde{u}_{n+1} satisfying

$$(5.2) \quad 2 < \tilde{x}_{n+1} \leq \tilde{y}_{n+1} \leq \tilde{z}_{n+1}.$$

Thus either the conclusion of Theorem 5.4 holds or this iterative procedure defines an infinite sequence u_n satisfying (5.2). By (5.2), the largest coordinate equals z_n and by Lemma 5.6, z_1, z_2, \dots is a decreasing sequence. Since $u_{n+1} \sim u_n$,

$$\kappa(u_1) = \cdots = \kappa(u_n) = \kappa(u_{n+1}) = \cdots$$

and we denote this common value by k . Let

$$\mu = 2\sqrt{k-2} > 0.$$

Lemma 5.8. *For each n , $z_{n+3} < z_n - \mu$.*

Proof. By Lemma 5.6, $z'_n < y_n$. Since we assume the sequence does not terminate, $z'_n > 2$. Two possibilities may occur, depending on whether $z'_n > x_n$ or $z'_n \leq x_n$. In either case the largest coordinate of u_{n+1} (assuming the linear normalization (5.2)) is the middle coordinate of u_n , that is

$$z_{n+1} = y_n.$$

Suppose first that $z'_n > x_n$. Since $2 < x_n < z'_n < y_n$, the next term in the sequence is

$$u_{n+1} = \begin{bmatrix} x_{n+1} \\ y_{n+1} \\ z_{n+1} \end{bmatrix} = \begin{bmatrix} x_n \\ z'_n \\ y_n \end{bmatrix}$$

and $y_{n+1} = z'_n$. Hence $z'_n < y_n \leq z_n$. By (5.1),

$$\begin{aligned} z_n - z_{n+2} &= z_n - y_{n+1} \\ &= z_n - z'_n > \mu \end{aligned}$$

by Lemma 5.6. Therefore

$$z_n - z_{n+3} > z_n - z_{n+2} > \mu.$$

Finally suppose that $z'_n < x_n$. Since $2 < z'_n < x_n < y_n$, the next term in the sequence is

$$u_{n+1} = \begin{bmatrix} x_{n+1} \\ y_{n+1} \\ z_{n+1} \end{bmatrix} = \begin{bmatrix} z'_n \\ x_n \\ y_n \end{bmatrix}$$

and $z_{n+1} = y_n$, $y_{n+1} = x_n$ and $x_{n+1} = z'_n$. Applying the iterative procedure to u_{n+1} , we distinguish two cases. The first case occurs when $z'_{n+1} < x_{n+1}$ as above; then

$$z_{n+1} - z_{n+3} > \mu.$$

and

$$z_n - z_{n+3} > z_{n+1} - z_{n+3} > \mu$$

as desired. The second case occurs when

$$u_{n+2} = \begin{bmatrix} x_{n+2} \\ y_{n+2} \\ z_{n+2} \end{bmatrix} = \begin{bmatrix} z'_{n+1} \\ x_{n+1} \\ y_{n+1} \end{bmatrix}.$$

Then $z_{n+3} = y_{n+2} = x_{n+1} = z'_n$ and by Lemma 5.6 implies

$$z_n - z_{n+3} = z_n - z'_n > \mu$$

as desired, completing the proof of Lemma 5.8 in both cases. \square

The proof of Theorem 5.4 easily follows from Lemma 5.8. First,

$$z = z_0 > z_1 > z_2 > \cdots > z_n > \cdots$$

and Lemma 5.8 implies that $z_{3N} < z_0 - N\mu$. Thus if $N > z_0/(3\mu)$, the value of z_{3N} will be negative, and the sequence does actually terminate. This contradiction completes the proof of Theorem 5.4.

APPENDIX: ELEMENTS OF THE MODULAR GROUP

We describe in more detail the automorphisms Γ and interpret them geometrically in terms of mapping classes of M .

Horowitz[18] determined the group $\text{Aut}(\mathbb{C}^3, \kappa)$ of polynomial mappings $\mathbb{C}^3 \rightarrow \mathbb{C}^3$ preserving κ . We have already observed that the *linear automorphisms* form the semidirect product $\Sigma \rtimes \mathfrak{S}_3$ of the group $\Sigma \cong \mathbb{Z}/2 \times \mathbb{Z}/2$ of sign-changes (see §2.2.1) and the symmetric group \mathfrak{S}_3 consisting of permutations of the coordinates x, y, z . Horowitz proved that the automorphism group of (\mathbb{C}^3, κ) is generated by the linear automorphism group

$$\text{Aut}(\mathbb{C}^3, \kappa) \cap \text{GL}(3, \mathbb{C}) = \Sigma \rtimes \mathfrak{S}_3$$

and the *quadratic reflection*:

$$\begin{bmatrix} x \\ y \\ z \end{bmatrix} \longrightarrow \begin{bmatrix} yz - x \\ y \\ z \end{bmatrix},$$

This group is commensurable with $\text{Out}(\pi)$.

We denote this group by Γ ; it is isomorphic to a semidirect product

$$\Gamma \cong \text{PGL}(2, \mathbb{Z}) \rtimes (\mathbb{Z}/2 \oplus \mathbb{Z}/2)$$

where $\text{PGL}(2, \mathbb{Z})$ is the quotient of $\text{GL}(2, \mathbb{Z})$ by the elliptic involution (see below) and $(\mathbb{Z}/2 \oplus \mathbb{Z}/2)$ is the group Σ of sign-changes.

A-4. The elliptic involution. However, $\text{Out}(\pi)$ does not act effectively on \mathbb{C}^3 . To describe elements of $\text{Out}(\pi)$, we use the isomorphism $h : \text{Out}(\pi) \rightarrow \text{GL}(2, \mathbb{Z})$ discussed in §1.2. The kernel of the homomorphism

$$\text{Out}(\pi) \longrightarrow \text{Aut}(\mathbb{C}^3, \kappa)$$

is generated by $h^{-1}(-I)$.

The elliptic involution is a nontrivial mapping class which acts trivially on the character variety. This phenomenon is due to the *hyperellipticity* of the one-holed torus (as in [14], §10.2). The automorphism

ε of π given by:

$$(5.3) \quad \begin{aligned} X &\longmapsto YX^{-1}Y^{-1} \sim X^{-1} \\ Y &\longmapsto (YX)Y^{-1}(X^{-1}Y^{-1}) \sim Y^{-1} \\ XY &\longmapsto X^{-1}Y^{-1} \sim (XY)^{-1} \end{aligned}$$

preserves $K = [X, Y]$ and acts on the homology by the element

$$h(\varepsilon_*) = -I = \begin{bmatrix} -1 & 0 \\ 0 & -1 \end{bmatrix} \in \mathrm{GL}(2, \mathbb{Z})$$

generating the center of $\mathrm{GL}(2, \mathbb{Z})$. Furthermore ε acts identically on the characters (x, y, z) . Thus the homomorphism

$$\mathrm{GL}(2, \mathbb{Z}) \xrightarrow{h^{-1}} \mathrm{Out}(\pi) \longrightarrow \mathrm{Aut}(\mathbb{C}^3, \kappa)$$

factors through $\mathrm{PGL}(2, \mathbb{Z}) := \mathrm{GL}(2, \mathbb{Z})/\{\pm \mathbb{I}\}$.

Note, however, that $\varepsilon^2 = \iota_{K^{-1}}$. The automorphism

$$\begin{aligned} \iota_{YX} \circ \varepsilon : \pi &\longrightarrow \pi \\ X &\longmapsto X^{-1} \\ Y &\longmapsto Y^{-1} \\ XY &\longmapsto X^{-1}Y^{-1} \sim (XY)^{-1} \end{aligned}$$

has order two in $\mathrm{Aut}(\pi)$ but does not preserve K .

A-5. The symmetric group. Next we describe the automorphisms of π which correspond to permutations of the three trace coordinates x, y, z . Permuting the two generators X, Y gives:

$$\begin{aligned} X &\longmapsto Y \\ Y &\longmapsto X \\ XY &\longmapsto YX \sim XY \end{aligned}$$

This automorphism $P_{(12)}$ sends $K \longrightarrow K^{-1}$. It acts on characters by:

$$(P_{(12)})_* : \begin{bmatrix} x \\ y \\ z \end{bmatrix} \longmapsto \begin{bmatrix} y \\ x \\ z \end{bmatrix}$$

and on the homology by:

$$\begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}.$$

Another transposition of the character space is defined by the involution

$$\begin{aligned} P_{(13)} : X &\longmapsto Y^{-1}X^{-1} \\ Y &\longmapsto XYX^{-1} \sim Y \\ XY &\longmapsto X^{-1} \end{aligned}$$

which maps $K \longmapsto K^{-1}$. It acts on characters by:

$$(P_{(13)})_* : \begin{bmatrix} x \\ y \\ z \end{bmatrix} \longmapsto \begin{bmatrix} z \\ y \\ x \end{bmatrix}$$

and on the homology by:

$$\begin{bmatrix} -1 & 0 \\ -1 & 1 \end{bmatrix}.$$

The involution

$$\begin{aligned} P_{(23)} : X &\longmapsto Y^{-1}XY \\ Y &\longmapsto Y^{-1}X^{-1} \\ XY &\longmapsto Y^{-1} \end{aligned}$$

maps $K \longmapsto K^{-1}$ and acts on characters by:

$$(P_{(23)})_* : \begin{bmatrix} x \\ y \\ z \end{bmatrix} \longmapsto \begin{bmatrix} x \\ z \\ y \end{bmatrix}$$

and on the homology by:

$$\begin{bmatrix} 1 & -1 \\ 0 & -1 \end{bmatrix}.$$

The composition $P_{(13)} \circ P_{(12)}$ will be denoted $P_{(123)}$ since the composition of the transposition (12) with the transposition (13) in \mathfrak{S}_3 equals the 3-cycle (123). Applying this composition, we obtain a 3-cycle of automorphisms

$$\begin{aligned} X &\longmapsto XYX^{-1} && \longmapsto XYX^{-1}Y^{-1}(XY)^{-1} \\ Y &\longmapsto Y^{-1}X^{-1} && \longmapsto (XY)X(XY)^{-1} \\ XY &\longmapsto XYX^{-1}Y^{-1}X^{-1} && \longmapsto (XY)(X^{-1}Y^{-1}X)(XY)^{-1} \end{aligned}$$

which preserve K , although $P_{(123)}^3$ equals the elliptic involution ε , not the identity. The action $P_{(123)}$ on characters is:

$$\begin{bmatrix} x \\ y \\ z \end{bmatrix} \xrightarrow{P_{(123)}} \begin{bmatrix} z \\ x \\ y \end{bmatrix} \xrightarrow{P_{(123)}} \begin{bmatrix} y \\ z \\ x \end{bmatrix}$$

which is the *inverse* of the permutation of coordinates given by (123). The action on homology is given by the respective matrices:

$$h(P_{(123)}) = \begin{bmatrix} 0 & -1 \\ 1 & -1 \end{bmatrix}, \quad h(P_{(132)}) = \begin{bmatrix} -1 & 1 \\ -1 & 0 \end{bmatrix}.$$

In summary, we have the following correspondence between permutations of the coordinates and elements of $\mathrm{PGL}(2, \mathbb{Z})$ given by $\tau \mapsto h(P_\tau)$:

$$\begin{aligned} (12) &\mapsto \pm \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \\ (13) &\mapsto \pm \begin{bmatrix} 1 & 0 \\ 1 & -1 \end{bmatrix} \\ (23) &\mapsto \pm \begin{bmatrix} -1 & 1 \\ 0 & 1 \end{bmatrix} \\ (132) &\mapsto \pm \begin{bmatrix} 1 & -1 \\ 1 & 0 \end{bmatrix} \\ (123) &\mapsto \pm \begin{bmatrix} 0 & 1 \\ -1 & 1 \end{bmatrix} \end{aligned}$$

The isomorphism $\mathrm{GL}(2, \mathbb{Z}/2) \cong \mathfrak{S}_3$ relates to the differential (at the origin) of the mappings in the image of $\mathrm{Out}(\pi) \rightarrow \mathrm{Aut}(\mathbb{C}^3, \kappa)$ as follows. The origin is the only isolated point in the $\mathrm{Out}(\pi)$ -invariant set $\kappa^{-1}(-2) \cap \mathbb{R}^3$, and is thus fixed by all of $\mathrm{Out}(\pi)$. (The origin corresponds to the quaternion representation (2.10); see §2.6.) Therefore taking the differential at the origin gives a representation

$$\begin{aligned} \mathrm{Out}(\pi) &\longrightarrow \mathrm{GL}(3, \mathbb{C}) \\ \phi &\longmapsto d_{(0,0,0)}\phi_* \end{aligned}$$

whose image is \mathfrak{S}_3 . In particular it identifies with the composition

$$\mathrm{Out}(\pi) \xrightarrow{h} \mathrm{GL}(2, \mathbb{Z}) \longrightarrow \mathrm{GL}(2, \mathbb{Z}/2) \cong \mathfrak{S}_3$$

where the last arrow denotes reduction modulo 2. These facts can be checked by direct computation.

A-6. A quadratic reflection. Here is a mapping class corresponding to a reflection preserving (x, y) . The automorphism Q_z of π given by:

$$\begin{aligned} X &\longmapsto (XYX^{-1})X(XY^{-1}X^{-1}) \sim X \\ Y &\longmapsto (XYX^{-1})Y^{-1}(XY^{-1}X^{-1}) \sim Y^{-1} \\ XY &\longmapsto (XYX^{-1})XY^{-1}(XY^{-1}X^{-1}) \sim XY^{-1} \end{aligned}$$

maps $K \mapsto K^{-1}$ and

$$(Q_z)_* : \begin{bmatrix} x \\ y \\ z \end{bmatrix} \mapsto \begin{bmatrix} x \\ y \\ xy - z \end{bmatrix}$$

$$h(Q_z) = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}.$$

Similarly the other quadratic reflections are:

$$Q_y : \begin{cases} X & \mapsto (XY)X(Y^{-1}X^{-1}) \sim X \\ Y & \mapsto (XY)X^{-1}Y^{-1}X^{-1}(Y^{-1}X^{-1}) \sim X^{-1}Y^{-1}X^{-1} \\ XY & \mapsto Y^{-1}X^{-1} = (XY)^{-1} \end{cases}$$

which induces

$$(Q_y)_* : \begin{bmatrix} x \\ y \\ z \end{bmatrix} \mapsto \begin{bmatrix} x \\ xz - y \\ z \end{bmatrix}, \quad h(Q_y) = \begin{bmatrix} 1 & -2 \\ 0 & -1 \end{bmatrix}$$

and

$$Q_x : \begin{cases} X & \mapsto (XY)Y^2X(Y^{-1}X^{-1}) \sim Y^2X \\ Y & \mapsto (XYX^{-1})Y^{-1}(XY^{-1}X^{-1}) \sim Y^{-1} \\ XY & \mapsto (XY^2)(XY)(Y^{-2}X^{-1}) \sim XY, \end{cases}$$

which induces

$$(Q_x)_* : \begin{bmatrix} x \\ y \\ z \end{bmatrix} \mapsto \begin{bmatrix} yz - x \\ y \\ z \end{bmatrix}, \quad h(Q_x) = \begin{bmatrix} 1 & 0 \\ 2 & -1 \end{bmatrix}.$$

The square of each of these three reflections is the identity element of $\text{Aut}(\pi)$. These reflections correspond to the generators (2.6) of the level-2 congruence subgroup $\text{GL}(2, \mathbb{Z})_{(2)}$.

A-7. Another involution. The automorphism ν

$$\begin{aligned} X &\mapsto Y^{-1} \\ Y &\mapsto YXY^{-1} \sim X \\ XY &\mapsto XY^{-1} \end{aligned}$$

preserves K , satisfies $\nu^2 = \varepsilon$ and acts by:

$$\nu_* : \begin{bmatrix} x \\ y \\ z \end{bmatrix} \mapsto \begin{bmatrix} y \\ x \\ xy - z \end{bmatrix}, \quad h(\nu_*) = \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix},$$

the composition of the transposition $P_{(12)}$ and the quadratic reflection Q_z . Note that $(P_{(12)})_*$ and $(Q_z)_*$ commute in $\text{Aut}(\mathbb{C}^3, \kappa)$.

A-8. **A Dehn twist.** The automorphism τ

$$\begin{aligned} X &\longmapsto XY \\ Y &\longmapsto Y \\ XY &\longmapsto XY^2 \end{aligned}$$

preserves K , and acts by:

$$\tau_* : \begin{bmatrix} x \\ y \\ z \end{bmatrix} \longmapsto \begin{bmatrix} xy - z \\ y \\ x \end{bmatrix}, \quad h(\tau) = \begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix},$$

the composition $P_{(13)} \circ Q_z$.

Similarly the Dehn twist τ_X around X

$$\begin{aligned} X &\longmapsto X \\ Y &\longmapsto YX \\ XY &\longmapsto XYX \end{aligned}$$

preserves K , and acts by:

$$\begin{bmatrix} x \\ y \\ z \end{bmatrix} \longmapsto \begin{bmatrix} x \\ xy - z \\ y \end{bmatrix}, \quad h(\tau) = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix},$$

the composition $P_{(23)} \circ Q_z$.

REFERENCES



1. Abikoff, W., "The Real Analytic Theory of Teichmüller Space," Lecture Notes in Mathematics **820** (1980) Springer-Verlag, Berlin and New York.
2. Benedetto, R. and Goldman, W., *The Topology of the Relative Character Varieties of a Quadruply-Punctured, Sphere*, Experimental Mathematics (1999) **8:1**, 85–104.
3. Bers, L. and Gardiner, F., *Fricke spaces*, Adv. Math. **62** (1986), 249–284.
4. Brumfiel, G. and Hilden, H., "SL(2)-representations of finitely presented groups," Contemp. Math. **187** Amer. Math. Soc., Providence, RI (1995).
5. Buser, P., "Geometry and Spectra of Compact Riemann Surfaces," Progress in Mathematics **106**, Birkhäuser Boston (1992).
6. Culler, M. and Shalen, P., *Varieties of group representations and splittings of 3-manifolds*, Ann. Math. **117** (1983), 109–146.
7. Feres, R., "Dynamical Systems and Semisimple Groups," Cambridge Tracts in Pure Mathematics **126**, Cambridge University Press (1998).
8. Fricke, R., *Über die Theorie der automorphen Modulgruppen*, Nachr. Akad. Wiss. Göttingen (1896), 91–101.
9. ——— and Klein, F., *Vorlesungen der Automorphen Funktionen*, Teubner, Leipzig, Vol. I (1897), Vol. II (1912).
10. Gilman, J. and Maskit, B., *An algorithm for 2-generator Fuchsian groups*, Michigan Math. J. **38** (1) (1991), 13–3.



11. Goldman, W., *Discontinuous groups and the Euler class*, Doctoral dissertation, University of California, Berkeley (1980).
12. ———, *Geometric structures and varieties of representations*, in “The Geometry of Group Representations,” Contemp. Math. **74**, Amer. Math. Soc. (1988), 169–198.
13. ———, *Topological components of spaces of representations*, Inv. Math. **93** (3) (1988), 557–607.
14. ———, *Ergodic theory on moduli spaces*, Ann. Math. **146** (1997), 1–33.
15. ———, *An exposition of results of Fricke*, (in preparation).
16. ——— and Neumann, W., *Action of the automorphism group of $x^2 + y^2 + x^2 - xyz - 2$ on homology*, (in preparation).
17. Harvey, W. J., *Spaces of Discrete Groups*, in “Discrete Groups and Automorphic Functions,” Academic Press (1977), 295–347.
18. Horowitz, R.D., *Induced automorphisms on Fricke characters of Free groups*, Trans. A.M.S. **208** (1975), 41–50.
19. Kern-Isberner, G. and Rosenberger, G., *Über Diskretheitsbedingungen und die Diophantische Gleichung $ax^2 + by^2 + cz^2 = dxyz$* , Archiv. Math. (Basel) **34** (1980), no. 6, 481–493.
20. Imayoshi, Y. and Taniguchi, M., “An Introduction to Teichmüller spaces,” Springer-Verlag Tokyo Berlin Heidelberg New York (1992).
21. Lubotzky, A. and Magid, A., *Varieties of representations of finitely generated groups*, Mem. A.M.S. **336** (58) (1985).
22. Lyndon, R. and Schupp, R., “Combinatorial Group Theory,” Springer-Verlag (1977), Springer-Verlag Berlin Heidelberg New York.
23. Magnus, W., *Rings of Fricke characters and automorphism groups of free groups*, Math. Zeit. **170** (1980), 91–103.
24. ———, Karrass, A. and Solitar, D., “Combinatorial Group Theory: Presentations of groups in terms of generators and relations,” (1976) Dover Publications, New York.
25. Margulis, G., “Discrete Subgroups of Semisimple Groups,” Ergebnisse der Math. **17**, Springer-Verlag Berlin-Heidelberg-New York (1991).
26. McOwen, R., *Prescribed curvature and singularities of conformal metrics on Riemann surfaces*, J. Math. Anal. Appl. **177** (1993), no. 1, 287–298.
27. Morgan, J. W. and Shalen, P. B., *Valuations, trees and degenerations of hyperbolic structures I*, Ann. Math. **120** (1984), 401–476.
28. Moore, C.C. *Ergodicity of flows on homogeneous spaces*, Amer. J. Math. **88** (1966), 154–178.
29. Nag, S., “The Complex Analytic Theory of Teichmüller Spaces,” Can. Math. Soc. Ser. of Monographs and Advanced Texts, (1988) John Wiley & Sons, New York, Chichester, Brisbane, Toronto, Singapore.
30. Nielsen, J., *Die Isomorphismen der allgemeinen unendlichen Gruppe mit zwei Erzeugenden*, Math. Ann. **71** (1918), 385–397.
31. ———, *Untersuchungen zur Topologie der geschlossenen zweiseitigen Flächen I*, Acta Math. **50** (1927), 189–358.
32. Previte, J. and Xia, E., *Topological dynamics on moduli spaces I*, Pac. J. Math. **193** (2000), no. 2, 397–417.
33. ———, *Topological dynamics on moduli space. II*, Trans. A. M. S. **354** (2002), no. 6, 2475–2494.



- 34. Procesi, C., *Invariant theory of n by n matrices*, Adv. Math. **19** (1976), 306–381.
- 35. Stantchev, G., *Dynamics of the modular group acting on $GL(2, \mathbb{R})$ -characters of a once-punctured torus*, Doctoral Dissertation, University of Maryland (2002).
- 36. Stillwell, J., “The Dehn-Nielsen theorem,” Appendix to *Papers on Group Theory and Topology*, by Max Dehn, Springer Verlag (1987) Berlin Heidelberg New York.
- 37. Troyanov, M., *Prescribing curvature on compact surfaces with conical singularities*, Trans. Amer. Math. Soc. **324** (2) (1991), 793–821.
- 38. Zimmer, R., “Ergodic theory and semisimple groups,” (1984) Monographs in Mathematics **81** Birkhäuser Boston Basel Stuttgart.



MATHEMATICS DEPARTMENT, UNIVERSITY OF MARYLAND, COLLEGE PARK,
MD 20742 USA

E-mail address: `wmg@math.umd.edu`

FIGURE 1. Level set $\kappa = -2.1$ FIGURE 2. Level set $\kappa = 1.9$

FIGURE 3. Level set $\kappa = 1.9$ FIGURE 4. Level set $\kappa = 2.1$

FIGURE 5. Level set $\kappa = 2.1$ FIGURE 6. Level set $\kappa = 4$

FIGURE 7. Level set $\kappa = 4$ FIGURE 8. Level set $\kappa = 19$





FIGURE 9. A convex embedded quadrilateral

FIGURE 10. When $\overline{p_3p_4}$ meets $\overline{p_1p_2}$



FIGURE 11. When $\overline{p_2p_3}$ meets $\overline{p_4p_1}$ 

FIGURE 12. A nonconvex embedded quadrilateral




FIGURE 13. Fundamental hexagons for a hyperbolic three-holed sphere