

THE GYSIN SEQUENCE FOR \mathbb{S}^1 -ACTIONS ON STRATIFIED PSEUDOMANIFOLDS

G. PADILLA

To grandmother Cira and her sisters, in loving memory.

ABSTRACT. For any stratified pseudomanifold X and any action of the unit circle \mathbb{S}^1 on X preserving the stratification and the local structure; the orbit space X/\mathbb{S}^1 is also a stratified pseudomanifold. For each perversity \bar{q} in X the orbit map $\pi : X \rightarrow X/\mathbb{S}^1$ induces a Gysin sequence relating the \bar{q} -intersection cohomologies of X and X/\mathbb{S}^1 . The third term of this sequence can be given by means of a spectral sequence on X/\mathbb{S}^1 whose second term is the cohomology of the fixed points' set $X^{\mathbb{S}^1}$ with values on a constructible sheaf. The above statements generalize a previous work on stratified pseudomanifolds with length 1.

FOREWORD

A stratified pseudomanifold is a topological space X with two features: the stratification and the local conical behavior. The stratification is a decomposition of X in a family of manifolds, called strata, endowed with a partial order of incidence. The union of open strata is a dense smooth manifold called the regular part, its complement Σ is the singular part of X . The local conical behavior is given by the existence of charts, the trivial model being a product $U \times c(L)$ where U is a smooth manifold and $c(L)$ is the cone of a compact stratified pseudomanifold L with lower length, we say that L is a link of U .

When \mathbb{S}^1 acts on X preserving the stratification and the local structure, then the orbit space X/\mathbb{S}^1 is again a stratified pseudomanifold. The orbit map $\pi : X \rightarrow X/\mathbb{S}^1$ preserves the strata and, for each perversity \bar{q} in X , it induces a long exact sequence, the Gysin sequence

$$\cdots \rightarrow H_{\bar{q}}^i(X) \rightarrow H^i(\mathcal{G}_{\bar{q}}(X/\mathbb{S}^1)) \xrightarrow{\partial} H_{\bar{q}}^{i+1}(X/\mathbb{S}^1) \xrightarrow{\pi^*} H_{\bar{q}}^{i+1}(X) \rightarrow \cdots$$

which relates the \bar{q} -intersection cohomologies of X and X/\mathbb{S}^1 . The connecting homomorphism ∂ depends on the Euler class $\varepsilon \in H_{\frac{\bar{q}}{2}}^2(X/\mathbb{S}^1)$; it vanishes if and only if there is a

Date: April 24/2003.

1991 Mathematics Subject Classification. 35S35; 55N33.

Key words and phrases. Intersection Cohomology, Stratified Pseudomanifolds.

foliation on the regular part of X transverse to the orbits of the action [18].

The third complex $\mathcal{G}_{\bar{q}}^*(X/\mathbb{S}^1)$ in the above expression is the Gysin term induced by the action. Its cohomology $H^*(\mathcal{G}_{\bar{q}}(X/\mathbb{S}^1))$ depends on basic cohomological data of two types: local and global. There is a second long exact sequence

$$\cdots \rightarrow H^i(\mathcal{G}_{\bar{q}}(X/\mathbb{S}^1)) \rightarrow H^i(\mathcal{L}\mathbf{ow}_{\bar{q}}(X/\mathbb{S}^1)) \xrightarrow{\partial'} H_{\bar{q}-\bar{e}}^{i+1}(X/\mathbb{S}^1) \xrightarrow{\iota} H^{i+1}(\mathcal{G}_{\bar{q}}(X/\mathbb{S}^1)) \rightarrow \cdots$$

where \bar{e} is a perversity in X/\mathbb{S}^1 vanishing on the mobile strata. The residual term $H^*(\mathcal{L}\mathbf{ow}_{\bar{q}}(X/\mathbb{S}^1))$ is calculated through a spectral sequence whose second term $E_2 = H^j(X^{\mathbb{S}^1}, \mathcal{P}^i)$ is the cohomology of the fixed points' set $X^{\mathbb{S}^1}$ with values on a graduated constructible sheaf \mathcal{P}^* . For each fixed point $x \in X$, the group \mathbb{S}^1 acts on the link L of the stratum containing x and the stalks

$$\mathcal{P}_x^i = \begin{cases} H^i(\mathcal{L}\mathbf{ow}_{\bar{q}}(L/\mathbb{S}^1)) & i \leq \bar{q}(x) - 3 \\ \ker\{\partial' : H^{\bar{q}(x)-1}(\mathcal{L}\mathbf{ow}_{\bar{q}}(L/\mathbb{S}^1)) \rightarrow H_{\bar{q}-\bar{e}}^{\bar{q}(x)}(L/\mathbb{S}^1)\} & i = \bar{q}(x) - 2 \\ \ker\{\partial : H^{\bar{q}(x)-1}(\mathcal{G}_{\bar{q}}(L/\mathbb{S}^1)) \rightarrow H_{\bar{q}}^{\bar{q}(x)+1}(L/\mathbb{S}^1)\} & i = \bar{q}(x) - 1 \\ 0 & i \geq \bar{q}(x) \end{cases}$$

are related to the Gysin sequence and the residual term of L .

Henceforth, when we write the word *manifold* we are talking about a smooth differential manifold of class C^∞ .

1. STRATIFIED SPACES

Recall the definition of a stratified spaces. The reader will find in [16] a detailed exposition.

1.1. Stratified Spaces: Let X be a Hausdorff, paracompact, 2nd countable topological space. A **stratification** of X is a locally finite partition \mathcal{S} whose elements are called **strata**, and satisfy:

- (1) Each stratum with the induced topology is a connected manifold.
- (2) For any two strata $S, S' \in \mathcal{S}$; if $S \cap \overline{S'}$ then $S \subset \overline{S'}$. In that case we say that S is **in the border of S'** , and we write $S \leq S'$.
- (3) There are open strata on X , all of them having the same dimension.

We say that X is a **stratified space** whenever it has some stratification \mathcal{S} . ♣

The open strata are also called **regular**. The **regular part** of X is the union of regular strata, it is a dense open manifold. A **singular** stratum is a non-regular stratum. The **singular part** $\Sigma \subset X$ is the complement of the regular part.

The border relationship (2) is a partial order. A stratum is maximal (resp. minimal) if and only if it is regular (resp. closed). Since the stratification \mathcal{S} is locally finite, strict order chains

$$S_0 < S_1 < \dots < S_l$$

on \mathcal{S} are always finite. The **length** of X is the supreme of the integers l such that there is a strict order chain as above; we write it $\text{len}(X)$.

Examples 1.2. Here there are some examples of stratified spaces.

- (1) For each manifold, the family of its connected components is a stratification.
- (2) For each stratified space X and each manifold M , the canonical stratification of $M \times X$ is

$$\{S \times S' : S \text{ is a connected component of } M \text{ and } S' \text{ is a stratum of } X'\}$$

- (3) For each compact stratified space L , the **cone of** L is the quotient space

$$c(L) = L \times [0, \infty) / L \times \{0\}$$

We write $[p, r]$ for the equivalence class of a point (p, r) , and \star for the equivalence class of $L \times \{0\}$, which we call the **vertex** of the cone. The family

$$\{\star\} \sqcup \{S \times \mathbb{R}^+ : S \text{ is a stratum of } L\}$$

is a stratification of $c(L)$. By convention we state $c(\emptyset) = \{\star\}$. The **radius** of the cone is the function $\rho : c(L) \rightarrow [0, \infty)$ given by $\rho[p, r] = r$. For each $\epsilon > 0$ we write $c_\epsilon(L) = \rho^{-1}[0, \epsilon)$, it is also a stratified space. ♣

Let X be a stratified space. For each paracompact subspace $Y \subset X$ the **induced partition** is

$$\mathcal{S}_{Y/X} = \{C : C \text{ a connected component of } Y \cap S, S \text{ a stratum of } X\}$$

If this family is a stratification of Y , then we say that Y is a **stratified subspace** of X .

A function $\alpha : X \rightarrow X'$ between two stratified spaces is a **morphism** (resp. **isomorphism**) if

- (1) α is a continuous function (resp. homeomorphism).
- (2) α preserves the regular part, i.e. $\alpha(X - \Sigma) \subset (X' - \Sigma')$.
- (3) α sends smoothly (resp. diffeomorphically) strata into strata.

In particular, α is an **embedding** if $\alpha(X) \subset Y$ is open and $\alpha : X \rightarrow \alpha(X)$ is an isomorphism. For instance, the inclusion $c_\epsilon(L) \xrightarrow{\iota} c(L)$ is an embedding, and the change of radius

$$f : c(L) \rightarrow c_\epsilon(L) \quad [p, r] \mapsto [p, \epsilon \cdot \arctan(r)/\pi]$$

is an isomorphism.

1.3. Stratified actions Take a stratified space X , a compact abelian Lie group G and a continuous effective action

$$\Phi : G \times X \rightarrow X$$

We will write $\Phi(g, x) = gx$, $B = X/G$ and $\pi : X \rightarrow B$ for the orbit map. We will say that Φ is **stratified** whenever

(1) The action $\Phi : G \times X \rightarrow X$ is a morphism, its restriction to $X - \Sigma$ is free.

(2) For each singular stratum S in X , the points of S have all the same isotropy G_S .

By convention, a stratum S in X is **mobile** (resp. **fixed**) if $G_S \neq \mathbb{S}^1$ (resp. $G_S = \mathbb{S}^1$).

Lemma 1.4. *The orbit space of a stratified action is a stratified space.*

Proof. By §1.3-(1), for each $g \in G$ the function $\Phi_g : X \rightarrow X$ given by $\Phi_g(x) = gx$ is an isomorphism. By §1.3-(2), for each stratum S of X the restriction $\pi : GS \rightarrow \pi(S)$ is a smooth locally trivial fiber bundle with fiber G/G_S . Since G is compact; B is a Hausdorff, paracompact, 2nd countable space [2]. The family

$$(1) \quad \mathcal{S}_B = \{\pi(S) : S \text{ is a stratum in } X\}$$

is a locally finite partition of B ; the border relationship passes from X to B through the orbit map. Hence \mathcal{S}_B is a stratification of B . \square

The family \mathcal{S}_B in equation (1) is the **stratification of B induced by the action Φ** . The orbit map is a morphism by construction.

2. STRATIFIED PSEUDOMANIFOLDS AND UNFOLDINGS

Stratified pseudomanifolds were introduced by Goresky and MacPherson in order to extend the Poincaré duality to the family of stratified spaces. The reader will find in [1] and [6] a detailed exposition of the subject.

2.1. Stratified Pseudomanifolds Let X be a stratified space, S a stratum of X . A **chart** of S in X is an embedding

$$\alpha : U \times c(L) \rightarrow X$$

where $c(L)$ is the cone of a compact stratified space, $U \subset S$ is open in S and $\alpha(u, \star) = u$ for each $u \in U$. Notice that $\text{len}(L) < \text{len}(X)$.

The definition of stratified pseudomanifolds is made by induction on the length: we say that X is a **stratified pseudomanifold** if for each stratum S there is a family of charts,

$$\mathcal{A}_S = \{\alpha : U_\alpha \times c(L) \rightarrow X\}_\alpha$$

such that $\{U_\alpha\}_\alpha$ is an open cover of S , and L is a compact stratified pseudomanifold which only depends on S , we call it a **link** of S . \clubsuit

For any stratified pseudomanifold the link of the regular strata is the empty set. Any open subset of a stratified pseudomanifold is a stratified pseudomanifold.

Examples 2.2. Here there are some examples of stratified pseudomanifolds.

- (1) Every manifold with the canonical stratification is a stratified pseudomanifold.
- (2) For each stratified pseudomanifold X and each manifold M , the product $M \times X$ is a stratified pseudomanifold.
- (3) If L is a compact stratified pseudomanifold then $c(L)$ is a stratified pseudomanifold.

Recall the definition of an unfolding, we use it in order to define the intersection cohomology of a stratified pseudomanifold by means of differential forms. For an introduction to the unfoldings and their properties, see [21].

2.3. Unfoldings Let X be a stratified pseudomanifold. An **unfolding** of X is a manifold \tilde{X} ; a surjective, proper, continuous function

$$\mathbf{L} : \tilde{X} \rightarrow X$$

and a family of unfoldings $\{\mathbf{L}_L : \tilde{L} \rightarrow L\}_L$ of the links of X ; satisfying:

- (1) The restriction $\mathbf{L} : \mathbf{L}^{-1}(X - \Sigma) \rightarrow X - \Sigma$ is a smooth trivial finite covering.
- (2) Each $z \in \mathbf{L}^{-1}(\Sigma)$ has an **unfoldable chart**; i.e., a commutative diagram

$$\begin{array}{ccc} U \times \tilde{L} \times \mathbb{R} & \xrightarrow{\tilde{\alpha}} & \tilde{X} \\ \downarrow c & & \downarrow \mathbf{L} \\ U \times c(L) & \xrightarrow{\alpha} & X \end{array}$$

where

- (a) α is a chart.
- (b) $\tilde{\alpha}$ is a diffeomorphism onto $\mathbf{L}^{-1}(\text{Im}(\alpha))$.
- (c) Left vertical arrow is $c(u, \tilde{p}, t) = (u, [\mathbf{L}_L(\tilde{p}), |t|])$ for each $u \in U$, $\tilde{p} \in \tilde{L}$, $t \in \mathbb{R}$.

We say that X is **unfoldable** when it has an unfolding.

Let $\mathbf{L} : \tilde{X} \rightarrow X$, $\mathbf{L}' : \tilde{X}' \rightarrow X'$ be two unfoldings. A morphism $\alpha : X \rightarrow X'$ is said to be **unfoldable** if there is some smooth function $\tilde{\alpha} : \tilde{X} \rightarrow \tilde{X}'$ such that the following square

$$\begin{array}{ccc} \tilde{X} & \xrightarrow{\tilde{\alpha}} & \tilde{X}' \\ \downarrow \mathbf{L} & & \downarrow \mathbf{L}' \\ X & \xrightarrow{\alpha} & X' \end{array}$$

is commutative. ♣

2.4. Examples These are some examples of unfoldings.

- (1) For each manifold M the identity $\iota : M \rightarrow M$ is an unfolding.
- (2) If $\mathbf{L} : \tilde{X} \rightarrow X$ is an unfolding then the product $\iota \times \mathbf{L} : M \times \tilde{X} \rightarrow M \times X$ is also an unfolding for any manifold M .
- (3) If $\mathbf{L}_L : \tilde{L} \rightarrow L$ is an unfolding on a compact stratified pseudomanifold L ; then the

arrow $c : \tilde{L} \times \mathbb{R} \rightarrow c(L)$ given by the rule $c(\tilde{p}, t) = [L_L(\tilde{p}), |t|]$ is an unfolding (cf. the left vertical arrow in diagram §2.3-(2)).

Lemma 2.5. *Let $L : \tilde{X} \rightarrow X$ be an unfolding. Then*

- (1) *The restriction $L : L^{-1}(A) \rightarrow A$ is an unfolding for each open subset $A \subset X$.*
- (2) *The restriction $L : L^{-1}(S) \rightarrow S$ is a smooth locally trivial fiber bundle with fiber \tilde{L} , for each singular stratum S with link L .*

3. INTERSECTION COHOMOLOGY

Between the various ways for defining the intersection cohomology; the reader can see [6] for a definition in pl -stratified pseudomanifolds; [7], [11], for a definition with sheaves; [13] for an approach with L^2 -cohomology; [4] for an exposition in Thom-Mather spaces. In this article, we use the DeRham-like definition of intersection cohomology as it is exposed in [21].

3.1. Lifiable Forms Let's fix an unfolding $L : \tilde{X} \rightarrow X$. A form $\omega \in \Omega^*(X - \Sigma)$ is **lifiable** if there is a form $\tilde{\omega} \in \Omega^*(\tilde{X})$ such that $L^*(\omega) = \tilde{\omega}$ on $L^{-1}(X - \Sigma)$. If such an $\tilde{\omega}$ exists then it is unique by density; we call it the **lifting** of ω . If ω, η are liftable forms then $d\omega$ is also liftable and we get the following equalities: $\widetilde{d\omega} = d\tilde{\omega}$, $\widetilde{\omega + \eta} = \tilde{\omega} + \tilde{\eta}$, $\widetilde{\omega \wedge \eta} = \tilde{\omega} \wedge \tilde{\eta}$. The liftable forms constitute a differential graded commutative algebra.

3.2. Intersection Cohomology Let $p : M \rightarrow B$ be a surjective submersion. A smooth vector field ξ in M is **vertical** if it is tangent to the fibers of p . Write i_ξ for the contraction by ξ . The **perverse degree** $\|\omega\|_B$ of a differential form $\omega \in \Omega(M)$ on B is the first integer m such that, for each vertical vector fields ξ_0, \dots, ξ_m ;

$$i_{\xi_0} \cdots i_{\xi_m}(\omega) = 0$$

Since contractions are antiderivatives of degree -1 , for each $\omega, \nu \in \Omega(M)$

$$(2) \quad \|\omega + \nu\|_B \leq \max \{ \|\omega\|_B, \|\nu\|_B \} \quad \|\omega \wedge \nu\|_B \leq \|\omega\|_B + \|\nu\|_B$$

By convention $\|0\|_B = -\infty$.

We define the DeRham-like intersection cohomology of X by means of liftable differential forms and an additional parameter which controls their behavior when approaching to Σ . This new parameter is a map \bar{q} which sends each singular stratum S in X to an integer $\bar{q}(S) \in \mathbb{Z}$; we call it a **perversity** in X .

For instance, for each integer $n \in \mathbb{Z}$ we denote by \bar{n} the constant perversity assigning n to any singular stratum. The top perversity in X is defined by $\bar{t}(S) = \text{codim}(S) - 2$ for each singular stratum S .

Fix a perversity \bar{q} . A \bar{q} -**form** on X is a liftable form ω on $X - \Sigma$ satisfying

$$\max\{\|\omega\|_S, \|d\omega\|_S\} \leq \bar{q}(S) \quad \forall S \text{ singular stratum}$$

where, with a little abuse of language, we denote by $\|\omega\|_S$ the perverse degree of the restriction $\tilde{\omega}|_{\mathbf{L}^{-1}(S)}$ with respect to the submersion $\mathbf{L} : \mathbf{L}^{-1}(S) \rightarrow S$. The \bar{q} -forms define a differential subcomplex $\Omega_{\bar{q}}^*(X)$ whose cohomology $H_{\bar{q}}^*(X)$ is the \bar{q} -**intersection cohomology** of X .

3.3. Some properties of intersection cohomology

- (a) $H_{\bar{q}}^*(X)$ does not depend on the particular choice of an unfolding, for any perversity \bar{q} .
- (b) If $\bar{q} > \bar{t}$ then $H_{\bar{q}}^*(X) = H^*(X - \Sigma)$ is the DeRham cohomology of $X - \Sigma$.
- (c) If $\bar{q} < \bar{0}$ then $H_{\bar{q}}^*(X) = H^*(X, \Sigma)$ is the relative cohomology of the pair.
- (d) If X is a manifold and $\bar{0} \leq \bar{q} \leq \bar{t}$ then $H_{\bar{q}}^*(X)$ coincides with the DeRham cohomology $H^*(X)$.
- (e) For any two perversities \bar{p}, \bar{q} ; the wedge product of the forms takes into account the perversities in the following way:

$$H_{\bar{p}}^i(X) \times H_{\bar{q}}^j(X) \xrightarrow{\wedge} H_{\bar{p}+\bar{q}}^{i+j}(X)$$

In particular, the $\bar{0}$ -intersection cohomology $H_{\bar{0}}^*(X)$ is a differential graded algebra and $H_{\bar{q}}^*(X)$ is an $H_{\bar{0}}(X)$ -module for any perversity \bar{q} . A **controlled** form is a $\bar{0}$ -form.

4. MODELLED ACTIONS

Now we introduce the family of action which we will use all along this work, we call them modelled actions.

4.1. Modelled Actions: Let $\Phi : G \times X \rightarrow X$ be a stratified action. We say that Φ is **modelled** if it satisfies conditions **MAI** and **MAII** stated below.

MAI For each singular stratum S there is a modelled action $\Psi : G_S \times L \rightarrow L$ of the isotropy subgroup on the link of S .

A **modelled unfolding** of X is an unfolding $\mathbf{L} : \tilde{X} \rightarrow X$ in the usual sense, together with a free smooth action $\tilde{\Phi} : G \times \tilde{X} \rightarrow \tilde{X}$ such that \mathbf{L} is equivariant and:

- (1) For each link L of X the induced unfolding $\mathbf{L}_L : \tilde{L} \rightarrow L$ is modelled.
- (2) For each singular stratum S and each $z \in \mathbf{L}^{-1}(S)$ there is a **modelled chart**; i.e., an unfoldable chart as in §2.3-(2), such that

- (a) The diagram §2.3-(2) is G_S -equivariant. Here the action of G_S on $U \times c(L)$ is given by the rule $g(u, [p, r]) = (u, [gp, r])$. The free action of G_S on $U \times \tilde{L} \times \mathbb{R}$ is defined as well.
- (b) The transformations of G are cone-preserving: For each $u \in U$, $g \in G$; if $\Phi_g(\alpha(\{u\} \times c(L))) \cap \text{Im}(\alpha) \neq \emptyset$ then the arrow

$$\alpha^{-1}\Phi_g\alpha|_u: \{u\} \times c(L) \rightarrow \{gu\} \times c(L)$$

is an (unfoldable) isomorphism and preserves the radius $\rho: U \times c(L) \rightarrow [0, \infty)$.

MAII X has a modelled unfolding. ♣

Remark 4.2. If $\text{len}(X) = 0$ then $\Sigma = \emptyset$ and the conditions **MAI**, **MAII** are trivial. If $\text{len}(X) = 1$ then **MAI** is trivial again; condition **MAII** can be simplified taking into account the existence of an equivariant normalization of the action; see [14] and [15].

Remark 4.3. Modelled actions constitute a category of actions. There are modelled morphisms, and they preserve the Euler class of the orbit space. We will have to wait until §5.8 before we can describe them in a precise way.

Examples 4.4. Here there are some examples of modelled actions:

- (1) If $\Psi: G \times L \rightarrow L$ is a modelled action with a modelled unfolding $L_L: \tilde{L} \rightarrow L$; then for any manifold U the induced action

$$\Phi: G \times U \times c(L) \rightarrow U \times c(L) \quad g(u, [p, r]) = (u, [gp, r])$$

is modelled. The canonical unfolding $c: U \times \tilde{L} \times \mathbb{R} \rightarrow U \times c(L)$ given in diagram §2.3-(2) is a modelled unfolding.

- (2) Let X be a Thom-Mather space. Any stratified action $\Phi: G \times X \rightarrow X$ preserving the tubular neighborhoods is a modelled action (see [23]).
- (3) If X is a manifold and Φ is a smooth effective action; then X can be endowed with the decomposition in orbit types. This decomposition is a stratification and X inherits an equivariant Thom-Mather structure. By example (1) Φ is a modelled action.

The main feature of modelled actions is that they preserve the category of unfoldable pseudomanifolds.

Proposition 4.5. *For each modelled action $\Phi: G \times X \rightarrow X$*

- (1) *The orbit space $B = X/G$ is a pseudomanifold.*
- (2) *The induced map $L_B: \tilde{B} = \tilde{X}/G \rightarrow B$ given by the rule $L_B(\tilde{\pi}(x)) = \pi(L(x))$ is an unfolding.*
- (3) *The orbit map $\pi: X \rightarrow B$ is an unfoldable morphism.*

Proof. Notice that, by §1.4, B is already a stratified space and $\pi: X \rightarrow B$ is a morphism.

(1) We verify the existence of charts in B . Proceed by induction on $l = \text{len}(X)$, for $l = 0$ it's trivial. Take some singular stratum S with link L . Applying induction, by **MAI** the quotient L/G_S is a stratified pseudomanifold. Fix a modelled chart on S

$$\begin{array}{ccc} U \times \tilde{L} \times \mathbb{R} & \xrightarrow{\tilde{\alpha}} & \tilde{X} \\ \downarrow^c & & \downarrow^{\tilde{L}} \\ U \times c(L) & \xrightarrow{\alpha} & X \end{array}$$

whose existence is given by **MAII**. Assume that $U = WV$ where $W \subset G$ is a contractible open neighborhood of $1 \in G$, V a slice in S . Write $\pi_L : L \rightarrow L/G_S$ for the orbit map. Since α is G_S -equivariant, the function

$$\beta : V \times c(L/G_S) \rightarrow B \quad \beta(y, [\pi_L(p), r]) = \pi\alpha(y, [p, r])$$

is well defined, we will show that it is an embedding:

- **β is injective:** Because V is a slice in S and by condition §4.1-(2.b), the transformations of G are cone-preserving.
- **β is continuous:** Because $B, L/G_S$ have the respective quotient topologies.
- **β is open:** Let $A \subset V \times c(L/G_S)$ be an open subset, $z \in A$. Take a compact neighborhood $z \in K \subset A$. Since $\beta : K \rightarrow \beta(K)$ is a continuous bijection from a compact space onto a Hausdorff space, it is a homeomorphism. There is an open $V' \subset V$ and $\epsilon > 0$ such that

$$z \in A' = V' \times c_\epsilon(L/G_S) \subset K \subset A$$

So $\beta : A' \rightarrow \beta(A')$ is a homeomorphism. We claim that

$$\beta(A') = \pi(\alpha(WV' \times c_\epsilon(L)))$$

the second set is open because $WV' \subset U$ is open in S and the orbit map π is open. In order to show the above equality take a point $\pi(\alpha(wv, [p, r])) \in \pi(\alpha(WV' \times c_\epsilon(L)))$. Then,

$$w^{-1}\alpha(wv, \star) = w^{-1}(wv) = v \in \text{Im}(\alpha)$$

By condition §4.1-(2.b), $\alpha^{-1}\Phi_w^{-1}\alpha : wv \times c(L) \rightarrow v \times c(L)$ is an isomorphism and preserves the radius. So

$$\pi(\alpha(wv, [p, r])) = \pi(\alpha(v, [p', r])) = \beta(v, [\pi_L(p'), r])$$

This proves that $\beta(A') \supset \pi(\alpha(WV' \times c_\epsilon(L)))$. The other inclusion is straightforward.

- **β is an embedding:** On $V \times \{\star\}$ the restriction $\beta : V \times \{\star\} \rightarrow \pi(V)$ given by $\beta(y, \star) \rightarrow \pi(y)$ is a diffeomorphism. For each stratum $R \subset L$ there is a stratum $S' \subset X$ such that $\alpha(V \times R \times \mathbb{R}^+) \subset S'$. Since α is G_S -equivariant, we get the following commutative diagram

$$\begin{array}{ccc} V \times G_S R \times \mathbb{R}^+ & \xrightarrow{\alpha} & GS' \\ \downarrow^{1 \times \pi_L} & & \downarrow^{\pi} \\ V \times \pi_L(R) \times \mathbb{R}^+ & \xrightarrow{\beta} & \pi(S') \end{array}$$

where the vertical arrows are submersions. So β is smooth on $V \times \pi_L(R) \times \mathbb{R}^+$ because α is smooth on $V \times G_S R \times \mathbb{R}^+$. The same argument can be applied to the inverse β^{-1} . Up to a change of variable, we can assume that $\beta(v, \star) = v \forall v$.

(2) By **MAII**, the function L_B is well defined because L is equivariant. If $\Sigma = \emptyset$ the proof is immediate, because \tilde{X} is a smooth equivariant finite trivial covering of X . In general, if $\Sigma \neq \emptyset$ then by the above remark L_B satisfies §2.3-(1).

Now we will prove that the charts given in the first step of this proof are unfoldable; this will show §2.3-(2). We apply induction on $l = \text{len}(X)$. For $l = 0$ there is nothing to do. Take a singular stratum S with link L . By induction, the G_S -equivariant unfolding $L_L : \tilde{L} \rightarrow L$ induces an unfolding $L_{L/G_S} : \tilde{L}/G_S \rightarrow L/G_S$. For each modelled chart α as in the first step of this proof, the lifted $\tilde{\alpha}$ satisfies a smooth-like property analogous to §4.1-(2.b). Hence, the arrow

$$\tilde{\beta} : V \times \tilde{L}/G_S \times \mathbb{R} \rightarrow \tilde{\pi}(\text{Im}(\tilde{\alpha})) \quad \tilde{\beta}(y, \tilde{\pi}_L(\tilde{p}), t) = \tilde{\pi}\tilde{\alpha}(y, \tilde{p}, t)$$

is well defined, injective and a smooth embedding onto an open subset of \tilde{B} . Notice that $\tilde{\beta}$ is the lifting corresponding to the map β given above. In consequence

$$\begin{array}{ccc} V \times \tilde{L}/G_S \times \mathbb{R} & \xrightarrow{\tilde{\beta}} & \tilde{B} \\ \downarrow^c & & \downarrow L_B \\ V \times c(L/G_S) & \xrightarrow{\beta} & B \end{array}$$

is an unfoldable chart in B . The details are left to the reader.

(3) It is immediate from the first two statements. □

5. INVARIANT FORMS

Some results of this § were taken of [9], [12]; these references deal with smooth non-free circle actions on manifolds, but the same proofs still hold in our context. The usual case of a smooth free circle action can be seen for instance in [8].

From now on, we fix a pseudomanifold X , a perversity \bar{q} in X , a modelled action $\Phi : \mathbb{S}^1 \times X \rightarrow X$ and a modelled unfolding $L : \tilde{X} \rightarrow X$.

5.1. Invariant forms A \bar{q} -form ω on X is **invariant** if for each $g \in \mathbb{S}^1$ the equation $g^*(\omega) = \omega$ holds. Since

$$\begin{array}{ccc} \tilde{X} & \xrightarrow{\Phi_g} & \tilde{X} \\ \downarrow L & & \downarrow L \\ X & \xrightarrow{\Phi_g} & X \end{array}$$

is an unfoldable isomorphism, $g^* : \Omega_{\bar{q}}^*(X) \rightarrow \Omega_{\underline{q}}^*(X)$ is an isomorphism of differential complexes. Invariant \bar{q} -forms define a differential complex, denoted $\mathbb{I}\Omega_{\bar{q}}^*(X)$. The inclusion

$$\iota : \mathbb{I}\Omega_{\bar{q}}^*(X) \rightarrow \Omega_{\bar{q}}^*(X)$$

induces an isomorphism in cohomology.

Next we will study the algebraic decomposition of an invariant \bar{q} -form. This decomposition depends on the existence of an unfoldable invariant riemannian metric and an unitary smooth vector field tangent to the orbits in the regular part.

5.2. Definitions The **Fundamental vector field** on X is the smooth vector field \mathcal{C} defined on $X - \Sigma$ by the rule

$$\mathcal{C}_x = d\Phi_x\left(\frac{\partial}{\partial g}\right) \Big|_{g=1}$$

The fundamental vector field \mathcal{C} never vanishes because $X - \Sigma$ has no fixed points. The lifted action $\tilde{\Phi} : \mathbb{S}^1 \times \tilde{X} \rightarrow \tilde{X}$ defines a fundamental vector field $\tilde{\mathcal{C}}$ on \tilde{X} . Notice that $\mathbb{L}_*(\tilde{\mathcal{C}}) = \mathcal{C}$ on $\mathbb{L}^{-1}(X - \Sigma)$.

An **unfoldable metric** on X is an invariant riemannian metric μ on $X - \Sigma$ such that there is an invariant riemannian metric $\tilde{\mu}$ on \tilde{X} satisfying:

- (1) $\mathbb{L}^*(\mu) = \tilde{\mu}$ in $\mathbb{L}^{-1}(X - \Sigma)$.
- (2) $\mu\langle \mathcal{C}, \mathcal{C} \rangle = \tilde{\mu}\langle \tilde{\mathcal{C}}, \tilde{\mathcal{C}} \rangle = 1$.
- (3) For each mobile stratum S and each vertical vector field ν with respect to the submersion $\mathbb{L}^{-1}(S) \xrightarrow{\mathbb{L}} S$; we have $\tilde{\mu}\langle \tilde{\mathcal{C}}, \nu \rangle = 0$.

For each modelled action in X there is an unfoldable metric; the reader will find a proof in [9].

Given an unfoldable metric μ on X ; the **characteristic form** induced by μ is the invariant 1-form χ defined by the rule $\chi(v) = \mu\langle \mathcal{C}, v \rangle$.

Lemma 5.3. *The characteristic form χ satisfies*

$$\|\chi\|_S = \begin{cases} 1 & S \text{ a fixed stratum} \\ 0 & S \text{ a mobile stratum} \end{cases}$$

Proof. By §5.2-(1), the characteristic form χ on $X - \Sigma$ lifts to the characteristic form $\tilde{\chi}$ on \tilde{X} . The perverse degree of χ is immediate from §5.2-(2) and (3). □

Each unfoldable metric μ in X induces an algebraic decomposition of the invariant forms. This decomposition is important in order to give a suitable presentation of the elements composing the Gysin Sequence of X .

5.4. Decomposition of an Invariant Form A form η on $X - \Sigma$ is **basic** if one of the following equivalent statements holds:

- (a) η is invariant and $\iota_{\mathcal{C}}(\eta) = 0$.
- (b) There is a unique differential form θ on $B - \Sigma = \pi(X - \Sigma)$ such that $\eta = \pi^*(\theta)$.

Notice also that if η is basic then

$$0 = L_{\mathcal{C}}(\eta) = d\iota_{\mathcal{C}}(\eta) + \iota_{\mathcal{C}}d(\eta) = \iota_{\mathcal{C}}d(\eta)$$

where $L_{\mathcal{C}}$ is the Lie derivative with respect to the fundamental vector field.

For each invariant form $\omega \in \mathrm{I}\Omega_{\bar{q}}^*(X - \Sigma)$ there are $\nu \in \Omega^*(B - \Sigma)$ and $\theta \in \Omega^{*-1}(B - \Sigma)$ satisfying

$$\omega = \pi^*(\nu) + \chi \wedge \pi^*(\theta)$$

The above expression is the **decomposition** of ω . The forms ν, θ are uniquely determined by the following equations

$$\pi^*(\theta) = \iota_{\mathcal{C}}(\omega) \quad \pi^*(\nu) = \omega - \chi \wedge \iota_{\mathcal{C}}(\omega)$$

When ω is a liftable form then

$$\tilde{\omega} = \tilde{\pi}^*(\tilde{\nu}) + \tilde{\chi} \wedge \tilde{\pi}^*(\tilde{\theta})$$

So ν, θ lift respectively to $\tilde{\nu}, \tilde{\theta}$. ♣

The first object involved in the Gysin sequence of X is the orbit map $\pi : X \rightarrow B$. For each perversity \bar{q} in X , π induces a well defined morphism in intersection cohomology

$$(3) \quad \pi^* : H_{\bar{q}}^*(B) \rightarrow H_{\bar{q}}^*(X)$$

As we shall see later, the Gysin sequence is a long exact sequence containing this map. Now we prove that this arrow makes sense; it is enough to study the behavior of π with respect to the perverse degree of an invariant \bar{q} -form and its algebraic decomposition.

Lemma 5.5. *Take a perversity \bar{q} on X , write also \bar{q} for the perversity induced on B in the obvious way. Then the arrow*

$$\pi^* : \Omega_{\bar{q}}^*(B) \rightarrow \mathrm{I}\Omega_{\bar{q}}^*(X)$$

is well defined. What's more, for each invariant form $\omega = \pi^(\nu) + \chi \wedge \pi^*(\theta)$ and each singular stratum S , we have*

$$\|\omega\|_S = \max\{\|\nu\|_{\pi(S)}, \|\chi\|_S + \|\theta\|_{\pi(S)}\}$$

Proof. See [9]. □

Since π^* commutes with the differential d ; the arrow (3) is well defined.

The second object that appears in the Gysin sequence of X is an intersection cohomology class in B uniquely determined by the action Φ ; we call it the Euler class. Up to this point, the situation is the analogous of the smooth case.

5.6. The Euler class of a modelled action Take an unfoldable metric μ on X , χ the characteristic form induced by μ . The differential form $d\chi$ is basic, so there is a unique form e on $B - \Sigma$ such that

$$d\chi = \pi^*(e)$$

This e is the **Euler form** induced by the action Φ and the metric μ . Since μ is unfoldable, e lifts to the Euler form \tilde{e} on \tilde{B} induced by the metric $\tilde{\mu}$.

The **Euler class** is the intersection cohomology class $\varepsilon = [e] \in H_{\bar{e}}^2(B)$ of the Euler form e with respect to a perversity \bar{e} in B called the **Euler perversity**. This \bar{e} is defined by induction on the length. For each singular stratum S in X

- (1) If S is mobile then $\bar{e}(\pi(S)) = 0$.
- (2) If S is fixed with link L and the Euler class $\varepsilon_L \in H_{\bar{e}_L}^2(L/S^1)$ vanishes, then $\bar{e}(\pi(S)) = 1$.
- (3) If S is fixed with link L and $\varepsilon_L \in H_{\bar{e}_L}^2(L/S^1) \neq 0$, then $\bar{e}(\pi(S)) = 2$. In this case we will say that S is a **perverse** stratum.

These properties determine the Euler perversity \bar{e} in a unique way. Following [15], the Euler class vanishes if and only if there is a foliation \mathcal{F} on $X - \Sigma$ transverse to the orbits of the action. ♣

Next we show that the Euler class is well defined.

Proposition 5.7. *There is an invariant unfoldable metric μ such that the Euler form e induced by μ belongs to $\Omega_{\bar{e}}^2(B)$.*

Proof. By §5.2-(3); for any unfoldable metric μ on X and any mobile stratum S we have $\|e\|_{\pi(S)} = 0$. So we restrict our attention to the fixed strata. We must give an unfoldable metric μ such that, for each fixed stratum S in X ,

$$(4) \quad \|e\|_{\pi(S)} = 2 \Leftrightarrow S \text{ is a perverse stratum}$$

Any unfoldable metric μ satisfying this property will be called a *good metric*. We show the existence of goods metrics by induction on $l = \text{len}(X)$; for $l = 0$ it is immediate. By inductive hypothesis, for each fixed stratum S in X , the link L has a good metric μ_L .

• **Construction of a global good metric μ from a family of local ones:**
 We give an invariant open cover $\mathcal{U} = \{X_\alpha\}_\alpha$ of X , and a family $\{\mu_\alpha\}_\alpha$ of unfoldable metrics such that each μ_α is a good metric in X_α .

- (a) The complement of the fixed points' set $X_0 = X - X^{S^1}$ belongs to \mathcal{U} . We take on X_0 an unfoldable metric μ_0 .

(b) For each fixed stratum S we take a family of modelled charts

$$\alpha : U_\alpha \times c(L) \rightarrow X$$

as in §4.1-(2); such that $\{U_\alpha\}_\alpha$ is a good cover of S . We put $X_\alpha = \text{Im}(\alpha)$ and take

$$\mu_\alpha = \alpha^{-*}(\mu_{U_\alpha} + \mu_L + dr^2)$$

where μ_{U_α} (resp. μ_L) is a riemannian (resp. good) metric in U_α (resp. in L). So μ_α is a good metric in X_α .

Fix an invariant controlled partition of the unity $\{\rho_\alpha\}_\alpha$ subordinated to \mathcal{U} . Define

$$(5) \quad \mu = \sum_{\alpha} \rho_\alpha \mu_\alpha$$

• **Goodness of μ on a fixed stratum S :** We verify the property (4) on S .

(\Rightarrow) Write χ, e (resp. χ_α, e_α) for the characteristic form and the Euler form induced by μ on X (resp. by μ_α on X_α). Notice that

$$(6) \quad d\chi = \sum_{\alpha} (d\rho_\alpha) \wedge \chi_\alpha + \sum_{\alpha} \rho_\alpha d\chi_\alpha$$

In the above expression, the first sum of the right side has perverse degree 1 (see §5.3). Recall that, by §5.5, $\|e\|_{\pi(S)} = \|d\chi\|_S$. If $\|d\chi\|_S = 2$ then, by equation (6),

$$\|d\chi_\alpha\|_{S \cap X_\alpha} = \|e_\alpha\|_{\pi(S \cap X_\alpha)} = 2$$

for some X_α intersecting S . So $\varepsilon_L \neq 0$ because μ_α is a good metric.

(\Leftarrow) In the rest of this proof we use some local properties of intersection cohomology. In particular, we use the step cohomology of a product $U \times c(L/\mathbb{S}^1)$ as it is defined in [10]. In section §7 the reader will find more details.

Assume that $\|e\|_{\pi(S)} < 2$ and take some $X_\alpha = \text{Im}(\alpha) \in \mathcal{U}$, the image of a modelled chart α on S . Write $B_\alpha = \pi(X_\alpha) \cong U_\alpha \times c(L/\mathbb{S}^1)$; so that $\|e|_{B_\alpha}\|_{U_\alpha} < 2$.

Consider the short exact sequence of step intersection cohomology

$$0 \rightarrow \Omega_{\mathbb{T}}^*(B_\alpha) \xrightarrow{i} \Omega_{\mathbb{Z}}^*(B_\alpha) \xrightarrow{pr} \Omega_{\mathbb{Z}/1}^*(B_\alpha) \rightarrow 0$$

which induces the long exact sequence

$$\dots \rightarrow H_{\mathbb{T}}^2(B_\alpha) \rightarrow H_{\mathbb{Z}}^2(B_\alpha) \xrightarrow{pr^*} H_{\mathbb{Z}/1}^2(B_\alpha) \xrightarrow{d} H_{\mathbb{T}}^3(B_\alpha) \rightarrow \dots$$

The inclusion $\iota_\epsilon : L/\mathbb{S}^1 \rightarrow U_\alpha \times c(L/\mathbb{S}^1)$ given by $p \mapsto (x_0, [p, \epsilon])$, induces the isomorphism

$$\iota_\epsilon^* : H_{\mathbb{Z}/1}^2(U_\alpha \times c(L/\mathbb{S}^1)) \xrightarrow{\cong} H_{\mathbb{Z}}^2(L/\mathbb{S}^1)$$

where $x_0 \in U_\alpha$ and $\epsilon > 0$. By the above remarks, $(\alpha \iota_\epsilon)^{-*}(\varepsilon_L) = pr^*[e|_{B_\alpha}] = 0$; so $\varepsilon_L = 0$.

□

5.8. Modelled morphisms Let $\Phi : \mathbb{S}^1 \times X \rightarrow X$ be a modelled action. A **perverse point** in X is a point of a perverse stratum. We will write X^{perv} for the set of perverse points, which is the union of the perverse strata, and $X^{\mathbb{S}^1}$ as usual for the set of fixed points. Let $\mathbb{S}^1 \times Y \rightarrow Y$ be any other modelled action. An unfoldable morphism

$$\begin{array}{ccc} \tilde{X} & \xrightarrow{\tilde{\alpha}} & \tilde{Y} \\ \mathbf{L} \downarrow & & \downarrow \mathbf{L}' \\ X & \xrightarrow{\alpha} & Y \end{array}$$

is said to be **modelled** if and only if each arrow in the above diagram is equivariant and

- (1) $\alpha^{-1}(Y^{\mathbb{S}^1}) \subset X^{\mathbb{S}^1}$.
- (2) $\alpha^{-1}(Y^{perv}) \subset X^{perv}$.

In other words, a modelled morphism is an equivariant unfoldable morphism which preserves the classification of the strata. For instance, any modelled chart of a fixed stratum in X is a modelled embedding in this new sense.

Proposition 5.9 (Functoriality of the Euler class). *The Euler class is preserved by modelled morphisms: If $\alpha : X \rightarrow Y$ is a modelled morphism, then $\alpha^*(\varepsilon_Y) = \varepsilon_X$.*

Proof. Write $\bar{\varepsilon}_X, \bar{\varepsilon}_Y$ for the Euler perversities in the orbit spaces B_X, B_Y . Each modelled morphism $\alpha : X \rightarrow Y$ induces an unfoldable morphism in the orbit spaces

$$\begin{array}{ccc} \tilde{B}_X & \xrightarrow{\tilde{\alpha}} & \tilde{B}_Y \\ \mathbf{L} \downarrow & & \downarrow \mathbf{L}' \\ B_X & \xrightarrow{\alpha} & B_Y \end{array}$$

which we still write α with a little abuse of notation. In order to see that the arrow

$$(7) \quad \alpha^* : \Omega_{\bar{\varepsilon}_Y}^*(B_Y) \rightarrow \Omega_{\bar{\varepsilon}_X}^*(B_X)$$

makes sense, we consider a third perversity $\alpha^*[\bar{\varepsilon}_Y]$ such that the arrow

$$\alpha^* : \Omega_{\bar{\varepsilon}_Y}^*(B_Y) \rightarrow \Omega_{\alpha^*[\bar{\varepsilon}_Y]}^*(B_X)$$

is well defined and $\alpha^*[\bar{\varepsilon}_Y] \leq \bar{\varepsilon}_X$. Then $\Omega_{\alpha^*[\bar{\varepsilon}_Y]}^*(B_X) \subset \Omega_{\bar{\varepsilon}_X}^*(B_X)$ and the map (7) is the composition with the inclusion.

The perversity $\alpha^*[\bar{\varepsilon}_Y]$ in B_X is given by the rule

$$\alpha^*[\bar{\varepsilon}_Y](\pi(S)) = \bar{\varepsilon}_Y(\pi(R))$$

for any singular strata S, R respectively in X, Y ; such that $\alpha(S) \subset R$. In this situation, we only need to show that

$$\alpha^*[\bar{e}_Y](\pi(S)) \leq \bar{e}_X(\pi(S))$$

If S is mobile then R is mobile since α is equivariant, so $\bar{e}_Y(\pi(R)) = \bar{e}_X(\pi(S)) = 0$ and the inequality holds. By the other hand, if S is fixed then it is a consequence of §5.8, since α preserves the classification of the strata. \square

6. THE GYSIN SEQUENCE

Take a stratified pseudomanifold X , a modelled action $\Phi : \mathbb{S}^1 \times X \rightarrow X$ with orbit space $B = X/\mathbb{S}^1$, and a perversity \bar{q} in X . The orbit map $\pi : X \rightarrow B$ preserves the strata and the perverse degree (see §5.5). Passing to the intersection cohomology we get a map

$$\pi^* : H_{\bar{q}}^*(B) \rightarrow H_{\bar{q}}^*(X)$$

which is a string of a long exact sequence; the Gysin sequence of X induced by the action. The third complex in the Gysin sequence is the Gysin term; its cohomology depends at the same time on global and local basic data. Global data concerns the Euler class $\varepsilon \in H_{\bar{q}}^*(B)$ induced by the action Φ , while local data concerns the Euler classes of the links of the perverse strata.

For instance, if $\Sigma = \emptyset$ then $\pi : X \rightarrow B$ is a smooth \mathbb{S}^1 -principal fiber bundle; we get the Gysin sequence by integrating along the fibers. If X is a manifold and Φ is a smooth non-free effective action, then $\Sigma \neq \emptyset$ and B is not a manifold anymore, but a stratified pseudomanifold. There is a Gysin sequence relating the DeRham cohomology of X with the intersection cohomology of B [9]. Something analogous happens for modelled actions on stratified pseudomanifolds with length ≤ 1 , cf. [15].

6.1. The Gysin Sequence Fix a modelled action $\Phi : \mathbb{S}^1 \times X \rightarrow X$ on a stratified pseudomanifold X , and perversity \bar{q} in X . Write still \bar{q} for the obvious perversity induced on B . The orbit map $\pi : X \rightarrow B$ induces a short exact sequence

$$0 \rightarrow \Omega_{\bar{q}}^{*+1}(B) \xrightarrow{\pi^*} \mathrm{I}\Omega_{\bar{q}}^{*+1}(X) \xrightarrow{pr} \mathcal{G}_{\bar{q}}^*(B) \rightarrow 0$$

The complex $\mathcal{G}_{\bar{q}}^*(B)$ is the **Gysin term**. There is a long exact sequence

$$(8) \quad \cdots \rightarrow H_{\bar{q}}^{i+1}(X) \xrightarrow{pr^*} H^i(\mathcal{G}_{\bar{q}}(B)) \xrightarrow{\partial} H_{\bar{q}}^{i+2}(B) \xrightarrow{\pi^*} H_{\bar{q}}^{i+2}(X) \rightarrow \cdots$$

This is the **Gysin sequence** of X . ♣

When the singular part of X is the empty set, then Φ is a free smooth action and $\pi : X \rightarrow B$ is a smooth principal fiber bundle with group \mathbb{S}^1 ; so (8) is the usual Gysin sequence. The cohomology of the Gysin term is $H^*(\mathcal{G}_{\bar{q}}(B)) = H^*(B)$ is the DeRham cohomology and the connecting homomorphism is the multiplication by the Euler class

(see [3], [8]). When X has a nonempty singular part then, by §3.3, for big perversities $\bar{q} > \bar{t}$ the sequence (8) is the usual Gysin sequence of $X - \Sigma$ in DeRham cohomology; and for negative perversities $\bar{q} < \bar{0}$ it is the Gysin sequence in relative cohomology. In general; we could naively conjecture that $H^*(\mathcal{G}_{\bar{q}}(B)) = H_{\bar{q}}^*(B)$ and the connecting morphism of the Gysin sequence to be the multiplication by the Euler class. As we will see, the real life is richer and more complicated.

Although the Gysin term is a quotient complex, it can be written by means of basic differential forms. The **characteristic perversity** in B induced by the action is

$$\bar{\chi}(\pi(S)) = \|\chi\|_S = \begin{cases} 1 & S \text{ a fixed stratum} \\ 0 & S \text{ a mobile stratum} \end{cases}$$

Lemma 6.2. *For each perversity $\bar{0} \leq \bar{q} \leq \bar{t}$ in X , the Gysin term $\mathcal{G}_{\bar{q}}^*(B)$ is isomorphic to the following complex*

$$\left\{ \begin{array}{l} \theta \in \Omega_{\bar{q}-\bar{\chi}}^*(B) / \exists \nu \in \Omega^*(B - \Sigma) : \\ \begin{array}{l} (1) \nu \text{ is liftable.} \\ (2) \max\{\|\nu\|_S, \|d\nu + e \wedge \theta\|_S\} \leq \bar{q}(S) \forall S \text{ perverse stratum} \end{array} \end{array} \right\}$$

Under this identification, the connecting homomorphism is

$$\partial : H^i(\mathcal{G}_{\bar{q}}(B)) \rightarrow H_{\bar{q}}^i(B) \quad \partial[\theta] = [d\nu + e \wedge \theta]$$

Proof. The restriction $\pi : X - \Sigma \rightarrow B - \Sigma$ is a \mathbb{S}^1 -principal fiber bundle. Consider the morphism of integration along the orbits

$$\oint = (-1)^{i-1} \pi^{-*} i_C : \mathbb{I}\Omega_{\bar{q}}^i(X) \rightarrow \Omega_{\bar{q}-\bar{\chi}}^{i-1}(B)$$

defined by

$$\oint \omega = (-1)^{i-1} \theta \quad \omega = \pi^*(\nu) + \chi \wedge \pi^*(\theta) \in \mathbb{I}\Omega_{\bar{q}}^i(X - \Sigma)$$

This morphism commutes with the differential.

The Gysin term $\mathcal{G}_{\bar{q}}^*(B) = \mathbb{I}\Omega_{\bar{q}}^{i+1}(X) / \pi^*(\Omega_{\bar{q}}^{i+1}(B))$ is a quotient with differential operator $\bar{d}(\bar{\omega}) = \overline{d\omega}$, where $\bar{\omega}$ is the equivalence class of a differential form $\omega \in \mathbb{I}\Omega_{\bar{q}}^i(X)$. The integration along the orbits passes well to this quotient,

$$\oint \mathcal{G}_{\bar{q}}^*(B) \rightarrow \Omega_{\bar{q}-\bar{\chi}}^*(B) \quad \bar{\omega} \mapsto \oint(\omega)$$

The complex given in the statement of §6.2 is the image of the above arrow. The connecting homomorphism arises as usual, from the Snake's Lemma. \square

In some cases, the cohomology of the Gysin term is closer of our naive conjecture.

Proposition 6.3. *If X has no perverse strata then the Euler class belongs to $H_{\bar{\chi}}^2(B)$ and, for each perversity $\bar{0} \leq \bar{q} \leq \bar{t}$, the Gysin sequence (8) becomes*

$$\dots \rightarrow H_{\bar{q}}^{i+1}(X) \xrightarrow{f} H_{\bar{q}-\bar{e}}^i(B) \xrightarrow{\varepsilon} H_{\bar{q}}^{i+2}(B) \xrightarrow{\pi^*} H_{\bar{q}}^{i+2}(X) \rightarrow \dots$$

where the connecting homomorphism ε is the multiplication by the Euler Class. If additionally X has no fixed strata, then the Euler class belongs to $H_{\bar{0}}^2(B)$ and the above sequence becomes

$$\dots \rightarrow H_{\bar{q}}^{i+1}(X) \xrightarrow{f} H_{\bar{q}}^i(B) \xrightarrow{\varepsilon} H_{\bar{q}}^{i+2}(B) \xrightarrow{\pi^*} H_{\bar{q}}^{i+2}(X) \rightarrow \dots$$

Proof. By §6.2 and the definition of $\bar{\chi}$, \bar{e} ; the Gysin term is an intermediate complex

$$(9) \quad \Omega_{\bar{q}-\bar{e}}^*(B) \subset \mathcal{G}_{\bar{q}}^*(B) \subset \Omega_{\bar{q}-\bar{\chi}}^*(B)$$

Now X has no perverse strata iff $\bar{\chi} = \bar{e}$ and the extremes in the above inequality are identical. For the connecting homomorphism we take the formula in §6.2 with $\nu = 0$. \square

Corollary 6.4. *If the Euler class $\varepsilon \in H_{\bar{e}}^2(B)$ vanishes then*

$$H_{\bar{q}}^*(X) = H_{\bar{q}}^{*-1}(B) \oplus H_{\bar{q}-\bar{e}}^*(B)$$

for each perversity $\bar{0} \leq \bar{q} \leq \bar{t}$. If additionally X has no fixed strata, then

$$H_{\bar{q}}(X) = H_{\bar{q}}(B) \otimes H(\mathbb{S}^1)$$

i.e., X is a cohomological product for intersection cohomology.

Proof. If the Euler class vanishes then X has no perverse strata. \square

6.5. Residual approximations Now let's assume that X has perverse strata. We will define the residual terms; those terms allows us to measure the difference between $H^*(\mathcal{G}_{\bar{q}}(B))$ and the intersection cohomology of B .

Consider the inclusions in the inequality (9). These inclusions provide the following short exact sequences

$$\begin{aligned} 0 \rightarrow \Omega_{\bar{q}-\bar{e}}^*(B) \hookrightarrow \mathcal{G}_{\bar{q}}^*(B) \xrightarrow{pr} \mathfrak{L}\mathfrak{ow}_{\bar{q}}^*(B) \rightarrow 0 \\ 0 \rightarrow \mathcal{G}_{\bar{q}}^*(B) \hookrightarrow \Omega_{\bar{q}-\bar{\chi}}^*(B) \xrightarrow{pr} \mathfrak{U}\mathfrak{pp}_{\bar{q}}^*(B) \rightarrow 0 \end{aligned}$$

We call $\mathfrak{L}\mathfrak{ow}_{\bar{q}}^*(B)$ (resp. $\mathfrak{U}\mathfrak{pp}_{\bar{q}}^*(B)$) the **lower residue** (resp. **upper residue**). The induced long exact sequences

$$(10) \quad \dots \rightarrow H_{\bar{q}-\bar{e}}^i(B) \rightarrow H^i(\mathcal{G}_{\bar{q}}(B)) \xrightarrow{pr} H^i(\mathfrak{L}\mathfrak{ow}_{\bar{q}}(B)) \xrightarrow{\partial'} H_{\bar{q}-\bar{e}}^{i+1}(B) \rightarrow \dots$$

$$(11) \quad \dots \rightarrow H^i(\mathcal{G}_{\bar{q}}(B)) \rightarrow H_{\bar{q}-\bar{\chi}}^i(B) \xrightarrow{pr} H^i(\mathfrak{U}\mathfrak{pp}_{\bar{q}}(B)) \xrightarrow{\partial''} H^{i+1}(\mathcal{G}_{\bar{q}}(B)) \rightarrow \dots$$

are the **residual approximations**. Next consider the cokernel

$$0 \rightarrow \Omega_{\frac{q-e}{q}}^*(B) \hookrightarrow \Omega_{\frac{q-\bar{x}}{q}}^*(B) \xrightarrow{pr} \Omega_{\frac{q-\bar{x}}{q-e}}^*(B) \rightarrow 0$$

its cohomology $H_{\frac{q-\bar{x}}{q-e}}^*(B)$ is called the **step intersection cohomology** of B [10]. The residual approximations are related by the long exact sequences

$$\begin{aligned} \dots \rightarrow H_{\frac{q-e}{q}}^i(B) \rightarrow H_{\frac{q-\bar{x}}{q}}^i(B) \rightarrow H_{\frac{q-\bar{x}}{q-e}}^i(B) \xrightarrow{d} H_{\frac{q-e}{q}}^{i+1}(B) \rightarrow \dots \\ \dots \rightarrow H^i(\mathcal{L}ow_{\frac{q}{q}}(B)) \rightarrow H_{\frac{q-\bar{x}}{q-e}}^i(B) \rightarrow H^i(\mathcal{U}pp_{\frac{q}{q}}(B)) \rightarrow H^i(\mathcal{L}ow_{\frac{q}{q}}(B)) \rightarrow \dots \end{aligned}$$

These sequences can be arranged in a commutative exact diagram; called the **Gysin braid**

$$\begin{array}{ccccccc} & & \curvearrowright & & \curvearrowright & & \curvearrowright \\ & & \downarrow & & \downarrow & & \downarrow \\ H_{\frac{q-e}{q}}^i(B) & & & H_{\frac{q-\bar{x}}{q}}^i(B) & & H^i(\mathcal{U}pp_{\frac{q}{q}}(B)) & & H^{i+1}(\mathcal{L}ow_{\frac{q}{q}}(B)) \\ & \searrow & & \swarrow & & \searrow & & \swarrow \\ & & H^i(\mathcal{G}_{\frac{q}{q}}(B)) & & H_{\frac{q-\bar{x}}{q-e}}^i(B) & & H^{i+1}(\mathcal{G}_{\frac{q}{q}}(B)) & \\ & \swarrow & & \searrow & & \swarrow & & \searrow \\ H^{i-1}(\mathcal{U}pp_{\frac{q}{q}}(B)) & & H^i(\mathcal{L}ow_{\frac{q}{q}}(B)) & & H_{\frac{q-e}{q}}^{i+1}(B) & & H_{\frac{q-\bar{x}}{q}}^{i+1}(B) \\ & & \curvearrowleft & & \curvearrowleft & & \curvearrowleft \end{array}$$

7. THE GYSIN THEOREM

We devote the rest of this work to calculate the residual cohomologies. The final goal is to relate $H^*(\mathcal{G}_{\frac{q}{q}}(B))$ with basic local cohomological data by means of the residual approximations; so we start this § with the local properties of the residues. An introduction to presheaves, sheaves and Čech cohomology can be found in [3], [5]; some results of this § were taken from [21].

Recall that a **presheaf** \mathcal{P} on X is **complete** (or it is a **sheaf**) iff, for any open cover $\mathcal{U} = \{X_{\alpha}\}_{\alpha}$ of X , the augmented Čech differential complex

$$(12) \quad 0 \rightarrow \mathcal{P}(X) \xrightarrow{\delta} C^0(\mathcal{U}, \mathcal{P}) \xrightarrow{\delta} C^1(\mathcal{U}, \mathcal{P}) \xrightarrow{\delta} C^2(\mathcal{U}, \mathcal{P}) \xrightarrow{\delta} \dots$$

is exact, where $C^j(\mathcal{U}, \mathcal{P}) = \prod_{\alpha_0 < \dots < \alpha_j} \mathcal{P}(X_{\alpha_0} \cap \dots \cap X_{\alpha_j})$ and δ is given coordinatewise by

the alternating sum of the restrictions. Notice that $(C^*(\mathcal{U}, \mathcal{P}), \delta)$ is a cohomological resolution of $\mathcal{P}(X)$.

For each perversity \bar{q} , the complex of \bar{q} -forms $\Omega_{\bar{q}}^*(-)$ is a presheaf on X (and also on B). The complex $\mathbb{I}\Omega_{\bar{q}}^*(-)$ of invariant \bar{q} -forms is a presheaf on X but it is defined in the topology of invariant open sets; we can also see it as a presheaf on B up to a composition with the orbit map. The complexes $\mathcal{G}_{\bar{q}}^*(-)$, $\mathfrak{L}\mathfrak{ow}_{\bar{q}}^*(-)$, and $\mathfrak{U}\mathfrak{pp}_{\bar{q}}^*(-)$ are presheaves on B .

Because of the existence of controlled invariant partitions of the unity, all these examples are sheaves. We will study their cohomological properties on the charts of B and X .

Lemma 7.1. *Let $\Phi : \mathbb{S}^1 \times X \rightarrow X$ be a modelled action. Consider on $\mathbb{R} \times X$ the (obvious) modelled action trivial in \mathbb{R} . Then the projection $pr : \mathbb{R} \times X \rightarrow X$ induces the following isomorphisms*

$$\begin{aligned} H_{\bar{q}}^i(\mathbb{R} \times X) &= H_{\bar{q}}^i(X) & H^i(\mathcal{G}_{\bar{q}}(\mathbb{R} \times B)) &= H^i(\mathcal{G}_{\bar{q}}(B)) \\ H^i(\mathfrak{L}\mathfrak{ow}_{\bar{q}}(\mathbb{R} \times B)) &= H^i(\mathfrak{L}\mathfrak{ow}_{\bar{q}}(B)) & H^i(\mathfrak{U}\mathfrak{pp}_{\bar{q}}(\mathbb{R} \times B)) &= H^i(\mathfrak{U}\mathfrak{pp}_{\bar{q}}(B)) \end{aligned}$$

Proof. See [15]. □

Proposition 7.2. *Let $\Psi : \mathbb{S}^1 \times L \rightarrow L$ be a modelled action on a compact stratified pseudomanifold L . For each perversity $\bar{0} \leq \bar{q} \leq \bar{t}$ and each $\epsilon > 0$ the map $\iota_{\epsilon} : L \rightarrow c(L)$ given by $p \mapsto [p, \epsilon]$ induces the following isomorphisms*

$$(13) \quad H_{\bar{q}}^i(c(L/\mathbb{S}^1)) = \begin{cases} H_{\bar{q}}^i(L/\mathbb{S}^1) & i \leq \bar{q}(\star) \\ 0 & i > \bar{q}(\star) \end{cases}$$

Also

$$(14) \quad H^i(\mathcal{G}_{\bar{q}}(c(L/\mathbb{S}^1))) = \begin{cases} H^i(\mathcal{G}_{\bar{q}}(L/\mathbb{S}^1)) & i \leq \bar{q}(\star) - 2 \\ \ker [\partial : H^{\bar{q}(\star)-1}(\mathcal{G}_{\bar{q}}(L/\mathbb{S}^1)) \rightarrow H_{\bar{q}}^{\bar{q}(\star)+1}(L/\mathbb{S}^1)] & i = \bar{q}(\star) - 1 \\ 0 & i \geq \bar{q}(\star) \end{cases}$$

where ∂ is the connecting homomorphism of the Gysin sequence on L .

Proof. For the first isomorphism see [21]. For the second one, we get a commutative diagram with exact horizontal rows

$$\begin{array}{ccccccc} \rightarrow & H_{\bar{q}}^{i+1}(c(L/\mathbb{S}^1)) & \xrightarrow{\pi^*} & H_{\bar{q}}^{i+1}(c(L)) & \rightarrow & H^i(\mathcal{G}_{\bar{q}}(c(L/\mathbb{S}^1))) & \xrightarrow{\partial} & H_{\bar{q}}^{i+2}(c(L/\mathbb{S}^1)) & \rightarrow \\ & \downarrow & & \downarrow & & \downarrow & & \downarrow & \\ \rightarrow & H_{\bar{q}}^{i+1}(L/\mathbb{S}^1) & \xrightarrow{\pi^*} & H_{\bar{q}}^{i+1}(L) & \rightarrow & H^i(\mathcal{G}_{\bar{q}}(L/\mathbb{S}^1)) & \xrightarrow{\partial} & H_{\bar{q}}^{i+2}(L/\mathbb{S}^1) & \rightarrow \end{array}$$

where the vertical arrows are induced by $\iota_{\epsilon} : L \rightarrow c(L)$ and $\iota_{\epsilon} : L/\mathbb{S}^1 \rightarrow c(L/\mathbb{S}^1)$. For $i \leq \bar{q}(\star) - 2$ we have enough vertical isomorphisms. By the Five Lemma

$$H^i(\mathcal{G}_{\bar{q}}(c(L/\mathbb{S}^1))) = H^i(\mathcal{G}_{\bar{q}}(L/\mathbb{S}^1))$$

For $i = \bar{q}(\star) - 1$ the two left vertical arrows are isomorphisms. In the upper right coin we get $H_{\bar{q}}^{\bar{q}(\star)+1}(c(L/\mathbb{S}^1)) = 0$. So

$$H^{\bar{q}(\star)-1}(\mathcal{G}_{\bar{q}}(c(L/\mathbb{S}^1))) = \text{coker}(\pi^*) = \ker \left[H^{\bar{q}(\star)-1}(\mathcal{G}_{\bar{q}}(L/\mathbb{S}^1)) \xrightarrow{\partial} H_{\bar{q}}^{\bar{q}(\star)+1}(L/\mathbb{S}^1) \right]$$

For $i \geq \bar{q}(\star)$ the upper horizontal row has four zeros, thus $H^i(\mathcal{G}_{\bar{q}}(c(L/\mathbb{S}^1))) = 0$. \square

Corollary 7.3. *In the same situation of §7.2; if the vertex is not perverse then the Gysin sequence of $c(L)$ is the Gysin sequence of L truncated in dimension $i = \bar{q}(\star) - 1$.*

Proposition 7.4. *In the same situation of §7.2; if the vertex is a perverse stratum then the map $\iota_\epsilon : L \rightarrow c(L)$ induces the following isomorphisms*

$$(15) \quad H^i(\mathfrak{L}\mathfrak{om}_{\bar{q}}(c(L/\mathbb{S}^1))) = \begin{cases} H^i(\mathfrak{L}\mathfrak{om}_{\bar{q}}(L/\mathbb{S}^1)) & i \leq \bar{q}(\star) - 3 \\ \ker\{\partial' : H^{\bar{q}(\star)-2}(\mathfrak{L}\mathfrak{om}_{\bar{q}}(L/\mathbb{S}^1)) \rightarrow H_{\bar{q}-\bar{\epsilon}}^{\bar{q}(\star)-1}(L/\mathbb{S}^1)\} & i = \bar{q}(\star) - 2 \\ \ker\{\partial : H^{\bar{q}(\star)-1}(\mathcal{G}_{\bar{q}}(L/\mathbb{S}^1)) \rightarrow H_{\bar{q}}^{\bar{q}(\star)+1}(L/\mathbb{S}^1)\} & i = \bar{q}(\star) - 1 \\ 0 & i \geq \bar{q}(\star) \end{cases}$$

And

$$(16) \quad H^i(\mathfrak{U}\mathfrak{pp}_{\bar{q}}(c(L/\mathbb{S}^1))) = \begin{cases} H^i(\mathfrak{U}\mathfrak{pp}_{\bar{q}}(L/\mathbb{S}^1)) & i \leq \bar{q}(\star) - 3 \\ \ker[\partial'' \partial : H^{\bar{q}(\star)-2}(\mathfrak{U}\mathfrak{pp}_{\bar{q}}(L/\mathbb{S}^1)) \rightarrow H_{\bar{q}}^{\bar{q}(\star)+1}(L/\mathbb{S}^1)] & i = \bar{q}(\star) - 2 \\ \frac{H_{\bar{q}-\bar{\chi}}^{\bar{q}(\star)-1}(L/\mathbb{S}^1)}{j^*(\ker^{\bar{q}(\star)-1}(\partial))} & i = \bar{q}(\star) - 1 \\ 0 & i \geq \bar{q}(\star) \end{cases}$$

Proof. We get the following commutative diagrams

$$(14) \quad \begin{array}{ccccccc} \rightarrow & H_{\bar{q}-\bar{\epsilon}}^i(c(L/\mathbb{S}^1)) & \xrightarrow{j^*} & H^i(\mathcal{G}_{\bar{q}}(c(L/\mathbb{S}^1))) & \rightarrow & H^i(\mathfrak{L}\mathfrak{om}_{\bar{q}}(c(L/\mathbb{S}^1))) & \xrightarrow{\partial'} & H_{\bar{q}-\bar{\epsilon}}^{i+1}(c(L/\mathbb{S}^1)) & \rightarrow \\ & \downarrow & & \downarrow & & \downarrow & & \downarrow & \\ \rightarrow & H_{\bar{q}-\bar{\epsilon}}^i(L/\mathbb{S}^1) & \xrightarrow{j^*} & H^i(\mathcal{G}_{\bar{q}}(L/\mathbb{S}^1)) & \rightarrow & H^i(\mathfrak{L}\mathfrak{om}_{\bar{q}}(L/\mathbb{S}^1)) & \xrightarrow{\partial'} & H_{\bar{q}-\bar{\epsilon}}^{i+1}(L/\mathbb{S}^1) & \rightarrow \end{array}$$

and

$$(15) \quad \begin{array}{ccccccc} \rightarrow & H^i(\mathcal{G}_{\bar{q}}(c(L/\mathbb{S}^1))) & \xrightarrow{j^*} & H_{\bar{q}-\bar{\chi}}^i(c(L/\mathbb{S}^1)) & \rightarrow & H^i(\mathfrak{U}\mathfrak{pp}_{\bar{q}}(c(L/\mathbb{S}^1))) & \xrightarrow{\partial''} & H^{i+1}(\mathcal{G}_{\bar{q}}(c(L/\mathbb{S}^1))) & \rightarrow \\ & \downarrow & & \downarrow & & \downarrow & & \downarrow & \\ \rightarrow & H^i(\mathcal{G}_{\bar{q}}(L/\mathbb{S}^1)) & \xrightarrow{j^*} & H_{\bar{q}-\bar{\chi}}^i(L/\mathbb{S}^1) & \rightarrow & H^i(\mathfrak{U}\mathfrak{pp}_{\bar{q}}(L/\mathbb{S}^1)) & \xrightarrow{\partial''} & H^{i+1}(\mathcal{G}_{\bar{q}}(L/\mathbb{S}^1)) & \rightarrow \end{array}$$

where the horizontal rows are residual approximations and the vertical arrows are induced by the maps $\iota_\epsilon : L \rightarrow c(L)$ and $\iota_\epsilon : L/\mathbb{S}^1 \rightarrow c(L/\mathbb{S}^1)$.

Since the vertex is a perverse stratum $\bar{\chi}(\star) = 1$ and $\bar{\epsilon}(\star) = 2$. In the above diagrams the case $i \leq \bar{q}(\star) - 3$ is direct from the Five Lemma and the case $i \geq \bar{q}(\star)$ is straightforward.

We verify the cases $i = \bar{q}(\star) - 2$ and $i = \bar{q}(\star) - 1$ proceeding in two steps.

• **Lower residue:** For $i = \bar{q}(\star) - 2$, by §7.2 the two left vertical arrows in diagram (14) are isomorphisms. In the upper right corner we get $H_{\bar{q}-\bar{e}}^{\bar{q}(\star)-1}(c(L/\mathbb{S}^1)) = 0$. So

$$H^{\bar{q}(\star)-2}(\mathcal{L}\mathbf{ow}_{\bar{q}}(c(L/\mathbb{S}^1))) = \ker^{\bar{q}(\star)-2}[\partial' : H^{\bar{q}(\star)-2}(\mathcal{L}\mathbf{ow}_{\bar{q}}(L/\mathbb{S}^1)) \rightarrow H_{\bar{q}-\bar{e}}^{\bar{q}(\star)-1}(L/\mathbb{S}^1)]$$

For $i = \bar{q}(\star) - 1$ the upper corners are zeros. By §7.2 and the exactness of the upper horizontal row,

$$H^{\bar{q}(\star)-1}(\mathcal{L}\mathbf{ow}_{\bar{q}}(c(L/\mathbb{S}^1))) = H^{\bar{q}(\star)-1}(\mathcal{G}_{\bar{q}}(c(L/\mathbb{S}^1))) = \ker^{\bar{q}(\star)}(\partial)$$

• **Upper residue:** For $i = \bar{q}(\star) - 2$; by §7.2 the left vertical arrows in the diagram (15) are isomorphisms. Hence

$$\iota_\epsilon : H^{\bar{q}(\star)-2}(\mathcal{U}\mathbf{pp}_{\bar{q}}(c(L/\mathbb{S}^1))) \rightarrow H^{\bar{q}(\star)-2}(\mathcal{U}\mathbf{pp}_{\bar{q}}(L/\mathbb{S}^1))$$

is injective. We get a commutative exact diagram

$$\begin{array}{ccccccc} & & 0 & & 0 & & 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 \rightarrow & \text{coker}^{\bar{q}(\star)-2}(j^\star, c(L/\mathbb{S}^1)) & \rightarrow & H^{\bar{q}(\star)-2}(\mathcal{U}\mathbf{pp}_{\bar{q}}(c(L/\mathbb{S}^1))) & \rightarrow & \ker^{\bar{q}(\star)-1}(j^\star, c(L/\mathbb{S}^1)) & \rightarrow 0 \\ & \downarrow & & \downarrow^{\iota_\epsilon} & & \downarrow & \\ 0 \rightarrow & \text{coker}^{\bar{q}(\star)-2}(j^\star, L/\mathbb{S}^1) & \rightarrow & H^{\bar{q}(\star)-2}(\mathcal{U}\mathbf{pp}_{\bar{q}}(L/\mathbb{S}^1)) & \rightarrow & \ker^{\bar{q}(\star)-1}(j^\star, L/\mathbb{S}^1) & \rightarrow 0 \\ & \downarrow & & \downarrow & & \downarrow & \\ & 0 & \rightarrow & \text{coker}^{\bar{q}(\star)-2}(\iota_\epsilon) & \rightarrow & \frac{\ker^{\bar{q}(\star)-1}(j^\star, L/\mathbb{S}^1)}{\ker^{\bar{q}(\star)-1}(j^\star, c(L/\mathbb{S}^1))} & \rightarrow 0 \\ & & & \downarrow & & \downarrow & \\ & & & 0 & & 0 & \end{array}$$

So $H^{\bar{q}(\star)-2}(\mathcal{U}\mathbf{pp}_{\bar{q}}(c(L/\mathbb{S}^1)))$ is the kernel of the map

$$\bar{\tau}_\epsilon : H^{\bar{q}(\star)-2}(\mathcal{U}\mathbf{pp}_{\bar{q}}(c(L/\mathbb{S}^1))) \rightarrow \frac{\ker^{\bar{q}(\star)-1}(j^\star, L/\mathbb{S}^1)}{\ker^{\bar{q}(\star)-1}(j^\star, c(L/\mathbb{S}^1))} = \frac{\text{Im}^{\bar{q}(\star)-1}(\partial', L/\mathbb{S}^1)}{\text{Im}^{\bar{q}(\star)-1}(\partial', L/\mathbb{S}^1) \cap \ker^{\bar{q}(\star)-1}(\partial)}$$

In the last equality we used §7.2, the exactness of the upper approximation and the fact that the third vertical arrow in the diagram (15) is injective. So we can identify the image of the third vertical arrow with $\ker^{\bar{q}(\star)-1}(\partial)$, the kernel of the connecting homomorphism of the Gysin sequence on L . We deduce that $\ker(\bar{\tau}_\epsilon)$ is the kernel of the composition

$$H^{\bar{q}(\star)-2}(\mathcal{U}\mathbf{pp}_{\bar{q}}(L/\mathbb{S}^1)) \xrightarrow{\partial''} H^{\bar{q}(\star)-1}(\mathcal{G}_{\bar{q}}(L/\mathbb{S}^1)) \xrightarrow{\partial} H_{\bar{q}}^{\bar{q}(\star)+1}(L/\mathbb{S}^1)$$

For $i = \bar{q}(\star) - 1$ the first left vertical arrow in the diagram (15) is injective, the second is an isomorphism. In the upper right corner we get $H^{\bar{q}(\star)}(\mathcal{U}\mathbf{pp}_{\bar{q}}(c(L/\mathbb{S}^1))) = 0$. We obtain

the exact commutative diagram

$$\begin{array}{ccccccc} 0 \rightarrow & \text{coker}^{\bar{q}^{(*)-1}}(j, c(L/\mathbb{S}^1)) & \xrightarrow{\cong} & H^{\bar{q}^{(*)-1}}(\mathfrak{Upp}_{\bar{q}}(c(L/\mathbb{S}^1))) & \rightarrow & 0 \\ & \downarrow \bar{\iota}_\epsilon & & \downarrow \iota_\epsilon & & \downarrow \\ 0 \rightarrow & \text{coker}^{\bar{q}^{(*)-1}}(j, L/\mathbb{S}^1) & \rightarrow & H^{\bar{q}^{(*)-1}}(\mathfrak{Upp}_{\bar{q}}(L/\mathbb{S}^1)) & \rightarrow & \ker^{\bar{q}^{(*)}}(j, L/\mathbb{S}^1) \rightarrow 0 \end{array}$$

Notice that

$$\ker^{\bar{q}^{(*)-1}}(\iota_\epsilon) \cong \ker(\bar{\iota}_\epsilon) = \frac{\text{Im}^{\bar{q}^{(*)-1}}(j^*, L/\mathbb{S}^1)}{\iota_\epsilon(\text{Im}^{\bar{q}^{(*)-1}}(j^*, c(L/\mathbb{S}^1)))} = \frac{\text{Im}^{\bar{q}^{(*)-1}}(j^*, L/\mathbb{S}^1)}{j^*(\text{Im}^{\bar{q}^{(*)-1}}(\iota_\epsilon))} = \frac{\text{Im}^{\bar{q}^{(*)-1}}(j^*, L/\mathbb{S}^1)}{j^*(\ker^{\bar{q}^{(*)-1}}(\partial))}$$

Also

$$\text{Im}^{\bar{q}^{(*)-1}}(\iota_\epsilon) = \text{coker}^{\bar{q}^{(*)-1}}(j^*, L/\mathbb{S}^1) = \frac{H_{\bar{q}-\bar{x}}^{\bar{q}^{(*)-1}}(L/\mathbb{S}^1)}{\text{Im}^{\bar{q}^{(*)-1}}(j^*, L/\mathbb{S}^1)}$$

So we get a short exact sequence

$$0 \rightarrow \frac{\text{Im}^{\bar{q}^{(*)-1}}(j^*, L/\mathbb{S}^1)}{j^*(\ker^{\bar{q}^{(*)-1}}(\partial))} \rightarrow H^{\bar{q}^{(*)-1}}(\mathfrak{Upp}_{\bar{q}}(c(L/\mathbb{S}^1))) \rightarrow \frac{H_{\bar{q}-\bar{x}}^{\bar{q}^{(*)-1}}(L/\mathbb{S}^1)}{\text{Im}^{\bar{q}^{(*)-1}}(j^*, L/\mathbb{S}^1)} \rightarrow 0$$

We deduce that

$$H^{\bar{q}^{(*)-1}}(\mathfrak{Upp}_{\bar{q}}(c(L/\mathbb{S}^1))) = \frac{H_{\bar{q}-\bar{x}}^{\bar{q}^{(*)-1}}(L/\mathbb{S}^1)}{j^*(\ker^{\bar{q}^{(*)-1}}(\partial))}$$

This finishes the proof. \square

Remarks 7.5. Statements §7.2, §7.3 and §7.4 imply that

- (1) For each fixed stratum S in X and each unfoldable chart $\beta : U \times c(L/\mathbb{S}^1) \rightarrow B$, we can calculate the cohomology of the Gysin term and the residues on the open $\text{Im}(\beta)$.
- (2) According to [1], $\mathcal{G}_{\bar{q}}^*(-)$ is a constructible sheaf on B . Also $\mathfrak{L}ow_{\bar{q}}^*(-)$ and $\mathfrak{Upp}_{\bar{q}}^*(-)$ are constructible sheaves on B with support on the perverse points' set $X^{per\bar{v}}$.

Theorem 7.6 (The Gysin Theorem). *Let X be a stratified pseudomanifold, \bar{q} a perversity in X , $\bar{0} \leq \bar{q} \leq \bar{t}$. For each modelled action $\mathbb{S}^1 \times X \rightarrow X$ there are two long exact sequences relating the intersection cohomology of X and B : the Gysin sequence*

$$\dots \rightarrow H_{\bar{q}}^{i+1}(X) \rightarrow H^i(\mathcal{G}_{\bar{q}}(B)) \xrightarrow{\partial} H_{\bar{q}}^{i+2}(B) \xrightarrow{\pi^*} H_{\bar{q}}^{i+2}(X) \rightarrow \dots$$

induced by the orbit map $\pi : X \rightarrow B$, and the lower approximation

$$\dots \rightarrow H^i(\mathcal{G}_{\bar{q}}(B)) \rightarrow H^i(\mathfrak{L}ow_{\bar{q}}(B)) \xrightarrow{\partial'} H_{\bar{q}-\bar{e}}^{i+1}(B) \rightarrow H^{i+1}(\mathcal{G}_{\bar{q}}(B)) \rightarrow \dots$$

induced by the inclusion $\Omega_{\bar{q}-\bar{e}}^(B) \xrightarrow{\iota} \mathcal{G}_{\bar{q}}^*(B)$. These sequences satisfy*

- (1) *If X has no perverse strata then $\mathcal{G}_{\bar{q}}^*(B) = \Omega_{\bar{q}-\bar{x}}^*(B) = \Omega_{\bar{q}-\bar{e}}^*(B)$, $\mathfrak{L}ow_{\bar{q}}^*(B) = 0$ and the connecting homomorphism of the Gysin sequence is the multiplication by the Euler Class $\varepsilon \in H_{\bar{x}}^2(B)$.*

- (2) If X has perverse strata, then $H^*(\mathcal{L}\mathbf{ow}_{\bar{q}}(B))$ is calculated through a spectral sequence in B , whose second term

$$E_2^{ij} = H^j(X^{perv}, \mathcal{P}^i)$$

is the cohomology of the perverse points' set X^{perv} with values on a locally constant graduated constructible presheaf \mathcal{P}^* . For each fixed point $x \in X$ the stalks

$$\mathcal{P}_x^i = \begin{cases} H^i(\mathcal{L}\mathbf{ow}_{\bar{q}}(L/\mathbb{S}^1)) & i \leq \bar{q}(S) - 3 \\ \ker\{\partial' : H^{\bar{q}(S)-2}(\mathcal{L}\mathbf{ow}_{\bar{q}}(L/\mathbb{S}^1)) \rightarrow H_{\frac{\bar{q}-\varepsilon}{\bar{q}}}^{\bar{q}(S)-1}(L/\mathbb{S}^1)\} & i = \bar{q}(S) - 2 \\ \ker\{H^{\bar{q}(S)-1}(\mathcal{G}_{\bar{q}}(L/\mathbb{S}^1)) \xrightarrow{\partial} H_{\bar{q}}^{\bar{q}(S)+1}(L/\mathbb{S}^1)\} & i = \bar{q}(S) - 1 \\ 0 & i \geq \bar{q}(S) \end{cases}$$

depend on the Gysin sequence and the residual approximation induced by the action of \mathbb{S}^1 on of the link L of the stratum containing x .

Proof. Statement (1) has been already proved in the preceding sections. Statement (2) arises from the usual spectral sequence induced by a double complex; see for instance [3], [5]. The double complex we take is the residual Čech double complex

$$(C^j(\mathcal{U}, \mathcal{L}\mathbf{ow}_{\bar{q}}^i(-)), \delta, d)$$

induced by an invariant open cover $\mathcal{U} = \{B_\alpha\}_\alpha$ of B ; where δ is the Čech differential induced by the restrictions, and d is the usual differential operator. We define \mathcal{U} as follows: First take the complement of the fixed points' set $X_0 = X - X^{\mathbb{S}^1}$; we ask $B_0 = \pi(X_0)$ to be in \mathcal{U} . Second, for each fixed point $x \in X$ we take a modelled chart

$$\alpha : U_\alpha \times c(L) \rightarrow X$$

in the stratum S containing x ; such that $x \in U_\alpha$. We ask the U_α 's intersecting S to be a good cover of S . We take

$$B_\alpha = \pi(\text{Im}(\alpha)) \cong U_\alpha \times c(L/\mathbb{S}^1)$$

Since the sheaf $\mathcal{L}\mathbf{ow}_{\bar{q}}^*(-)$ vanishes identically on B_0 ; the second term of the spectral sequence is

$$E_2^{ij} = H_\delta^j H_d^i(\mathcal{U}, \mathcal{L}\mathbf{ow}_{\bar{q}}^i(-)) = H^j(\mathcal{U}, \mathcal{H}\mathcal{L}\mathbf{ow}_{\bar{q}}^i(-)) = H^j(X^{perv}, \mathcal{H}\mathcal{L}\mathbf{ow}_{\bar{q}}^i(-))$$

So $\mathcal{P}^* = \mathcal{H}\mathcal{L}\mathbf{ow}_{\bar{q}}^*(-)$ is the desired presheaf. The remarks on the stalks are immediate from §7.1, §7.2. \square

7.7. Exceptional actions A modelled action $\Phi : \mathbb{S}^1 \times X \rightarrow X$ is **exceptional** if the links of X have no perverse strata; i.e., if any perverse stratum of X is a closed (minimal) stratum.

Corollary 7.8. *For any exceptional action $\Phi : \mathbb{S}^1 \times X \rightarrow X$ we have*

$$(17) \quad H^*(\mathcal{L}\mathbf{ow}_{\bar{q}}(B)) = \prod_S H^*(S, \mathcal{I}\mathbf{m}_{\bar{q}}(\varepsilon_L))$$

where S runs over the perverse strata and $H^*(S, \mathcal{I}\mathbf{m}_{\bar{q}}(\varepsilon_L))$ is the cohomology of S with values on a locally constant presheaf with stalk

$$\ker\{\varepsilon_L : H_{\bar{q}-\bar{x}}^{\bar{q}(S)-1}(L/\mathbb{S}^1) \rightarrow H_{\bar{q}}^{\bar{q}(S)+1}(L/\mathbb{S}^1)\}$$

the kernel of the multiplication by the Euler class $\varepsilon_L \in H_{\bar{x}}^2(L/\mathbb{S}^1)$ of the link L of S .

Proof. If the link L of a perverse stratum S has no perverse strata, then the stalk $\mathcal{H}^i \mathcal{L}\mathbf{ow}_{\bar{q}}(-)$ vanishes for $i \neq \bar{q}(S) - 1$ (so it is a single presheaf). The equality (17) is straightforward, since the perverse strata are disjoint closed subsets. \square

ACKNOWLEDGMENTS

We would like to thank the accurate observations of M. Saralegi. While writing this article, the author received the financial support of the CDCH-Universidad Central de Venezuela and the hospitality of the staff in the Math Department-Université D’Artois.

REFERENCES

- [1] BOREL, A. & SPALTENSTEIN, N. "Sheaf theoretic intersection cohomology (Bern, 1983)" in *Intersection Cohomology-Swiss seminars*. (Bern, 1983). Progress in Mathematics Vol.50, 1-22. Birkhäuser. Boston (1984).
- [2] BREDON, G. *Introduction to Compact Transformation Groups*. Pure and Applied Mathematics Vol.46. Academic Press. New York (1972).
- [3] BOTT, R. & LU T. *Differential Forms in Algebraic Topology*. Graduate Texts in Mathematics Vol.82. Springer-Verlag. New York-Heidelberg- Berlin (1982).
- [4] BRYLINSKI, J.L. *Equivariant Intersection Cohomology*. *Contemporary Math.* **132** (1992), 5-32.
- [5] GODEMENT, R. *Topologie Algébrique et Thorie des Faisceaux*. Hermann. Paris (1958).
- [6] GORESKEY, M. & MACPHERSON, R. *Intersection Homology Theory*. *Topology* **19**, 135-162 (1980).
- [7] GORESKEY, M. & MACPHERSON, R. *Intersection Homology II*. *Invent. Math.* **71**, 77-129 (1983).
- [8] GREUB, W.; HALPERIN, S. & VANSTONE, R. *Conexions, curvature and cohomology*. Pure and Applied Mathematics Vol.47. Academic Press. New York (1972).
- [9] HECTOR, G. & SARALEGI, M. *Intersection Cohomology of \mathbb{S}^1 -actions*. *Trans. Amer. Math. Soc.* **338**, 263-288 (1983).
- [10] KING, H. *Intersection homology and homology of manifolds*. *Topology* **21** (1982), 229-234.
- [11] MAC PHEARSON, R. *Intersection Homology and Perverse Sheaves*. Colloquium Lectures, Annual Meeting of the Amer. Math. Soc. San Francisco (1991).
- [12] MASA, X.; MACIAS E. & ALVAREZ J. *Analysis and Geometry in Foliated Manifolds*. World Scientific. Santiago de Compostela (1994).
- [13] NAGASE, M. *L^2 -cohomology and intersection cohomology of stratified spaces*. *Duke Math. J.* **50**, 329-368 (1983).
- [14] PADILLA, G. *On normal stratified pseudomanifolds*. (2002) To appear in *Extracta Math.*

- [15] PADILLA, G. *Intersection cohomology of S^1 -actions on pseudomanifolds*. (2003) Submitted to *Indag. Math.*
- [16] PFLAUM, M. *Analytic and Geometric study of Stratified Spaces*. Lecture Notes in Mathematics Vol. **1768**. Springer. Berlin (2001).
- [17] ROYO, J. *The Euler Class for Riemannian Flows*, C. R. Acad. Sci. Paris, t.332, Serie I, pp. 45–50 (2001).
- [18] ROYO, J. *The Gysin Sequence for Riemannian Flows*, Contemporary Mathematics v.288, pp. 415–419 (2001).
- [19] SARALEGI, M. *Cohomologie d'Intersection des Actions Toriques Simples*. *Indag. Math.* **33**, 389–417 (1996).
- [20] SARALEGI, M. *A Gysin Sequence for Semifree Actions of S^3* . *Proc. Amer. Math. Soc.* **118**, 1335–1345 (1993).
- [21] SARALEGI, M. *Homological Properties of Stratified Spaces*. *Illinois J. Math.* **38**, 47–70 (1994).
- [22] SARALEGI, M. *The Euler class for flows of isometries*. *Research Notes in Math.* **131**, 25–28 (1989).
- [23] THOM, R. *Ensembles et morphismes stratifiés*. *Bull. Amer. Math. Soc.* **75**, 240–284 (1969).

UNIVERSIDAD CENTRAL DE VENEZUELA- ESCUELA DE MATEMÁTICA. CARACAS 1010.

E-mail address: gabrielp@euler.ciens.ucv.ve