

# ESTIMATES OF AUTOMORPHIC FUNCTIONS

JOSEPH BERNSTEIN AND ANDRE REZNIKOV

**ABSTRACT.** We present a new method to estimate trilinear period for automorphic representations of  $SL_2(\mathbb{R})$ . The method is based on the uniqueness principle in representation theory. We show how to separate the exponentially decaying factor in the triple period from the essential automorphic factor which behaves polynomially. We also describe a general method which gives an estimate on the average of the automorphic factor and thus prove a convexity bound for the triple period.

## 1. INTRODUCTION

**1.1. Maass forms.** Let  $Y$  be a compact Riemann surface with a Riemannian metric of constant curvature  $-1$  and the associated volume element  $dv$ . The corresponding Laplace-Beltrami operator is non-negative and has purely discrete spectrum on the space  $L^2(Y, dv)$  of functions on  $Y$ . We will denote by  $0 = \mu_0 < \mu_1 \leq \mu_2 \leq \dots$  its eigenvalues of and by  $\phi_i = \phi_{\mu_i}$  the corresponding eigenfunctions (normalized to have  $L^2$  norm one). In the theory of automorphic forms the functions  $\phi_{\mu_i}$  are called automorphic functions or **Maass forms** (after H. Maass).

The study of Maass forms plays an important role in analytic number theory.

We are interested in their analytic properties and will present a new method how to bound some important quantities arising from  $\phi_i$ .

**1.2. Triple products.** For any three Maass forms  $\phi_i, \phi_j, \phi_k$  we define the following triple product or triple period:

$$(1) \quad c_{ijk} = \int_Y \phi_i \phi_j \phi_k dv .$$

We would like to bound coefficients  $c_{ijk}$  as a function of eigenvalues  $\mu_i, \mu_j, \mu_k$ . In particular we would like to find bounds for these coefficients when one or more of these indexes tend to infinity.

**1.3. Motivation.** First of all we would like to explain why this problem is interesting. The explanation goes back to pioneering works of Rankin and Selberg (see [Ra], [Se]). They discovered that in special cases triple products as above give rise to automorphic  $L$ -functions. That allowed them to obtain analytic continuation and effective bounds for

these  $L$ -functions and as an application to obtain bounds on Fourier coefficients of cusp forms towards Ramanujan conjecture.

Since then the Rankin-Selberg method had many generalizations. Recently, for  $Y$  arising from the full modular group  $SL_2(\mathbb{Z})$  and for cuspidal functions  $\phi$ , Watson (see [Wa]) proved the following beautiful formula:

$$(2) \quad \left| \int_Y \phi_i \phi_j \phi_k dv \right|^2 = G(\lambda_i, \lambda_j, \lambda_k) \frac{L(1/2, \phi_i \otimes \phi_j \otimes \phi_k)}{L(1, \phi_i, Ad) L(1, \phi_j, Ad) L(1, \phi_k, Ad)}.$$

Here  $\lambda_t$  is a natural parameter of an eigenfunction  $\phi_t$  which is related to the eigenvalue by  $\mu_t = \frac{1-\lambda_t^2}{4}$ . Functions  $L(s, \phi_i \otimes \phi_j \otimes \phi_k)$ ,  $L(s, \phi_t, Ad)$  are appropriate automorphic  $L$ -functions associated with  $\phi_i$ , and the function  $G(\lambda_i, \lambda_j, \lambda_k)$  is some explicit rational expression in the ordinary  $\Gamma$ -functions. The relation (2) could be viewed as a far reaching generalization of the original Rankin-Selberg formula and was motivated by a work [HK] by Harris and Kudla on a conjecture of Jacquet.

**1.4. Results.** In this paper we will consider the following problem. We fix two Maass forms  $\phi = \phi_\tau$ ,  $\phi' = \phi_{\tau'}$  as above and consider coefficients defined by the triple period as above:

$$(3) \quad c_i = \int_Y \phi \phi' \phi_i dv$$

as  $\{\phi_i = \phi_{\lambda_i}\}$  run over the basis of Maass forms.

Thus we see from (2) that estimates of coefficients  $c_i$  are equivalent to estimates of the corresponding  $L$ -functions. One would like to have a general method to estimate the coefficients  $c_i$  and similar quantities. This problem was raised by Selberg in his celebrated paper [Se].

Let us understand what kind of bounds on the left hand side of (2) one would like to have in order to estimate effectively  $L$ -functions involved on the right hand side of (2) (or at least the ratio of  $L$ -functions).

We note first that one expects that  $c_i$  have exponential decay in  $|\lambda_i|$  as  $i$  goes to  $\infty$ . Namely, general experience from the analytic theory of automorphic  $L$ -functions tells us that  $L$ -functions have at most polynomial growth when  $|\lambda_i| \rightarrow \infty$ . Hence, analyzing the function  $G(\lambda)$ , one would expect from (2) and the Stirling formula for the asymptotics of  $\Gamma$ -function that the normalized coefficients

$$(4) \quad b_i = |c_i|^2 \exp\left(\frac{\pi}{2} |\lambda_i|\right)$$

have at most polynomial growth in  $|\lambda_i|$ , and hence  $c_i$  decay exponentially. However, it is difficult to see from the definition of the coefficients  $c_i$  that they have exponential decay and it is not clear what should be the rate of this decay.

The fact that an exponential decay with correct exponent holds for a general Riemann surface was first shown in papers by Good and Sarnak (see [Go] and [Sa1]). Both proofs used ingenious analytic continuation of automorphic functions in the *variable* parameter.

In this paper we will explain how to separate naturally the exponential decay from a polynomial growth in coefficients  $c_i$  using representation theory. We also prove the following

**Theorem.** *There exists an effectively computable constant  $A$  such that the following bound holds for arbitrary  $T > 0$*

$$(5) \quad \sum_{T \leq |\lambda_i| \leq 2T} b_i \leq A .$$

**1.5. A conjecture.** The estimate in the theorem is tight but when we try to use it to get a bound for an individual term  $b_i$  we get only an inequality

$$(6) \quad b_i \leq A .$$

According to Weyl's law there are approximately  $cT^2$  eigenvalues  $\mu_i$  with  $\lambda_i$  between  $T$  and  $2T$ , so the individual bound for the coefficient  $b_i$  is definitely not tight. We would like to make the following conjecture concerning the size of coefficients  $b_i$ :

**Conjecture.** *For any  $\varepsilon > 0$  there exists a constant  $C_\varepsilon > 0$  such that*

$$b_i \leq C_\varepsilon |\lambda_i|^{-2+\varepsilon} ,$$

*as  $|\lambda_i| \rightarrow \infty$ .*

For  $Y$  arising from congruence subgroups this conjecture is consistent with the Lindelöf conjecture for appropriate automorphic  $L$ -functions (see [BR1], [Sa2] and [Wa] for more details). We note that the bound in the Theorem above corresponds to the so-called **convexity bound**.

**1.6. The method.** The first proof of the (slightly weaker) version of Theorem 1.4 appeared in [BR1]. It was based on the analytic continuation of representations from a real group to a complex group (generalizing methods of [Sa1]). The method based on the analytic continuation was extended in [KS] to the case of higher rank groups. While it gives bounds which are tight for general representations it was not able, so far, to cover cases relevant to  $L$ -functions.

The proof we present here is based on the uniqueness of triple product in representation theory. It has an advantage that it could be generalized to higher rank groups and gives bounds which are consistent with the theory of  $L$ -functions. The present method also could be applied to  $p$ -adic groups (unlike methods of [BR1]).

We describe now the general ideas behind our new proof. It is based on ideas from representation theory. Namely, we use the fact that every automorphic form  $\phi$  generates

an automorphic representation of the group  $G = PGL_2(\mathbb{R})$ ; this means that starting from  $\phi$  we produce a smooth irreducible representation of the group  $G$  in a space  $V$  and its realization  $\nu : V \rightarrow C^\infty(X)$  in the space of smooth functions on the automorphic space  $X = \Gamma \backslash G$ .

The triple product  $c_i = \int_Y \phi \phi' \phi_i dv$  extends to a  $G$ -equivariant trilinear form on the corresponding automorphic representations  $l^{aut} : V \otimes V' \otimes V_i \rightarrow \mathbb{C}$ , where  $V = V_\tau, V' = V_{\tau'}, V_i = V_{\lambda_i}$ .

Then we use a general result from representation theory that such  $G$ -equivariant trilinear form is unique up to a scalar. This implies that the automorphic form  $l^{aut}$  is proportional to an explicit "model" form  $l^{mod}$  which we describe using explicit realizations of representations of the group  $G$ ; it is important that this last form carries no arithmetic information.

Thus we can write  $l^{aut} = a_i \cdot l^{mod}$  for some constant  $a_i$  and hence  $c_i = l^{aut}(e_\tau \otimes e_{\tau'} \otimes e_{\lambda_i}) = a_i \cdot l^{mod}(e_\tau \otimes e_{\tau'} \otimes e_{\lambda_i})$ , where  $e_\tau, e_{\tau'}, e_{\lambda_i}$  are  $K$ -invariant unit vectors in the automorphic representations  $V, V', V_i$  corresponding to the automorphic forms  $\phi, \phi'$  and  $\phi_i$ .

It turns out that in the last formula the proportionality coefficient  $a_i$  carries an important "automorphic" information while the second factor carries no arithmetic information and can be computed in terms of  $\Gamma$ -functions using explicit realizations of representations  $V_\tau, V_{\tau'}$  and  $V_{\lambda_i}$ . This second factor is responsible for the exponential decay, while the first factor  $a_i$  has a polynomial behavior in parameter  $\lambda_i$ .

In order to bound the quantities  $a_i$  we use the fact that they appear as coefficients in the spectral decomposition of the diagonal Hermitian form  $H_\Delta$  on the space  $E = V_\tau \otimes V_{\tau'}$  (see 4.2, 4.3). This gives us an inequality  $\sum |a_i|^2 H_i \leq H_\Delta$ , where  $H_i$  is an Hermitian form on  $E$  induced by the model trilinear form  $l^{mod} : V \otimes V' \otimes V_i \rightarrow \mathbb{C}$  above.

Using the geometric properties of the diagonal form and simple explicit estimates of forms  $H_i$  we establish the convexity bound for the coefficients  $a_i$ .

It is known that the uniqueness principle plays central role in the theory of automorphic functions (see [PS]). The impact that the uniqueness has on the analytic behavior of automorphic functions is yet another manifestation of this principle.

**Acknowledgments.** We would like to thank Peter Sarnak for many fruitful discussions. We also would like to thank Yuri Neretin for discussions on trilinear functionals.

Research was partially supported by EC TMR network "Algebraic Lie Representations", grant no. ERB FMRX-CT97-0100, by BSF grant, Minerva Foundation and by the Excellency Center "Group Theoretic Methods in the Study of Algebraic Varieties" of the Israel Science Foundation, the Emmy Noether Institute for Mathematics (the Center of Minerva Foundation of Germany).

The paper was written during our stay at IHES and is an extended version of the talk given by the first author at Cartier Colloquium in June 2002.

## 2. REPRESENTATION THEORETIC SETTING

We recall the standard connection of the above setting with the representation theory (see [G6]).

**2.1. Automorphic functions and automorphic representations.** Let us describe the geometric construction which allows one to pass from analysis on a Riemann surface to representation theory.

Let  $\mathbb{H}$  be the upper half plane with the hyperbolic metric of constant curvature  $-1$ . The group  $SL_2(\mathbb{R})$  acts on  $\mathbb{H}$  by fractional linear transformations. This action allows to identify the group  $PSL_2(\mathbb{R})$  with the group of all orientation preserving motions of  $\mathbb{H}$ . For reasons explained below we would like to work with the group  $G$  of all motions of  $\mathbb{H}$ ; this group is isomorphic to  $PGL_2(\mathbb{R})$ . Hence throughout the paper we denote  $G = PGL_2(\mathbb{R})$ .

Let us fix a discrete co-compact subgroup  $\Gamma \subset G$  and set  $Y = \Gamma \backslash \mathbb{H}$ . We consider the Laplace operator on the Riemann surface  $Y$  and denote by  $\mu_i$  its eigenvalues and by  $\phi_i$  the corresponding normalized eigenfunctions.

The case when  $\Gamma$  acts freely on  $\mathbb{H}$  precisely corresponds to the case discussed in the introduction (this follows from the uniformization theorem for the Riemann surface  $Y$ ). Our results hold for general co-compact subgroup  $\Gamma$  (and in fact, with slight modifications, for any lattice  $\Gamma \subset G$ ).

We will identify the upper half plane  $\mathbb{H}$  with  $G/K$ , where  $K = PO(2)$  is a maximal compact subgroup of  $G$  (this follows from the fact that  $G$  acts transitively on  $\mathbb{H}$  and the stabilizer in  $G$  of the point  $z_0 = i \in \mathbb{H}$  coincides with  $K$ ).

We denote by  $X$  the compact quotient  $\Gamma \backslash G$  (we call it the automorphic space). In the case when  $\Gamma$  acts freely on  $\mathbb{H}$  one can identify the space  $X$  with the bundle of unit tangent vectors to the Riemann surface  $Y = \Gamma \backslash \mathbb{H}$ .

The group  $G$  acts on  $X$  (from the right) and hence on the space of functions on  $X$ . We fix the unique  $G$ -invariant measure  $\mu_X$  on  $X$  of total mass one. Let  $L^2(X) = L^2(X, d\mu_X)$  be the space of square integrable functions and  $(\Pi_X, G, L^2(X))$  the corresponding unitary representation. We will denote by  $P_X$  the Hermitian form on  $L^2(X)$  given by the scalar product. We denote by  $\|\cdot\|_X$  or simply  $\|\cdot\|$  the corresponding norm and by  $\langle f, g \rangle_X$  the corresponding scalar product.

The identification  $Y = \Gamma \backslash \mathbb{H} \simeq X/K$  induces the embedding  $L^2(Y) \subset L^2(X)$ . We will always identify the space  $L^2(Y)$  with the subspace of  $K$ -invariant functions in  $L^2(X)$ .

Let  $\phi$  be a normalized eigenfunction of the Laplace-Beltrami operator on  $Y$ . Consider a closed  $G$ -invariant subspace  $L_\phi \subset L^2(X)$  generated by  $\phi$  under the action of  $G$ . It is well-known that  $(\pi, L) = (\pi_\phi, L_\phi)$  is an irreducible unitary representation of  $G$  (see [G6]).

Usually it is more convenient to work with the space  $V = L^\infty$  of smooth vectors in  $L$ . The unitary Hermitian form  $P$  on  $V$  is  $G$ -invariant.

A representation  $(\pi, G, V)$  equipped with a positive  $G$ -invariant Hermitian form  $P$  we will call a **smooth pre-unitary representation**; this simply means that  $V$  is the space of smooth vectors in the unitary representation obtained from  $V$  by completion with respect to  $P$ .

Thus starting with an automorphic function  $\phi$  we constructed an irreducible smooth pre-unitary representation  $(\pi, V)$ . In fact we constructed this space together with a canonical morphism  $\nu : V \rightarrow C^\infty(X)$  since  $C^\infty(X)$  is the smooth part of  $L^2(X)$ .

**Definition.** A smooth pre-unitary representation  $(\pi, G, V)$  equipped with a  $G$ -morphism  $\nu : V \rightarrow C^\infty(X)$  we will call an  **$X$ -enhanced representation**.

In this note we will assume that the morphism  $\nu$  is normalized, i.e. it carries the standard  $L^2$  Hermitian form  $P_X$  on  $C^\infty(X)$  into Hermitian form  $P$  on  $V$ .

Thus starting with an automorphic function  $\phi$  we constructed

- (i) An  $X$ -enhanced irreducible pre-unitary representation  $(\pi, V, \nu)$ ,
- (ii) A  $K$ -invariant unit vector  $e_V \in V$  (this vector is just our function  $\phi$ ).

Conversely, suppose we are given an irreducible smooth pre-unitary  $X$ -enhanced representation  $(\pi, V, \nu)$  of the group  $G$  and a  $K$ -fixed unit vector  $e_V \in V$ . Then the function  $\phi = \nu(e_V) \in C^\infty(X)$  is  $K$ -invariant and hence can be considered as a function on  $Y$ . The fact that the representation  $(\pi, V)$  is irreducible implies that  $\phi$  is an automorphic function.

Thus we have established a natural correspondence between Maass forms  $\phi$  and tuples  $(\pi, V, \nu, e_V)$ , where  $(\pi, V, \nu)$  is an  $X$ -enhanced irreducible smooth pre-unitary representation and  $e_V \in V$  is a unit  $K$ -invariant vector.

**2.2. Decomposition of the representation  $(\Pi_X, G, L^2(X))$ .** It is well known that in case when  $X$  is compact the representation  $(\Pi_X, G, L^2(X))$  decomposes into a direct (infinite) sum

$$(7) \quad L^2(X) = \oplus_j (\pi_j, L_j)$$

of irreducible unitary representations of  $G$  (all representations appear with finite multiplicities (see [G6])). Let  $(\pi, L)$  be one of these irreducible "automorphic" representations and  $V = L^\infty$  its smooth part. By definition  $V$  is given with a  $G$ -equivariant isometric morphism  $\nu : V \rightarrow C^\infty(X)$ , i.e.  $V$  is an  $X$ -enhanced representation.

If  $V$  has a  $K$ -invariant vector it corresponds to a Maass form. There are other spaces in this decomposition which correspond to discrete series representations. Since they are not related to Maass forms we will not study them in more detail.

**2.3. Representations of  $PGL_2(\mathbb{R})$ .** All irreducible unitary representations of  $G$  are classified. For simplicity we consider those with a nonzero  $K$ -fixed vector (so called representations of class one) since only these representations arise from Maass forms.

These are the representations of the principal and the complementary series and the trivial representation.

We will use the following standard explicit model for irreducible smooth representations of  $G$ .

For every complex number  $\lambda$  consider the space  $V_\lambda$  of smooth even homogeneous functions on  $\mathbb{R}^2 \setminus 0$  of homogeneous degree  $\lambda - 1$  (which means that  $f(ax, ay) = |a|^{\lambda-1} f(x, y)$  for all  $a \in \mathbb{R} \setminus 0$ ). The representation  $(\pi_\lambda, V_\lambda)$  is induced by the action of the group  $GL_2(\mathbb{R})$  given by  $\pi_\lambda(g)f(x, y) = f(g^{-1}(x, y))|\det g|^{(\lambda-1)/2}$ . This action is trivial on the center of  $GL_2(\mathbb{R})$  and hence defines a representation of  $G$ . The representation  $(\pi_\lambda, V_\lambda)$  is called **representation of the generalized principal series**.

When  $\lambda = it$  is purely imaginary the representation  $(\pi_\lambda, V_\lambda)$  is pre-unitary; the  $G$ -invariant scalar product in  $V_\lambda$  is given by  $\langle f, g \rangle_{\pi_\lambda} = \frac{1}{2\pi} \int_{S^1} f \bar{g} d\theta$ . These representations are called representations of **the principal series**.

When  $\lambda \in (-1, 1)$  the representation  $(\pi_\lambda, V_\lambda)$  is called a representation of the complementary series. These representations are also pre-unitary, but the formula for the scalar product is more complicated (see [G6]).

All these representations have  $K$ -invariant vectors. We fix a  $K$ -invariant unit vector  $e_\lambda \in V_\lambda$  to be a function which is one on the unit circle in  $\mathbb{R}^2$ .

Representations of the principal and the complimentary series exhaust all nontrivial irreducible pre-unitary representations of  $G$  of class one.

In what follows we will do necessary computations for representation of the principal series. Computations for the complementary series are a little more involved but essentially the same (compare with [BR1], section 5.5, where similar computations are described in detail).

Suppose we are given a class one  $X$ -enhanced representation  $\nu : V_\lambda \rightarrow C^\infty(X)$ ; we assume  $\nu$  to be an isometric embedding. Such  $\nu$  gives rise to an eigenfunction of the Laplacian on the Riemann surface  $Y = X/K$  as before. Namely, if  $e_\lambda \in V_\lambda$  is a unit  $K$ -fixed vector then the function  $\phi = \nu(e_\lambda)$  is a normalized eigenfunction of the Laplacian on the space  $Y = X/K$  with the eigenvalue  $\mu = \frac{1-\lambda^2}{4}$ . This explains why  $\lambda$  is a natural parameter to describe Maass forms.

**2.4. Triple products.** We introduce now our main tool.

**2.4.1. Automorphic triple products.** Suppose we are given three  $X$ -enhanced representations of  $G$

$$\nu_j : V_j \rightarrow C^\infty(X), \quad j = 1, 2, 3.$$

We define the  $G$ -invariant trilinear form  $l_{\pi_1, \pi_2, \pi_3}^{aut} : V_1 \otimes V_2 \otimes V_3 \rightarrow \mathbb{C}$ , by formula

$$(8) \quad l_{\pi_1, \pi_2, \pi_3}^{aut}(v_1 \otimes v_2 \otimes v_3) = \int_X \phi_{v_1}(x) \phi_{v_2}(x) \phi_{v_3}(x) d\mu_X ,$$

where  $\phi_{v_j} = \nu_j(v_j) \in C^\infty(X)$  for  $v_j \in V_j$ .

In particular, the triple periods  $c_i$  in (3) can be expressed in terms of this form as  $c_i = l_{\pi, \pi', \pi_i}^{aut}(e_\tau \otimes e_{\tau'} \otimes e_{\lambda_i})$ , where  $e_\lambda \in V_\lambda$  is the  $K$ -fixed unit vector.

**2.4.2. Uniqueness of triple products.** The central fact about invariant trilinear functionals is the following uniqueness result:

**Theorem.** *Let  $(\pi_j, V_j)$ ,  $j = 1, 2, 3$ , be three irreducible smooth admissible representations of  $G$ . Then  $\dim \text{Hom}_G(V_1 \otimes V_2 \otimes V_3, \mathbb{C}) \leq 1$ .*

*Remark.* The uniqueness statement was proven by Oksak in [O] for the group  $SL(2, \mathbb{C})$  and the proof could be adopted for  $PGL_2(\mathbb{R})$  as well (see also [Mo] and [Lo]). For the  $p$ -adic  $GL(2)$  more refined results were obtained by Prasad (see [P]). He also proved the uniqueness when at least one representation is a discrete series representation of  $GL_2(\mathbb{R})$ .

There is no uniqueness of trilinear functionals for representations of  $SL_2(\mathbb{R})$  (the space is two-dimensional). This is the reason why we prefer to work with  $PGL_2(\mathbb{R})$ .

For  $SL_2(\mathbb{R})$  one has the following uniqueness statement instead. Let  $(\pi, V)$  and  $(\sigma, W)$  be two irreducible smooth pre-unitary representations of  $SL_2(\mathbb{R})$  of class one. Then the space of  $SL_2(\mathbb{R})$ -invariant trilinear functionals on  $V \otimes V \otimes W$  which are symmetric in the first two variables is one-dimensional. This is the correct uniqueness result needed if one wants to work with  $SL_2(\mathbb{R})$ ; this was implicitly done in [Re2], where the second author missed the absence of the uniqueness for  $SL_2(\mathbb{R})$ . We take an opportunity to correct this gap.

We note however, that the absence of uniqueness does not pose any problem for the method we present. All what is really needed for our method is the fact that the space of invariant functionals is finite dimensional.

### 3. TRIPLE PRODUCTS: EXPONENTIAL DECAY

We now explain our method how to bound coefficients  $c_i$ . It is based on the uniqueness of trilinear functionals.

**3.1. Model triple products.** Let  $(\pi, V)$  and  $(\pi', V')$  be automorphic representations corresponding to Maass forms  $\phi$  and  $\phi'$ . Any Maass form  $\phi_i$  gives us an automorphic representation  $(\pi_i, V_{\lambda_i})$  and hence defines a trilinear functional

$$l_{\pi, \pi', \pi_i}^{aut} : V \otimes V' \otimes V_{\lambda_i} \rightarrow \mathbb{C} .$$



In 5.1 we use an explicit model for representations  $\pi_1, \pi_2, \pi_3$  to construct a model invariant trilinear functional which is given by an explicit formula. We call it the **model triple product** and denote by  $l_{\pi_1, \pi_2, \pi_3}^{mod}$ .

By the uniqueness principle for representations  $\pi, \pi', \pi_i$  there exists a constant  $a_i = a_{\pi, \pi', \pi_i}$  such that:

$$(9) \quad l_{\pi, \pi', \pi_i}^{aut} = a_i \cdot l_{\pi, \pi', \pi_i}^{mod}.$$

**3.2. Exponential decay.** This gives a formula for the triple products  $c_i$

$$(10) \quad c_i = l_{\lambda_i}^{aut}(e_\tau \otimes e_{\tau'} \otimes e_{\lambda_i}) = a_i \cdot l_{\lambda_i}^{mod}(e_\tau \otimes e_{\tau'} \otimes e_{\lambda_i}).$$

Here we denoted  $l_{\lambda_i}^{aut} = l_{\pi, \pi', \pi_i}^{aut}$ ,  $l_{\lambda_i}^{mod} = l_{\pi, \pi', \pi_i}^{mod}$  and  $e_\lambda$  is the unit  $K$ -fixed vector in the representation  $V_\lambda$ .

The model triple product  $l_{\lambda_i}^{mod}(e_\tau \otimes e_{\tau'} \otimes e_{\lambda_i})$  constructed in 5.1 is given by an explicit integral. In Appendix A we evaluate this integral by a direct computation in the model. It turns out that it has an exponential decay in  $|\lambda|$  which explains the exponential decay of coefficients  $c_i$ . Namely, we prove the following

**Proposition.** *Set  $k(\lambda) := |l_{\lambda}^{mod}(e_\tau \otimes e_{\tau'} \otimes e_\lambda)|^2$ . Then there exists a constant  $c > 0$  such that*

$$k_\lambda = c \exp(-\frac{\pi}{2}|\lambda|) \cdot |\lambda|^{-2}(1 + O(|\lambda|^{-1}))$$

as  $|\lambda| \rightarrow \infty$ .

#### 4. TRIPLE PRODUCTS: POLYNOMIAL BOUNDS

We explain now how to obtain bounds on the coefficients  $a_i$  (note that these coefficients encode deep arithmetic information - values of  $L$ -functions).

Our method is based on the fact that these coefficients appear in the spectral decomposition of some geometrically defined Hermitian form on the space  $E$  which is essentially the tensor product of spaces  $V$  and  $V'$ .

More precisely, denote by  $L$  and  $L'$  the Hilbert completions of spaces  $V$  and  $V'$ , consider the unitary representation  $(\Pi, G \times G, L \otimes L')$  of the group  $G \times G$  and denote by  $E$  its smooth part; so  $E$  is a smooth completion of  $V \otimes V'$ .

Denote by  $\mathcal{H}(E)$  the (real) vector space of continuous Hermitian forms on  $E$  and by  $\mathcal{H}^+(E)$  the cone of nonnegative Hermitian forms.

We will describe several classes of Hermitian forms on  $E$ ; some of them have spectral description, others are described geometrically.

**4.1. Hermitian forms corresponding to trilinear functionals.** Let  $W$  be a smooth pre-unitary admissible representation of  $G$ . Any  $G$ -invariant functional  $l : V \otimes V' \otimes W \rightarrow \mathbb{C}$  defines a  $G$ -intertwining morphism  $T^l : V \otimes V' \rightarrow W^*$  which extends to a  $G$ -morphism

$$(11) \quad T^l : E \rightarrow \bar{W}.$$

where we identify the complex conjugate space  $\bar{W}$  with the smooth part of the space  $W^*$ .

The standard Hermitian form (scalar product)  $P_W$  on the space  $W$  induces the Hermitian form  $\bar{P}$  on  $\bar{W}$ . Using the operator  $T^l$  we define the Hermitian form  $H^l$  on the space  $E$  by  $H^l = (T^l)^*(\bar{P})$ , i.e.  $H^l(u) = \bar{P}(T^l(u))$  for  $u \in E$ .

We note that if the representation of  $G$  in the space  $W$  is irreducible then starting with the Hermitian form  $H^l$  we can reconstruct the space  $W$ , the functional  $l$  and the morphism  $T^l$  uniquely up to an isomorphism.

Let us introduce a special notation for the particular case we are interested in. For any number  $\lambda \in i\mathbb{R}$  consider the representation of the principal series  $W = V_\lambda$ , choose the model trilinear functional  $l^{mod} : V \otimes V' \otimes V_\lambda \rightarrow \mathbb{C}$  described in 5.1 and denote the corresponding Hermitian form on  $E$  by  $H_\lambda^{mod}$ .

**4.2. Diagonal form  $H_\Delta$ .** Consider the space  $C^\infty(X \times X)$ . The diagonal  $\Delta : X \rightarrow X \times X$  gives rise to the restriction morphism  $r_\Delta : C^\infty(X \times X) \rightarrow C^\infty(X)$ . We define a nonnegative Hermitian form  $H_\Delta$  on  $C^\infty(X \times X)$  by  $H_\Delta = (r_\Delta)^*(P_X)$ , i.e.

$$H_\Delta(u) = P_X(r_\Delta(u)) = \int_X |r_\Delta(u)|^2 d\mu_X \text{ for } u \in C^\infty(X \times X).$$

We call  $H_\Delta$  the diagonal form.

More generally, if  $L$  is a closed subspace of  $L^2(X)$  and  $pr_L : L^2(X) \rightarrow L$  the orthogonal projection onto  $L$  we can define a Hermitian form  $P_L$  on  $C^\infty(X)$  by  $P_L = (pr_L)^*(P_X)$  and consider the induced Hermitian form  $H_L = (r_\Delta)^*(P_L)$  on  $C^\infty(X \times X)$ .

Clearly the correspondence  $L \mapsto H_L$  is additive (which means that  $H_{L+L'} = H_L + H_{L'}$  if  $L$  and  $L'$  are orthogonal) and monotone.

**4.3. First basic inequality.** Let us realize the space  $E = V \otimes V'$  as a  $G \times G$ -invariant subspace of  $C^\infty(X \times X)$ . We consider the restrictions of the Hermitian forms  $H_\Delta, H_L$  discussed above to the space  $E$  and will denote them by the same symbols.

**Claim.** *Let  $\phi_{\lambda_i}$  be a Maass form. Consider the  $G$ -invariant subspace  $L_i \subset L^2(X)$  generated by  $\phi_{\lambda_i}$  and its complex conjugate  $\bar{L}_i \subset L^2(X)$ .*

*Then on the space  $E$  the Hermitian form  $H_{\bar{L}_i}$  coincides with the form  $H_{\lambda_i}^{aut}$  corresponding to the automorphic trilinear form  $l = l_{\pi, \pi', \pi_i}^{aut} : V \otimes V' \otimes V_{\lambda_i} \rightarrow \mathbb{C}$ .*

Indeed, if we identify the space  $\bar{L}_i$  with  $L_i^*$  then the operator  $pr_{\bar{L}_i} \circ r_\Delta : E \rightarrow \bar{L}_i$  coincides with the operator  $T^l$  corresponding to the automorphic trilinear form  $l = l_{\pi, \pi', \pi_i}^{aut}$ .

This claim implies the first basic inequality

$$(12) \quad \sum_{\lambda_i} |a_i|^2 H_{\lambda_i}^{mod} \leq H_{\Delta} .$$

Indeed, by the uniqueness principle (9) we have:

$$(13) \quad H_{\lambda_i}^{aut} = |a_i|^2 \cdot H_{\lambda_i}^{mod} ,$$

where  $a_i = a_{\pi, \pi', \pi_i}$  are as in (9).

Since all the spaces  $\bar{L}_i$  are orthogonal we have  $\sum_i H_{\lambda_i}^{aut} \leq H_{\Delta}$  which proves the first basic inequality.

**4.4. Second basic inequality.** We would like to use the inequality (12) to bound the coefficients  $a_i$ . In order to do this we have to establish some bounds for the diagonal form  $H_{\Delta}$ .

The group  $G \times G$  naturally acts on the space of Hermitian forms on  $C^\infty(X \times X)$  - we denote this action by  $\Pi$ . We extend this action to the action of the algebra  $H(G) = C_c^\infty(G \times G, \mathbb{R})$  of smooth real valued functions with compact support. Note that if  $h \in H(G)$  is a nonnegative function then the operator  $\Pi(h)$  preserves the cone of positive forms.

We have then the second basic inequality

**Claim.** *Let  $h \in H(G)$  be a non-negative function. Then there exists a constant  $C$ , depending on  $h$ , such that we have  $\Pi(h)H_{\Delta} \leq C \cdot P_{X \times X}$ , where  $P_{X \times X}$  is the standard  $L^2$  Hermitian form on the space  $C^\infty(X \times X)$ .*

*Proof.* Let  $u \in C^\infty(X \times X)$  and  $f = |u|^2$ . Then  $P_{X \times X}(u) = \langle \mu, f \rangle$  and  $\Pi(h)H_{\Delta}(u) = \langle \mu', f \rangle$ , where  $\mu = \mu_{X \times X}$  and  $\mu' = \Pi(h)(\Delta_*(\mu_X))$  are two measures on  $X \times X$ .

Since the measure  $\mu'$  is smooth it is bounded by  $C\mu$ . □

Note that the bound in the claim is essentially tight. Namely if the function  $h$  has large enough support then we also have a bound in the opposite direction.

**4.5. Positive functionals.** We can now prove that the coefficients  $a_i$  have at most polynomial growth in  $|\lambda_i|$ .

We start with the inequality (12) of *non-negative forms*. We want to produce out of it an inequality for coefficients  $a_i$ . There is a standard way to do this by means of positive functionals on the space of Hermitian forms  $\mathcal{H}(E)$ .

**Definition.** A **positive functional** on the space  $\mathcal{H}(E)$  is an additive map  $\rho : \mathcal{H}(E)^+ \rightarrow \mathbb{R}^+ \cup \infty$ .

It is easy to see that the positive functional  $\rho$  is automatically monotone and homogeneous (i.e.  $\rho(H) \leq \rho(H')$  if  $H \leq H'$  and  $\rho(aH) = a\rho(H)$  for  $a > 0$ ).

**Example.** Any vector  $u \in E$  gives us an elementary positive functional  $\rho_u$  defined by  $\rho_u(H) = H(u)$ .

Fix a positive functional  $\rho$  and consider the weight function  $h(\lambda) = \rho(H_\lambda^{mod})$ . Then from the first basic inequality (12) we can deduce the following inequality for a weighted sum of coefficients  $|a_i|^2$ :

$$\sum_i h(\lambda_i) |a_i|^2 \leq \rho(H_\Delta) .$$

**4.6. Test functional  $\rho_T$ .** For any real  $T$  we construct in 5.2 the positive "test" functional  $\rho_T$  on  $\mathcal{H}(E)$  with the properties described in the proposition below. Let us fix automorphic representations  $V, V' \subset C^\infty(X)$ ,  $E = V \otimes V' \subset C^\infty(X \times X)$  as above.

**Proposition.** *We can find a constant  $C$  which depends only on  $G$  and  $\Gamma$  and a constant  $T_0$  which depends on  $V$  and  $V'$  such that for any  $T \geq T_0$  there exists a positive functional  $\rho_T$  on  $\mathcal{H}(E)$  satisfying*

$$(14) \quad \rho_T(H_\Delta) \leq CT^2 ,$$

$$(15) \quad h_T(\lambda) := \rho_T(H_\lambda^{mod}) \geq 1 \text{ for any } |\lambda| \leq 2T .$$

**4.7. Proof of Theorem 1.4.** Consider the inequality  $\sum_i |a_i|^2 \rho_T(H_{\lambda_i}^{mod}) \leq \rho_T(H_\Delta)$ .

The right hand side  $\rho_T(H_\Delta)$  is bounded by  $CT^2$ . In the left hand side we can leave only terms with  $|\lambda_i| \leq 2T$ . Thus we arrive at inequality

$$(16) \quad \sum_{|\lambda_i| \leq 2T} |a_i|^2 \leq CT^2 .$$

This gives the desired bound for  $\sum |a_i|^2$ .

According to Proposition 3.2 there exists a constant  $b$  such that  $b_i T^2 \leq b |a_i|^2$  for  $T \leq |\lambda_i| \leq 2T$ . This shows that  $\sum_{T < |\lambda_i| < 2T} b_i \leq A$  for some constant  $A$ , which finishes the proof of Theorem 1.4.

**4.7.1. A conjecture.** One can show (see [Re1]) that the mean-value result in (16) is essentially sharp. One expects that for  $T \leq |\lambda_i| \leq 2T$  all terms in the sum (16) are at most of order  $T^\varepsilon$  for any  $\varepsilon > 0$ . Hence, we have established a sharp bound on the average and a rather weak bound for each term. This is a typical situation which one often encounters in the analytic theory of  $L$ -functions, the so-called convexity bound. The major problem

hence is to find a method which would allow us to obtain better bound for a single term or for a short interval – the so-called subconvexity bounds.

We would like to make the following conjecture concerning the size of coefficients  $a_{\pi, \pi', \pi_i}$  which is equivalent to Conjecture 1.5:

**Conjecture.** *For fixed  $\pi, \pi'$  and for any  $\varepsilon > 0$  there exists  $C_\varepsilon > 0$  independent of  $\lambda_i$  such that*

$$|a_{\pi, \pi', \pi_{\lambda_i}}| \leq C_\varepsilon |\lambda_i|^\varepsilon ,$$

as  $|\lambda_i| \rightarrow \infty$ .

## 5. CONSTRUCTION OF MODEL TRILINEAR FUNCTIONALS AND OF TEST FUNCTIONALS

**5.1. Model trilinear functionals.** For every  $\lambda \in \mathbb{C}$  we denote by  $(\pi_\lambda, V_\lambda)$  the smooth class one representation of the generalized principle series of the group  $G = PGL_2(\mathbb{R})$  described in 2.3. We will use the realization of  $(\pi_\lambda, V_\lambda)$  in the space of smooth homogeneous functions on  $\mathbb{R}^2 \setminus 0$  of homogeneous degree  $\lambda - 1$ .

For explicit computations it is often convenient to pass from plane model to a circle model. Namely, the restriction of functions in  $V_\lambda$  to the unit circle  $S^1 \subset \mathbb{R}^2$  defines an isomorphism of the space  $V_\lambda$  with the space  $C^\infty(S^1)^{even}$  of even smooth functions on  $S^1$  so we can think about vectors in  $V_\lambda$  as functions on  $S^1$ .

In this section we describe the **model** invariant trilinear functional using the geometric models. Namely for given three complex numbers  $\lambda_j, j = 1, 2, 3$ , we construct explicitly nontrivial trilinear functional  $l^{mod} : V_{\lambda_1} \otimes V_{\lambda_2} \otimes V_{\lambda_3} \rightarrow \mathbb{C}$  by means of its kernel.

**5.1.1. Kernel of  $l^{mod}$ .** Let  $\omega(\xi, \eta) = \xi_1\eta_2 - \xi_2\eta_1$  be  $SL_2(\mathbb{R})$ -invariant of a pair of vectors  $\xi, \eta \in \mathbb{R}^2$ . We set

$$(17) \quad K_{\lambda_1, \lambda_2, \lambda_3}(s_1, s_2, s_3) = |\omega(s_2, s_3)|^{(\alpha-1)/2} |\omega(s_1, s_3)|^{(\beta-1)/2} |\omega(s_1, s_2)|^{(\gamma-1)/2}$$

for  $s_1, s_2, s_3 \in \mathbb{R}^2 \setminus 0$ , where  $\alpha = \lambda_1 - \lambda_2 - \lambda_3, \beta = -\lambda_1 + \lambda_2 - \lambda_3, \gamma = -\lambda_1 - \lambda_2 + \lambda_3$ .

The kernel function  $K_{\lambda_1, \lambda_2, \lambda_3}(s_1, s_2, s_3)$  satisfies two main properties:

- (1)  $K$  is invariant with respect to the diagonal action of  $SL_2(\mathbb{R})$ .
- (2)  $K$  is homogeneous of degree  $-1 - \lambda_j$  in each variable  $s_j$ .

Hence if  $f_j$  are homogeneous functions of degree  $-1 + \lambda_j$ , then the function

$$F(s_1, s_2, s_3) = f_1(s_1)f_2(s_2)f_3(s_3)K_{\lambda_1, \lambda_2, \lambda_3}(s_1, s_2, s_3) ,$$

is homogeneous of degree  $-2$  in each variable  $s_j \in \mathbb{R}^2 \setminus 0$ .

5.1.2. *Functional  $l^{mod}$ .* To define the model trilinear functional  $l^{mod}$  we notice that on the space  $\mathcal{V}$  of functions of homogeneous degree  $-2$  on  $\mathbb{R}^2 \setminus 0$  there exists a natural  $SL_2(\mathbb{R})$ -invariant functional  $\mathfrak{L} : \mathcal{V} \rightarrow \mathbb{C}$ . It is given by the formula  $\mathfrak{L}(f) = \int_{\Sigma} f d\sigma$  where the integral is taken over any closed curve  $\Sigma \subset \mathbb{R}^2 \setminus 0$  which goes around 0 and the measure  $d\sigma$  on  $\Sigma$  is given by the area element inside of  $\Sigma$  divided by  $\pi$ ; his last normalization factor is chosen so that  $\mathfrak{L}(Q^{-1}) = 1$  for the standard quadratic form  $Q$  on  $\mathbb{R}^2$ .

Applying  $\mathfrak{L}$  separately to each variable  $s_i \in \mathbb{R}^2 \setminus 0$  of the function  $F(s_1, s_2, s_3)$  above we obtain the  $G$ -invariant functional

$$(18) \quad l_{\pi_1, \pi_2, \pi_3}^{mod}(f_1 \otimes f_2 \otimes f_3) := \langle \mathfrak{L} \otimes \mathfrak{L} \otimes \mathfrak{L}, F \rangle .$$

In the circle model this functional is expressed by the following integral:

$$(19) \quad l_{\pi_1, \pi_2, \pi_3}^{mod}(f_1 \otimes f_2 \otimes f_3) = (2\pi)^{-3} \iiint f_1(x) f_2(y) f_3(z) K_{\lambda_1, \lambda_2, \lambda_3}(x, y, z) dx dy dz,$$

where  $x, y, z$  are the standard angular parameters on the circle.

*Remark.* The integral defining the trilinear functional is often divergent and the functional should be defined using regularization of this integral. There are standard procedures how to make such a regularization (see e.g. [G1]).

Fortunately in the case of unitary representations all integrals converge absolutely so we will not discuss the regularization procedure.

**5.2. Construction of test functionals.** In this section we will present a construction of a family of test functionals  $\rho_T$  on the space  $\mathcal{H}(E)$ .

Fix smooth irreducible pre-unitary representations of class one  $V = V_{\tau}, V = V_{\tau'}$  and denote by  $E$  the smooth completion of  $V \otimes V'$  as in section 4. We will do the computations only for representations of the principal series; complementary series are treated similarly.

For computations we will identify the spaces  $V$  and  $V'$  with  $C^{\infty}(S^1)^{even}$ .

Our aim is to prove the following

**Proposition.** *There exist constants  $T_0, C, c > 0$  such that for any  $T \geq T_0$  there exists a positive functional  $\rho$  on  $\mathcal{H}(E)$  satisfying*

$$(20) \quad \rho(H_{\Delta}) \leq CT^2 ,$$

$$(21) \quad h_T(\lambda) := \rho(H_{\lambda}^{mod}) \geq c \text{ for any } |\lambda| \leq 2T .$$

The functional  $\rho_T = c^{-1}\rho$  is the required test functional in 4.6.

**5.3. Proof of Proposition 5.2.** We will construct a functional  $\rho$  as an integral of elementary functionals. Namely we find a positive function  $h \in H(G) \subset C^{\infty}(G \times G)$  and a

vector  $u \in E$  and define  $\rho(H) = \rho_u(\Pi(h)(H))$ , where  $\rho_u$  is the elementary functional on the space  $\mathcal{H}(E)$  corresponding to the vector  $u$ .

**5.3.1. Construction of function  $h$ .** We construct the function  $h$  independent of parameter  $T$ . Let  $D_1 \subset SL_2(\mathbb{R}) \subset G$  be the subset of matrices  $g$  with  $\|g\| \leq 2$ . We consider the subset  $D = D_1 \times D_1 \subset G \times G$  and choose a positive function  $h \in H(G \times G) = C_c^\infty(G \times G)$  which is  $\geq 1$  on the subset  $D$  and is supported in some neighborhood of  $D$ . We also assume that the function  $h$  is invariant under left and right translations by elements of the maximal compact subgroup  $K \times K$ .

**5.3.2. Construction of vector  $u$ .** Let us identify the space  $E = V \otimes V'$  with the space of smooth functions  $C^\infty(S^1 \times S^1)^{even}$ . Let  $S$  be a disc in  $S^1 \times S^1$  of radius  $(100T)^{-1}$ . We construct  $u$  as a smooth non-negative real valued function on  $S^1 \times S^1$  supported in  $S$  such that

- (i)  $\int u dx dy = 1$ ,
- (ii)  $\|u\|_{L^2}^2 \leq 10^5 T^2$ .

We would like to show that the functional  $\rho$  constructed in 5.3 satisfies conditions formulated in Proposition 5.2.

**5.3.3. Geometric bound.** We have

$$\rho(H_\Delta) = \rho_u(\Pi(h)(H_\Delta)) \leq C' \rho_u(P_E) = C' P_E(u) \leq CT^2$$

see 4.4, 5.3.2.

**5.3.4. Spectral bound.** First we would like to give another description of the Hermitian form  $H_\lambda^{mod}$ . Consider the model trilinear functional  $l = l_{\pi, \pi', \pi_i}^{mod}$  described in 5.1 and the corresponding operator  $T^l : E \rightarrow \bar{V}_\lambda$ .

We will identify the space  $V_\lambda$  with the space  $C^\infty(S^1)^{even}$ . Fix some point  $z \in S^1$ . Then the  $\delta$  functional  $\delta_z$  at this point defines a functional  $f$  on  $E$ . We denote by  $P_f$  the corresponding Hermitian form on  $E$ ,  $P_f(u) = |\langle f, u \rangle|^2$ .

Since the scalar product on the space  $\bar{V}_\lambda \simeq C^\infty(S^1)^{even}$  is given by the standard integral we see that the standard Hermitian form  $P_{\bar{V}_\lambda}$  is an average over the compact group  $K$  of the forms  $\pi(k)(P_{\delta_z})$ . This implies that  $H_\lambda^{mod} = \int_K (\Pi(k, k)(P_f)) dk$ .

Since we assumed the function  $h \in H(G \times G)$  to be  $K \times K$ -invariant we see that

$$\rho(H_\lambda^{mod}) = \rho_u(\Pi(h)(P_f))$$

Thus we see that in order to prove a lower bound for  $\rho(H_\lambda^{mod})$  it is enough to establish a lower bound for  $\rho_u(\Pi(g)P_f) := |\langle \Pi(g)f, u \rangle|^2$ .

Namely, the desired lower bound follows from the following

**Lemma.** *Let  $T_0$  be large enough. Then there exists an open non-empty subset  $D_0 \subset D$  such that for  $T \geq T_0$  and for  $g \in D_0$  we have  $|\langle \Pi(g)f, u \rangle| \geq 1/2$ .*

*Proof.* Let  $x, y$  be parameters on circles  $S^1$  describing  $V$  and  $V'$ . As follows from the definition (19) the functional  $f$  is given by the function  $f = f(x, y)$  on  $S^1 \times S^1$  described by formula

$$f(x, y) = |\sin(y - z)|^{(\alpha-1)/2} |\sin(x - z)|^{(\beta-1)/2} |\sin(x - y)|^{(\gamma-1)/2} .$$

Let  $D_0 \subset D$  be the subset of elements  $g \in D$  such that the function  $\Pi(g)(f)$  restricted to the subset  $S \subset S^1 \times S^1$  has absolute value  $\leq 10$ . It is easy to see that for large  $T$  this set is non empty and of some measure bounded below.

On the other hand, for  $g$  in this set we see that the gradient of the function  $\Pi(g)(f)$  on the subset  $S$  is bounded by  $3T$ . Hence the lower bound on  $|\langle P(g)f, u \rangle|$  for  $g \in D_0$  follows from the following easy claim

**Claim.** *Let  $S$  be a set with a measure  $\nu$  and  $u, h$  be two measurable functions on  $S$ . Let us assume that*

(i)  *$u$  is real valued positive function and  $\int u d\nu = 1$ .*

(ii)  *$\sup |h(s)| \geq 1$  and the variation  $\text{Var}(h) := \sup |h(s) - h(s')|$  is bounded by  $1/2$ .*

*Then  $|\int h u d\nu| \geq 1/2$ .*

□

**5.4. Construction of test functionals via Sobolev norms.** In this section we outline another, slightly more conceptual, construction of test functionals. This construction uses the notion of Sobolev norms on representation spaces (see [BR2]).

**5.4.1. Sobolev norms.** Let  $G$  be a Lie group and  $(\pi, G, V)$  a smooth pre-unitary representation. Then we can construct a family of positive definite Hermitian forms on the space  $V$  as follows.

Fix a basis  $\{X_j | j = 1, \dots, r\}$  of the Lie algebra  $\mathfrak{g}$  of the group  $G$ . Then for any natural number  $l$  and any  $T > 0$  we define a Hermitian form  $Q_{l,T}$  on  $V$  by

$$Q_{l,T}(v) = \sum_{\nu} T^{2(l-|\nu|)} P(X^{\nu}(v)) .$$

Here the sum is over all multi indexes  $\nu = (n_1, \dots, n_r)$  with the norm  $|\nu| := \sum n_j$  bounded by  $l$  and  $P = P_V$  is the Hermitian form defining the unitary structure on  $V$ .

**5.4.2. Positive functionals defined by forms.** Every positive definite Hermitian form  $Q$  on  $V$  defines a positive functional  $\rho_Q$  on  $\mathcal{H}(V)$  by  $\rho_Q(H) = \text{tr}(H|Q)$ . Here  $\text{tr}(H|Q)$  denotes



the relative trace of forms  $H$  and  $Q$ ; by definition it is equal to the square of the Hilbert-Schmidt norm of the identity operator on  $V$  considered as a morphism of pre-Hilbert spaces  $(V, Q) \rightarrow (V, H)$ . This notion is discussed in detail in [BR2].

**5.4.3. Construction of Sobolev test functionals.** Let us apply these constructions to the representation  $(\Pi, G \times G, E)$  discussed in 4.

Fix  $l$  and  $T$ , consider the Sobolev Hermitian form  $Q = Q_{l,T}$  on the space  $E$  and define the positive functional  $\rho$  on  $\mathcal{H}(E)$  to be  $\rho = \rho_Q$ .

**Proposition.** *Suppose  $l \geq 2$ . Then*

- (i)  $\rho(H_\Delta) \leq CT^{2-2l}$ ,
- (ii) *There exists  $c > 0$  such that  $\rho(H_\lambda^{mod}) \geq cT^{-2l}$  for  $|\lambda| \leq 2T$ .*

This gives another proof of Proposition 4.6.

**5.4.4. Sketch of the proof of Proposition 5.4.3.** (i) Since the representation  $\Pi$  is continuous with respect to the form  $Q_{l,T}$  the second basic inequality 4.4 implies that  $\rho(H_\Delta) \leq C'\rho(P_E)$ . The proof of the inequality  $\rho(P_E) \leq C''T^{2-2l}$  is the same as in [BR2], section 4.

In order to prove (ii) it is enough to find a vector  $u \in E$  such that  $Q_{l,T}(u) \leq T^{2l}$  and  $|\langle f, u \rangle| \geq c$ , where  $f = f_z$  is the function described in 5.3.4. We can take a function  $u \in C^\infty(S^1 \times S^1)$  of the form  $u = \phi f$  where  $\phi$  is a smooth cut-off function which equals 0 around singularities of the function  $f$ .

We leave details to the reader.

## APPENDIX A.

**A.1. Computation of  $l^{mod}$  for  $K$ -fixed vectors.** In this appendix we prove the Proposition 3.2 which describes the asymptotic behavior of the function  $k_\lambda$ .

One can prove this proposition applying the stationary phase method directly to the integral (19). To do this we need to consider the complexification of the functions  $e_\lambda(s_i)$  and the function  $K_{\lambda_1, \lambda_2, \lambda_3}(s_1, s_2, s_3)$  in the variables  $s_i$  and move contour of integration towards the singularities of the complexified integral. This could be done either in a classical language or using analytic continuation of representations in the spirit of [BR1].

**A.2. Computation of the integral.** We prefer to prove this proposition in a different way. Namely we explicitly compute the value of the model functional on the unit vectors in terms of  $\Gamma$ -functions and then prove the proposition by applying Stirling formulas for asymptotic behavior of  $\Gamma$ -functions.

Let  $\pi_{\lambda_i}$ ,  $i = 1, 2, 3$  be three representations of the generalized principal series and  $e_{\lambda_i}$  be the corresponding  $K$ -fixed unit vectors (they correspond to function 1 in the circle model). Set  $A(\lambda_1, \lambda_2, \lambda_3) := l_{\pi_{\lambda_1}, \pi_{\lambda_2}, \pi_{\lambda_3}}^{mod}(e_{\lambda_1} \otimes e_{\lambda_2} \otimes e_{\lambda_3})$ .

In sections A.4, A.5 we explicitly compute the function  $A(\lambda_1, \lambda_2, \lambda_3)$  (see the final expression in A.5) .

**A.3. Gaussian.** We would like to compute our integral by comparing it with Gaussian integrals which are much easier to manipulate with.

Namely, suppose we are given a finite-dimensional Euclidean vector space  $L$ . In this case we introduce the Gaussian probability measure  $G$  on  $L$  by  $dG = \pi^{-\dim L/2} \exp(-Q)dl$ , where  $Q$  is the quadratic form which defines the Euclidean structure on  $L$  and  $dl$  is the standard Euclidean measure on  $L$ .

We are interested in the quantities  $\langle f, G \rangle = \langle f, G \rangle_L := \int f dG$  for various (usually homogeneous) functions  $f$  on  $L$ . The main properties of the Gaussian which we use are the following:

(i) **Normalization.**  $\langle 1, G \rangle = 1$ .

(ii) **Product formula.** Suppose that the Euclidean space  $L$  is a product of Euclidean spaces  $L_1$  and  $L_2$ . Then the Gaussian measure  $G$  on  $L$  is the product of Gaussian measures  $G_1$  and  $G_2$  on  $L_1$  and  $L_2$ . In particular, if a function  $f$  decomposes as a product of functions  $f_1$  and  $f_2$  on  $L_1$  and  $L_2$  we have  $\langle f, G \rangle = \langle f_1, G_1 \rangle \langle f_2, G_2 \rangle$ .

The following integrals are classical

**Proposition.**

*Let  $L = \mathbb{R}^n$  be the standard Euclidean space.*

*(i) Let  $r$  denote the radius function on  $L$ . Then  $\langle r^s, G \rangle = \Gamma((s+n)/2)/\Gamma(n/2)$ .*

*(ii) Let  $h$  be a linear functional on  $L$ . Then  $\langle |h|^s, G \rangle = ||h||^s \Gamma((s+1)/2)/\Gamma(1/2)$ .*

*(iii) Let  $L$  be the space  $M_{2,2}$  of  $2 \times 2$  matrices with the standard Euclidean structure. Then  $\langle |\det|^s, G \rangle = \Gamma((s+1)/2)\Gamma(s/2+1)/\Gamma(1/2)$ .*

*Proof.* In (i) passing to spherical coordinates we get the integral

$$2c \int r^{s+n-1} \exp(-r^2) dr = c \int u^{(s+n)/2} \exp(-u) du / u = c \Gamma((s+n)/2) .$$

The normalization at  $s = 0$  defines the constant  $c = \Gamma(n/2)$ .

The proof of (ii) is reduced to the one variable case using product formula and then it follows from (i).

In (iii) we can write  $L$  as a product of two column spaces  $L_1$  and  $L_2$ . Then we have

$$\begin{aligned} \langle |\det|^s, G \rangle &= \int |\omega(x, y)|^s dG_1(x) dG_2(y) = \int (\int |\omega(x, y)|^s dG_1(x)) dG_2(y) = \\ &\Gamma((s+1)/2)/\Gamma(1/2) \cdot \int |y|^s dG_2(y) = \Gamma((s+1)/2)\Gamma(s/2+1)/\Gamma(1/2) \end{aligned}$$

since  $\Gamma(1) = 1$ . □

**A.4. Reduction 1.** The Proposition A.3 allows us to write the integrals which we would like to compute as some Gaussian integrals.

**Corollary.** *For any function  $h \in V_{-\lambda}$  we have*

$$\langle h, G \rangle = \Gamma((1 - \lambda)/2) \cdot \mathfrak{L}(h \cdot e_\lambda) .$$

Indeed, after averaging  $h$  with respect to the action of  $SO(2)$  we can assume that it is proportional to the function  $e_{-\lambda}$ . Then the formula follows from A.3(i).

Using this corollary we can rewrite the integral for the function  $A(\lambda_1, \lambda_2, \lambda_3)$ .

**Proposition.** *Consider the Euclidean space  $L = \mathbb{R}^2 \times \mathbb{R}^2 \times \mathbb{R}^2$  and define the function  $B(\lambda_1, \lambda_2, \lambda_3)$  by Gaussian integral  $B(\lambda_1, \lambda_2, \lambda_3) := \langle K_{\lambda_1, \lambda_2, \lambda_3}(s_1, s_2, s_3), G \rangle$ . Then*

$$B(\lambda_1, \lambda_2, \lambda_3) = A(\lambda_1, \lambda_2, \lambda_3) \cdot \Gamma((1 - \lambda_1)/2) \Gamma((1 - \lambda_2)/2) \Gamma((1 - \lambda_3)/2) .$$

**A.5. Reduction 2.** Let us rewrite the integral defining the function  $B$ . First, we identify the Euclidean space  $L$  in section A.4 with the space  $M_{2,3}$  of  $2 \times 3$  matrices. We consider the Euclidean space  $W \approx \mathbb{R}^3$  and define the map  $\nu : M_{2,3} \rightarrow W$  using  $2 \times 2$  minors. Let us define the function  $f$  on  $W$  by formula

$$f(w_1, w_2, w_3) = |w_1|^{(\alpha-1)/2} |w_2|^{(\beta-1)/2} |w_3|^{(\gamma-1)/2} .$$

We can write  $K_{\lambda_1, \lambda_2, \lambda_3} = \nu^*(f)$  and hence  $B(\lambda_1, \lambda_2, \lambda_3) = \langle \nu^*(f), G \rangle$

(here  $\alpha = \lambda_1 - \lambda_2 - \lambda_3, \beta = -\lambda_1 + \lambda_2 - \lambda_3, \gamma = -\lambda_1 - \lambda_2 + \lambda_3$  as in 5.1.1).

Now we will use the following general lemma which we prove in section A.6.

**Lemma.** *Let  $h$  be a homogeneous function on the space  $W$  of homogeneous degree  $s$ . Then*

$$\langle \nu^*(h), G \rangle_L = \langle h, G \rangle_W \cdot \Gamma(s/2 + 1) .$$

From this lemma we see that the computation of the function  $B(\lambda_1, \lambda_2, \lambda_3)$  is reduced to the computation of the function  $C(\alpha, \beta, \gamma) := \langle f, G \rangle$ .

Since the Gaussian  $G$  on  $W$  is a direct product of three one dimensional Gaussians and the function  $f$  is a product of functions depending only on one coordinate we deduce that the integral  $\langle f, G \rangle$  is a product of three one dimensional integrals which can be computed using A.3.

Thus we obtain  $C(\alpha, \beta, \gamma) = \Gamma((\alpha + 1)/4) \Gamma((\beta + 1)/4) \Gamma((\gamma + 1)/4)$ .

The final expression for the function  $A(\lambda_1, \lambda_2, \lambda_3)$  is

$$A(\lambda_1, \lambda_2, \lambda_3) = \frac{\Gamma((\alpha+1)/4)\Gamma((\beta+1)/4)\Gamma((\gamma+1)/4)\Gamma((\delta+1)/4)}{\Gamma((1-\lambda_1)/2)\Gamma((1-\lambda_2)/2)\Gamma((1-\lambda_3)/2)},$$

where  $\alpha = \lambda_1 - \lambda_2 - \lambda_3$ ,  $\beta = -\lambda_1 + \lambda_2 - \lambda_3$ ,  $\gamma = -\lambda_1 - \lambda_2 + \lambda_3$ ,  $\delta = -\lambda_1 - \lambda_2 - \lambda_3$ .

**A.6. Proof of Lemma A.5.** Consider the natural actions of the group  $SO(3)$  on the Euclidean spaces  $M_{2,3} \approx W \times W$  and  $W$ ; these actions preserve Gaussian measures.

The map  $\nu : M_{2,3} \rightarrow W$  is  $SO(3)$ -equivariant; it is nothing else than the exterior product map  $W \times W \rightarrow \bigwedge^2(W) = W^* = W$ . Hence we can replace the function  $h$  by its average with respect to the action of the group  $SO(3)$ , i.e. up to some constant by a function  $h = r^s$ . This shows that  $\langle \nu^*(h), G \rangle = a(s) \langle h, G \rangle$ , where  $a(s)$  depends on  $s$  but not on  $h$ .

In order to compute the function  $a(s)$  we can consider the identity above for the function  $h(w) = |w_3|^s$ . According to Proposition A.3(ii) we have  $\langle h, G \rangle = \Gamma((s+1)/2)/\Gamma(1/2)$ .

On the other hand it is clear that the function  $\nu^*(h)$  depends only on four variables and hence the integral  $\langle \nu^*(h), G \rangle$  coincides with the integral  $\langle h', G \rangle$  over the space  $M_{2,2}$  of  $2 \times 2$  matrices, where  $h'(m) = |\det(m)|^s$ .

From Proposition A.3(iii) we deduce that  $a(s) = \Gamma(s/2 + 1)$ .

**A.7. Proof of Proposition 3.2.** According to Stirling formulas for any fixed  $\sigma$  and large  $t$  we have  $\Gamma(\sigma + it) = \sqrt{2\pi} \exp(-\frac{\pi}{2}|t|) |t|^{\sigma-1/2} (1 + O(|t|^{-1}))$ .

This and the explicit formula for the function  $A(\lambda_1, \lambda_2, \lambda_3)$  implies the proposition.

## REFERENCES

- [BR1] J. Bernstein, A. Reznikov, Analytic continuation of representations, *Ann. Math.*, 150(1999), 329–352.
- [BR2] J. Bernstein, A. Reznikov, Sobolev norms of automorphic functionals, *IMRN* 2002:40 (2002), 2155–2174.
- [G1] I. Gelfand, G. Shilov, Generalized Functions. vol. 1, Academic Press, 1964.
- [G5] I. Gelfand, M. Graev, N. Vilenkin, Generalized Functions. vol. 5, Academic Press, 1966.
- [G6] I. Gelfand, M. Graev, I. Piatetski-Shapiro, Representation Theory and Automorphic Forms. Saunders, 1969.
- [Go] A. Good, Cusp forms and eigenfunctions of Laplacian, *Math. Ann.*, 255 (1981), 523–548.
- [HK] M. Harris, S. Kudla, The central critical value. *Ann. of Math.* 133, (1991), 605–672.
- [KS] B. Kröetz, R. Stanton, Holomorphic extension of representations, to appear in *Ann. of Math.*
- [L] S. Lang,  $SL_2(\mathbb{R})$ . Springer GTM105, 1985.
- [Lo] H. Loke, Trilinear forms of  $GL(2)$ , *Pacific J. Math.* 197 (2001), no. 1, 119–144.
- [Mo] V. Molchanov, Tensor products of unitary representations of the three-dimensional Lorentz group, *Izv. Akad. Nauk SSSR Ser. Mat.* 43 (1979), no. 4, 860–891.
- [O] A. Oksak, Trilinear Lorenz invariant forms. *Comm. Math. Phys.* 29 (1973), 189–217.
- [P] D. Prasad, Trilinear forms for representations of  $GL(2)$ , *Composito Math.*, 75(1990), 1–46.
- [PS] I. Piatetsky-Shapiro, Euler subgroups, in *Lie groups and their representations*, Halsted, (1975), 597–620.

- [Ra] R. Rankin, Contributions to the theory of Ramanujan's function  $\tau(n)$ , Proc. Camb. Philos. Soc. 35 (1939), 357–372.
- [Re1] A. Reznikov, Non-vanishing of periods of automorphic functions, Forum Math. 13, No.4 (2001), 485–493.
- [Re2] A. Reznikov, Laplace-Beltrami operator on a Riemann surface and equidistribution of measures, Comm. Math. Phys. 222 (2001), no. 2, 249–267.
- [Sa1] P. Sarnak, Integrals of products of eigenfunctions, IMRN, no. 6, (1994), 251–260.
- [Sa2] P. Sarnak, Arithmetical Quantum Chaos, Israeli Math. Conference Proc., vol. 8, 1995.
- [Se] A. Selberg On the estimation of Fourier coefficients, in Collected works, Springer-Verlag, New York (1989), 506–520.
- [Wa] T. Watson, Thesis, Princeton, 2001.

TEL AVIV UNIVERSITY, RAMAT AVIV, ISRAEL

*E-mail address:* `bernstei@math.tau.ac.il`

BAR ILAN UNIVERSITY, RAMAT-GAN, ISRAEL

*E-mail address:* `reznikov@math.biu.ac.il`