

On the Lie's Theorem for Lie Color Algebras

Shouchuan Zhang ^{a, b}

a. Department of Mathematics, Hunan University

Changsha 410082, P.R. China

b. Department of Mathematics, University of Queensland

Brisbane 4072, Australia

Abstract

We show that the Lie's Theorem holds for Lie color algebras with a torsion-free abelian group G . We give an example to show that the torsion-free condition is necessary.

0 Introduction

Since Kac's work [1] on the classification of Lie superalgebras, Lie color algebras have attracted much attention of mathematicians and mathematical physicists. For instance, Scheunert [2] studied Lie color algebras and obtained the PBW basis and Ado theorem of these algebras. Montgomery et al [3] [4] studied the properties of Lie color algebras determined by braided algebras, H -Lie algebras (or generalized Lie algebras) and (H, r) -Lie algebras. Majid [5][6] introduced the basic theory of Lie braided algebras and studied the transmutation and bosonisation of Lie braided bialgebras.

As generalized algebraic structures, Lie color algebras have found many physical applications, e.g. in the construction of the Yang-Baxter equation and integrable models [9][10]. In this letter we obtain the Lie's Theorem for Lie color algebras. We show that the Lie's Theorem for Lie color algebras holds provided that the abelian group G is torsion-free. We give an example to show that the torsion-free condition is necessary. Our results should shed new light on the study of structures of Lie color algebras.

1 Main Results

The concepts of nilpotent elements, ad-nilpotent elements, nilpotent Lie color algebras and solvable Lie color algebras are similar to ones in Lie algebras.

Let G be an abelian group with a skew symmetric bicharacter r . If $V = \sum_{g \in G} \oplus V_g$ is a graded vector space over field F with $\dim V_g = n_g < \infty$ and $A_{gh} = \text{Hom}_F(V_h, V_g)$ for any $g, h \in G$, then $A = \sum \{A_{ij} \mid i, j \in G\}$ is an algebra graded by G with $A_g = \sum_{i=j+g} A_{ij}$ for any $g \in G$. A is a Lie color algebra under the bracket operation $[a, b] = ab - r(|b|, |a|)ba$ for any homogeneous elements $a, b \in A$. Here $|a|$ stands for the degree of the element a . This Lie color algebra is referred to as general linear Lie color algebra, written as $gl(\{n_g\}, F)$ or $gl(\{V_g\}, F)$. Its Lie color subalgebras are called linear Lie color algebras. In fact, $gl(\{n_g\}, F) = \{f \in \text{End}_k V \mid \ker f \text{ is finite codimensional}\}$. When G is finite, we may view $gl(\{n_g\}, F)$ as a block matrix algebra over F . When $G = 0$, we denote $gl(\{n_g\}, F)$ by $gl(n, F)$, in this case it is the ungraded general linear Lie algebra.

Lemma 1.1 *Let L be a Lie color algebra and N be a Lie color subalgebra of $gl(\{V_g\}, F)$. Then, (i) $(ad X)^m(Y) = \sum_{i+j=m} k_{ij} X^i Y X^j$ for any homogeneous elements $X, Y \in N, k_{ij} \in F$, and natural number m .*

(ii) *If $X \in N$ is nilpotent homogeneous, then $ad X$ is nilpotent too.*

(iii) *For any homogeneous $x, y, z \in L$, $[[x, y], z] = [x[y, z]] + r(|z|, |y|)[[x, z], y]$, i.e. $ad [x, y] = [ad x, ad y]$. In other word, ad is a color representation of Lie color algebra L .*

(iv) *$Z(L)$ is a color ideal of Lie color algebra L . Here $Z(L) = \{x \in L \mid [x, y] = 0 \text{ for any } y \in L\}$, called the center of L .*

Proof. (i) It can be obtained by induction on natural number m .

(ii) It immediately follows from (i).

(iii) It is trivial.

(iv) We only need show that $Z(L)$ is G -graded. For any $x \in Z(L)$ and for any homogeneous element $y \in L$, since $[x, y] = 0$, we have that $\sum_{g \in G} [x_g, y] = 0$, which implies that $[x_g, y] = 0$. Consequently, $x_g \in Z(L)$. \square

Theorem 1.2 *Let L be a finite-dimensional Lie color subalgebra of $gl(\{V_g\}, F)$. If every homogeneous element in L is nilpotent and $V \neq 0$, then there exists a non-zero homogeneous element $v \in V$ such that $Lv = 0$.*

Proof. We use induction on $\dim L = n$. The case $\dim L = 0$ is obvious. Let K be a color subalgebra of L with $K \neq L$. For any $X \in K$, it is clear that there exists an F -linear map $\phi(X)$ from L/K to L/K such that the below diagram becomes commutative.

$$\begin{array}{ccc}
 & ad X & \\
 L & \longrightarrow & L \\
 \pi \downarrow & \searrow & \downarrow \pi \\
 L/K & \longrightarrow & L/K \\
 & \phi(X) &
 \end{array}$$

i.e. for any $Y \in L$, $\phi(X)(Y + K) = [X, Y] + K$. It is easy to verify that ϕ is a Lie color algebra homomorphism from K to $gl(\{L_g/K_g\}, F)$. The induction hypothesis guarantees

existence of a homogeneous element $X_0 + K \neq K$ in L/K such that $\phi(X)(X_0 + K) = K$, i.e. $[X, X_0] + K = K$ and $[X, X_0] \subseteq K$ for any $X \in K$. Thus $K + FX_0$ is a Lie color subalgebra of L with $\dim(K + FX_0) = \dim K + 1$. Consequently, there exist a Lie color subalgebra K and a homogeneous element $z \in L$ such that $L = K + Fz$. Obviously, K is a color ideal of L .

By induction, $W = \{v \in V \mid Kv = 0\} \neq 0$. Now we show that W is a graded space. For any homogeneous element $0 \neq x \in K$, $0 \neq w \in W$, we see that $x(w) = \sum_{g \in G} x(w_g) = 0$ implies $x(w_g) = 0$ since x is homogeneous. Thus $w_g \in W$ for any $g \in G$, which implies that W is a graded space. Since K is a G -graded ideal, W is stable under L : for three homogeneous elements $x \in L, y \in K, w \in W$, we see that $y(x(w)) = [y, x](w) + r(|x|, |y|)x(y(w)) = 0$. Since z is nilpotent on W there exists a non-zero $v \in W$ such that $z(w) = 0$ and $\sum_{g \in G} z(w_g) = 0$, which implies $z(w_g) = 0$ for any $g \in G$. Consequently, there exists a non-zero homogeneous $w_g \in W$ such that $z(w_g) = 0$. Finally, $L(w_g) = 0$, as desired. \square In [7], Scheunert stated that the Engel's theorem is true by the usual proof. Here for completeness, we have written down the theorem and provided a short proof.

Theorem 1.3 (See [7])(Engel's Theorem) *Let L be a finite-dimensional Lie color algebra. If all homogeneous elements of L is ad-nilpotent, then L is nilpotent.*

Proof. It is clear that $ad L = \{ad x \mid x \in L\} \subseteq gl(\{L_g\}, F)$ satisfies the hypothesis of Theorem 1.2. Thus there exists a homogeneous element $0 \neq x$ in L for which $[L, x] = 0$, i.e. $Z(L) \neq 0$. Now $L/Z(L)$ evidently consists of ad-nilpotent elements and has smaller dimension than L . Using induction on $\dim L$, we find that $L/Z(L)$ is nilpotent, which implies L is nilpotent. \square

Lemma 1.4 *Let $A \in gl(\{n_g\}, F)_u$ with torsion-free $u \neq 0$. Then*

(i) *If A has an eigenvalue, then A must have a zero eigenvalue and has homogeneous eigenvector of eigenvalue zero.*

(ii) *every eigenvalue of A is zero.*

(iii) *A is nilpotent.*

Proof. (i) If λ is an eigenvalue of A , then there exists an eigenvector $v \in V$ such that $A(v) = \lambda v$, i.e.

$$\sum_{g \in G} Av_g = \sum_{g \in G} \lambda v_g.$$

Thus $A(v_g) = \lambda v_{g+u}$ for any $g \in G$. Assume $v_{g_0} \neq 0$. There exists a natural number m such that $v_{g_0+mu} \neq 0$ and $v_{g_0+(m+1)u} = 0$. Since $A(v_{g_0+mu}) = \lambda v_{g_0+(m+1)u} = 0$, which implies that A has a zero eigenvalue and this eigenvector is homogeneous.

(ii) If λ is a non-zero eigenvalue of A , then there exists an eigenvector $v \in V$ such that $A(v) = \lambda v$, i.e.

$$\sum_{g \in G} Av_g = \sum_{g \in G} \lambda v_g.$$

Thus $A(v_g) = \lambda v_{g+u}$ for any $g \in G$. Assume $v_{g_0} \neq 0$. There exists a natural number m such that $V_{g_0-mu} \neq 0$ but $v_{g_0-(m+1)u} = 0$. Since $A(v_{g_0-(m+1)u}) = \lambda v_{g_0-mu} = 0$, $\lambda = 0$. We get a contradiction.

(iii) follows from part (ii). \square

Theorem 1.5 *Let L be a finite-dimensional solvable Lie color subalgebra of $gl(\{V_g\}, F)$ over algebraically closed field F and all of homogeneous elements in $[L, L]$ be nilpotent. If $V \neq 0$ and G is a torsion-free abelian group, then V contains a common homogeneous eigenvector for all the endomorphisms in L .*

Proof. Use induction on $\dim L$. The case $\dim L = 0$ is trivial. We attempt to imitate the proof of Theorem 1.2. The idea is (1) to locate a color ideal K of codimension one, (2) to show by induction that common homogeneous eigenvectors exist for K , (3) to verify that L stabilizes a space consisting of such eigenvectors, and (4) to find in that space homogeneous eigenvector for a single homogeneous $x \in L$ satisfying $L = K + Fz$.

Step (1) is easy. Since L is solvable, $[L, L] \neq 0$ and L properly includes $[L, L]$. $L/[L, L]$ being abelian, any G -graded subspace is automatically an ideal. Take a G -graded subspace of codimension one, then its inverse image K is a color ideal of codimension one in L . Set $L = K + Fz$ for some homogeneous element $z \in L$.

For step (2), use induction to find a common homogeneous eigenvector $v \in V$ for K . This means that for $x \in K$, $x(v) = \lambda(x)v$, $\lambda : K \rightarrow F$ is some linear function. Fix the λ and denote by W the color subspace generated by

$$\{w \in V \mid w \text{ is homogeneous and } x(w) = \lambda(x)w, \text{ for all } x \in K\}; \text{ so } W \neq 0.$$

Step (3) consists in showing that L leaves W invariant. Assuming for the moment that this is done, proceed to step (4); write $L = K + Fz$ and use the fact that F is algebraically closed and Lemma 1.4 to find a homogeneous eigenvector $v_0 \in W$ (for some eigenvector of z). Then v_0 is obviously a common eigenvector for L (and λ can be extended to a linear function on L such that $x(v_0) = \lambda(x)v_0, x \in L$).

It remains to show that L stabilizes W . Let homogeneous $w \in W$, homogeneous $x \in L$. To test whether or not $x(w)$ lies in W , we must take arbitrary $y \in K$ and examine $y(x(w)) = [y, x](w) + r(|x|, |y|)x(y(w)) = \lambda([y, x])(w) + r(|x|, |y|)\lambda(y)x(w)$. Thus we have to prove that $\lambda([y, x]) = 0$. However, since $[y, x]$ nilpotent, $\lambda([y, x]) = 0$. We complete the proof. \square

Corollary 1.6 (*Lie's Theorem*) *Let L be a finite-dimensional solvable Lie color subalgebra of $gl(\{V_g\}, F)$ and all of homogeneous elements in $[L, L]$ nilpotent. If G is a torsion-free abelian group, then L stabilizes some color flag in V (in other words, the matrices of L relative to a suitable homogeneous basis of V are uppertriangular).*

The Lie's theorem is one of the main results in the present letter. The condition that G is torsion-free is necessary for the Lie's theorem to be true as is seen from the following example.

Example 1.7 Let $G = \mathbf{Z}_3$ with trivial bicharacter $r(x, y) = 1$ for $x, y \in \mathbf{Z}_3$, $\{e_1, e_2, e_3\}$ the natural basis of \mathbf{F}^3 , $V_i = Fe_i$ for $i = 1, 2, 3$ and a linear transformation $A = \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{pmatrix}$. Let $L = FA$ be a linear Lie color algebra in $gl(\{V_g\}, F)$. Obviously, the degree of A is 2 and $[L, L] = 0$. In other word, L is a finite-dimensional solvable color subalgebra of $gl(\{V_g\}, F)$ and all homogeneous elements in $[L, L]$ are nilpotent. But the matrices of linear transformation A relative to any homogeneous basis of V can not be upper triangular.

Remark: In the case of Lie algebras, the other conditions in Corollary 1.6 imply the condition that every homogeneous element in $[L, L]$ is nilpotent by [8, Corollary 4.1 A]. In other word, our Corollary 1.6 implies the Lie's theorem for (ungraded) Lie algebras.

Corollary 1.8 Let L be a finite-dimensional solvable Lie color subalgebra of $gl(\{n_g\}, F)$ and all homogeneous elements in $[L, L]$ be nilpotent. If G is a torsion-free abelian group, then there exists a chain of color ideals of L , $0 = L_0 \subseteq L_1 \subseteq \dots \subseteq L_n = L$ such that $\dim L_i = i$.

Acknowledgement : The work is supported by Australian Research Council. S.C.Z thanks the Departemnt of Mathematics, University of Queensland for hospitality.

References

- [1] V. G. Kac, Adv. Math., **26** (1977), 8–96.
- [2] M. Scheunert, J. Math. Phys. **20** (1979), 712-720.
- [3] Y. Bahturin, D. Fishman and S. Montgomery, J. Alg. **96** (1996), 27-48.
- [4] Y. Bahturin, D. Fischman and S. Montgomery, J. Alg. **236** (2001), 246-276.
- [5] S. Majid, J. Geom. Phys. **13** (1994), 307–356.
- [6] S. Majid, Pacific J. Math. **192** (2000)2, 329–356.
- [7] M. Scheunert, Lie superalgebras, Springer Lecture Notes in Math. **716**, 1979.
- [8] J. E. Humphreys, Introduction to Lie algebras and representation theory, Graduate Texts in Mathematics 9, Springer-Verlag, 1972.

- [9] D.S. McAnally and A.J. Bracken, *Bull. Austral. Math. Soc.* **55** (1997), 425–428.
- [10] D.S. McAnally, in *Group22: Proc. of the XXII Int. Coll. on Group Theor. Methods in Phys.*, pp428-432, 1998, eds: S.P. Corney, R. Delbourgo and P.D. Jarvis.