

REGULAR POINTS IN AFFINE SPRINGER FIBERS

MARK GOESKY, ROBERT KOTTWITZ, AND ROBERT MACPHERSON

1. INTRODUCTION

Let G be a connected reductive group over \mathbf{C} with Lie algebra \mathfrak{g} . We put $F = \mathbf{C}((\epsilon))$ and $\mathcal{O} = \mathbf{C}[[\epsilon]]$. Let $X = X_G$ denote the affine Grassmannian $G(F)/G(\mathcal{O})$. For $u \in \mathfrak{g}(F)$ we write X^u for the affine Springer fiber

$$X^u = \{g \in G(F)/G(\mathcal{O}) : \text{Ad}(g^{-1})(u) \in \mathfrak{g}(\mathcal{O})\}.$$

studied by Kazhdan and Lusztig in [KL88].

For $x = gG(\mathcal{O}) \in X^u$ the $G(\mathcal{O})$ -orbit (for the adjoint action) of $\text{Ad}(g^{-1})(u)$ in $\mathfrak{g}(\mathcal{O})$ depends only on x , and its image under $\mathfrak{g}(\mathcal{O}) \rightarrow \mathfrak{g}(\mathbf{C})$ is a well-defined $G(\mathbf{C})$ -orbit in $\mathfrak{g}(\mathbf{C})$. We say that $x \in X^u$ is *regular* if the associated orbit is regular in $\mathfrak{g}(\mathbf{C})$. (Recall that an element of $\mathfrak{g}(\mathbf{C})$ is regular if the nilpotent part of its Jordan decomposition is a principal nilpotent element in the centralizer of the semisimple part of its Jordan decomposition.) We write X_{reg}^u for the (Zariski open) subset of regular elements in X^u .

From now on we assume that u is regular semisimple with centralizer T , a maximal torus in G over F . Assume further that u is *integral*, by which we mean that X^u is non-empty. Kazhdan and Lusztig [KL88] show that X^u is then a locally finite union of projective algebraic varieties, and in Cor. 1 of §4 of [KL88] they show that the open subset X_{reg}^u of X^u is non-empty (and hence dense in at least one irreducible component of X^u). The action of $T(F)$ on X clearly preserves the subsets X^u and X_{reg}^u . Bezrukavnikov [Bez96] proved that X_{reg}^u forms a single orbit under $T(F)$. (Actually Kazhdan-Lusztig and Bezrukavnikov consider only topologically nilpotent elements u , but the general case can be reduced to their special case by using the topological Jordan decomposition of u .)

The goal of this paper is to characterize regular elements in X^u (for integral regular semisimple u as above). When T is elliptic (in other words, F -anisotropic modulo the center of G) the characterization gives no new information. At the other extreme, in the split case, the characterization gives a clear picture of what it means for a point in X^u to be regular.

We will now state our characterization in the split case, leaving the more technical general statement to the next section (see Theorem 1). Fix a split maximal torus $A \subset G$ over \mathbf{C} and denote by \mathfrak{a} its Lie algebra. We identify the affine

Date: August 7, 2003.

1991 Mathematics Subject Classification. Primary: 22E50; Secondary: 20G25, 22E67.

Key words and phrases. Affine Springer fiber, fixed point set, affine Grassmannian.

M.G. was partially supported by NSF Grant DMS-0139986.

R.K. was partially supported by NSF Grant DMS-0245639.

Grassmannian $A(F)/A(\mathcal{O})$ for A with the cocharacter lattice $X_*(A)$, the cocharacter μ corresponding to the class of $\mu(\epsilon)$ in $A(F)/A(\mathcal{O})$. For any Borel subgroup $B = AN$ containing A (N denoting the unipotent radical of B) there is a well-known retraction $r_B : X \rightarrow X_*(A)$ defined using the Iwasawa decomposition $G(F) = N(F)A(F)G(\mathcal{O})$: the fiber of r_B over $\mu \in X_*(A)$ is $N(F)\mu(\epsilon)G(\mathcal{O})/G(\mathcal{O})$. The family of cocharacters $r_B(x)$ (B ranging through all Borel subgroups containing A) has been studied by Arthur [Art76, Lemma 3.6]; it is the volume of the convex hull of these points that arises as the weight factor for (fully) weighted orbital integrals for elements in $A(F)$. In particular Arthur shows that for $x \in X$ and any pair B, B' of adjacent Borel subgroups containing A , there is a unique non-negative integer $n(x, B, B')$ such that

$$(1.0.1) \quad r_B(x) - r_{B'}(x) = n(x, B, B') \cdot \alpha_{B, B'}^\vee$$

where $\alpha_{B, B'}$ is the unique root of A that is positive for B and negative for B' .

The main result of this paper (in the split case) is that for $x \in X^u$

$$(1.0.2) \quad n(x, B, B') \leq \text{val } \alpha_{B, B'}(u)$$

for every pair B, B' of adjacent Borel subgroups containing A , and that $x \in X^u$ is regular if and only if all the inequalities (1.0.2) are actually equalities.

2. STATEMENTS

2.1. Notation. We write \mathfrak{g} for the Lie algebra of G and follow the same convention for groups denoted by other letters.

Choose an algebraic closure \overline{F} of F and let $\Gamma = \text{Gal}(\overline{F}/F)$. We write G_F for the F -group obtained from G by extension of scalars from \mathbf{C} to F .

As before we use $\mu \mapsto \mu(\epsilon)$ to identify the cocharacter group $X_*(A)$ with $A(F)/A(\mathcal{O})$. By means of this identification the canonical surjection $A(F) \rightarrow A(F)/A(\mathcal{O})$ can be viewed as a surjection

$$(2.1.1) \quad A(F) \rightarrow X_*(A).$$

Let $\Lambda = \Lambda_G$ denote the quotient of the coweight lattice $X_*(A)$ by the coroot lattice (the subgroup of $X_*(A)$ generated by the coroots of A in G). Up to canonical isomorphism Λ is independent of the choice of A ; moreover when defining Λ we could replace A by any maximal torus T in G_F . There is a canonical surjective homomorphism

$$(2.1.2) \quad G(F) \rightarrow \Lambda,$$

characterized by the following two properties: it is trivial on the image of $G_{\text{sc}}(F)$ in $G(F)$ (where G_{sc} denotes the simply connected cover of the derived group of G), and its restriction to $A(F)$ coincides with the composition of (2.1.1) and the canonical surjection $X_*(A) \rightarrow \Lambda$.

Recall that X denotes the affine Grassmannian $G(F)/G(\mathcal{O})$ for G . The homomorphism (2.1.2) is trivial on $G(\mathcal{O})$ and hence induces a canonical surjection

$$(2.1.3) \quad \nu_G : X \rightarrow \Lambda,$$

whose fibers are the connected components of X .

2.2. Parabolic subgroups. We will be concerned with parabolic subgroups P of G containing A . Such a parabolic subgroup has a unique Levi subgroup M containing A , and we refer to M as *the* Levi component of P .

As usual, by a Levi subgroup of G , we mean a Levi subgroup of some parabolic subgroup of G . Let M be a Levi subgroup of G containing A . We write $\mathcal{P}(M)$ for the set of parabolic subgroups of G that contain A and have Levi component M . Thus any $P \in \mathcal{P}(M)$ can be written as $P = MN$ where $N = N_P$ denotes the unipotent radical of P . As usual there is a notion of adjacency: two parabolic subgroups $P = MN$ and $P' = MN'$ in $\mathcal{P}(M)$ are said to be *adjacent* if there exists (a unique) parabolic subgroup $Q = LU$ containing both P and P' such that the semisimple rank of L is one greater than the semisimple rank of M . Thus $U = N \cap N'$, and, moreover, if L is chosen so that $L \supset A$, then

$$\mathfrak{l} = \mathfrak{m} \oplus (\mathfrak{n} \cap \bar{\mathfrak{n}}') \oplus (\mathfrak{n}' \cap \bar{\mathfrak{n}})$$

where \bar{N} denotes the unipotent radical of the parabolic subgroup $\bar{P} = M\bar{N}$ opposite to P (and where \bar{N}' is opposite to N').

Given adjacent P, P' in $\mathcal{P}(M)$ we define an element $\beta_{P,P'} \in \Lambda_M$ (the coweight lattice for A modulo the coroot lattice for M) as follows. Consider the collection of elements in Λ_M obtained from coroots α^\vee where α ranges through the set of roots of A in $\mathfrak{n} \cap \bar{\mathfrak{n}}'$. We define $\beta_{P,P'}$ to be the unique element in this collection such that all other members in the collection are positive integral multiples of $\beta_{P,P'}$. Note that although Λ_M may have torsion elements, the elements in our collection lie in the kernel of the canonical map from Λ_M to Λ_G , and this kernel is torsion-free. Thus any member of our collection can be written uniquely as a positive integer times $\beta_{P,P'}$. Note also that $\beta_{P',P} = -\beta_{P,P'}$. In case $M = A$, so that P, P' are Borel subgroups, $\beta_{P,P'}$ is the unique coroot of A that is positive for P and negative for P' .

2.3. Retractions from X to X_M . The inclusion of $M(F)$ into $G(F)$ induces an inclusion of the affine Grassmannian X_M for M into the affine Grassmannian X for G . Let $P \in \mathcal{P}(M)$ and let X_P denote the set $P(F)/P(\mathcal{O})$. The canonical inclusion of P in G induces a bijection i from X_P to X , and the canonical surjection $P \rightarrow M$ induces a canonical surjective map p (of sets) from X_P to X_M . We define the retraction $r_P = r_P^G : X \rightarrow X_M$ as the composed map $p \circ i^{-1}$. Given $x \in X$ we often denote by x_P the image of x under the retraction r_P .

These retractions satisfy the following transitivity property. Suppose that $L \supset M$ are Levi subgroups containing A , and suppose further that $P \in \mathcal{P}(M)$ and $Q \in \mathcal{P}(L)$ satisfy $Q \supset P$. Let P_L denote the parabolic subgroup $P \cap L$ in L . Then

$$(2.3.1) \quad r_P^G = r_{P_L}^L \circ r_Q^G.$$

Moreover, for any $x \in X$ the element $\nu_M(x_P)$ maps to $\nu_L(x_Q)$ under the canonical surjection $\Lambda_M \rightarrow \Lambda_L$, and in particular $\nu_M(x_P) \mapsto \nu_G(x)$ under $\Lambda_M \rightarrow \Lambda_G$.

2.4. Definition of $n(x, P, P')$. A point $x \in X$ determines points $\nu_M(x_P)$ in Λ_M , one for each $P \in \mathcal{P}(M)$. This family of points arises in the definition of the weighted orbital integrals occurring in Arthur's work. A basic fact [Art76] about this family of points is that whenever P, P' are adjacent parabolic subgroups in $\mathcal{P}(M)$, there is a (unique) non-negative integer $n(x, P, P')$ such that

$$(2.4.1) \quad \nu_M(x_P) - \nu_M(x_{P'}) = n(x, P, P') \cdot \beta_{P,P'}.$$

2.5. Fixed point sets X^u . Let $u \in \mathfrak{g}(F)$. Define a subset X^u of X by

$$X^u = \{g \in G(F)/G(\mathcal{O}) : \text{Ad}(g^{-1})(u) \in \mathfrak{g}(\mathcal{O})\}.$$

2.6. Conjugacy classes associated to fixed points. Let $u \in \mathfrak{g}(F)$. Suppose that the coset $x = gG(\mathcal{O})$ lies in X^u . The image of $\text{Ad}(g^{-1})(u)$ under the canonical surjection $\mathfrak{g}(\mathcal{O}) \rightarrow \mathfrak{g}(\mathbf{C})$ gives a well-defined $G(\mathbf{C})$ -conjugacy class $\bar{u}_G(x)$ (for the adjoint action) in $\mathfrak{g}(\mathbf{C})$.

As above let M be a Levi subgroup of G and let $P \in \mathcal{P}(M)$. Now suppose that $u \in \mathfrak{m}(F)$ and that $x \in X^u$. Choose $p \in P(F)$ such that $x = pG(\mathcal{O})$; thus x_P is the coset $mM(\mathcal{O})$, where m denotes the image of p under the canonical homomorphism from P onto M . Of course $\text{Ad}(p^{-1})(u)$ lies in $\mathfrak{p}(\mathcal{O})$, and its image in $\mathfrak{p}(\mathbf{C})$ gives a well-defined $P(\mathbf{C})$ -conjugacy class $\bar{u}_P(x)$ in $\mathfrak{p}(\mathbf{C})$. It follows that x_P lies in X_M^u (as was first noted by Kazhdan-Lusztig [KL88]), and also that $\bar{u}_P(x)$ maps to $\bar{u}_G(x)$ (respectively, $\bar{u}_M(x_P)$) under the map on conjugacy classes induced by $\mathfrak{p}(\mathbf{C}) \hookrightarrow \mathfrak{g}(\mathbf{C})$ (respectively, $\mathfrak{p}(\mathbf{C}) \rightarrow \mathfrak{m}(\mathbf{C})$).

2.7. Review of regular elements. An element $u \in \mathfrak{g}(\mathbf{C})$ is *regular* if the nilpotent part of its Jordan decomposition is a principal nilpotent element in the centralizer of the semisimple part of its Jordan decomposition, or, equivalently, if the set of Borel subalgebras containing u is finite. It is well-known that the set of regular elements in $\mathfrak{g}(\mathbf{C})$ is open.

As above let M be a Levi subgroup of G and let $P \in \mathcal{P}(M)$. Suppose that u is a regular element in $\mathfrak{g}(\mathbf{C})$ that happens to lie in $\mathfrak{p}(\mathbf{C})$. Then the image u_M of u in $\mathfrak{m}(\mathbf{C})$ is regular in $\mathfrak{m}(\mathbf{C})$.

2.8. Regular points in X^u . We say that $x \in X^u$ is *regular* if the associated conjugacy class $\bar{u}_G(x) \in \mathfrak{g}(\mathbf{C})$ consists of regular elements. We denote by X_{reg}^u the set of regular elements in X^u ; the subset X_{reg}^u is open in X^u .

As above let M be a Levi subgroup of G and let $P \in \mathcal{P}(M)$. Suppose that $u \in \mathfrak{m}(F)$. We have already seen that r_P maps X^u into X_M^u , and that the conjugacy class in $\mathfrak{g}(\mathbf{C})$ associated to $x \in X^u$ is compatible with the conjugacy class in $\mathfrak{m}(\mathbf{C})$ associated to the retracted point $x_P \in X_M^u$, compatible in the sense that there is a conjugacy class in $\mathfrak{p}(\mathbf{C})$ that maps to both of them. Therefore x_P is regular in X_M^u if x is regular in X^u .

2.9. Set-up for the main result. As before let M denote a Levi subgroup of G containing A . We now assume that u is an integral regular semisimple element of $\mathfrak{g}(F)$ that happens to lie in $\mathfrak{m}(F)$. (It is equivalent to assume that the centralizer T of u is contained in M_F .) For each pair $P = MN$, $P' = MN'$ of adjacent parabolic subgroups in $\mathcal{P}(M)$ we are going to define a non-negative integer $n(u, P, P')$. This collection of integers measures how far X^u sticks out from X_M^u .

As before we need the parabolic subgroups $\bar{P} = M\bar{N}$ and $\bar{P}' = M\bar{N}'$ opposite to P and P' respectively. Let α be a root of T in $N \cap \bar{N}'$. Since T , N and N' are defined over F , the group $\text{Gal}(\bar{F}/F)$ preserves the set of roots of T in $N \cap \bar{N}'$. Let F_α denote the field of definition of α , so that $\text{Gal}(\bar{F}/F_\alpha)$ is the stabilizer of α in $\text{Gal}(\bar{F}/F)$. For any finite extension F' of F (e.g. F_α) we normalize the valuation $\text{val}_{F'}$ on F' so that a uniformizing element in F' has valuation 1, or, equivalently, so that ϵ has valuation $[F' : F]$. There exists a unique positive integer m_α such that the image of the element α^\vee in Λ_M is equal to $m_\alpha \cdot \beta_{P, P'}$, where $\beta_{P, P'}$ is the element of Λ_M defined above. Note that m_α depends only on the orbit of α under

the Galois group; here we use that the Galois group acts on the cocharacter group of T through the Weyl group of M , so that any two elements in the Galois orbit of α^\vee have the same image in Λ_M . Finally we define $n(u, P, P')$ as the sum

$$(2.9.1) \quad n(u, P, P') = \sum \text{val}_{F_\alpha}(\alpha(u)) \cdot m_\alpha,$$

where the sum is taken over a set of representatives α of the orbits of $\text{Gal}(\overline{F}/F)$ on the set of roots of T in $N \cap \overline{N}'$. In the special case that $M = A$ (and hence $T = A$) $n(u, P, P')$ is equal to $\text{val}_F(\alpha(u))$, where α is the unique root of A that is positive for P and negative for P' .

Theorem 1. *Let M and u be as above, and let $x \in X^u$. Recall that $x_P \in X_M^u$ for all $P \in \mathcal{P}(M)$.*

(a) *For every pair $P, P' \in \mathcal{P}(M)$ of adjacent parabolic subgroups*

$$n(x, P, P') \leq n(u, P, P').$$

(b) *The point x is regular in X^u if and only if the following two conditions hold:*

- (i) *the point x_P is regular in X_M^u for all $P \in \mathcal{P}(M)$, and*
- (ii) *for every pair $P, P' \in \mathcal{P}(M)$ of adjacent parabolic subgroups*

$$n(x, P, P') = n(u, P, P').$$

3. PROOFS

3.1. The case of $SL(2)$. The key step in proving our main theorem is to verify it for $SL(2)$, where it reduces to a computation that can be found in [Lan80]. To keep things self-contained we reproduce the calculation here. Let A, B, \bar{B} denote the diagonal, upper triangular and lower triangular subgroups of $SL(2)$ respectively, and let α be the unique root of A that is positive for B . Of course $\beta_{B, \bar{B}} = \alpha^\vee$. Let $x \in X$ and let $u = \begin{bmatrix} c & 0 \\ 0 & -c \end{bmatrix}$ for non-zero $c \in \mathcal{O}$. Note that $n(u, B, \bar{B}) = \text{val}_F(c)$. We will show that $x \in X^u$ if and only if $n(x, B, \bar{B}) \leq n(u, B, \bar{B})$, and that $x \in X_{\text{reg}}^u$ if and only if $n(x, B, \bar{B}) = n(u, B, \bar{B})$.

The difference $\nu_A(x_B) - \nu_A(x_{\bar{B}})$ and the sets X^u and X_{reg}^u are invariant under the action of $A(F)$ on X , so it is enough to consider x of the form $x = gG(\mathcal{O})$ with $g = \begin{bmatrix} 1 & 0 \\ t & 1 \end{bmatrix}$. (Note that for this reason our calculations apply just as well to any group whose semisimple rank is 1.) For such x we have $\nu_A(x_B) = 0$. If $t \in \mathcal{O}$, then $\nu_A(x_B) = 0$. If $t \notin \mathcal{O}$, then $\begin{bmatrix} 0 & -1 \\ 1 & t^{-1} \end{bmatrix} \in G(\mathcal{O})$ and thus

$$\begin{bmatrix} 1 & 0 \\ t & 1 \end{bmatrix} = \begin{bmatrix} t^{-1} & 1 \\ 0 & t \end{bmatrix} \begin{bmatrix} 0 & -1 \\ 1 & t^{-1} \end{bmatrix} \in \begin{bmatrix} t^{-1} & 1 \\ 0 & t \end{bmatrix} \cdot G(\mathcal{O}),$$

which shows that $\nu_A(x_B) = \text{val}_F(t^{-1}) \cdot \alpha^\vee$. We conclude that $n(x, B, \bar{B})$ equals 0 if $t \in \mathcal{O}$ and equals $\text{val}_F(t^{-1})$ if $t \notin \mathcal{O}$. In any case $n(x, B, \bar{B})$ is a non-negative integer.

For x, u as above we have

$$\text{Ad}(g^{-1})u = \begin{bmatrix} c & 0 \\ -2ct & -c \end{bmatrix}.$$

Therefore $x \in X^u \iff ct \in \mathcal{O} \iff n(x, B, \bar{B}) \leq n(u, B, \bar{B})$. Moreover $x \in X_{\text{reg}}^u \iff ct \in \mathcal{O}^\times$ or $(c \in \mathcal{O}^\times \text{ and } t \in \mathcal{O}) \iff n(x, B, \bar{B}) = n(u, B, \bar{B})$.

3.2. Review of $n(x, P, P')$. We need to review Arthur's proof of the existence of the non-negative integers $n(x, P, P')$. We begin with the case $M = A$. Let $x \in X$. We must check that for any two adjacent Borel subgroups $P, P' \in \mathcal{P}(A)$ there is a (unique) non-negative integer $n(x, P, P')$ such that

$$\nu_A(x_P) - \nu_A(x_{P'}) = n(x, P, P') \cdot \alpha^\vee,$$

where α is the unique root of A that is positive for P and negative for P' . For this we consider the unique parabolic subgroup Q containing P and P' whose Levi component L has semisimple rank 1. By transitivity of retractions we have

$$(3.2.1) \quad \nu_A(x_P) - \nu_A(x_{P'}) = \nu_A(y_B) - \nu_A(y_{\bar{B}})$$

where $y = x_Q$ and $B = L \cap P, \bar{B} = L \cap P'$. This reduces us to the case in which G has semisimple rank 1, which has already been done. For future use we note that (3.2.1) can be reformulated as the equality

$$n(x, P, P') = n(y, B, \bar{B}).$$

Again let $x \in X$. Now we check that for any Levi subgroup $M \supset A$ and any adjacent parabolic subgroups $P = MN, P' = MN'$ in $\mathcal{P}(M)$ there is a (unique) non-negative integer $n(x, P, P')$ such that

$$\nu_M(x_P) - \nu_M(x_{P'}) = n(x, P, P') \cdot \beta_{P, P'}.$$

Fix a Borel subgroup B_M in M and let B (respectively, B') be the inverse image of B_M under $P \rightarrow M$ (respectively, $P' \rightarrow M$); thus B, B' are Borel subgroups containing A .

Now choose a minimal gallery of Borel subgroups $B = B_0, B_1, B_2, \dots, B_l = B'$ joining B to B' , and for $i = 1, \dots, l$ let α_i be the unique root of A that is positive for B_{i-1} and negative for B_i . Then

$$\nu_A(x_B) - \nu_A(x_{B'}) = \sum_{i=1}^l n(x, B_{i-1}, B_i) \cdot \alpha_i^\vee.$$

Note that $\{\alpha_1, \dots, \alpha_l\}$ is precisely the set of roots of A in $\mathfrak{n} \cap \bar{\mathfrak{n}}'$ and that for each i there exists a (unique) positive integer m_i such that the image of α_i^\vee in Λ_M is equal to $m_i \cdot \beta_{P, P'}$. Applying the canonical surjection $\Lambda_A \rightarrow \Lambda_M$ to the previous equation, we find that (see 2.3)

$$\nu_M(x_P) - \nu_M(x_{P'}) = n(x, P, P') \cdot \beta_{P, P'}.$$

where $n(x, P, P')$ is the non-negative integer

$$\sum_{i=1}^l m_i \cdot n(x, B_{i-1}, B_i).$$

3.3. Proof of part of the main theorem in case $A = T$. Let $u \in \mathfrak{a}(\mathcal{O})$ and assume that u is regular in $\mathfrak{g}(F)$. Let $x \in X^u$.

Let M be a Levi subgroup of G containing A . We are now going to prove the first main assertion in our theorem, namely that for any pair of adjacent $P, P' \in \mathcal{P}(M)$ there is an inequality

$$n(x, P, P') \leq n(u, P, P').$$

Let B, B', B_0, \dots, B_l and α_i, m_i ($i = 1, \dots, l$) be as in 3.2. Then by definition

$$n(u, P, P') = \sum_{i=1}^l m_i \cdot \text{val}_F(\alpha_i(u)).$$

Let M_i be the Levi subgroup containing A whose root system is $\{\pm\alpha_i\}$, and let B'_{i-1}, B'_i denote the Borel subgroups in M_i obtained by intersecting B_{i-1}, B_i with M_i . Let Q_i be the unique parabolic subgroup in $\mathcal{P}(M_i)$ such that Q_i contains B_{i-1} and B_i . We showed in 3.2 that

$$n(x, P, P') = \sum_{i=1}^l m_i \cdot n(x, B_{i-1}, B_i)$$

and that

$$n(x, B_{i-1}, B_i) = n(y_i, B'_{i-1}, B'_i),$$

where $y_i = x_{Q_i} \in X_{M_i}^u$. Since M_i has semisimple rank 1, we know that

$$n(y_i, B'_{i-1}, B'_i) \leq \text{val}_F(\alpha_i(u)).$$

This completes the proof of the first main assertion.

Now suppose that x is regular in X^u . Then each point $y_i \in X_{M_i}^u$ above is regular in $X_{M_i}^u$, and therefore from the rank 1 case (see 3.1) we know that

$$n(y_i, B'_{i-1}, B'_i) = \text{val}_F(\alpha_i(u)).$$

We conclude that if x is regular in X^u , then

$$n(x, P, P') = n(u, P, P'),$$

which is another of the assertions in our theorem.

3.4. Proof of the rest of the main theorem in case $M = A = T$. We continue with $u \in \mathfrak{a}(\mathcal{O})$ and $x \in X^u$ as before, but for the moment we only consider the case $M = A$. We assume that

$$(3.4.1) \quad n(x, P, P') = \text{val}_F(\alpha_{P, P'}(u))$$

for all adjacent Borel subgroups $P, P' \in \mathcal{P}(A)$, where $\alpha_{P, P'}$ is the unique root of A that is positive for P and negative for P' . We want to prove that x is regular in X^u . To do so we must first select a suitable Borel subgroup $B \in \mathcal{P}(A)$.

Let $u_0 \in \mathfrak{a}(\mathbf{C})$ denote the image of u under $\mathfrak{a}(\mathcal{O}) \rightarrow \mathfrak{a}(\mathbf{C})$, and let M denote the centralizer of u_0 in G . Thus M is a Levi subgroup of G containing A , and we choose $P \in \mathcal{P}(M)$. Then we obtain a suitable Borel subgroup by taking any $B \in \mathcal{P}(A)$ such that $B \subset P$. For any B -simple root α we denote by B_α the unique Borel subgroup in $\mathcal{P}(A)$ that is adjacent to B and for which α is negative, and we write P_α for the unique parabolic subgroup containing B and B_α such that the semisimple rank of the Levi component M_α of P_α is 1. Consider the element (well-defined up to $B(\mathbf{C})$ -conjugacy) $v := \bar{u}_B(x) \in \mathfrak{b}(\mathbf{C})$ defined in 2.6. The equation (3.4.1) plus the semisimple rank 1 theory implies that the points $x_{P_\alpha} \in X_{M_\alpha}^u$ are

regular, and this in turn implies (see 2.6) that for every B -simple root α the image of the element v under $\mathfrak{b}(\mathbf{C}) \hookrightarrow \mathfrak{p}_\alpha(\mathbf{C}) \twoheadrightarrow \mathfrak{m}_\alpha(\mathbf{C})$ is regular in $\mathfrak{m}_\alpha(\mathbf{C})$. Moreover it is evident that the image of v under the canonical surjection $\mathfrak{b}(\mathbf{C}) \twoheadrightarrow \mathfrak{a}(\mathbf{C})$ is equal to u_0 . Using only these facts, we now check that v is regular in $\mathfrak{g}(\mathbf{C})$ (and hence that x is regular in X^u).

Let $v = v_s + v_n$ be the Jordan decomposition of v , with v_s semisimple and v_n nilpotent. Since it is harmless to replace v by any $B(\mathbf{C})$ -conjugate, we may assume without loss of generality that $v_s \in \mathfrak{a}(\mathbf{C})$. Then, since $v_s \mapsto u_0$ under $\mathfrak{b}(\mathbf{C}) \twoheadrightarrow \mathfrak{a}(\mathbf{C})$, it follows that $v_s = u_0$. Since v_n commutes with $v_s = u_0$, it lies in $\mathfrak{m}(\mathbf{C})$, and we must check that v_n is a principal nilpotent element in $\mathfrak{m}(\mathbf{C})$. As v_n lies in the Borel subalgebra $(\mathfrak{b} \cap \mathfrak{m})(\mathbf{C})$ of $\mathfrak{m}(\mathbf{C})$, it is enough to check that the projection of v_n into each simple root space of $(\mathfrak{b} \cap \mathfrak{m})(\mathbf{C})$ is non-zero, and this follows from the statement (proved above) that the image of v under $\mathfrak{b}(\mathbf{C}) \hookrightarrow \mathfrak{p}_\alpha(\mathbf{C}) \twoheadrightarrow \mathfrak{m}_\alpha(\mathbf{C})$ is regular in $\mathfrak{m}_\alpha(\mathbf{C})$ for every simple root α of A in M .

3.5. End of the proof of the main theorem in case $A = T$. We continue with $u \in \mathfrak{a}(\mathcal{O})$ and $x \in X^u$ as above. Let M be any Levi subgroup containing A . It remains to prove that if x_P is regular in X_M^u for all $P \in \mathcal{P}(M)$ and if

$$(3.5.1) \quad n(x, P, P') = n(u, P, P')$$

for every adjacent pair $P, P' \in \mathcal{P}(M)$, then x is regular in X^u . We have already proved this in case $M = A$, and now we want to reduce the general case to this special case.

The equality (3.5.1) is equivalent to the equality

$$(3.5.2) \quad \nu_M(x_P) - \nu_M(x_{P'}) = n(u, P, P') \cdot \beta_{P, P'}.$$

Fix $P \in \mathcal{P}(M)$ and sum (3.5.2) over the set of neighboring pairs in a minimal gallery joining P to its opposite $\bar{P} \in \mathcal{P}(M)$. Doing this yields the equality

$$(3.5.3) \quad \nu_M(x_P) - \nu_M(x_{\bar{P}}) = \sum_{\alpha \in R_N} \text{val}_F(\alpha(u)) \cdot \pi_M(\alpha^\vee),$$

where $\pi_M : X_*(A) \rightarrow \Lambda_M$ is the canonical surjection and R_N is the set of roots of A in \mathfrak{n} .

Fix a Borel subgroup B_M in M containing A and let B (respectively, B_1) be the Borel subgroups in $\mathcal{P}(A)$ obtained as the inverse image of B_M under $P \rightarrow M$ (respectively, $\bar{P} \rightarrow M$). Then (3.5.3) implies (see 2.3) that

$$\nu_A(x_B) - \nu_A(x_{B_1}) \equiv \sum_{\alpha \in R_N} \text{val}_F(\alpha(u)) \cdot \alpha^\vee$$

modulo the coroot lattice for M . Since R_N is also the set of roots that are positive on B and negative on B_1 , it follows that

$$\nu_A(x_B) - \nu_A(x_{B_1}) = \sum_{\alpha \in R_N} j_\alpha \cdot \alpha^\vee$$

for some integers j_α such that $0 \leq j_\alpha \leq \text{val}_F(\alpha(u))$. (To prove this pick a minimal gallery joining B to B_1 and use the inequality stated in the main theorem for each neighboring pair in the gallery.) Comparing this equality with the congruence, we see that the linear combination

$$(3.5.4) \quad \sum_{\alpha \in R_N} (\text{val}_F(\alpha(u)) - j_\alpha) \cdot \alpha^\vee$$

maps to 0 in Λ_M .

We get a basis for $\Lambda_M \otimes \mathbf{R}$ by taking the elements $\beta_{P,P'}$ as P' varies through the set of parabolic subgroups in $\mathcal{P}(M)$ adjacent to P . Moreover for any $\alpha \in R_N$ the image $\pi_M(\alpha^\vee)$ of α^\vee in Λ_M is a non-negative linear combination of basis elements $\beta_{P,P'}$ (with at least one non-zero coefficient). Therefore the fact that (3.5.4) maps to 0 in Λ_M means that

$$(3.5.5) \quad \nu_A(x_B) - \nu_A(x_{B_1}) = \sum_{\alpha \in R_N} \text{val}_F(\alpha(u)) \cdot \alpha^\vee$$

By hypothesis $x_{\bar{P}}$ is regular. Therefore (transitivity of retractions plus the part of our theorem we have already proved) for all adjacent Borel subgroups $B_1, B_2 \in \mathcal{P}(A)$ such that $B_1, B_2 \subset \bar{P}$ we have

$$\nu_A(x_{B_1}) - \nu_A(x_{B_2}) = \text{val}_F(\alpha_{B_1, B_2}(u)) \cdot \alpha_{B_1, B_2}^\vee,$$

where α_{B_1, B_2} denotes the unique root that is positive on B_1 and negative on B_2 . Summing these equalities over neighboring pairs in a minimal gallery joining B_1 to \bar{B} , we find that

$$\nu_A(x_{B_1}) - \nu_A(x_{\bar{B}}) = \sum_{\alpha \in R_M^+} \text{val}_F(\alpha(u)) \cdot \alpha^\vee,$$

where R_M^+ denotes the set of roots of A in B_M . Adding this last equality to (3.5.5), we see that

$$(3.5.6) \quad \nu_A(x_B) - \nu_A(x_{\bar{B}}) = \sum_{\alpha \in R^+} \text{val}_F(\alpha(u)) \cdot \alpha^\vee.$$

Now consider any minimal gallery $B = B_0, B_1, \dots, B_l = \bar{B}$ joining B to \bar{B} . Then

$$(3.5.7) \quad \nu_A(x_B) - \nu_A(x_{\bar{B}}) = \sum_{i=1}^l n(x, B_{i-1}, B_i) \cdot \alpha_i^\vee,$$

where α_i is the unique root that is positive for B_{i-1} and negative for B_i . We know that $n(x, B_{i-1}, B_i) \leq \text{val}_F(\alpha_i(u))$ for all i . Subtracting (3.5.7) from (3.5.6), we find that 0 is a non-negative linear combination of positive roots; therefore each coefficient in this linear combination is 0, which means that

$$n(x, B_{i-1}, B_i) = \text{val}_F(\alpha_i(u))$$

for $i = 1, \dots, l$.

Now consider any pair B', B'' of adjacent Borel subgroups in $\mathcal{P}(A)$. After reversing the order of B', B'' if necessary we can find a minimal gallery as above and an index i such that $(B_{i-1}, B_i) = (B', B'')$. Therefore

$$(3.5.8) \quad n(x, B', B'') = \text{val}_F(\alpha(u)),$$

where α is the unique root that is positive on B' and negative on B'' . Since both sides of (3.5.8) remain unchanged when B', B'' are switched, we see that (3.5.8) holds for any adjacent pair B', B'' . By what we have already done, it follows that x is regular in X^u .

3.6. Proof of the main theorem in general. Now let M be any Levi subgroup of G containing A , and let u be an integral regular semisimple element of $\mathfrak{g}(F)$ that happens to lie in $\mathfrak{m}(F)$. Let $T = \text{Cent}_{G_F}(u)$, a maximal torus in M_F . We choose a finite extension F'/F that splits T .

We normalize the valuation $\text{val}_{F'}$ on F' so that uniformizing elements in F' have valuation 1. Thus $\text{val}_{F'}(\epsilon) = [F' : F]$. We write X' for the set $G(F')/G(\mathcal{O}_{F'})$. The inclusion $G(F) \hookrightarrow G(F')$ induces a canonical injection $X \hookrightarrow X'$.

For any $P \in \mathcal{P}(M)$ the diagram

$$\begin{array}{ccc} X & \xrightarrow{r_P} & X_M \\ \downarrow & & \downarrow \\ X' & \xrightarrow{r'_P} & X'_M \end{array}$$

commutes, where the horizontal maps are retractions and the vertical maps are the canonical injections. Moreover the diagram

$$\begin{array}{ccc} X & \xrightarrow{\nu_G} & \Lambda_G \\ \downarrow & & \downarrow \\ X' & \xrightarrow{\nu'_G} & \Lambda_G \end{array}$$

commutes, where the left vertical map is the canonical injection and the right vertical map is multiplication by $e := [F' : F]$.

For any $x \in X^u$ the image of x in X' lies in $(X')^u$, and x is regular in X^u if and only if x is regular in $(X')^u$. Indeed the conjugacy class $\bar{u}_G(x)$ attached to u and x is the same for X and X' .

The torus T is conjugate under $M(F')$ to A , so our theorem is true for T over F' . Therefore for $x \in X^u$ and adjacent $P = MN, P' = MN' \in \mathcal{P}(M)$

$$(3.6.1) \quad e \cdot n(x, P, P') \leq \sum_{\alpha \in R_N \cap R_{N'}} \text{val}_{F'}(\alpha(u)) \cdot m_\alpha,$$

and x is regular in X^u if and only if all of these inequalities are equalities. (As before R_N denotes the set of roots of A in \mathfrak{n} ; the positive integers m_α were defined in 2.9.) Dividing by e , and noting that the term indexed by α depends only on the Γ -orbit of α , we find that (3.6.1) is equivalent to the inequality

$$n(x, P, P') \leq n(u, P, P').$$

This completes the proof of the theorem.

REFERENCES

- [Art76] J. Arthur, *The characters of discrete series as orbital integrals*, Invent. Math. **32** (1976), 205–261.
- [Bez96] R. Bezrukavnikov, *The dimension of the fixed point set on affine flag manifolds*, Math. Res. Lett. **3** (1996), 185–189.
- [KL88] D. Kazhdan and G. Lusztig, *Fixed point varieties on affine flag manifolds*, Israel J. Math. **62** (1988), 129–168.
- [Lan80] R. P. Langlands, *Base change for $GL(2)$* , Ann. of Math. Studies 96, Princeton University Press, 1980.

M. GORESKY, SCHOOL OF MATHEMATICS, IAS, PRINCETON, NJ 08540
E-mail address: `goresky@ias.edu`

R. KOTTWITZ, DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CHICAGO, 5734 UNIVERSITY
AVE., CHICAGO, IL 60637
E-mail address: `kottwitz@math.uchicago.edu`

R. MACPHERSON, SCHOOL OF MATHEMATICS, IAS, PRINCETON, NJ 08540
E-mail address: `rdm@ias.edu`