MINIMAL RATIONAL CURVES IN MODULI SPACES OF STABLE BUNDLES

XIAOTAO SUN

Introduction

Let C be a smooth projective curve of genus $g \geq 2$ and \mathcal{L} be a line bundle on C of degree d. Assume that $r \geq 2$ is an integer coprime with d. Let $M := \mathcal{U}_C(r, \mathcal{L})$ be the moduli space of stable vector bundles on C of rank r and with the fixed determinant \mathcal{L} . It is well-known that M is a smooth projective Fano variety with Picard number 1. For any projective curve in M, we can define its degree with respect to the ample anti-canonical line bundle $-K_M$. A natural question raised by Jun-Muk Hwang (see Question 1 in [Hw]) is to determine all rational curves of minimal degree passing through a generic point of M. In this short note, we prove the following theorem

Main Theorem. Assume that $g \geq 4$. Then any rational curve of M passing through a generic point has degree at least 2r. It has degree 2r if and only if it is a Hecke curve.

Our idea is simple and the proof is elementary. If E is a vector bundle on $X = C \times \mathbb{P}^1$ that induces the morphism of \mathbb{P}^1 to M. Then a simple computation shows that its degree equals to the second Chern class of $\mathcal{E}nd(E)$. If the restriction of E to the generic fiber of ruled surface $f: X \to C$ is semistable, then one sees easily that $c_2(\mathcal{E}nd(E))$ is at least 2r, and it is 2r if and only if $c_2(E) = 1$ (after tensoring E by suitable line bundle pulling back from \mathbb{P}^1). This will force E to be an extension

$$0 \to f^*V \xrightarrow{i} E \xrightarrow{\phi} \mathcal{O}_{\{p\} \times \mathbb{P}^1}(-1) \to 0,$$

where V is a bundle on C. That is, after performing elementary transformation on E once along one fiber, E becomes a pullback of a vector bundle on C. For any $x \in \mathbb{P}^1$, restricting above sequence to $C \times \{x\}$ and denote $E|_{C \times \{x\}}$ by E_x , we have

$$0 \to V \xrightarrow{i_x} E_x \xrightarrow{\phi_x} \mathcal{O}_{\{p\} \times \mathbb{P}^1}(-1)_x \to 0.$$

Let $\iota_x: V_p \to E_x|_p = E_{(p,x)}$ be the homomorphism between the fibers at p induced by the sheaf map i_x . Then the Hecke modifications $\{(\widetilde{W}^{ker(\iota_x)})^{\vee}; x \in \mathbb{P}^1\}$ of V along $\{ker(\iota_x) \subset V_p; x \in \mathbb{P}^1\}$ are exactly $\{E_x; x \in \mathbb{P}^1\}$. Thus the given curve is a Hecke curve by definition.

m +1 4 cm v

The work is supported by a grant of NFSC for outstanding young researcher at contract number 10025103.

If the restriction of E to the generic fiber is not semistable, then using the so called relative Harder-Narasimhan filtration we are able to prove that $c_2(\mathcal{E}nd(E)) > 2r$. Here we also need the condition that the rational curve passes through the generic point of M which corresponds a (1,1)-stable bundle when $g \geq 4$.

§1 Hecke curves

Since we are discussing the question raised in [Hw], we copy the definition of Hecke curves in [Hw] and adopt notation there. Given two nonnegative integers k, ℓ , a vector bundle W of rank r and degree d on C is (k,ℓ) -stable, if, for each proper subbundle W' of W, we have

$$\frac{deg(W') + k}{rk(W')} < \frac{deg(W) + k - \ell}{r}.$$

The usual stability is equivalent to (0,0)-stability. The dual bundle of a (k,ℓ) -stable bundle is (ℓ,k) -stable.

Lemma 1.1 ([NR]). If $g \ge 4$, a generic point $[W] \in M$ corresponds to a (1,1)-stable bundle W.

Lemma 1.2 ([NR]). Let $0 \to V \to W \to \mathcal{O}_p \to 0$ be an exact sequence, where \mathcal{O}_p is the 1-dimensional skyscraper sheaf at $p \in C$. If W is (k,ℓ) -stable, then V is $(k,\ell-1)$ -stable.

Let $[W] \in M$ be a generic point corresponding to a (1,1)-stable bundle W over C. We will use V^{\vee} to denote the dual vector bundle (or dual vector space) of a vector bundle V (or a vector space V). Let $\mathbb{P}(W)$ be the projective bundle consisting of lines through the origin on each fiber. For $p \in C$ and $\zeta \in \mathbb{P}(W_p^{\vee})$, define a vector bundle W^{ζ} by

$$(1.1) 0 \to W^{\zeta} \to W \to (W_p/\zeta^{\perp}) \otimes \mathcal{O}_p \to 0$$

where ζ^{\perp} denotes the hyperplane in W_p annihilated by ζ . Let $\iota: W_p^{\zeta} \to W_p$ be the homomorphism between the fibers at p induced by the sheaf injection $W^{\zeta} \to W$. The kernel $ker(\iota)$ of ι is a 1-dimensional subspace of W_p^{ζ} and its annihilator $ker(\iota)^{\perp}$ is a hyperplane in $(W^{\zeta})_p^{\vee}$. Let \mathcal{H} be a line in $\mathbb{P}(W_p^{\zeta})$ containing the point $[ker(\iota)]$. For each point $[l] \in \mathcal{H}$ corresponding to a 1-dimensional subspace $l \subset W_p^{\zeta}$, define a vector bundle \widetilde{W}^l by

$$(1.2) 0 \to \widetilde{W}^l \to (W^\zeta)^\vee \to ((W^\zeta)_p^\vee/l^\perp)) \otimes \mathcal{O}_p \to 0$$

where $l^{\perp} \subset (W^{\zeta})_p^{\vee}$ is the hyperplane annihilating l. This bundle \widetilde{W}^l is stable for each $[l] \in \mathcal{H}$ by Lemma 1.2. It is easy to check that for $l = ker(\iota)$,

$$(1.3) \widetilde{W}^{ker(\iota)} \cong W^{\vee}.$$

Thus $\{(\widetilde{W}^l)^{\vee}; l \in \mathcal{H}\}$ defines a rational curve passing through $[W] \in M$. A rational curve on M constructed in this way is called a Hecke curve. By using [NR], it can be shown that a Hecke curve is smooth and has degree 2r with respect to $-K_M$. There is an equivalent description for $(\widetilde{W}^l)^{\vee}$. For any vector bundle, say W^{ζ} , on C and a subspace K of the fiber W_p^{ζ} $(p \in C)$. According to [NS], there are two canonical constructions called Hecke modifications (see Remark 2.4 of [NS]). For any $[l] \in \mathcal{H}$, the stable bundle $(\widetilde{W}^l)^{\vee}$ is in fact obtained from W^{ζ} by performing the second Hecke modification along the subspace $l \subset W_p^{\zeta}$.

§2 Proof of the Main Theorem

For any rational curve $\mathbb{P}^1 \subset M$ through a general point of M, let E be the vector bundle on $X := C \times \mathbb{P}^1$, which induces the embedding $\mathbb{P}^1 \subset M$. Let $\pi : X = C \times \mathbb{P}^1 \to \mathbb{P}^1$ be the projection and $\mathbb{E} \subset \mathcal{E}nd(E)$ be the subbundle of trace free. Then, since $\pi_*(\mathbb{E}) = 0$, we have $T_M|_{\mathbb{P}^1} = R^1\pi_*\mathbb{E}$ and, by using Leray spectral sequence and Riemann-Roch theorem,

$$-\chi(\mathbb{E}) = \chi(R^1 \pi_* \mathbb{E}) = -K_M \cdot \mathbb{P}^1 + (r^2 - 1)(g - 1).$$

By using $\chi(\mathbb{E}) = deg(ch(\mathbb{E}) \cdot td(T_X))_2$, noting $c_1(\mathbb{E}) = c_1(\mathcal{E}nd(E)) = 0$, we get

$$(2.1) -K_M \cdot \mathbb{P}^1 = c_2(\mathbb{E}) = 2rc_2(E) - (r-1)c_1(E)^2 := \Delta(E).$$

Let $f: X = C \times \mathbb{P}^1 \to C$ be the projection. Then, for any torsion free sheaf E on the ruled surface X, its restriction to a generic fiber $f^{-1}(\xi) = X_{\xi}$ has the form

$$E|_{X_{\xi}} = \bigoplus_{i=1}^{n} \mathcal{O}_{X_{\xi}}(\alpha_i)^{\oplus r_i}, \quad \alpha_1 > \cdots > \alpha_n.$$

The $\alpha = (\alpha_1^{\oplus r_1}, ..., \alpha_n^{\oplus r_n})$ is called the generic splitting type of E. In our case, tensoring E by $\pi^*\mathcal{O}(-\alpha_n)$, we can (and we will) assume that $\alpha_n = 0$. Any such E admits a relative Hardar-Narasimhan filtration

$$0 = E_0 \subset E_1 \subset \cdots \subset E_n = E$$

of which the quotient sheaves $F_i = E_i/E_{i-1}$ are torsion free with generic splitting type $(\alpha_i^{\oplus r_i})$ respectively. Then it is easy to see that

$$2c_2(E) = 2\sum_{i=1}^n c_2(F_i) + 2\sum_{i=1}^n c_1(E_{i-1})c_1(F_i)$$
$$= 2\sum_{i=1}^n c_2(F_i) + c_1(E)^2 - \sum_{i=1}^n c_1(F_i)^2.$$

Thus
$$\Delta(E) = 2r \sum_{i=1}^{n} c_2(F_i) + c_1(E)^2 - r \sum_{i=1}^{n} c_1(F_i)^2$$
.

Let $F'_i = F_i \otimes \pi^* \mathcal{O}_{\mathbb{P}^1}(-\alpha_i)$ (i = 1, ..., n), thus they have generic splitting type $(0^{\oplus r_i})$ respectively. Let $c_1(F_i) = f^* \mathcal{O}_C(d_i) + \pi^* \mathcal{O}_{\mathbb{P}^1}(r_i\alpha_i)$, where $\mathcal{O}_C(d_i)$, $\mathcal{O}_{\mathbb{P}^1}(r_i\alpha_i)$ are divisors of degree d_i , $r_i\alpha_i$ on C, \mathbb{P}^1 respectively. Here we remark that for any torsion free sheaf F_i on X we have $c_1(F_i)|_{f^{-1}(\bullet)} = c_1(F_i|_{f^{-1}(\bullet)})$ for general points on C (resp. $c_1(F_i)|_{\pi^{-1}(\bullet)} = c_1(F_i|_{\pi^{-1}(\bullet)})$ for general points on \mathbb{P}^1). Therefore d_i are the degrees of F_i on the general fiber of π respectively. Without confusion, we denote the degree of F_i (resp. E_i) on the generic fiber of π by $deg(F_i)$ (resp. $deg(E_i)$). Consequently, $\mu(E_i)$, $\mu(E)$ denote the slope of restrictions of E_i , E to the generic fiber of π respectively. Note that

$$c_2(F_i') = c_2(F_i) - (r_i - 1)c_1(F_i)\pi^*\mathcal{O}_{\mathbb{P}^1}(\alpha_i) = c_2(F_i) - (r_i - 1)d_i\alpha_i,$$

 $c_1(F_i)^2 = 2r_i d_i \alpha_i$ and $c_1(E)^2 = 2d \sum_{i=1}^n r_i \alpha_i$, we have

$$\Delta(E) = 2r \left(\sum_{i=1}^{n} c_2(F_i') + \mu(E) \sum_{i=1}^{n} r_i \alpha_i - \sum_{i=1}^{n} d_i \alpha_i \right).$$

Let $rk(E_i)$ denote the rank of E_i , note that $r_i = rk(E_i) - rk(E_{i-1})$ and $d_i = deg(E_i) - deg(E_{i-1})$, we have

(2.2)
$$\Delta(E) = 2r \left(\sum_{i=1}^{n} c_2(F_i') + \sum_{i=1}^{n-1} (\mu(E) - \mu(E_i))(\alpha_i - \alpha_{i+1})rk(E_i) \right).$$

Lemma 2.1. Any torsion free sheaf \mathcal{E} of rank r on a ruled surface, with generic splitting type $(0^{\oplus r})$, must have $c_2(\mathcal{E}) \geq 0$.

Proof. The argument is in fact contained in the proof of Lemma 1.4 of [GL]. One can choose a divisor D on C such that $f^*\mathcal{O}_C(D) \subset \mathcal{E}$ and $\mathcal{E}/f^*\mathcal{O}_C(D)$ is torsion free. Since $\mathcal{E}/f^*\mathcal{O}_C(D)$ has generic splitting type $(0^{\oplus (r-1)})$, by induction hypothesis on rank, we can assume that $c_2(\mathcal{E}/f^*\mathcal{O}_C(D)) \geq 0$. Hence

$$c_2(\mathcal{E}) = c_2(\mathcal{E}/f^*\mathcal{O}_C(D)) + f^{-1}(D)(c_1(\mathcal{E}) - f^{-1}(D)) = c_2(\mathcal{E}/f^*\mathcal{O}_C(D)) \ge 0.$$

Proposition 2.2. If $g \ge 4$, then $\Delta(E) \ge 2r$. The equality holds if and only if E has generic splitting type $(0^{\oplus r})$ and $c_2(E) = 1$.

Proof. Since the rational curve passes through a generic point $[W] \in M$, then the bundle $E|_{\pi^{-1}([W])} \cong W$ is (1,1)-stable. Thus, using the inequality (2.2), we have

$$\Delta(E) \ge 2r \left(\sum_{i=1}^{n} c_2(F_i') + \sum_{i=1}^{n-1} (\alpha_i - \alpha_{i+1}) \right),$$

the equality holds if and only if n = 1.

By Lemma 2.1, if n > 1, we have

$$\Delta(E) > 2r \sum_{i=1}^{n-1} (\alpha_i - \alpha_{i+1}) \ge 2r.$$

If n = 1, then $\Delta(E) = 2rc_2(E) \ge 2r$ (note that we have assumed $\alpha_n = 0$). The equality $\Delta(E) = 2r$ holds if and only if $c_2(E) = 1$.

From now on, we assume that E has generic splitting type $(0^{\oplus r})$. If E has a jumping line $X_p = f^{-1}(p)$ $(p \in C)$, i.e.,

$$E|_{X_p} = \bigoplus_{i=1}^n \mathcal{O}_{X_p}(\beta_i)^{\oplus r_i}, \quad \beta_1 > \dots > \beta_n$$

with the type $(\beta_1^{\oplus r_1}, \dots, \beta_n^{\oplus r_n})$ different from $(0^{\oplus r})$. Then we can perform the elementary transformation on E along X_p by taking F to be the kernel of the (unique surjective) homomorphism $\phi: E \to E|_{X_p} \to \mathcal{O}_{X_p}(\beta_n)^{\oplus r_n}$. Clearly,

$$(2.3) 0 \to F \to E \xrightarrow{\phi} \mathcal{O}_{X_p}(\beta_n)^{\oplus r_n} \to 0.$$

An easy calculation yields

Lemma 2.3.
$$c_1(F) = c_1(E) - r_n X_p$$
 and $c_2(F) = c_2(E) + r_n \beta_n$.

Proof. By the exact sequence (2.3), the computation is straightforward.

Lemma 2.4. If $c_2(E) = 1$ and E has generic splitting type $(0^{\oplus r})$, then E has exactly one jumping line X_p and the elementary transformation F along X_p is isomorphic to f^*V for a vector bundle V over C.

Proof. The E has at least one jumping line. Otherwise, E will be a pullback of a vector bundle over C, which is impossible. At any jumping line X_p , with splitting type $(\beta_1^{\oplus r_1}, \dots, \beta_n^{\oplus r_n})$, we must have $\beta_n < 0$. Hence, by Lemma 2.3 and Lemma 2.1, E has a unique jumping line X_p with $\beta_n = -1$ and $r_n = 1$. Then F has no jumping line, thus $F = f^*V$ for a vector bundle V over C.

Therefore by Proposition 2.2 and Lemma 2.4, if $\Delta(E) = 2r$, we have

$$(2.4) 0 \to f^*V \to E \xrightarrow{\phi} \mathcal{O}_{X_n}(-1) \to 0.$$

Proposition 2.5. If $g \geq 4$ and $\Delta(E) = 2r$, then the rational curve is a Hecke curve.

Proof. For any $x \in \mathbb{P}^1$, let E_x denote $E|_{C \times \{x\}}$. Restrict the sequence (2.4) to $\pi^{-1}(x) = C \times \{x\}$, we get

$$(2.5) 0 \to V \to E_x \xrightarrow{\phi_x} \mathcal{O}_{X_n}(-1)_x \to 0.$$

Since the curve passes through the generic point $[W] \in M$, there is a $x_0 \in \mathbb{P}^1$ such that $E_{x_0} \cong W$ is (1,1)-stable by Lemma 1.1. Then V is a (1,0)-stable bundle by Lemma 1.2 (thus when $x = x_0$, the sequence (2.5) corresponds to the sequence (1.1) and V corresponds to W^{ζ} in the definition of Hecke curves, see $\delta 1$)

Let $\iota_x: V_p \to E_x|_p = E_{(p,x)}$ be the homomorphism between the fibers at p induced by the sheaf injection $V \to E_x$ in sequence (2.5). Then the kernel $ker(\iota_x)$ is a 1-dimensional subspace of V_p . When x moves on \mathbb{P}^1 , these $[ker(\iota_x)] \in \mathbb{P}(V_p)$ form a line $\mathcal{H} \subset \mathbb{P}(V_p)$. It is easy to check that, as the same as (1.3), for any $x \in \mathbb{P}^1$

$$\widetilde{W}^{ker(\iota_x)} \cong E_x^{\vee}.$$

Thus $\{(\widetilde{W}^{ker(\iota_x)})^{\vee}; [ker(\iota_x)] \in \mathcal{H}\}$ defines the given rational curve. That is, the given rational curve is a Hecke curve.

Theorem 2.6. Assume that $g \ge 4$. Then any rational curve of M passing through the generic point of M has at least degree 2r with respect to $-K_M$. It has degree 2r if and only if it is a Hecke curve.

Proof. By (2.1), the degree $-K_M \cdot \mathbb{P}^1$ equals to $\Delta(E)$. Then, by Proposition 2.2, it has degree at least 2r. If it has degree 2r, then by Proposition 2.5 it must be a Hecke curve. It was known that any Hecke curve has degree 2r. We are done

REFERENCES

- [GL] D. Gieseker and J. Li, Moduli of high rank vector bundles over surfaces, J. Amer. Math. Soc. 9 (1996), 107–151.
- [Hw] J.-M. Hwang, Hecke curves on the moduli space of vector bundles over an algebraic curve, Proceedings of the Symposium Algebraic Geometry in East Asia, Kyoto (2001), 155–164.
- [NR] M. S. Narasimhan and S. Ramanan, Geometry of Hecke cycles I., in C. P. Ramanujama tribute. Springer Verlag, (1978), 291–345.
- [NS] D.S. Nagaraj and C.S. Seshadri, Degenerations of the moduli spaces of vector bundles on curves I, Proc. Indian Acad. Sci.(Math. Sci.) 107 (1997), 101–137.

INSTITUTE OF MATHEMATICS, CHINESE ACADEMY OF SCIENCES, BEIJING 100080, CHINA *E-mail address*: xsun@math08.math.ac.cn

DEPARTMENT OF MATHEMATICS, THE UNIVERSITY OF HONG KONG, POKFULAM ROAD, HONG KONG

E-mail address: xsun@maths.hku.hk