

# On singular Artin monoids

EDDY GODELLE AND LUIS PARIS

February 8, 2021

## Abstract

In this paper we study some combinatorial aspects of the singular Artin monoids. Firstly, we show that a singular Artin monoid  $SA$  can be presented as a semidirect product of a graph monoid with its associated Artin group  $A$ . Such a decomposition implies that a singular Artin monoid embeds in a group. Secondly, we give a solution to the word problem for the FC type singular Artin monoids. Afterwards, we show that FC type singular Artin monoids have the FRZ property. Briefly speaking, this property says that the centralizer in  $SA$  of any non-zero power of a standard singular generator  $\tau_s$  coincides with the centralizer of any non-zero power of the corresponding non-singular generator  $\sigma_s$ . Finally, we prove Birman's conjecture, namely, that the desingularization map  $\eta : SA \rightarrow \mathbb{Z}[A]$  is injective, for right-angled singular Artin monoids.

**AMS Subject Classification:** Primary 20F36.

## 1 Introduction

Let  $S$  be a finite set. Recall that a *Coxeter matrix* over  $S$  is a matrix  $M = (m_{st})_{s,t \in S}$  indexed by the elements of  $S$  and such that  $m_{ss} = 1$  for all  $s \in S$ , and  $m_{st} = m_{ts} \in \{2, 3, 4, \dots, +\infty\}$  for all  $s, t \in S$ ,  $s \neq t$ . A Coxeter matrix  $M = (m_{st})$  is usually represented by its *Coxeter graph*,  $\Gamma$ , which is defined as follows.  $S$  is the set of vertices of  $\Gamma$ , two vertices  $s, t$  are joined by an edge if  $m_{st} \geq 3$ , and this edge is labelled by  $m_{st}$  if  $m_{st} \geq 4$ . If  $a, b$  are two letters and  $m \in \mathbb{Z}_{\geq 2}$ , then we denote by  $w(m : a, b)$  the word  $\dots bab$  of length  $m$ . We take an abstract set  $\mathcal{S} = \{\sigma_s; s \in S\}$  in one-to-one correspondence with  $S$ , and we define the *Artin group* associated to  $\Gamma$  to be the group  $A = A_\Gamma$  presented by

$$A = A_\Gamma = \langle \mathcal{S} \mid w(m_{st} : \sigma_s, \sigma_t) = w(m_{st} : \sigma_t, \sigma_s) \text{ for } s, t \in S, s \neq t \text{ and } m_{st} < +\infty \rangle.$$

The *Coxeter group* associated to  $\Gamma$  is the quotient  $W = W_\Gamma$  of  $A$  by the relations  $\sigma_s^2 = 1$ ,  $s \in S$ .

Take  $X \subset S$  and put  $\mathcal{S}_X = \{\sigma_s; s \in X\}$ . We denote by  $A_X$  the subgroup of  $A$  generated by  $\mathcal{S}_X$ , and by  $W_X$  the subgroup of  $W$  generated by  $\mathcal{S}_X$ . Let  $\Gamma_X$  be the full

subgraph of  $\Gamma$  generated by  $X$ . Then  $A_X$  is the Artin group associated to  $\Gamma_X$  (see [32] and [34]), and  $W_X$  is the Coxeter group associated to  $\Gamma_X$  (see [10]). The subgroup  $A_X$  is called *standard parabolic subgroup* of  $A$ , and  $W_X$  is called *standard parabolic subgroup* of  $W$ .

We say that  $\Gamma$  (or  $A$ ) is of *spherical type* if  $W$  is finite, that  $\Gamma$  (or  $A$ ) is *right-angled* if  $m_{st} \in \{2, +\infty\}$  for all  $s, t \in S$ ,  $s \neq t$ , and that  $\Gamma$  (or  $A$ ) is of *type FC* if, for all  $X \subset S$ , either  $W_X$  is finite, or there exist  $s, t \in X$ ,  $s \neq t$ , such that  $m_{st} = +\infty$ . Note that the spherical type Artin groups as well as the right-angled Artin groups are both FC type Artin groups. The number  $n = |S|$  is called the *rank* of  $A$ .

The first (non-abelian) example of Artin group which has appeared in the literature is certainly the braid group  $\mathcal{B}_n$  introduced by Artin [3] in 1925. One of the most important works in the subject is a paper by Garside [23] where the word problem and the conjugacy problem for  $\mathcal{B}_n$  are solved. Garside's ideas have been extended to all spherical type Artin groups by Brieskorn, Saito [12], and Deligne [17] in 1972. These two papers, [12] and [17], are the foundation of the theory of Artin groups, and, more specifically, of the spherical type Artin groups. Right-angled Artin groups are also known as *graph groups* or as *free partially commutative groups*. They have been widely studied, and their applications extend to various domains like parallel computation, random walks, and cohomology of groups. We mention, for example, the paper [8] where a group which is  $FP_2$  but not finitely presented is constructed as a subgroup of some right-angled Artin group. Artin groups of type FC have been introduced by Charney and Davis [15] in 1995 in their study of the  $K(\pi, 1)$ -problem for complements of infinite hyperplane arrangements associated to reflection groups.

In the same way as the braid group  $\mathcal{B}_n$  has been extended to the singular braid monoid  $S\mathcal{B}_n$  (see [5] and [9]), the Artin groups can be extended to the singular Artin monoids as follows. Take a new abstract set  $\mathcal{T} = \{\tau_s; s \in S\}$  in one-to-one correspondence with  $S$ , and define the *singular Artin monoid* associated to  $\Gamma$  to be the monoid  $SA = SA_\Gamma$  presented as a monoid by the generating set  $\mathcal{S} \cup \mathcal{S}^{-1} \cup \mathcal{T}$ , where  $\mathcal{S}^{-1} = \{\sigma_s^{-1}; s \in S\}$ , and by the relations

$$\begin{aligned} \sigma_s \sigma_s^{-1} &= \sigma_s^{-1} \sigma_s = 1, & \text{for } s \in S, \\ \sigma_s \tau_s &= \tau_s \sigma_s, & \text{for } s \in S, \\ w(m_{st} : \sigma_s, \sigma_t) &= w(m_{st} : \sigma_t, \sigma_s), & \text{for } s, t \in S, s \neq t, \text{ and } m_{st} < +\infty, \\ w(m_{st} - 1 : \sigma_s, \sigma_t) \tau_s &= \tau_{s \wedge t} w(m_{st} - 1 : \sigma_s, \sigma_t), & \text{for } s, t \in S, s \neq t, \text{ and } m_{st} < +\infty, \\ \tau_s \tau_t &= \tau_t \tau_s, & \text{for } s, t \in S, s \neq t, \text{ and } m_{st} = 2, \end{aligned}$$

where  $s \wedge t = s$  if  $m_{st}$  is even, and  $s \wedge t = t$  if  $m_{st}$  is odd. Observe that we have an epimorphism  $\theta : SA \rightarrow A$  which sends  $\sigma_s^{\pm 1}$  to  $\sigma_s^{\pm 1}$  and  $\tau_s$  to  $\sigma_s$  for all  $s \in S$ , and that this epimorphism has a section  $\iota : A \rightarrow SA$  which sends  $\sigma_s^{\pm 1}$  to  $\sigma_s^{\pm 1}$  for all  $s \in S$ . In particular,  $A$  embeds in  $SA$ .

The combinatorial study of the singular Artin monoids (of spherical type) has been initiated by Corran [16] with techniques inspired from [12].

The purpose of the present paper is to study different combinatorial aspects of the monoid  $SA$ .

Firstly, in Section 2, we prove that  $SA$  can be decomposed as a semidirect product of a so-called graph monoid with the Artin group  $A$ . A consequence of this decomposition shall be that  $SA$  embeds in a group. Note that this last result has been previously proved by Basset [7] and Keyman [30] with completely different proofs.

Sections 3 and 4 concern only singular Artin monoids of type FC. We solve the word problem for  $SA$  in Section 3. In [21], Fenn, Rolfsen, and Zhu proved that the centralizer in the singular braid monoid of a standard singular generator  $\tau_j$  is equal to the centralizer of any non-zero power of  $\tau_j$ , and that this centralizer coincides with the centralizer of any non-zero power of  $\sigma_j$ . This property, which we like to call *FRZ property*, is of importance in the study of singular braids, and, in particular, in the proof of Birman's conjecture for braid groups (see [36]). In Section 4, we extend the FRZ property to all singular Artin monoids of type FC.

Define the *desingularization map* as the multiplicative homomorphism  $\eta : SA \rightarrow \mathbb{Z}[A]$  which sends  $\sigma_s^{\pm 1}$  to  $\sigma_s^{\pm 1}$  and  $\tau_s$  to  $\sigma_s - \sigma_s^{-1}$  for all  $s \in S$ . One of the main questions in the subject, known as *Birman's conjecture*, is to determine whether the desingularization map is injective. This is known to be true for braid groups [36] and for rank 2 Artin groups [20]. We prove Birman's conjecture for right-angled Artin groups in Section 5.

The last section is dedicated to some questions for which we do not have any significant result but that deserve to be mentioned.

## 2 Semidirect product structure

Our purpose in this section is to determine a decomposition of  $SA$  as a semidirect product of a so-called graph monoid with the Artin group  $A$ . A first consequence of this decomposition is that a singular Artin monoid embeds in a group. This decomposition shall be also used in Sections 3 and 4 to solve the word problem and to prove the "FRZ property" for singular Artin monoids of type FC.

Let  $G$  be a (standard) graph, let  $V$  be its set of vertices, and let  $E = E(G)$  be its set of edges. Define the *graph monoid* of  $G$  to be the monoid  $\mathcal{M}(G)$  presented as a monoid by

$$\mathcal{M}(G) = \langle V \mid uv = vu \text{ if } \{u, v\} \in E \rangle^+.$$

The first (standard) graph that we shall consider is the graph  $\hat{\Omega}$  defined by the following data.

- $\hat{Y} = \{\alpha\tau_s\alpha^{-1}; \alpha \in A \text{ and } s \in S\}$  is the set of vertices of  $\hat{\Omega}$ ;
- $\{\hat{u}, \hat{v}\}$  is an edge of  $\hat{\Omega}$  if  $\hat{u}\hat{v} = \hat{v}\hat{u}$  in  $SA$ .

**Proposition 2.1.** *We have  $SA = \mathcal{M}(\hat{\Omega}) \rtimes A$ .*

**Proof.** We have a homomorphism  $f : \mathcal{M}(\hat{\Omega}) \rtimes A \rightarrow SA$  defined by  $f(\alpha) = \alpha \in SA$  for all  $\alpha \in A$ , and  $f(\hat{u}) = \hat{u} \in SA$  for all  $\hat{u} \in \hat{Y}$ . Conversely, One can easily verify using the presentation of  $SA$  that there is a homomorphism  $g : SA \rightarrow \mathcal{M}(\hat{\Omega}) \rtimes A$  which sends  $\sigma_s^{\pm 1}$  to  $\sigma_s^{\pm 1} \in A$ , and  $\tau_s$  to  $\tau_s \in \hat{Y}$ , for all  $s \in S$ . Obviously,  $f \circ g = \text{Id}$  and  $g \circ f = \text{Id}$ .  $\square$

**Corollary 2.2.** *SA embeds in a group.*

**Proof.** Let  $\mathcal{G}(\hat{\Omega})$  be the group presented by  $\mathcal{G}(\hat{\Omega}) = \langle \hat{\Upsilon} \mid \hat{u}\hat{v} = \hat{v}\hat{u} \text{ if } \{\hat{u}, \hat{v}\} \in E(\hat{\Omega}) \rangle$ . Then  $\mathcal{M}(\hat{\Omega})$  embeds in  $\mathcal{G}(\hat{\Omega})$  (see [19] and [18]), thus  $SA = \mathcal{M}(\hat{\Omega}) \rtimes A$  embeds in  $\mathcal{G}(\hat{\Omega}) \rtimes A$ .  $\square$

### 3 The word problem

Note that a solution to the word problem for  $SA$  will also give a solution to the word problem for the Artin group  $A$ . So, a reasonable approach would be to study the word problem for those singular Artin monoids whose associated Artin groups have known solutions to the word problem. In the case of Artin groups of type FC, a solution has been found by Altobelli [2]. Another observation is that a given element of a graph monoid  $\mathcal{M}(G)$  has finitely many representatives and these representatives can be easily listed. Other solutions to the word problem for  $\mathcal{M}(G)$  can be found in [13], [41], and [19]. Now, assume that  $A$  is of type FC and consider the decomposition  $SA = \mathcal{M}(\hat{\Omega}) \rtimes A$  of the previous section. By the above observations, in order to solve the word problem for  $SA$ , it suffices to find an algorithm which decides whether two elements  $\alpha\tau_s\alpha^{-1}$  and  $\beta\tau_t\beta^{-1}$  of  $\hat{\Upsilon}$  are equal, and, if not, whether they commute or not. Such an algorithm can be easily derived from Proposition 3.1 below together with Altobelli's solution to the word problem for  $A$ .

Define the graph  $\Omega$  as follows.

- $\Upsilon = \{\alpha\sigma_s\alpha^{-1}; \alpha \in A \text{ and } s \in S\}$  is the set of vertices of  $\Omega$ ;
- $\{u, v\}$  is an edge of  $\Omega$  if  $uv = vu$  in  $A$ .

**Proposition 3.1.** *Assume  $\Gamma$  to be of type FC. Then there exists an isomorphism  $\varphi : \hat{\Omega} \rightarrow \Omega$  which sends  $\alpha\tau_s\alpha^{-1}$  to  $\alpha\sigma_s\alpha^{-1}$  for all  $\alpha \in A$  and all  $s \in S$ .*

The remainder of the section is dedicated to the proof of Proposition 3.1.

Define the *Artin monoid* associated to  $\Gamma$  to be the monoid  $A^+ = A_\Gamma^+$  presented as a monoid by

$$A^+ = \langle \mathcal{S} \mid w(m_{st} : \sigma_s, \sigma_t) = w(m_{st} : \sigma_t, \sigma_s) \text{ for } s, t \in S, s \neq t, \text{ and } m_{st} < +\infty \rangle^+.$$

By [35], the natural homomorphism  $A^+ \rightarrow A$  which sends  $\sigma_s$  to  $\sigma_s$  for all  $s \in S$  is injective. We can define the *length function*  $\text{lg} : A^+ \rightarrow \mathbb{N}$  which associates to each element of  $A^+$  the length of any of its representatives with respect to the generating set  $\mathcal{S}$ . Since the defining relations of  $A^+$  are homogeneous, this function is well-defined and is a homomorphism of monoids. For the same reason, we can define a partial order  $\leq_R$  on  $A^+$  by setting  $a \leq_R b$  if there exists  $c \in A^+$  such that  $ca = b$ . Now, the following proposition is a mixture of several well-known facts on spherical type Artin groups.

**Proposition 3.2.** *Assume  $\Gamma$  to be of spherical type.*

(1) **(Brieskorn-Saito [12], Deligne [17]).**  $(A^+, \leq_R)$  is a lattice. The lattice operations of  $(A^+, \leq_R)$  are denoted by  $\wedge_R$  and  $\vee_R$ .

(2) **(Brieskorn-Saito [12], Deligne [17]).** Let  $s, t \in S$ ,  $s \neq t$ . Then  $\sigma_s \vee_R \sigma_t = w(m_{st} : \sigma_s, \sigma_t) = w(m_{st} : \sigma_t, \sigma_s)$ .

(3) **(Brieskorn-Saito [12], Deligne [17]).** Let  $\Delta = \vee_R \{\sigma_s; s \in S\}$ . Then there exists a permutation  $\mu : S \rightarrow S$  such that  $\mu^2 = \text{Id}$  and  $\Delta \sigma_s \Delta^{-1} = \sigma_{\mu(s)}$  for all  $s \in S$ .

(4) **(Brieskorn-Saito [12], Deligne [17]).** Each  $\alpha \in A$  can be written as  $\alpha = a\Delta^k$  with  $a \in A^+$  and  $k \in \mathbb{Z}$ .

(5) **(Charney [14]).** Each  $\alpha \in A$  can be uniquely written as  $\alpha = ab^{-1}$  with  $a, b \in A^+$  and  $a \wedge_R b = 1$ . Such an expression  $\alpha = ab^{-1}$  is called the Charney form of  $\alpha$ .

(6) **(Charney [14]).** Let  $\alpha \in A$  and  $u, v \in A^+$  such that  $\alpha = uv^{-1}$ . Then  $u = ac$  and  $v = bc$ , where  $c = u \wedge_R v$  and  $ab^{-1}$  is the Charney form of  $\alpha$ .  $\square$

Let  $s, t \in S$  and  $\omega \in A$ . We say that  $\omega$  is an *elementary positive  $(t, s)$ -ribbon* if either

- $s = t$ , and  $\omega = \sigma_s$ ; or
- $s = t$ , and there exists  $r \in S$  such that  $m_{sr}$  is even and  $\omega = w(m_{sr} - 1 : \sigma_s, \sigma_r)$ ; or
- $s \neq t$ ,  $m_{st}$  is odd, and  $\omega = w(m_{st} - 1 : \sigma_s, \sigma_t)$ .

Define an *elementary  $(t, s)$ -ribbon* to be either an elementary positive  $(t, s)$ -ribbon, or the inverse of an elementary positive  $(s, t)$ -ribbon. Note that, if  $\omega$  is an elementary  $(t, s)$ -ribbon, then  $\omega\sigma_s = \sigma_t\omega$  and  $\omega\tau_s = \tau_t\omega$ . We say that  $\omega$  is a  *$(t, s)$ -ribbon* if there exist a sequence  $s_0 = s, s_1, \dots, s_p = t$  in  $S$ , and a sequence  $\omega_1, \dots, \omega_p$  in  $A$ , such that  $\omega_i$  is an elementary  $(s_i, s_{i-1})$ -ribbon for all  $i = 1, \dots, p$ , and  $\omega = \omega_p \dots \omega_2 \omega_1$ . Clearly, if  $\omega$  is a  $(t, s)$ -ribbon, then  $\omega\sigma_s = \sigma_t\omega$  and  $\omega\tau_s = \tau_t\omega$ .

The key point in the proof of Proposition 3.1 is the following result which can be found in [25] (see also [24]).

**Proposition 3.3 (Godelle [25]).** *Assume  $\Gamma$  to be of type FC. Let  $s \in S$ ,  $X \subset S$ , and  $\alpha \in A$ . Then the followings are equivalent.*

- (1)  $\alpha\sigma_s\alpha^{-1} \in A_X$ .
- (2) There exists  $k \in \mathbb{Z} \setminus \{0\}$  such that  $\alpha\sigma_s^k\alpha^{-1} \in A_X$ .
- (3) There exist  $t \in X$ ,  $\omega \in A$ , and  $\beta \in A_X$ , such that  $\omega$  is a  $(t, s)$ -ribbon and  $\alpha = \beta\omega$ .

$\square$

**Corollary 3.4 (Godelle [25]).** *Assume  $\Gamma$  to be of type FC. Let  $s, t \in S$  and  $\alpha \in A$ . Then the followings are equivalent.*

- (1)  $\alpha\sigma_s\alpha^{-1} = \sigma_t$ .
- (2) There exists  $k \in \mathbb{Z} \setminus \{0\}$  such that  $\alpha\sigma_s^k\alpha^{-1} = \sigma_t^k$ .
- (3)  $\alpha$  is a  $(t, s)$ -ribbon.  $\square$

**Lemma 3.5.** *Assume  $\Gamma$  to be of type FC. Then there exists a bijection  $\varphi : \hat{\Upsilon} \rightarrow \Upsilon$  which sends  $\alpha\tau_s\alpha^{-1}$  to  $\alpha\sigma_s\alpha^{-1}$  for all  $\alpha \in A$  and all  $s \in S$ .*

**Proof.** We take  $\alpha, \beta \in A$  and  $s, t \in S$ , and we turn to prove the following equivalence.

$$\alpha\tau_s\alpha^{-1} = \beta\tau_t\beta^{-1} \quad \Leftrightarrow \quad \alpha\sigma_s\alpha^{-1} = \beta\sigma_t\beta^{-1}.$$

Assume first that  $\alpha\tau_s\alpha^{-1} = \beta\tau_t\beta^{-1}$ . Recall the epimorphism  $\theta : SA \rightarrow A$  which sends  $\sigma_s^{\pm 1}$  to  $\sigma_s^{\pm 1}$  and  $\tau_s$  to  $\sigma_s$  for all  $s \in S$ . Then

$$\alpha\sigma_s\alpha^{-1} = \theta(\alpha\tau_s\alpha^{-1}) = \theta(\beta\tau_t\beta^{-1}) = \beta\sigma_t\beta^{-1}.$$

Now, assume that  $\alpha\sigma_s\alpha^{-1} = \beta\sigma_t\beta^{-1}$ . By Corollary 3.4,  $\beta^{-1}\alpha$  is a  $(t, s)$ -ribbon, thus  $\beta^{-1}\alpha\tau_s = \tau_t\beta^{-1}\alpha$ , and therefore  $\alpha\tau_s\alpha^{-1} = \beta\tau_t\beta^{-1}$ .  $\square$

**Lemma 3.6.** *Assume  $\Gamma$  to be of spherical type. Then the bijection  $\varphi : \hat{\Upsilon} \rightarrow \Upsilon$  extends to an isomorphism  $\varphi : \hat{\Omega} \rightarrow \Omega$ .*

**Proof.** We take  $\alpha, \beta \in A$  and  $s, t \in S$ , and we turn to prove the following equivalence.

$$(\alpha\tau_s\alpha^{-1})(\beta\tau_t\beta^{-1}) = (\beta\tau_t\beta^{-1})(\alpha\tau_s\alpha^{-1}) \quad \Leftrightarrow \quad (\alpha\sigma_s\alpha^{-1})(\beta\sigma_t\beta^{-1}) = (\beta\sigma_t\beta^{-1})(\alpha\sigma_s\alpha^{-1}).$$

First, assume that  $(\alpha\tau_s\alpha^{-1})(\beta\tau_t\beta^{-1}) = (\beta\tau_t\beta^{-1})(\alpha\tau_s\alpha^{-1})$ . Then

$$(\alpha\sigma_s\alpha^{-1})(\beta\sigma_t\beta^{-1}) = \theta((\alpha\tau_s\alpha^{-1})(\beta\tau_t\beta^{-1})) = \theta((\beta\tau_t\beta^{-1})(\alpha\tau_s\alpha^{-1})) = (\beta\sigma_t\beta^{-1})(\alpha\sigma_s\alpha^{-1}).$$

Now, we assume that  $(\alpha\sigma_s\alpha^{-1})(\beta\sigma_t\beta^{-1}) = (\beta\sigma_t\beta^{-1})(\alpha\sigma_s\alpha^{-1})$ , and we prove that  $(\alpha\tau_s\alpha^{-1})(\beta\tau_t\beta^{-1}) = (\beta\tau_t\beta^{-1})(\alpha\tau_s\alpha^{-1})$ . Our proof is divided into 3 steps.

*Step 1:* Assume  $\beta = 1$ ,  $\alpha = a \in A^+$ , and  $a\sigma_s \wedge_R a = 1$ . So,  $a\sigma_s a^{-1}$  is a Charney form. Since  $\sigma_t a \sigma_s a^{-1} \sigma_t^{-1} = a\sigma_s a^{-1}$ , by Proposition 3.2, there exists  $c \in A^+$  such that  $\sigma_t a = ac$  and  $\sigma_t a \sigma_s = a\sigma_s c$ . The element  $c$  is clearly of length 1, namely,  $c = \sigma_r$  for some  $r \in S$ , and, by Corollary 3.4, the equality  $\sigma_t a = a\sigma_r$  implies that  $a$  is a  $(t, r)$ -ribbon. Moreover, we have  $a\sigma_r \sigma_s = \sigma_t a \sigma_s = a\sigma_s \sigma_r$ , thus  $\sigma_r \sigma_s = \sigma_s \sigma_r$ , therefore  $m_{sr} = 2$ . So,

$$\tau_t a \tau_s a^{-1} = a \tau_r \tau_s a^{-1} = a \tau_s \tau_r a^{-1} = a \tau_s a^{-1} \tau_t.$$

*Step 2:* Assume  $\beta = 1$  and  $\alpha = a \in A^+$ . We argue by induction on the length of  $a$ . The case  $a\sigma_s \wedge_R a = 1$  is treated in Step 1, thus we can suppose that  $a\sigma_s \wedge_R a \neq 1$ . In particular, there exists some  $r \in S$  such that  $\sigma_r \leq_R a\sigma_s \wedge_R a$ . Suppose  $r = s$ . Then  $a$  can be written as  $a = a_1\sigma_s$  with  $a_1 \in A^+$ , and, moreover,  $\sigma_t a_1 \sigma_s a_1^{-1} = a_1 \sigma_s a_1^{-1} \sigma_t$ . By the inductive hypothesis, it follows that

$$\tau_t a \tau_s a^{-1} = \tau_t a_1 \tau_s a_1^{-1} = a_1 \tau_s a_1^{-1} \tau_t = a \tau_s a^{-1} \tau_t.$$

Suppose  $r \neq s$ . We have  $\sigma_s, \sigma_r \leq_R a\sigma_s$ , thus, by Proposition 3.2,  $\sigma_s \vee_R \sigma_r = w(m_{sr} : \sigma_r, \sigma_s) \leq_R a\sigma_s$ , therefore  $\omega = w(m_{sr} - 1 : \sigma_s, \sigma_r) \leq_R a$ . Write  $u = s$  if  $m_{sr}$  is even, and  $u = r$  if  $m_{sr}$  is odd. Then  $\omega$  is an (elementary)  $(u, s)$ -ribbon and  $a$  can be written as

$a = a_1\omega$  with  $a_1 \in A^+$ . Moreover,  $\sigma_t a_1 \sigma_u a_1^{-1} = \sigma_t a \sigma_s a^{-1} = a \sigma_s a^{-1} \sigma_t = a_1 \sigma_u a_1^{-1} \sigma_t$ . By the inductive hypothesis, it follows that

$$\tau_t a \tau_s a^{-1} = \tau_t a_1 \tau_u a_1^{-1} = a_1 \tau_u a_1^{-1} \tau_t = a \tau_s a^{-1} \tau_t.$$

*Step 3:* General case. By Proposition 3.2, there exist  $a \in A^+$  and  $k \in \mathbb{Z}$  such that  $\beta^{-1}\alpha = a\Delta^k$ . Recall the permutation  $\mu : S \rightarrow S$  such that  $\Delta\sigma_r\Delta^{-1} = \sigma_{\mu(r)}$  for all  $r \in S$ . Let  $r = \mu^k(s)$ . Then  $\Delta^k$  is a  $(r, s)$ -ribbon (by Corollary 3.4), and, moreover,  $\sigma_t a \sigma_r a^{-1} = \sigma_t (\beta^{-1}\alpha\sigma_s\alpha^{-1}\beta) = (\beta^{-1}\alpha\sigma_s\alpha^{-1}\beta)\sigma_t = a\sigma_r a^{-1}\sigma_t$ . By Step 2, it follows that

$$\tau_t(\beta^{-1}\alpha\tau_s\alpha^{-1}\beta) = \tau_t a \tau_r a^{-1} = a \tau_r a^{-1} \tau_t = (\beta^{-1}\alpha\tau_s\alpha^{-1}\beta)\tau_s,$$

hence  $(\beta\tau_t\beta^{-1})(\alpha\tau_s\alpha^{-1}) = (\alpha\tau_s\alpha^{-1})(\beta\tau_t\beta^{-1})$ .  $\square$

In order to extend this result to all Artin groups of type FC (namely, in order to prove Proposition 3.1), we need one more preliminary result (on amalgamated products).

**Proposition 3.7 (Serre [39]).** *Let  $G = G_1 *_H G_2$  be the amalgamated product of two groups  $G_1$  and  $G_2$  over  $H$ . Let  $C_1$  and  $C_2$  be transversals of  $G_1/H$  and  $G_2/H$ , respectively, which contain 1. For all  $g \in G$  there exists a unique sequence  $(g_1, \dots, g_l, h)$  such that*

- $g = g_1 g_2 \dots g_l h$ ;
- $h \in H$ , and either  $g_i \in C_1 \setminus \{1\}$  or  $g_i \in C_2 \setminus \{1\}$  for all  $i = 1, \dots, l$ ;
- $g_i \in C_1 \Rightarrow g_{i+1} \in C_2$ , and  $g_i \in C_2 \Rightarrow g_{i+1} \in C_1$ , for all  $i = 1, \dots, l-1$ .  $\square$

The above sequence  $(g_1, \dots, g_l, h)$  is called the *amalgam normal form* of  $g$  (relative to the amalgamated product  $G_1 *_H G_2$ ). The number  $l$  is called the *amalgam norm* of  $g$  (relative to the amalgamated product  $G_1 *_H G_2$ ) and is denoted by  $l = |g|_*$ .

**Proof of Proposition 3.1.** We take  $\alpha, \beta \in A$  and  $s, t \in S$ , and we turn to prove the following equivalence.

$$(\alpha\tau_s\alpha^{-1})(\beta\tau_t\beta^{-1}) = (\beta\tau_t\beta^{-1})(\alpha\tau_s\alpha^{-1}) \quad \Leftrightarrow \quad (\alpha\sigma_s\alpha^{-1})(\beta\sigma_t\beta^{-1}) = (\beta\sigma_t\beta^{-1})(\alpha\sigma_s\alpha^{-1}).$$

The implication  $\Rightarrow$  can be proved exactly in the same manner as the implication  $\Rightarrow$  in the proof of Lemma 3.6. So, we assume that  $(\alpha\sigma_s\alpha^{-1})(\beta\sigma_t\beta^{-1}) = (\beta\sigma_t\beta^{-1})(\alpha\sigma_s\alpha^{-1})$ , and we prove that  $(\alpha\tau_s\alpha^{-1})(\beta\tau_t\beta^{-1}) = (\beta\tau_t\beta^{-1})(\alpha\tau_s\alpha^{-1})$ . We argue by induction on the rank of  $A$  (*i.e.* on the number of vertices of  $\Gamma$ ). Up to changing  $\alpha$  by  $\beta^{-1}\alpha$ , we can also assume that  $\beta = 1$ .

If  $m_{xy} < +\infty$  for all  $x, y \in S$ ,  $x \neq y$ , then  $A$  is of spherical type and, therefore, the equality  $(\alpha\tau_s\alpha^{-1})\tau_t = \tau_t(\alpha\tau_s\alpha^{-1})$  follows from Lemma 3.6. So, we can assume that there exist  $x, y \in S$ ,  $x \neq y$ , such that  $m_{xy} = +\infty$ . Let  $X = S \setminus \{x\}$  and  $Y = S \setminus \{y\}$ . Then  $A = A_X *_A_{X \cap Y} A_Y$ . We choose transversals  $C_X$  and  $C_Y$  of  $A_X/A_{X \cap Y}$  and  $A_Y/A_{X \cap Y}$ , respectively, which contain 1, we consider the amalgam normal form  $(\alpha_1, \dots, \alpha_l, \gamma)$  of  $\alpha$ , and we argue by induction on  $l = |\alpha|_*$ .

Assume  $l = 0$ . So,  $\alpha = \gamma \in A_{X \cap Y}$ . The fact that the amalgam normal forms of  $\sigma_t(\alpha\sigma_s\alpha^{-1})$  and  $(\alpha\sigma_s\alpha^{-1})\sigma_t$  are equal implies that either  $\sigma_s, \sigma_t \in A_X$  (namely  $s, t \in X$ ), or  $\sigma_s, \sigma_t \in A_Y$  (namely,  $s, t \in Y$ ). Say  $s, t \in X$ . We have  $s, t \in X$  and  $\alpha \in A_X$ , thus, by the inductive hypothesis,  $\tau_t(\alpha\tau_s\alpha^{-1}) = (\alpha\tau_s\alpha^{-1})\tau_t$ .

Now, we assume that  $l = |\alpha|_* \geq 1$ . Without loss of generality, we can assume that  $\alpha_l \in C_X$ . Then  $\alpha_1 \in C_Z$ , where  $Z = Y$  if  $l$  is even, and  $Z = X$  if  $l$  is odd. We consider 4 different cases.

*Case 1:*  $\alpha_l\gamma\sigma_s\gamma^{-1}\alpha_l^{-1} \in A_{X \cap Y}$ . By Proposition 3.3,  $\alpha_l\gamma$  can be written as  $\alpha_l\gamma = \gamma_1\omega$  where  $\omega$  is a  $(r, s)$ -ribbon for some  $r \in X \cap Y$ , and  $\gamma_1 \in A_{X \cap Y}$ . Let  $\alpha' = \alpha_1\alpha_2 \dots \alpha_{l-1}\gamma_1 = \alpha\omega^{-1}$ . We have  $(\alpha'\sigma_r\alpha'^{-1})\sigma_t = (\alpha\sigma_s\alpha^{-1})\sigma_t = \sigma_t(\alpha\sigma_s\alpha^{-1}) = \sigma_t(\alpha'\sigma_r\alpha'^{-1})$ , thus, by the inductive hypothesis (on the amalgam norm of  $\alpha$ ), we have  $(\alpha\tau_s\alpha^{-1})\tau_t = (\alpha'\tau_r\alpha'^{-1})\tau_t = \tau_t(\alpha'\tau_r\alpha'^{-1}) = \tau_t(\alpha\tau_s\alpha^{-1})$ .

*Case 2:*  $l = 1$ ,  $s \in X$ , and  $\alpha\sigma_s\alpha^{-1} \in A_X \setminus A_{X \cap Y}$ . The fact that the amalgam normal forms of  $\sigma_t(\alpha\sigma_s\alpha^{-1})$  and  $(\alpha\sigma_s\alpha^{-1})\sigma_t$  are equal implies that  $\sigma_t \in A_X$ , namely, that  $t \in X$ . We have  $s, t \in X$  and  $\alpha \in A_X$ , thus, by the inductive hypothesis (on the rank of  $A$ ), we have  $\tau_t(\alpha\tau_s\alpha^{-1}) = (\alpha\tau_s\alpha^{-1})\tau_t$ .

*Case 3:*  $l \geq 2$ ,  $s \in X$ , and  $\alpha_l\gamma\sigma_s\gamma^{-1}\alpha_l^{-1} \in A_X \setminus A_{X \cap Y}$ . Then the amalgam normal form of  $\alpha\sigma_s\alpha^{-1}$  has the form  $(\alpha_1, \dots, \alpha_{l-1}, \beta'_1, \beta'_2, \dots, \beta'_l, \gamma_1)$ , and  $\alpha_1, \beta'_i \in C_Z$ . The fact that the amalgam normal forms of  $\sigma_t(\alpha\sigma_s\alpha^{-1})$  and  $(\alpha\sigma_s\alpha^{-1})\sigma_t$  are equal implies that  $\sigma_t \in A_Z$ , namely, that  $t \in Z$ . Then the amalgam normal form of  $(\alpha\sigma_s\alpha^{-1})\sigma_t$  has either the form  $(\alpha_1, \dots, \alpha_{l-1}, \beta'_1, \beta'_2, \dots, \beta'_{l-1}, \gamma_2)$  if  $\beta'_l\gamma_1\sigma_t \in A_{X \cap Y}$ , or the form  $(\alpha_1, \dots, \alpha_{l-1}, \beta'_1, \beta'_2, \dots, \beta'_{l-1}, \beta''_l, \gamma_2)$  if  $\beta'_l\gamma_1\sigma_t \notin A_{X \cap Y}$ . Now,  $\sigma_t(\alpha\sigma_s\alpha^{-1})$  has also this amalgam normal form, thus  $\alpha_1 A_{X \cap Y} = \sigma_t \alpha_1 A_{X \cap Y}$ , namely,  $\alpha_1^{-1} \sigma_t \alpha_1 \in A_{X \cap Y}$ . By Proposition 3.3, we deduce that  $\alpha_1$  can be written as  $\alpha_1 = \omega\delta$ , where  $\omega$  is a  $(t, r)$ -ribbon for some  $r \in X \cap Y$ , and  $\delta \in A_{X \cap Y}$ . Let  $\alpha' = \delta\alpha_2 \dots \alpha_l\gamma = \omega^{-1}\alpha$ . We have  $\sigma_r(\alpha'\sigma_s\alpha'^{-1}) = (\alpha'\sigma_s\alpha'^{-1})\sigma_r$ , thus, by the inductive hypothesis (on the amalgam norm of  $\alpha$ ), we have  $\tau_r(\alpha'\tau_s\alpha'^{-1}) = (\alpha'\tau_s\alpha'^{-1})\tau_r$ , therefore  $\tau_t(\alpha\tau_s\alpha^{-1}) = (\alpha\tau_s\alpha^{-1})\tau_t$ .

*Case 4:*  $l \geq 1$  and  $s = x \notin X$ . Then the amalgam normal form of  $\alpha\sigma_s\alpha^{-1}$  has the form  $(\alpha_1, \dots, \alpha_{l-1}, \alpha_l, \beta'_0, \beta'_1, \dots, \beta'_l, \gamma_1)$ . Applying the same argument as in Case 3, we conclude that  $\tau_t(\alpha\tau_s\alpha^{-1}) = (\alpha\tau_s\alpha^{-1})\tau_t$ .  $\square$

## 4 The FRZ property

The aim of this section is to prove the following.

**Proposition 4.1.** *Assume  $\Gamma$  to be of type FC. Let  $\alpha \in SA$  and  $s, t \in S$ . Then the followings are equivalent.*

- (1)  $\alpha\sigma_s = \sigma_t\alpha$ .
- (2) There exists  $k \in \mathbb{Z} \setminus \{0\}$  such that  $\alpha\sigma_s^k = \sigma_t^k\alpha$ .

(3)  $\alpha\tau_s = \tau_t\alpha$ .

(4) There exists  $k \in \mathbb{N} \setminus \{0\}$  such that  $\alpha\tau_s^k = \tau_t^k\alpha$ .

The following lemma is a preliminary result to the proof of Proposition 4.1.

**Lemma 4.2.** *Let  $G$  be a graph, let  $V$  be its set of vertices, and let  $E$  be its set of edges. Let  $u_1, \dots, u_l, v, w \in V$  and  $p \in \mathbb{N} \setminus \{0\}$  such that  $v^p u_1 u_2 \dots u_l = u_1 u_2 \dots u_l w^p$  (in  $\mathcal{M}(G)$ ). Then  $v = w$ , and  $u_i v = v u_i$  for all  $i = 1, \dots, l$ .*

**Proof.** Observe that, if  $H$  is a full subgraph of  $G$ , then there is an epimorphism  $f_H : \mathcal{M}(H) \rightarrow \mathcal{M}(G)$  which sends the vertices of  $H$  to themselves and sends the other vertices of  $G$  to 1. First, applying this observation to the graph  $H = \{v\}$  with one vertex,  $v$ , and no edge, we deduce that  $v = w$ . Now, we argue by induction on  $l$ . Suppose that  $u_1 \neq v$  and that  $u_1$  and  $v$  are not joined by an edge. Let  $H$  be the full subgraph of  $G$  generated by  $\{u_1, v\}$ . Then  $\mathcal{M}(H)$  is the free monoid freely generated by  $\{u_1, v\}$ ,  $f_M(v^p u_1 \dots u_l)$  is a word which starts with  $v$ , and the word  $f_M(u_1 \dots u_l v^p)$  starts with  $u_1$ : a contradiction. So, either  $u_1 = v$  or  $\{u_1, v\} \in E$ , that is,  $u_1 v = v u_1$ . It follows also that  $v^p u_2 \dots u_l = u_2 \dots u_l v^p$ , and, by the inductive hypothesis, we conclude that  $u_i v = v u_i$  for all  $i = 2, \dots, l$ .  $\square$

**Proof of Proposition 4.1.** The implications (1) $\Rightarrow$ (2) and (3) $\Rightarrow$ (4) are obvious, thus it remains to prove (2) $\Rightarrow$ (3) and (4) $\Rightarrow$ (1).

*Proof of (2) $\Rightarrow$ (3).* We assume that  $\alpha\sigma_s^k = \sigma_t^k\alpha$  for some  $k \in \mathbb{Z} \setminus \{0\}$ . Recall the epimorphism  $\theta : SA \rightarrow A$  which sends  $\sigma_s^{\pm 1}$  to  $\sigma_s^{\pm 1}$  and  $\tau_s$  to  $\sigma_s$  for all  $s \in S$ . Let  $\alpha_0 = \theta(\alpha) \in A$ . Then  $\alpha_0\sigma_s^k = \sigma_t^k\alpha_0$ , thus, by Corollary 3.4,  $\alpha_0$  is a  $(t, s)$ -ribbon, therefore  $\alpha_0\tau_s = \tau_t\alpha_0$ . Let  $\beta = \alpha_0^{-1}\alpha$ . Then  $\beta\sigma_s^k = \sigma_s^k\beta$ . Recall the decomposition  $SA = \mathcal{M}(\hat{\Omega}) \rtimes A$ , and write  $\beta = u_1 u_2 \dots u_l \beta_0$  where  $u_1, \dots, u_l \in \hat{\Upsilon}$  and  $\beta_0 \in A$ . We have  $\sigma_s^k \beta \sigma_s^{-k} = u'_1 u'_2 \dots u'_l \beta'_0$ , where  $u'_i = \sigma_s^k u_i \sigma_s^{-k} \in \hat{\Upsilon}$  for all  $i = 1, \dots, l$ , and  $\beta'_0 = \sigma_s^k \beta_0 \sigma_s^{-k} \in A$ . The equality  $\sigma_s^k \beta \sigma_s^{-k} = \beta$  implies that  $\beta'_0 = \beta_0$ , and that there exists a permutation  $\chi \in \text{Sym}_l$  such that  $u'_i = u_{\chi(i)}$  for all  $i = 1, \dots, l$ . Firstly, the equality  $\beta'_0 = \sigma_s^k \beta_0 \sigma_s^{-k} = \beta_0$  implies by Corollary 3.4 that  $\beta_0$  is a  $(s, s)$ -ribbon, thus  $\beta_0\tau_s = \tau_s\beta_0$ . Now, let  $p$  be the order of  $\chi$ . Then  $\sigma_s^{pk} u_i \sigma_s^{-pk} = u_i$  for all  $i = 1, \dots, l$ . Take  $i \in \{1, \dots, l\}$  and write  $u_i = \gamma_i \tau_{r_i} \gamma_i^{-1}$  where  $\gamma_i \in A$  and  $r_i \in S$ . The element  $\sigma_s^{pk}$  commutes with  $\gamma_i \tau_{r_i} \gamma_i^{-1} = u_i$ , thus  $\sigma_s^{pk}$  commutes with  $\gamma_i \sigma_{r_i} \gamma_i^{-1} = \theta(\gamma_i \tau_{r_i} \gamma_i^{-1})$ , therefore  $\sigma_s$  commutes with  $\gamma_i \sigma_{r_i} \gamma_i^{-1}$  (by Corollary 3.4), and hence, by Proposition 3.1,  $\tau_s$  commutes with  $\gamma_i \tau_{r_i} \gamma_i^{-1} = u_i$ . This shows that  $\tau_s$  commutes with  $\beta = u_1 \dots u_l \beta_0$ , and we conclude that  $\alpha\tau_s = \alpha_0\beta\tau_s = \alpha_0\tau_s\beta = \tau_t\alpha_0\beta = \tau_t\alpha$ .

*Proof of (4) $\Rightarrow$ (1).* We assume that  $\alpha\tau_s^k = \tau_t^k\alpha$  for some  $k \in \mathbb{N} \setminus \{0\}$ . We write  $\alpha = u_1 u_2 \dots u_l \alpha_0$ , where  $u_1, \dots, u_l \in \hat{\Upsilon}$  and  $\alpha_0 \in A$ . Then the equality  $\alpha\tau_s^k = \tau_t^k\alpha$  implies that we have  $u_1 u_2 \dots u_l (\alpha_0 \tau_s \alpha_0^{-1})^k = \tau_t^k u_1 u_2 \dots u_l$  in  $\mathcal{M}(\hat{\Omega})$ . By Lemma 4.2, it follows that  $\alpha_0 \tau_s \alpha_0^{-1} = \tau_t$ , and  $\tau_t u_i = u_i \tau_t$  for all  $i = 1, \dots, l$ . The equality  $\alpha_0 \tau_s \alpha_0^{-1} = \tau_t$  implies that  $\alpha_0 \sigma_s \alpha_0^{-1} = \theta(\alpha_0 \tau_s \alpha_0^{-1}) = \theta(\tau_t) = \sigma_t$ . Now, take  $i \in \{1, \dots, l\}$  and write  $u_i = \gamma_i \tau_{r_i} \gamma_i^{-1}$  where  $\gamma_i \in A$  and  $r_i \in S$ . The element  $\tau_t$  commutes with  $u_i = \gamma_i \tau_{r_i} \gamma_i^{-1}$ ,

thus  $\sigma_t = \theta(\tau_t)$  commutes with  $\gamma_i \sigma_{r_i} \gamma_i^{-1} = \theta(u_i)$ , that is,  $\sigma_t \gamma_i \sigma_{r_i} \gamma_i^{-1} \sigma_t^{-1} = \gamma_i \sigma_{r_i} \gamma_i^{-1}$ , hence, by Lemma 3.5,  $\sigma_t u_i \sigma_t^{-1} = \sigma_t \gamma_i \tau_{r_i} \gamma_i^{-1} \sigma_t^{-1} = \gamma_i \tau_{r_i} \gamma_i^{-1} = u_i$ . This shows that  $\alpha \sigma_s = u_1 \dots u_l \alpha_0 \sigma_s = u_1 \dots u_l \sigma_t \alpha_0 = \sigma_t u_1 \dots u_l \alpha_0 = \sigma_t \alpha$ .  $\square$

## 5 Birman's conjecture

The purpose of this section is to prove Birman's conjecture for right-angled Artin groups. As these groups are the same as graph groups, for convenience, we shall use the terminology of graph groups and graph products of groups.

Let  $G$  be a graph, let  $V$  be its set of vertices, and let  $E$  be its set of edges. Take an (abstract) set  $\mathcal{S} = \{\sigma_u; u \in V\}$  in one-to-one correspondence with  $V$ , and define the *graph group* associated to  $G$  to be the group presented by

$$\mathcal{G}(G) = \langle \mathcal{S} \mid \sigma_u \sigma_v = \sigma_v \sigma_u \text{ for } \{u, v\} \in E \rangle.$$

Take another abstract set  $\mathcal{T} = \{\tau_u; u \in V\}$  in one-to-one correspondence with  $V$ , and define the *singular graph monoid* associated to  $G$  to be the monoid  $S\mathcal{G}(G)$  presented as a monoid by the generating set  $\mathcal{S} \cup \mathcal{S}^{-1} \cup \mathcal{T}$  and by the relations

$$\begin{aligned} \sigma_u \sigma_u^{-1} = \sigma_u^{-1} \sigma_u = 1, \quad \sigma_u \tau_u = \tau_u \sigma_u, & \quad \text{for } u \in V, \\ \sigma_u \sigma_v = \sigma_v \sigma_u, \quad \sigma_u \tau_v = \tau_v \sigma_u, \quad \tau_u \tau_v = \tau_v \tau_u, & \quad \text{for } \{u, v\} \in E. \end{aligned}$$

The *desingularization map* is defined in this context as the multiplicative homomorphism  $\eta : S\mathcal{G}(G) \rightarrow \mathbb{Z}[\mathcal{G}(G)]$  which sends  $\sigma_u^{\pm 1}$  to  $\sigma_u^{\pm 1}$  and  $\tau_u$  to  $\sigma_u - \sigma_u^{-1}$  for all  $u \in V$ .

**Theorem 5.1.** *The desingularization map  $\eta : S\mathcal{G}(G) \rightarrow \mathbb{Z}[\mathcal{G}(G)]$  is injective.*

We begin with some definitions and results on graph products of groups.

Suppose given a group (or a monoid)  $K_u$  for each  $u \in V$ . Then the *graph product* of the family  $\{K_u\}_{u \in V}$  along the graph  $G$  is the quotient  $\mathcal{G}(\{K_u\}_{u \in V}, G)$  of the free product  $*_{u \in V} K_u$  by the relations

$$g_u g_v = g_v g_u \quad \text{for } \{u, v\} \in E, \quad g_u \in K_u, \quad g_v \in K_v.$$

Note that, if  $K_u = \mathbb{N}$  for all  $u \in V$ , then  $\mathcal{G}(\{K_u\}, G) = \mathcal{M}(G)$ , if  $K_u = \mathbb{Z}$  for all  $u \in V$ , then  $\mathcal{G}(\{K_u\}, G) = \mathcal{G}(G)$ , and if  $K_u = \mathbb{Z} \times \mathbb{N}$  for all  $u \in V$ , then  $\mathcal{G}(\{K_u\}, G) = S\mathcal{G}(G)$ .

Let  $\mathcal{G}(\{K_u\}, G)$  be a graph product of groups (or monoids). Let  $g \in \mathcal{G}(\{K_u\}, G)$ . An *expression* for  $g$  is a sequence  $W = (g_1, g_2, \dots, g_l)$  such that

- there exists  $u_i \in V$  such that  $g_i \in K_{u_i} \setminus \{1\}$  for all  $i = 1, \dots, l$ ,
- $g = g_1 g_2 \dots g_l$ .

The *support* of  $W$  is the sequence  $\text{Supp}(W) = (u_1, u_2, \dots, u_l)$ , and the *length* of  $W$  is  $|W| = l$ . The minimal length for an expression for  $g$  is called the *syllable length* of  $g$  and is denoted by  $|g|_G$ . An expression  $W$  of  $g$  is called *reduced* if its length is equal to the length of  $g$ .

Let  $g \in \mathcal{G}(\{K_u\}, G)$ , let  $W = (g_1, \dots, g_l)$  be an expression for  $g$ , and let  $(u_1, u_2, \dots, u_l)$  be the support of  $W$ . Suppose there exists  $i \in \{1, \dots, l-1\}$  such that  $u_i = u_{i+1}$ , and put

$$W' = \begin{cases} (\dots, g_{i-1}, g_{i+2}, \dots) & \text{if } g_i g_{i+1} = 1, \\ (\dots, g_{i-1}, g_i g_{i+1}, g_{i+2}, \dots) & \text{if } g_i g_{i+1} \neq 1. \end{cases}$$

We say that  $W'$  is obtained from  $W$  via an *elementary M-operation of type I*. This operation shortens the length of an expression by 1 or 2. Suppose there exists  $i \in \{1, \dots, l-1\}$  such that  $\{u_i, u_{i+1}\} \in E$ , and put

$$W'' = (\dots, g_{i-1}, g_{i+1}, g_i, g_{i+2}, \dots).$$

We say that  $W''$  is obtained from  $W$  via an *elementary M-operation of type II*. This operation leaves the length of an expression unchanged. We say that an expression  $W$  is *M-reduced* if its length cannot be reduced applying a sequence of elementary M-operations.

The following proposition is essentially proved in [27] (see also [28] and [29]).

**Proposition 5.2 (Green [27]).** (1) *Let  $g \in \mathcal{G}(\{K_u\}, G)$ , and let  $W_1$  and  $W_2$  be two M-reduced expressions for  $g$ . Then  $W_1$  and  $W_2$  are related by a sequence of elementary M-operations of type II.*

(2) *Let  $g \in \mathcal{G}(\{K_u\}, G)$ , and let  $W$  be an expression for  $g$ . Then  $W$  is M-reduced if and only if  $W$  is reduced.*  $\square$

Now, assume that the set  $V$  of vertices is endowed with a total order,  $\leq$ . Let  $g \in \mathcal{G}(\{K_u\}, G)$ , and let  $W$  be a reduced expression for  $g$ . We say that  $W$  is a *normal form* for  $g$  if  $\text{Supp}(W)$  is the smallest support of a reduced expression for  $g$  with respect to the lexicographic order.

**Corollary 5.3.** (1) *Each element of  $\mathcal{G}(\{K_u\}, G)$  has a unique normal form.*

(2) *To be a normal form depends only on its support, namely, if  $W$  is the normal form for some  $g \in \mathcal{G}(\{K_u\}, G)$ , if  $W'$  is some expression for some  $g' \in \mathcal{G}(\{K_u\}, G)$ , and if  $\text{Supp}(W') = \text{Supp}(W)$ , then  $W'$  is the normal form for  $g'$ .*  $\square$

We return to the study of the graph group  $\mathcal{G}(G)$ . Consider the homomorphism  $\text{deg} : \mathcal{G}(G) \rightarrow \mathbb{Z}$  which sends  $\sigma_u$  to 1 for all  $u \in V$ . For  $n \in \mathbb{Z}$ , we put  $\mathcal{G}_n(G) = \{g \in \mathcal{G}(G); \text{deg}(g) \geq n\}$ , and we denote by  $\mathcal{A}_n$  the free  $\mathbb{Z}$ -module freely generated by  $\mathcal{G}_n(G)$ . Let  $\mathcal{A} = \mathbb{Z}[\mathcal{G}(G)]$ . Then  $\{\mathcal{A}_n\}_{n \in \mathbb{Z}}$  is a filtration of  $\mathcal{A}$  compatible with the multiplication, that is,

- $\mathcal{A}_n \subset \mathcal{A}_m$  if  $n \geq m$ ,
- $\mathcal{A}_p \mathcal{A}_q \subset \mathcal{A}_{p+q}$  for  $p, q \in \mathbb{Z}$ ,
- $1 \in \mathcal{A}_0$ .

Moreover, this filtration is separate, namely,

- $\bigcap_{n \in \mathbb{Z}} \mathcal{A}_n = \{0\}$ .

We denote by  $\tilde{\mathcal{A}}$  the completion of  $\mathcal{A}$ . For  $n \in \mathbb{Z}$ , we put  $\mathcal{G}^{(n)}(G) = \{g \in \mathcal{G}(G); \deg(g) = n\}$ , and we denote by  $\mathcal{A}^{(n)}$  the free  $\mathbb{Z}$ -module freely generated by  $\mathcal{G}^{(n)}(G)$ . Then each element of  $\tilde{\mathcal{A}}$  can be uniquely written as a formal series  $\sum_{n=k}^{+\infty} P_n$ , where  $k \in \mathbb{Z}$  (which may be negative), and  $P_n \in \mathcal{A}^{(n)}$  for all  $n \geq k$ .

Take a free abelian group  $K_u \simeq \mathbb{Z} \times \mathbb{Z}$  of rank 2 generated by  $\{\sigma_u, \tau_u\}$  for all  $u \in V$ , and write  $\tilde{\mathcal{G}} = \mathcal{G}(\{K_u\}, G)$ . Let  $\mathcal{U}(\tilde{\mathcal{A}})$  be the group of unities of  $\tilde{\mathcal{A}}$ . We have a homomorphism  $\tilde{\eta} : \tilde{\mathcal{G}} \rightarrow \mathcal{U}(\tilde{\mathcal{A}})$  defined by

$$\tilde{\eta}(\sigma_u) = \sigma_u, \quad \tilde{\eta}(\tau_u) = \sigma_u - \sigma_u^{-1}, \quad \text{for } u \in V.$$

Note that

$$\tilde{\eta}(\tau_u^{-1}) = - \sum_{n=0}^{+\infty} \sigma_u^{2n+1}, \quad \text{for } u \in V.$$

Note also that  $S\mathcal{G}(G)$  is a submonoid of  $\tilde{\mathcal{G}}$ , that  $\mathcal{A} = \mathbb{Z}[\mathcal{G}(G)]$  is a subalgebra of  $\tilde{\mathcal{A}}$ , and that the restriction of  $\tilde{\eta}$  to  $S\mathcal{G}(G)$  is the desingularization map  $\eta : S\mathcal{G}(G) \rightarrow \mathbb{Z}[\mathcal{G}(G)]$ . So, Theorem 5.1 is a consequence of the following.

**Proposition 5.4.** *The homomorphism  $\tilde{\eta} : \tilde{\mathcal{G}} \rightarrow \mathcal{U}(\tilde{\mathcal{A}})$  is injective.*

**Proof.** For  $u \in V$  and  $(p, q) \in \mathbb{Z} \times \mathbb{Z}$ ,  $(p, q) \neq (0, 0)$ , we write

$$(\sigma_u - \sigma_u^{-1})^p \sigma_u^q = \sum_{n=k}^{+\infty} c_{npq} \sigma_u^n.$$

Note that the numbers  $c_{npq}$  do not depend on  $u$ , but only on  $n, p$ , and  $q$ . Note also that there always exists some  $a \geq k$  (which may be negative) such that  $a \neq 0$  and  $C_{apq} \neq 0$ .

Let  $\tilde{g} \in \tilde{\mathcal{G}}$ ,  $\tilde{g} \neq 1$ . Let  $(\tau_{u_1}^{p_1} \sigma_{u_1}^{q_1}, \tau_{u_2}^{p_2} \sigma_{u_2}^{q_2}, \dots, \tau_{u_l}^{p_l} \sigma_{u_l}^{q_l})$  be the normal form for  $\tilde{g}$ . We have

$$\begin{aligned} \tilde{\eta}(\tilde{g}) &= (\sigma_{u_1} - \sigma_{u_1}^{-1})^{p_1} \sigma_{u_1}^{q_1} (\sigma_{u_2} - \sigma_{u_2}^{-1})^{p_2} \sigma_{u_2}^{q_2} \dots (\sigma_{u_l} - \sigma_{u_l}^{-1})^{p_l} \sigma_{u_l}^{q_l} \\ &= \sum_{n_1 \geq k_1, \dots, n_l \geq k_l} c_{n_1 p_1 q_1} c_{n_2 p_2 q_2} \dots c_{n_l p_l q_l} \sigma_{u_1}^{n_1} \sigma_{u_2}^{n_2} \dots \sigma_{u_l}^{n_l}. \end{aligned}$$

By the above observations, we can find  $a_1, \dots, a_l \in \mathbb{Z} \setminus \{0\}$  such that  $c_{a_i p_i q_i} \neq 0$  for all  $i = 1, \dots, l$ . We turn to show that  $\sigma_{u_1}^{n_1} \sigma_{u_2}^{n_2} \dots \sigma_{u_l}^{n_l} \neq \sigma_{u_1}^{a_1} \sigma_{u_2}^{a_2} \dots \sigma_{u_l}^{a_l}$  for any  $l$ -tuple  $(n_1, \dots, n_l)$  in  $\mathbb{Z}^l$  different from  $(a_1, \dots, a_l)$ . This implies that  $\sigma_{u_1}^{a_1} \sigma_{u_2}^{a_2} \dots \sigma_{u_l}^{a_l} \neq 1$ , and that the coefficient of  $\sigma_{u_1}^{a_1} \sigma_{u_2}^{a_2} \dots \sigma_{u_l}^{a_l}$  in  $\tilde{\eta}(\tilde{g})$  is  $c_{a_1 p_1 q_1} c_{a_2 p_2 q_2} \dots c_{a_l p_l q_l} \neq 0$ , thus that  $\tilde{\eta}(\tilde{g}) \neq 1$ .

The sequence  $(\tau_{u_1}^{p_1} \sigma_{u_1}^{q_1}, \dots, \tau_{u_l}^{p_l} \sigma_{u_l}^{q_l})$  is a normal form, thus, by Corollary 5.3, the sequence  $(\sigma_{u_1}^{a_1}, \sigma_{u_2}^{a_2}, \dots, \sigma_{u_l}^{a_l})$  is the normal form for  $\sigma_{u_1}^{a_1} \sigma_{u_2}^{a_2} \dots \sigma_{u_l}^{a_l}$ . Assume  $n_i \neq 0$  for all  $i = 1, \dots, l$ . Again, by Corollary 5.3, the sequence  $(\sigma_{u_1}^{n_1}, \sigma_{u_2}^{n_2}, \dots, \sigma_{u_l}^{n_l})$  is the normal form for  $\sigma_{u_1}^{n_1} \sigma_{u_2}^{n_2} \dots \sigma_{u_l}^{n_l}$ , thus  $\sigma_{u_1}^{n_1} \sigma_{u_2}^{n_2} \dots \sigma_{u_l}^{n_l} \neq \sigma_{u_1}^{a_1} \sigma_{u_2}^{a_2} \dots \sigma_{u_l}^{a_l}$  if  $(n_1, n_2, \dots, n_l) \neq (a_1, a_2, \dots, a_l)$ . Now, assume that there exists  $i \in \{1, \dots, l\}$  such that  $n_i = 0$ . Then

$$|\sigma_{u_1}^{n_1} \sigma_{u_2}^{n_2} \dots \sigma_{u_l}^{n_l}|_G < l = |(\sigma_{u_1}^{a_1}, \sigma_{u_2}^{a_2}, \dots, \sigma_{u_l}^{a_l})| = |\sigma_{u_1}^{a_1} \sigma_{u_2}^{a_2} \dots \sigma_{u_l}^{a_l}|_G,$$

hence  $\sigma_{u_1}^{n_1} \sigma_{u_2}^{n_2} \dots \sigma_{u_l}^{n_l} \neq \sigma_{u_1}^{a_1} \sigma_{u_2}^{a_2} \dots \sigma_{u_l}^{a_l}$ . □

## 6 Other questions

### 6.1 Topological interpretation

A *geometric braid* on  $n$  strings is a  $n$ -tuple  $\beta = (b_1, \dots, b_n)$  of smooth disjoint paths in  $\mathbb{C} \times [0, 1]$  such that

- the projection of  $b_k(t)$  on the second component is  $t$  for all  $k \in \{1, \dots, n\}$  and all  $t \in [0, 1]$ ,
- $b_k(0) = (k, 0)$  and  $b_k(1) = (\xi(k), 1)$ , where  $\xi$  is some permutation of  $\{1, \dots, n\}$ , for all  $k \in \{1, \dots, n\}$ .

The *braid group* on  $n$  strings, denoted by  $\mathcal{B}_n$ , is the group of isotopy classes of braids. Define a *singular braid* on  $n$  strings as a  $n$ -tuple  $\beta = (b_1, \dots, b_n)$  of smooth paths in  $\mathbb{C} \times [0, 1]$  such that

- the projection of  $b_k(t)$  on the second component is  $t$  for all  $k \in \{1, \dots, n\}$  and all  $t \in [0, 1]$ ,
- $b_k(0) = (k, 0)$  and  $b_k(1) = (\xi(k), 1)$ , where  $\xi$  is some permutation of  $\{1, \dots, n\}$ , for all  $k \in \{1, \dots, n\}$ ,
- the strings intersect transversely in finitely many double points.

The *singular braid monoid*, denoted by  $S\mathcal{B}_n$ , is the monoid of isotopy classes of singular braids. By [3] (see also [4]), the braid group  $\mathcal{B}_n$  is isomorphic to the Artin group  $A = A_\Gamma$  associated to the Coxeter graph  $\Gamma = A_{n-1}$ , and, by [9],  $S\mathcal{B}_n$  is the singular Artin monoid associated to  $\Gamma = A_{n-1}$ .

Some other Artin groups can be viewed as “geometric braid groups”. For instance, the Artin group associated to the Coxeter graph  $\Gamma = B_n$  is the braid group on  $n$  strings of the annulus, and the Artin group associated to the graph  $\Gamma = D_n$  is an index 2 subgroup of the braid group on  $n$  strings of the plane endowed with a singular point of degree 2 (see [1]). One can easily verify using the techniques of [26] that the singular braid monoid of the annulus coincides with the singular Artin monoid associated to  $\Gamma = B_n$ . However, we did not succeed to find any embedding of the singular Artin monoid associated to  $\Gamma = D_n$  into the singular braid monoid of the plane endowed with a singular point of degree 2, and we suspect that such an embedding does not exist.

In other respects, the spherical type Artin groups can be interpreted as fundamental groups of regular orbit spaces (see [11]), and the groups of type  $A_n$  and  $D_n$  as geometric monodromy groups of simple singularities (see [37]). In both cases, we do not know whether these topological interpretations can be extended to the singular Artin monoids, even for the singular Artin monoids of type  $A_n$  (namely, the singular braid monoids).

### 6.2 Vassiliev invariants

An *invariant* on  $A$  is a set-map  $v : A \rightarrow H$ , where  $H$  is some abelian group, or, equivalently, a homomorphism  $v : \mathbb{Z}[A] \rightarrow H$  of  $\mathbb{Z}$ -modules. Consider the homomorphism  $\text{ord} : SA \rightarrow \mathbb{N}$  which sends  $\sigma_s^{\pm 1}$  to 0 and  $\tau_s$  to 1 for all  $s \in S$ . For  $d \in \mathbb{N}$ , we put

$S_d A = \{\alpha \in SA; \text{ord}(\alpha) = d\}$ . So,  $S_d A$  is the set of elements of  $SA$  that have exactly  $d$  “singularities”. Recall the desingularization map  $\eta : SA \rightarrow \mathbb{Z}[A]$  which sends  $\sigma_s^{\pm 1}$  to  $\sigma_s^{\pm 1}$  and  $\tau_s$  to  $\sigma_s - \sigma_s^{-1}$  for all  $s \in S$ . Then define a *Vassiliev invariant* of type  $d$  as an invariant  $v : \mathbb{Z}[A] \rightarrow H$  which vanishes on  $\eta(S_{d+1}A)$ .

The following questions have been solved for braid groups (see [6], [31], and [33]), and remain open for the other Artin groups.

**Question 6.1.** *Do Vassiliev invariants separate the elements of  $A$ ? In other words, given two elements  $\alpha, \beta \in A$ ,  $\alpha \neq \beta$ , does there exist a Vassiliev invariant  $v : \mathbb{Z}[A] \rightarrow H$  such that  $v(\alpha) \neq v(\beta)$ ?*

By a *universal Vassiliev invariant* we mean an invariant  $Z : \mathbb{Z}[A] \rightarrow \mathcal{A}$  such that, if  $v : \mathbb{Z}[A] \rightarrow H$  is some Vassiliev invariant, then there exists a homomorphism  $u : \mathcal{A} \rightarrow H$  of  $\mathbb{Z}$ -modules such that  $v = u \circ Z$ .

**Question 6.2.** *Describe a universal Vassiliev invariant for  $A$  in terms of generators and relations.*

### 6.3 Conjugacy problem

Let  $\mathcal{M}$  be a monoid, and let  $\mathcal{G}$  be the group of unities of  $\mathcal{M}$ . We say that two elements  $\alpha, \beta \in \mathcal{M}$  are *conjugate* if there exists some  $\gamma \in \mathcal{G}$  such that  $\gamma\alpha\gamma^{-1} = \beta$ . A *solution to the conjugacy problem* in  $\mathcal{M}$  is an algorithm which decides whether two given elements  $\alpha, \beta \in \mathcal{M}$  are conjugate or not.

As for the word problem, a solution to the conjugacy problem for  $SA$  will give a solution to the conjugacy problem for  $A$  (which, by the way, is the group of unities of  $SA$ ). So, a reasonable approach is to study the conjugacy problem for those singular Artin monoids whose associated Artin groups have known solutions to the conjugacy problem. For instance, a solution to the conjugacy problem for Artin groups of spherical type can be found in [12] (see also [38] and [22]).

A solution to the conjugacy problem for singular braid monoids is given in [40]. We suspect that this algorithm can be extended to all spherical type singular Artin monoids. On the other hand, we do not know whether the techniques introduced in the present paper, more specifically, Propositions 2.1 and 3.1, can be also used to solve the conjugacy problem in  $SA$ .

On other respects, let  $\tilde{A} = \mathcal{G}(\hat{\Omega}) \rtimes A$  be the smallest group which contains  $SA = \mathcal{M}(\hat{\Omega}) \rtimes A$ . As far as we know, no solution to the conjugacy problem for  $\tilde{A}$  is known, even in the case where  $SA$  is the singular braid monoid.

## References

- [1] D. Allcock, *Braid pictures for Artin groups*, Trans. Amer. Math. Soc. **354** (2002), 3455–3474.

- [2] J.A. Altobelli, *The word problem for Artin groups of FC type*, J. Pure Appl. Algebra **129** (1998), 1–22.
- [3] E. Artin, *Theorie der Zöpfe*, Hamburg Abh. **4** (1925), 47–72.
- [4] E. Artin, *Theory of braids*, Ann. of Math. (2) **48** (1947), 101–126.
- [5] J.C. Baez, *Link invariants of finite type and perturbation theory*, Lett. Math. Phys. **26** (1992), 43–51.
- [6] D. Bar-Natan, *Vassiliev homotopy string link invariants*, J. Knot Theory Ramifications **4** (1995), 13–32.
- [7] G. Basset, *Quasi-commuting extensions of groups*, Comm. Algebra **28** (2000), 5443–5454.
- [8] M. Bestvina, N. Brady, *Morse theory and finiteness properties of groups*, Invent. Math. **129** (1997), 445–470.
- [9] J.S. Birman, *New points of view in knot theory*, Bull. Amer. Math. Soc. (N.S.) **28** (1993), 253–287.
- [10] N. Bourbaki, *Groupes et algèbres de Lie, Chapitres 4, 5 et 6*, Hermann, Paris, 1968.
- [11] E. Brieskorn, *Die Fundamentalgruppe des Raumes der regulären Orbits einer endlichen Komplexen Spiegelungsgruppe*, Invent. Math. **12** (1971), 57–61.
- [12] E. Brieskorn, K. Saito, *Artin-Gruppen und Coxeter-Gruppen*, Invent. Math. **17** (1972), 245–271.
- [13] P. Cartier, D. Foata, *Problèmes combinatoires de commutation et réarrangements*, Lecture Notes in Mathematics, No. 85, Springer-Verlag, Berlin-New York, 1969.
- [14] R. Charney, *Geodesic automation and growth functions for Artin groups of finite type*, Math. Ann. **301** (1995), 307–324.
- [15] R. Charney, M.W. Davis, *The  $K(\pi, 1)$ -problem for hyperplane complements associated to infinite reflection groups*, J. Amer. Math. Soc. **8** (1995), 597–627.
- [16] R. Corran, *A normal form for a class of monoids including the singular braid monoids*, J. Algebra **223** (2000), 256–282.
- [17] P. Deligne, *Les immeubles des groupes de tresses généralisés*, Invent. Math. **17** (1972), 273–302.
- [18] C. Duboc, *Commutations dans les monoïdes libres : un cadre théorique pour l'étude du parallélisme*, Ph. D. Thesis, Université de Rouen, 1986.

- [19] G. Duchamp, D. Krob, *Free partially commutative structures*, J. Algebra **156** (1993), 318–361.
- [20] J. East, *Birman’s conjecture is true for  $I_2(p)$* , preprint, 2003.
- [21] R. Fenn, D. Rolfsen, J. Zhu, *Centralisers in the braid group and singular braid monoid*, Enseign. Math. (2) **42** (1996), 75–96.
- [22] N. Franco, J. González-Meneses, *Conjugacy problem for braid groups and Garside groups*, J. Algebra **266** (2003), 112–132.
- [23] F.A. Garside, *The braid group and other groups*, Quart. J. Math. Oxford Ser. (2) **20** (1969), 235–254.
- [24] E. Godelle, *Normalisateur et groupe d’Artin de type sphérique*, J. Algebra **269** (2003), 263–274.
- [25] E. Godelle, *Parabolic subgroups of Artin groups of type FC*, Pacific J. Math. **208** (2003), 243–254.
- [26] J. González-Meneses, *Presentations for the monoids of singular braids on closed surfaces*, Comm. Algebra **30** (2002), 2829–2836.
- [27] E.R. Green, *Graph products of groups*, Ph. D. Thesis, The University of Leeds, 1990.
- [28] S. Hermiller, J. Meier, *Algorithms and geometry of graph products of groups*, J. Algebra **171** (1995), 230–257.
- [29] T. Hsu, D.T. Wise, *On linear and residual properties of graph products*, Michigan Math. J. **46** (1999), 251–259.
- [30] E. Keyman, *A class of monoids embeddable in a group*, Turkish J. Math. **25** (2001), 299–305.
- [31] T. Kohno, *Vassiliev invariants and de Rham complex on the space of knots*, Symplectic geometry and quantization (Sanda and Yokohama, 1993), 123–138, Contemp. Math., 179, Amer. Math. Soc., Providence, RI, 1994.
- [32] H. van der Lek, *The homotopy type of complex hyperplane complements*, Ph. D. Thesis, University of Nijmegen, 1983.
- [33] S. Papadima, *the universal finite-type invariant for braids, with integer coefficients*, Topology Appl. **118** (2002), 169–185.
- [34] L. Paris, *Parabolic subgroups of Artin groups*, J. Algebra **196** (1997), 369–399.
- [35] L. Paris, *Artin monoids inject in their groups*, Comment. Math. Helv. **77** (2002), 609–637.

- [36] L. Paris, *The proof of Birman's conjecture on singular braid monoids*, preprint, [ArXiv:math.GR/0306422](https://arxiv.org/abs/math/0306422).
- [37] B. Perron, J.P. Vannier, *Groupe de monodromie géométrique des singularités simples*, *Math. Ann.* **306** (1996), 231–245.
- [38] M. Picantin, *The conjugacy problem in small Gaussian groups*, *Comm. Algebra* **29** (2001), 1021–1039.
- [39] J.P. Serre, *Arbres, amalgames,  $SL_2$* , *Astérisque*, No. 46, Société Mathématique de France, Paris, 1977.
- [40] V.V. Vershinin, *On the singular braid monoid*, preprint, [ArXiv:math.GR/0309339](https://arxiv.org/abs/math/0309339).
- [41] G.X. Viennot, *Heaps of pieces. I. Basic definitions and combinatorial lemmas*, *Graph theory and its applications: East and West* (Jinan, 1986), 542–570, *Ann. New York Acad. Sci.*, 576, New York Acad. Sci., New York, 1989.

Eddy Godelle  
 Laboratoire de Mathématiques  
 Nicolas Oresme  
 Université de Caen  
 UMR 6139 du CNRS, BP 5186  
 14032 Caen cedex  
 France  
[eddy.godelle@math.unicaen.fr](mailto:eddy.godelle@math.unicaen.fr)

Luis Paris  
 Institut de Mathématiques  
 de Bourgogne  
 Université de Bourgogne  
 UMR 5584 du CNRS, BP 47870  
 21078 Dijon cedex  
 France  
[lparis@u-bourgogne.fr](mailto:lparis@u-bourgogne.fr)