

YETTER-DRINFELD CATEGORIES FOR QUASI-HOPF ALGEBRAS

D. BULACU, S. CAENEPEEL, AND F. PANAITTE

ABSTRACT. We show that all possible categories of Yetter-Drinfeld modules over a quasi-Hopf algebra H are isomorphic. We prove also that the category ${}^H_H\mathcal{YD}^{\text{fd}}$ of finite dimensional left Yetter-Drinfeld modules is rigid and then we compute explicitly the canonical isomorphisms in ${}^H_H\mathcal{YD}^{\text{fd}}$. Finally, we show that certain duals of H_0 , the braided Hopf algebra introduced in [6, 7], are isomorphic as braided Hopf algebras if H is a finite dimensional triangular quasi-Hopf algebra.

INTRODUCTION

Let H be a Hopf algebra with a bijective antipode. We can introduce left, right, left-right and right-left Yetter-Drinfeld modules over H , and it is well-known (see [2, 17]) that the corresponding categories ${}^H_H\mathcal{YD}$, \mathcal{YD}_H^H , ${}_H\mathcal{YD}^H$ and ${}^H\mathcal{YD}_H$ are isomorphic. These categories are also isomorphic to the center of the monoidal category ${}_H\mathcal{M}$ of left H -modules, and, if H is finite dimensional, to the category ${}_{D(H)}\mathcal{M}$ of left modules over the Drinfeld double $D(H)$. It also known that the category of finite dimensional Yetter-Drinfeld modules is rigid, that is, we have left and right duality in this category.

In [10], Dijkgraaf, Pasquier and Roche introduced the so-called “twisted double” of a finite group, which is a Hopf algebra-type object $D^\omega(G)$ associated to a pair (G, ω) , where G is a finite group and ω is a normalized 3-cocycle on G ; this object is *not* a Hopf algebra, but a quasi-Hopf algebra in the sense of Drinfeld [11]. The construction is similar to the quantum double, so it appears natural to try to define the quantum double of an arbitrary finite dimensional quasi-Hopf algebra, generalizing the Drinfeld double for Hopf algebras, and then to show that $D^\omega(G)$ is such a quantum double. This has been done first by Majid in [15]; he first computed the center of the monoidal category ${}_H\mathcal{M}$ of left H -modules over the quasi-Hopf algebra H (we will denote this center by ${}^H_H\mathcal{YD}$ and call its objects left Yetter-Drinfeld modules over H). Then he defined the quantum double $D(H)$ by an implicit Tannaka-Krein reconstruction procedure, in such a way that ${}_{D(H)}\mathcal{M} \cong {}^H_H\mathcal{YD}$. An explicit construction of the quantum double (as a so-called “diagonal crossed product”) has been given afterwards by Hausser and Nill in [12, 13]. They

1991 *Mathematics Subject Classification.* 16W30.

Key words and phrases. quasi-Hopf algebra, Yetter-Drinfeld module, braided monoidal category.

Research supported by the bilateral project “Hopf Algebras in Algebra, Topology, Geometry and Physics” of the Flemish and Romanian governments. This paper was finished while the first author was visiting the Vrije Universiteit Brussel, and he would like to thank VUB for its warm hospitality. The third author was also partially supported by the programmes SCOPES and EURROMMAT.

identified the category of left modules over their quantum double with the category $\frac{H^{\text{cop}}}{H^{\text{cop}}}\mathcal{YD}$ (we will denote this category by ${}_H\mathcal{YD}^H$ and call its objects left-right Yetter-Drinfeld modules over H). Now the question arises whether $\frac{H}{H}\mathcal{YD} \cong {}_H\mathcal{YD}^H$.

In Section 2, we will show that the categories $\frac{H}{H}\mathcal{YD}$ and ${}_H\mathcal{YD}^H$ (and also two other categories \mathcal{YD}_H^H and ${}^H\mathcal{YD}_H$ which we will introduce) are isomorphic, even in the situation where H is not finite dimensional. The isomorphisms can be constructed explicitly, and we even have isomorphisms $\frac{H}{H}\mathcal{YD} \cong {}_H\mathcal{YD}^H$ and $\mathcal{YD}_H^H \cong {}^H\mathcal{YD}_H$ of braided monoidal categories. As we will see, the proof is much more technical than in the classical case.

In Section 3, we show that the category $\frac{H}{H}\mathcal{YD}^{\text{fd}}$ of finite dimensional (left) Yetter-Drinfeld modules is rigid; the left and right duals are constructed explicitly. In an arbitrary rigid braided monoidal category \mathcal{C} , we have canonical isomorphisms $M \cong M^{**}$ and $(M \otimes N)^* \cong M^* \otimes N^*$. In Section 4, we will compute these isomorphisms in the cases $\mathcal{C} = \frac{H}{H}\mathcal{YD}^{\text{fd}}$. If we then specialize to finite dimensional left modules over a quasitriangular quasi-Hopf algebra, we recover some results from [8]. Let B be a braided Hopf algebra, that is a Hopf algebra in $\frac{H}{H}\mathcal{YD}^{\text{fd}}$. Then B^* and *B are also braided Hopf algebras. In Section 5, we study the special case where $B = H_0$, the braided Hopf algebra introduced in [6, 7]. In particular, we prove that *H_0 and H_0^* are isomorphic braided Hopf algebras if H is a finite dimensional triangular quasi-Hopf algebra. They are also isomorphic to the coproduct of the braided Hopf algebra \underline{H}^* introduced in [3].

1. PRELIMINARIES

1.1. Quasi-Hopf algebras. We work over a commutative field k . All algebras, linear spaces etc. will be over k ; unadorned \otimes means \otimes_k . Following Drinfeld [11], a quasi-bialgebra is a fourtuple $(H, \Delta, \varepsilon, \Phi)$ where H is an associative algebra with unit, Φ is an invertible element in $H \otimes H \otimes H$, and $\Delta : H \rightarrow H \otimes H$ and $\varepsilon : H \rightarrow k$ are algebra homomorphisms satisfying the identities

$$(1.1) \quad (id \otimes \Delta)(\Delta(h)) = \Phi(\Delta \otimes id)(\Delta(h))\Phi^{-1},$$

$$(1.2) \quad (id \otimes \varepsilon)(\Delta(h)) = h \otimes 1, \quad (\varepsilon \otimes id)(\Delta(h)) = 1 \otimes h,$$

for all $h \in H$. Φ has to be a normalized 3-cocycle, in the sense that

$$(1.3) \quad (1 \otimes \Phi)(id \otimes \Delta \otimes id)(\Phi)(\Phi \otimes 1) = (id \otimes id \otimes \Delta)(\Phi)(\Delta \otimes id \otimes id)(\Phi),$$

$$(1.4) \quad (id \otimes \varepsilon \otimes id)(\Phi) = 1 \otimes 1 \otimes 1.$$

The map Δ is called the coproduct or the comultiplication, ε the counit and Φ the reassociator. As for Hopf algebras [18] we denote $\Delta(h) = \sum h_1 \otimes h_2$, but since Δ is only quasi-coassociative we adopt the further convention

$$(\Delta \otimes id)(\Delta(h)) = \sum h_{(1,1)} \otimes h_{(1,2)} \otimes h_2, \quad (id \otimes \Delta)(\Delta(h)) = \sum h_1 \otimes h_{(2,1)} \otimes h_{(2,2)},$$

for all $h \in H$. We will denote the tensor components of Φ by capital letters, and the ones of Φ^{-1} by small letters, namely

$$\Phi = \sum X^1 \otimes X^2 \otimes X^3 = \sum T^1 \otimes T^2 \otimes T^3 = \sum V^1 \otimes V^2 \otimes V^3 = \dots$$

$$\Phi^{-1} = \sum x^1 \otimes x^2 \otimes x^3 = \sum t^1 \otimes t^2 \otimes t^3 = \sum v^1 \otimes v^2 \otimes v^3 = \dots$$

H is called a quasi-Hopf algebra if, moreover, there exists an anti-automorphism S of the algebra H and elements $\alpha, \beta \in H$ such that, for all $h \in H$:

$$(1.5) \quad \sum S(h_1)\alpha h_2 = \varepsilon(h)\alpha \quad \text{and} \quad \sum h_1\beta S(h_2) = \varepsilon(h)\beta,$$

$$(1.6) \quad \sum X^1\beta S(X^2)\alpha X^3 = 1 \quad \text{and} \quad \sum S(x^1)\alpha x^2\beta S(x^3) = 1.$$

The antipode of a quasi-Hopf algebra is unique up to a transformation $\alpha \mapsto U\alpha$, $\beta \mapsto \beta U^{-1}$, $S(h) \mapsto US(h)U^{-1}$, where $U \in H$ is invertible. The axioms for a quasi-Hopf algebra imply that $\varepsilon(\alpha)\varepsilon(\beta) = 1$, so, by rescaling α and β , we may assume without loss of generality that $\varepsilon(\alpha) = \varepsilon(\beta) = 1$ and $\varepsilon \circ S = \varepsilon$. The identities (1.2-1.4) also imply that

$$(1.7) \quad (\varepsilon \otimes id \otimes id)(\Phi) = (id \otimes id \otimes \varepsilon)(\Phi) = 1 \otimes 1 \otimes 1.$$

Together with a quasi-Hopf algebra $H = (H, \Delta, \varepsilon, \Phi, S, \alpha, \beta)$ we also have H^{op} , H^{cop} and $H^{\text{op,cop}}$ as quasi-Hopf algebras, where "op" means opposite multiplication and "cop" means opposite comultiplication. The quasi-Hopf structures are obtained by putting $\Phi_{\text{op}} = \Phi^{-1}$, $\Phi_{\text{cop}} = (\Phi^{-1})^{321}$, $\Phi_{\text{op,cop}} = \Phi^{321}$, $S_{\text{op}} = S_{\text{cop}} = (S_{\text{op,cop}})^{-1} = S^{-1}$, $\alpha_{\text{op}} = S^{-1}(\beta)$, $\beta_{\text{op}} = S^{-1}(\alpha)$, $\alpha_{\text{cop}} = S^{-1}(\alpha)$, $\beta_{\text{cop}} = S^{-1}(\beta)$, $\alpha_{\text{op,cop}} = \beta$ and $\beta_{\text{op,cop}} = \alpha$.

Recall that the definition of a quasi-Hopf algebra is "twist covariant" in the following sense. An invertible element $F \in H \otimes H$ is called a *gauge transformation* or *twist* if $(\varepsilon \otimes id)(F) = (id \otimes \varepsilon)(F) = 1$. If H is a quasi-Hopf algebra and $F = \sum F^1 \otimes F^2 \in H \otimes H$ is a gauge transformation with inverse $F^{-1} = \sum G^1 \otimes G^2$, then we can define a new quasi-Hopf algebra H_F by keeping the multiplication, unit, counit and antipode of H and replacing the comultiplication, reassociator and the elements α and β by

$$(1.8) \quad \Delta_F(h) = F\Delta(h)F^{-1},$$

$$(1.9) \quad \Phi_F = (1 \otimes F)(id \otimes \Delta)(F)\Phi(\Delta \otimes id)(F^{-1})(F^{-1} \otimes 1),$$

$$(1.10) \quad \alpha_F = \sum S(G^1)\alpha G^2, \quad \beta_F = \sum F^1\beta S(F^2).$$

It is well-known that the antipode of a Hopf algebra is an anti-coalgebra morphism. For a quasi-Hopf algebra, we have the following statement: there exists a gauge transformation $f \in H \otimes H$ such that

$$(1.11) \quad f\Delta(S(h))f^{-1} = (S \otimes S)(\Delta^{\text{op}}(h)), \quad \text{for all } h \in H,$$

where $\Delta^{\text{op}}(h) = \sum h_2 \otimes h_1$. The element f can be computed explicitly. First set

$$\sum A^1 \otimes A^2 \otimes A^3 \otimes A^4 = (\Phi \otimes 1)(\Delta \otimes id \otimes id)(\Phi^{-1}),$$

$$\sum B^1 \otimes B^2 \otimes B^3 \otimes B^4 = (\Delta \otimes id \otimes id)(\Phi)(\Phi^{-1} \otimes 1).$$

Then define $\gamma, \delta \in H \otimes H$ by

$$(1.12) \quad \gamma = \sum S(A^2)\alpha A^3 \otimes S(A^1)\alpha A^4 \quad \text{and} \quad \delta = \sum B^1\beta S(B^4) \otimes B^2\beta S(B^3).$$

Then f and f^{-1} are given by the formulae:

$$(1.13) \quad f = \sum (S \otimes S)(\Delta^{\text{op}}(x^1))\gamma\Delta(x^2\beta S(x^3)),$$

$$(1.14) \quad f^{-1} = \sum \Delta(S(x^1)\alpha x^2)\delta(S \otimes S)(\Delta^{\text{op}}(x^3)).$$

Moreover, f satisfies the following relations:

$$(1.15) \quad f\Delta(\alpha) = \gamma, \quad \Delta(\beta)f^{-1} = \delta.$$

Furthermore the corresponding twisted reassociator (see (1.9)) is given by

$$(1.16) \quad \Phi_f = \sum (S \otimes S \otimes S)(X^3 \otimes X^2 \otimes X^1).$$

In a Hopf algebra H , we obviously have the identity $\sum h_1 \otimes h_2 S(h_3) = h \otimes 1$, for all $h \in H$. We will need the generalization of this formula to the quasi-Hopf algebra setting. Following [12] and [13], we define

$$(1.17) \quad p_R = \sum p_R^1 \otimes p_R^2 = \sum x^1 \otimes x^2 \beta S(x^3)$$

$$(1.18) \quad q_R = \sum q_R^1 \otimes q_R^2 = \sum X^1 \otimes S^{-1}(\alpha X^3) X^2$$

$$(1.19) \quad p_L = \sum p_L^1 \otimes p_L^2 = \sum X^2 S^{-1}(X^1 \beta) \otimes X^3$$

$$(1.20) \quad q_L = \sum q_L^1 \otimes q_L^2 = \sum S(x^1) \alpha x^2 \otimes x^3$$

For all $h \in H$, we then have:

$$(1.21) \quad \sum \Delta(h_1) p_R [1 \otimes S(h_2)] = p_R [h \otimes 1]$$

$$(1.22) \quad \sum [1 \otimes S^{-1}(h_2)] q_R \Delta(h_1) = (h \otimes 1) q_R$$

$$(1.23) \quad \sum \Delta(h_2) p_L [S^{-1}(h_1) \otimes 1] = p_L (1 \otimes h)$$

$$(1.24) \quad \sum [S(h_1) \otimes 1] q_L \Delta(h_2) = (1 \otimes h) q_L$$

and

$$(1.25) \quad \sum \Delta(q_R^1) p_R [1 \otimes S(q_R^2)] = 1 \otimes 1$$

$$(1.26) \quad \sum [1 \otimes S^{-1}(p_R^2)] q_R \Delta(p_R^1) = 1 \otimes 1$$

$$(1.27) \quad \sum [S(p_L^1) \otimes 1] q_L \Delta(p_L^2) = 1 \otimes 1$$

$$(1.28) \quad \sum \Delta(q_L^2) p_L [S^{-1}(q_L^1) \otimes 1] = 1 \otimes 1$$

$$(1.29) \quad \begin{aligned} (q_R \otimes 1)(\Delta \otimes id)(q_R)\Phi^{-1} &= \sum [1 \otimes S^{-1}(X^3) \otimes S^{-1}(X^2)] \\ & [1 \otimes S^{-1}(f^2) \otimes S^{-1}(f^1)](id \otimes \Delta)(q_R \Delta(X^1)), \end{aligned}$$

$$(1.30) \quad \begin{aligned} \Phi^{-1}(id \otimes \Delta)(p_L)(1 \otimes p_L) &= \sum (\Delta \otimes id)(\Delta(X^3) p_L) \\ & (S^{-1}(X^2 g^2) \otimes S^{-1}(X^1 g^1) \otimes 1), \end{aligned}$$

$$(1.31) \quad \begin{aligned} (1 \otimes q_L)(id \otimes \Delta)(q_L)\Phi &= \sum (S(x^2) \otimes S(x^1) \otimes 1) \\ & (f \otimes 1)(\Delta \otimes id)(q_L \Delta(x^3)), \end{aligned}$$

where $f = \sum f^1 \otimes f^2$ is the twist defined in (1.13) with its inverse $f^{-1} = \sum g^1 \otimes g^2$ defined in (1.14).

1.2. Quasitriangular quasi-Hopf algebras. A quasi-Hopf algebra H is quasi-triangular if there exists an element $R \in H \otimes H$ such that

$$(1.32) \quad (\Delta \otimes id)(R) = \sum \Phi_{312} R_{13} \Phi_{132}^{-1} R_{23} \Phi,$$

$$(1.33) \quad (id \otimes \Delta)(R) = \sum \Phi_{231}^{-1} R_{13} \Phi_{213} R_{12} \Phi^{-1},$$

$$(1.34) \quad \Delta^{\text{op}}(h)R = R\Delta(h), \text{ for all } h \in H,$$

$$(1.35) \quad (\varepsilon \otimes id)(R) = (id \otimes \varepsilon)(R) = 1.$$

Here we use the following notation. If σ is a permutation of $\{1, 2, 3\}$, we set $\Phi_{\sigma(1)\sigma(2)\sigma(3)} = \sum X^{\sigma^{-1}(1)} \otimes X^{\sigma^{-1}(2)} \otimes X^{\sigma^{-1}(3)}$, and R_{ij} means R acting non-trivially in the i^{th} and j^{th} positions of $H \otimes H \otimes H$.

In [5] it is shown that R is invertible. The inverse of R is given by

$$(1.36) \quad R^{-1} = \sum X^1 \beta S(Y^2 R^1 x^1 X^2) \alpha Y^3 x^3 X_2^3 \otimes Y^1 R^2 x^2 X_1^3.$$

Furthermore, the element

$$(1.37) \quad u = \sum S(R^2 p^2) \alpha R^1 p^1$$

(with $p_R = \sum p^1 \otimes p^2$ defined as in (1.17)) satisfies $S^2(u) = u$, is invertible in H , and

$$(1.38) \quad u^{-1} = \sum X^1 R^2 p^2 S(S(X^2 R^1 p^1) \alpha X^3),$$

$$(1.39) \quad \varepsilon(u) = 1 \text{ and } S^2(h) = uhu^{-1},$$

for all $h \in H$. Consequently the antipode S is bijective, so, as in the Hopf algebra case, the assumptions about invertibility of R and bijectivity of S can be dropped. Moreover, the R -matrix $R = \sum R^1 \otimes R^2$ satisfies the identities (see [1], [13], [5]):

$$(1.40) \quad f_{21} R f^{-1} = (S \otimes S)(R),$$

$$(1.41) \quad \sum S(R^2) \alpha R^1 = S(\alpha)u,$$

where $f = \sum f^1 \otimes f^2$ is the twist defined in (1.13), and $f_{21} = \sum f^2 \otimes f^1$. In addition, a second formula for the inverse of R is

$$(1.42) \quad R^{-1} = \sum \tilde{q}_1^2 X^2 R^1 p^1 \otimes \tilde{q}_2^2 X^3 S^{-1}(\tilde{q}^1 X^1 R^2 p^2),$$

where $p_R = \sum p^1 \otimes p^2$ and $q_L = \sum \tilde{q}^1 \otimes \tilde{q}^2$ are the elements defined by (1.17) and (1.20). Finally, recall that a quasi-Hopf algebra (H, R) is called triangular if $R^{-1} = R_{21}$, where $R_{21} = \sum R^2 \otimes R^1$.

1.3. Braided Hopf algebras. A monoidal category is a category \mathcal{C} with objects U, V, W etc., a functor $\otimes : \mathcal{C} \times \mathcal{C} \rightarrow \mathcal{C}$ equipped with an associativity natural transformation consisting of functorial isomorphisms $a_{U,V,W} : (U \otimes V) \otimes W \rightarrow U \otimes (V \otimes W)$ satisfying a pentagonal identity, and a compatible unit object $\underline{1}$ and associated natural isomorphisms (the left and right unit constraints, $l_V : V \cong V \otimes \underline{1}$ and $r_V : V \cong \underline{1} \otimes V$). If \mathcal{C} and \mathcal{D} are monoidal categories, then, roughly speaking, we say that $\mathbf{T} : \mathcal{C} \rightarrow \mathcal{D}$ is a monoidal functor if it respects the tensor products (in the sense that for any two objects $U, V \in \mathcal{C}$ there exists a functorial isomorphism $\Psi_{U,V} : \mathbf{T}(U) \otimes \mathbf{T}(V) \rightarrow \mathbf{T}(U \otimes V)$ such that Ψ respects the associativity constraints), the unit object and the left and right unit constraints (for a complete definition see [16, p. 421]).

Let \mathcal{C} be a monoidal category. An object $V \in \mathcal{C}$ has a left dual if there exists an object V^* and morphisms $ev_V : V^* \otimes V \rightarrow \underline{1}$, $coev_V : \underline{1} \rightarrow V \otimes V^*$ in \mathcal{C} such that

$$(1.43) \quad l_V^{-1} \circ (id_V \otimes ev_V) \circ a_{V, V^*, V} \circ (coev_V \otimes id_V) \circ r_V = id_V,$$

$$(1.44) \quad r_{V^*}^{-1} \circ (ev_V \otimes id_{V^*}) \circ a_{V^*, V, V^*}^{-1} \circ (id_{V^*} \otimes coev_V) \circ l_{V^*} = id_{V^*}.$$

There exists a similar notion of right duality. An object $V \in \mathcal{C}$ has a right dual if there exist an object ${}^*V \in \mathcal{C}$ and morphisms $ev'_V : V \otimes {}^*V \rightarrow \underline{1}$, $coev'_V : \underline{1} \rightarrow {}^*V \otimes V$ in \mathcal{C} such that

$$(1.45) \quad r_V^{-1} \circ (ev'_V \otimes id_V) \circ a_{V, {}^*V, V}^{-1} \circ (id_V \otimes coev'_V) \circ l_V = id_V,$$

$$(1.46) \quad l_{{}^*V}^{-1} \circ (id_{{}^*V} \otimes ev'_V) \circ a_{{}^*V, V, {}^*V} \circ (coev'_V \otimes id_{{}^*V}) \circ r_{{}^*V} = id_{{}^*V}.$$

If every object in the category has a left and right dual, then we say that \mathcal{C} is a rigid monoidal category.

A braided category is a monoidal category equipped with a commutativity natural transformation consisting of functorial isomorphisms $c_{U, V} : U \otimes V \rightarrow V \otimes U$ compatible with the unit and the associativity constraints in a natural way (for a complete definition see [14], [16]). If \mathcal{C} and \mathcal{D} are braided categories then a monoidal functor $F : \mathcal{C} \rightarrow \mathcal{D}$ is braided if, for any pair (U, V) of objects of \mathcal{C} we have $F(c_{U, V}) \circ \Psi_{U, V} = \Psi_{V, U} \circ c_{F(U), F(V)}$.

Suppose that $(H, \Delta, \varepsilon, \Phi)$ is a quasi-bialgebra. If U, V, W are left H -modules, define $a_{U, V, W} : (U \otimes V) \otimes W \rightarrow U \otimes (V \otimes W)$ by

$$(1.47) \quad a_{U, V, W}((u \otimes v) \otimes w) = \Phi \cdot (u \otimes (v \otimes w)).$$

Then the category ${}_H\mathcal{M}$ of left H -modules becomes a monoidal category with tensor product \otimes given via Δ , associativity constraints $a_{U, V, W}$, unit k as a trivial H -module and the usual left and right unit constraints. If H is a quasi-Hopf algebra then the category of finite dimensional left H -modules is rigid; the left dual of V is V^* with the H -module structure given by $(h \cdot \varphi)(v) = \varphi(S(h) \cdot v)$, for all $v \in V$, $\varphi \in V^*$, $h \in H$ and with

$$(1.48) \quad ev_V(\varphi \otimes v) = \varphi(\alpha \cdot v), \quad coev_V(1) = \sum_{i=1}^n \beta \cdot v_i \otimes v^i,$$

where $\{v_i\}$ is a basis in V with dual basis $\{v^i\}$. The right dual *V of V is the same dual vector space now equipped with the left H -module structure given for all $h \in H$, $v \in V$ and linear map φ on V by $(h \cdot \varphi)(v) = \varphi(S^{-1}(h) \cdot v)$, and with

$$(1.49) \quad ev'_V(v \otimes \varphi) = \varphi(S^{-1}(\alpha) \cdot v), \quad coev'_V(1) = \sum_{i=1}^n v^i \otimes S^{-1}(\beta) \cdot v_i,$$

where we used the same notation as above.

Now let H be a quasitriangular quasi-Hopf algebra, with R -matrix $R = \sum R^1 \otimes R^2$. For two left H -modules U and V , we define $c_{U, V} : U \otimes V \rightarrow V \otimes U$ by

$$(1.50) \quad c_{U, V}(u \otimes v) = \sum R^2 \cdot v \otimes R^1 \cdot u$$

and then $({}_H\mathcal{M}, \otimes, k, a, \lambda, \rho, c)$ is a braided monoidal category (cf. [14] or [16]).

We are interested in general Hopf algebras in braided categories \mathcal{C} . The definition of a Hopf algebra B in \mathcal{C} is obtained in the obvious way in analogy with the standard axioms [18]. Thus, a bialgebra in \mathcal{C} is $(B, m, \eta, \Delta, \varepsilon)$ where B is an object in \mathcal{C} and the morphism $m : B \otimes B \rightarrow B$ gives a multiplication that is associative up to the isomorphism a . Similarly for the coassociativity of the comultiplication

$\Delta : B \rightarrow B \otimes B$. The identity in the algebra B is expressed as usual by $\eta : \mathbf{1} \rightarrow B$ such that $m \circ (\eta \otimes id) = m \circ (id \otimes \eta) = id$. The counit axiom is $(\varepsilon \otimes id) \circ \Delta = (id \otimes \varepsilon) \circ \Delta = id$. In addition, Δ is required to be an algebra morphism where $B \otimes B$ has the multiplication $m_{B \otimes B}$, defined as the composition

$$(1.51) \quad \begin{array}{ccccc} (B \otimes B) \otimes (B \otimes B) & \xrightarrow{a} & B \otimes (B \otimes (B \otimes B)) & \xrightarrow{B \otimes a^{-1}} & B \otimes ((B \otimes B) \otimes B) \\ & \searrow^{B \otimes c \otimes B} & \searrow^{B \otimes ((B \otimes B) \otimes B)} & \searrow^{B \otimes a} & \searrow^{B \otimes (B \otimes (B \otimes B))} \\ & \xrightarrow{a^{-1}} & (B \otimes B) \otimes (B \otimes B) & \xrightarrow{m \otimes m} & B \otimes B \end{array}$$

A Hopf algebra B is a bialgebra with a morphism $S : B \rightarrow B$ in \mathcal{C} (the antipode) satisfying the usual axioms $m \circ (S \otimes id) \circ \Delta = \eta \circ \varepsilon = m \circ (id \otimes S) \circ \Delta$.

Let \mathcal{C} be a braided category. Let $\overline{\mathcal{C}}$ be equal to \mathcal{C} as a monoidal category, with the mirror-reversed braiding $\overline{c}_{M,N} = c_{N,M}^{-1}$. For a Hopf algebra $B \in \mathcal{C}$ with bijective antipode, we define B^{op} , B^{cop} and $B^{\text{op,cop}}$ by

$$(1.52) \quad m_{B^{\text{op}}} = m_B \circ c_{B,B}^{-1}, \quad \Delta_{B^{\text{op}}} = \Delta_B, \quad S_{B^{\text{op}}} = S_B^{-1},$$

$$(1.53) \quad m_{B^{\text{cop}}} = m_B, \quad \Delta_{B^{\text{cop}}} = c_{B,B}^{-1} \circ \Delta_B, \quad S_{B^{\text{cop}}} = S_B^{-1},$$

$$(1.54) \quad m_{B^{\text{op,cop}}} = m_B \circ c_{B,B}, \quad \Delta_{B^{\text{op,cop}}} = c_{B,B}^{-1} \circ \Delta_B, \quad S_{B^{\text{op,cop}}} = S_B,$$

and the other structure morphisms remain the same as for B . It is well-known (see for example [2, 19]) that B^{op} and B^{cop} are Hopf algebras in $\overline{\mathcal{C}}$, and $B^{\text{op,cop}}$ is a Hopf algebra in \mathcal{C} .

2. YETTER-DRINFELD MODULES OVER A QUASI-HOPF ALGEBRA

In this Section we will investigate the relations between the various kinds of Yetter-Drinfeld modules (${}^H_H\mathcal{YD}$, ${}_H\mathcal{YD}^H$, etc.) over a quasi-Hopf algebra. First we recall from [15] the notion of left Yetter-Drinfeld module over a quasi-bialgebra.

Definition 2.1. Let H be a quasi-bialgebra with reassociator Φ . A left H -module M together with a left H -coaction

$$\lambda_M : M \rightarrow H \otimes M, \quad \lambda_M(m) = \sum m_{(-1)} \otimes m_{(0)}$$

is called a left Yetter-Drinfeld module if the following equalities hold, for all $h \in H$ and $m \in M$:

$$(2.1) \quad \begin{aligned} & \sum X^1 m_{(-1)} \otimes (X^2 \cdot m_{(0)})_{(-1)} X^3 \otimes (X^2 \cdot m_{(0)})_{(0)} \\ & = \sum X^1 (Y^1 \cdot m)_{(-1)_1} Y^2 \otimes X^2 (Y^1 \cdot m)_{(-1)_2} Y^3 \otimes X^3 \cdot (Y^1 \cdot m)_{(0)}, \end{aligned}$$

$$(2.2) \quad \sum \varepsilon(m_{(-1)}) m_{(0)} = m,$$

$$(2.3) \quad \sum h_1 m_{(-1)} \otimes h_2 \cdot m_{(0)} = \sum (h_1 \cdot m)_{(-1)} h_2 \otimes (h_1 \cdot m)_{(0)}.$$

The category of left Yetter-Drinfeld H -modules and k -linear maps that preserve the H -action and H -coaction is denoted by ${}^H_H\mathcal{YD}$. In [15] it is shown that ${}^H_H\mathcal{YD}$ is a pre-braided monoidal category. The forgetful functor ${}^H_H\mathcal{YD} \rightarrow {}_H\mathcal{M}$ is monoidal, and the coaction on the tensor product $M \otimes N$ of two Yetter-Drinfeld modules M and N is given by

$$(2.4) \quad \begin{aligned} \lambda_{M \otimes N}(m \otimes n) & = \sum X^1 (x^1 Y^1 \cdot m)_{(-1)} x^2 (Y^2 \cdot n)_{(-1)} Y^3 \\ & \otimes X^2 \cdot (x^1 Y^1 \cdot m)_{(0)} \otimes X^3 x^3 \cdot (Y^2 \cdot n)_{(0)}. \end{aligned}$$

The braiding is given by

$$(2.5) \quad c_{M,N}(m \otimes n) = \sum m_{(-1)} \cdot n \otimes m_{(0)}.$$

This braiding is invertible if H is a quasi-Hopf algebra [6], and its inverse is given by

$$(2.6) \quad c_{M,N}^{-1}(n \otimes m) = \sum \tilde{q}_1^2 X^2 \cdot (p^1 \cdot m)_{(0)} \otimes S^{-1}(\tilde{q}^1 X^1 (p^1 \cdot m)_{(-1)} p^2 S(\tilde{q}_2^2 X^3)) \cdot n$$

where $p_R = \sum p^1 \otimes p^2$ and $q_L = \sum \tilde{q}^1 \otimes \tilde{q}^2$ are the elements defined by (1.17) and (1.20), respectively.

Now, we will define left-right Yetter-Drinfeld modules for quasi-bialgebras. Namely, if H is a quasi-bialgebra, we define ${}_H\mathcal{YD}^H$ to be ${}_{H^{\text{cop}}}\mathcal{YD}$. More explicitly:

Definition 2.2. Let H be a quasi-bialgebra. A k -linear space M is called a left-right Yetter-Drinfeld module if M is a left H -module (denote the action by $h \otimes m \mapsto h \cdot m$) and H coacts on M to the right (denote the right coaction by $M \rightarrow M \otimes H$, $m \mapsto \sum m_{(0)} \otimes m_{(1)}$) such that for all $m \in M$ and $h \in H$ the following relations hold:

$$(2.7) \quad \begin{aligned} & \sum (x^2 \cdot m_{(0)})_{(0)} \otimes (x^2 \cdot m_{(0)})_{(1)} x^1 \otimes x^3 m_{(1)} \\ &= \sum x^1 \cdot (y^3 \cdot m)_{(0)} \otimes x^2 (y^3 \cdot m)_{(1)_1} y^1 \otimes x^3 (y^3 \cdot m)_{(1)_2} y^2, \end{aligned}$$

$$(2.8) \quad \sum \varepsilon(m_{(1)}) m_{(0)} = m,$$

$$(2.9) \quad \sum h_1 \cdot m_{(0)} \otimes h_2 m_{(1)} = \sum (h_2 \cdot m)_{(0)} \otimes (h_2 \cdot m)_{(1)} h_1.$$

The category ${}_H\mathcal{YD}^H$ has left-right Yetter-Drinfeld modules as objects; the morphisms intertwine the H -action and the corresponding H -coaction.

If H is a quasi-Hopf algebra then we will prove that the categories ${}_H\mathcal{YD}^H$ and ${}^H_H\mathcal{YD}$ are isomorphic. To this end, we first need some formulae.

Lemma 2.3. *Let H be a quasi-Hopf algebra and M an object of ${}_H\mathcal{YD}^H$. Then the following relations hold:*

$$(2.10) \quad \sum x^1 \tilde{p}^1 \otimes x^2 \tilde{p}_1^2 \otimes x^3 \tilde{p}_2^2 = \sum X_1^2 \tilde{p}^1 S^{-1}(X^1) \otimes X_2^2 \tilde{p}^2 \otimes X^3,$$

$$(2.11) \quad \sum q_1^1 x^1 \otimes q_2^1 x^2 \otimes q^2 x^3 = \sum X^1 \otimes q^1 X_1^2 \otimes S^{-1}(X^3) q^2 X_2^2,$$

$$(2.12) \quad \begin{aligned} & \sum (\tilde{p}^2 h \cdot m)_{(0)} \otimes (\tilde{p}^2 h \cdot m)_{(1)} \tilde{p}^1 \\ &= \sum h_{(2,1)} \cdot (\tilde{p}^2 \cdot m)_{(0)} \otimes h_{(2,2)} (\tilde{p}^2 \cdot m)_{(1)} \tilde{p}^1 S^{-1}(h_1), \end{aligned}$$

for all $h \in H$ and $m \in M$, where $q_R = \sum q^1 \otimes q^2$ and $p_L = \sum \tilde{p}^1 \otimes \tilde{p}^2$ are the elements defined by (1.18) and (1.19).

Proof. The relations (2.10) and (2.11) follow easily by (1.3) and (1.5). The equality (2.12) is an immediate consequence of (1.23) and (2.9), the details are left to the reader. \square

Proposition 2.4. *Let H be a quasi-Hopf algebra. We have a functor $F : {}_H\mathcal{YD}^H \rightarrow {}^H_H\mathcal{YD}$, defined as follows. For $M \in {}_H\mathcal{YD}^H$, $F(M) = M$ as a left H -module; the left H -coaction is given by*

$$(2.13) \quad \lambda_M(m) = \sum m_{(-1)} \otimes m_{(0)} = \sum q_1^1 x^1 S(q^2 x^3 (\tilde{p}^2 \cdot m)_{(1)} \tilde{p}^1) \otimes q_2^1 x^2 \cdot (\tilde{p}^2 \cdot m)_{(0)},$$

for all $m \in M$, where $q_R = \sum q^1 \otimes q^2$ and $p_L = \sum \tilde{p}^1 \otimes \tilde{p}^2$ are the elements defined by (1.18) and (1.19), and $\rho_M(m) = \sum m_{(0)} \otimes m_{(1)}$ is the right coaction of H on M . The functor F sends a morphism to itself.

Proof. The main thing to show is that $F(M)$ is an object of ${}^H_H\mathcal{YD}$. It is then easy to see that F behaves well at the level of the morphisms.

Write $q_R = \sum q^1 \otimes q^2 = \sum Q^1 \otimes Q^2$, $p_L = \sum \tilde{p}^1 \otimes \tilde{p}^2 = \sum \tilde{P}^1 \otimes \tilde{P}^2$, $f = \sum f^1 \otimes f^2$ and $f^{-1} = \sum g^1 \otimes g^2$. We prove first that (2.1) holds for $F(M)$. For all $m \in M$ we compute:

$$\begin{aligned}
& \sum X^1(Y^1 \cdot m)_{(-1)_1} Y^2 \otimes X^2(Y^1 \cdot m)_{(-1)_2} Y^3 \otimes X^3 \cdot (Y^1 \cdot m)_{(0)} \\
(2.13) \quad &= \sum X^1 q_{(1,1)}^1 x_1^1 S(q^2 x^3 (\tilde{p}^2 Y^1 \cdot m)_{(1)} \tilde{p}^1)_1 Y^2 \otimes X^2 q_{(1,2)}^1 x_2^1 \\
& \quad S(q^2 x^3 (\tilde{p}^2 Y^1 \cdot m)_{(1)} \tilde{p}^1)_2 Y^3 \otimes X^3 q_2^1 x^2 \cdot (\tilde{p}^2 Y^1 \cdot m)_{(0)} \\
(2.12, 1.1, 1.11) \quad &= \sum X^1 (q^1 Y_1^1)_{(1,1)} x_1^1 g^1 S(q_2^2 Y_{(2,2)}^1 x_2^3 (\tilde{p}^2 \cdot m)_{(1)_2} \tilde{p}_2^1) f^1 Y^2 \\
& \quad \otimes X^2 (q^1 Y_1^1)_{(1,2)} x_2^1 g^2 S(q_1^2 Y_{(2,1)}^1 x_1^3 (\tilde{p}^2 \cdot m)_{(1)_1} \tilde{p}_1^1) f^2 Y^3 \\
& \quad \otimes X^3 (q^1 Y_1^1)_{(2)} x^2 \cdot (\tilde{p}^2 \cdot m)_{(0)} \\
(1.1, 1.29) \quad &= \sum (q^1 Q_1^1 y^1)_1 X^1 x_1^1 g^1 S(Q^2 y^3 x_2^3 (\tilde{p}^2 \cdot m)_{(1)_2} \tilde{p}_2^1) \\
& \quad \otimes (q^1 Q_1^1 y^1)_{(2,1)} X^2 x_2^1 g^2 S(q^2 Q_2^1 y^2 x_1^3 (\tilde{p}^2 \cdot m)_{(1)_1} \tilde{p}_1^1) \\
& \quad \otimes (q^1 Q_1^1 y^1)_{(2,2)} X^3 x^2 \cdot (\tilde{p}^2 \cdot m)_{(0)} \\
(1.3, 2.9) \quad &= \sum (q^1 Q_1^1 y^1)_1 z^1 X^1 g^1 S(Q^2 y^3 z_2^3 x_2^3 (X_2^3 \tilde{p}^2 \cdot m)_{(1)_2} X_{(1,2)}^3 \tilde{p}_2^1) \\
& \quad \otimes (q^1 Q_1^1 y^1)_{(2,1)} z_1^2 x^1 X^2 g^2 S(q^2 Q_2^1 y^2 z_1^3 x_1^3 (X_2^3 \tilde{p}^2 \cdot m)_{(1)_1} X_{(1,1)}^3 \tilde{p}_1^1) \\
& \quad \otimes (q^1 Q_1^1 y^1)_{(2,2)} z_2^2 x^2 \cdot (X_2^3 \tilde{p}^2 \cdot m)_{(0)} \\
(1.30, 2.7) \quad &= \sum (q^1 Q_1^1 y^1)_1 z^1 S(Q^2 y^3 z_2^3 x_2^3 X^3 t^3 (\tilde{p}_2^2 \tilde{P}^2 \cdot m)_{(1)} \tilde{p}_1^2 \tilde{P}^1) \\
& \quad \otimes (q^1 Q_1^1 y^1)_{(2,1)} z_1^2 x^1 S(q^2 Q_2^1 y^2 z_1^3 x_1^3 X^2 (t^2 \cdot (\tilde{p}_2^2 \tilde{P}^2 \cdot m)_{(0)})_{(1)} t^1 \tilde{p}^1) \\
& \quad \otimes (q^1 Q_1^1 y^1)_{(2,2)} z_2^2 x^2 X^1 \cdot (t^2 \cdot (\tilde{p}_2^2 \tilde{P}^2 \cdot m)_{(0)})_{(0)} \\
(2.9, 2.10, 2.9) \quad &= \sum (q^1 Q_1^1 y^1)_1 z^1 S(Q^2 y^2 z_2^3 x_2^3 X^3 Y^3 (\tilde{P}^2 \cdot m)_{(1)} \tilde{P}^1) \\
& \quad \otimes (q^1 Q_1^1 y^1)_{(2,1)} z_1^2 x^1 Y^1 S(q^2 Q_2^1 y^2 z_1^3 x_1^3 X^2 Y_2^2 (\tilde{p}^2 \cdot (\tilde{P}^2 \cdot m)_{(0)})_{(1)} \tilde{p}^1) \\
& \quad \otimes (q^1 Q_1^1 y^1)_{(2,2)} z_2^2 x^2 X^1 Y_1^2 \cdot (\tilde{p}^2 \cdot (\tilde{P}^2 \cdot m)_{(0)})_{(0)} \\
(1.3) \quad &= \sum (q^1 Q_1^1)_1 y_1^1 z^1 S(Q^2 y^3 z_2^3 Y^3 (\tilde{P}^2 \cdot m)_{(1)} \tilde{P}^1) \\
& \quad \otimes (q^1 Q_1^1)_{(2,1)} (y_2^1 z^2 Y^1)_1 x^1 S(q^2 Q_2^1 y^2 z_1^3 Y^2 x^3 (\tilde{p}^2 \cdot (\tilde{P}^2 \cdot m)_{(0)})_{(1)} \tilde{p}^1) \\
& \quad \otimes (q^1 Q_1^1)_{(2,2)} (y_2^1 z^2 Y^1)_2 x^2 \cdot (\tilde{p}^2 \cdot (\tilde{P}^2 \cdot m)_{(0)})_{(0)} \\
(1.3, 1.1, 2.12) \quad &= \sum q_1^1 z^1 Q_1^1 y^1 S(Q^2 y^3 (\tilde{P}^2 \cdot m)_{(1)} \tilde{P}^1) \\
& \quad \otimes (q_2^1 z^2)_1 x^1 (Q_2^1)_1 S(q^2 z^3 x^3 (Q_2^1)_{(2,2)} (\tilde{p}^2 y^2 \cdot (\tilde{P}^2 \cdot m)_{(0)})_{(1)} \tilde{p}^1) \\
& \quad \otimes (q_2^1 z^2)_2 x^2 (Q_2^1)_{(2,1)} \cdot (\tilde{p}^2 y^2 \cdot (\tilde{P}^2 \cdot m)_{(0)})_{(0)} \\
(2.11, 2.12) \quad &= \sum X^1 Q_1^1 y^1 S(Q^2 y^3 (\tilde{P}^2 \cdot m)_{(1)} \tilde{P}^1) \\
& \quad \otimes q_1^1 X_{(1,1)}^2 x^1 S(q^2 X_2^2 x^3 (\tilde{p}^2 Q_2^1 y^2 \cdot (\tilde{P}^2 \cdot m)_{(0)})_{(1)} \tilde{p}^1) X^3 \\
& \quad \otimes q_2^1 X_{(1,2)}^2 x^2 \cdot (\tilde{p}^2 Q_2^1 y^2 \cdot (\tilde{P}^2 \cdot m)_{(0)})_{(0)}
\end{aligned}$$

$$(1.1, 2.12, 2.13) = \sum X^1 m_{(-1)} \otimes (X^2 \cdot m_{(0)})_{(-1)} X^3 \otimes (X^2 \cdot m_{(0)})_{(0)}.$$

Similar computations show that

$$\begin{aligned} & \sum (h_1 \cdot m)_{(-1)} h_2 \otimes (h_1 \cdot m)_{(0)} \\ (2.13) &= \sum q_1^1 x^1 S(q^2 x^3 (\tilde{p}^2 h_1 \cdot m)_{(1)} \tilde{p}^1) h_2 \otimes q_2^1 x^2 \cdot (\tilde{p}^2 h_1 \cdot m)_{(0)} \\ (2.12, 1.1) &= \sum (q^1 h_{(1,1)})_1 x^1 S(q^2 h_{(1,2)} x^3 (\tilde{p}^2 \cdot m)_{(1)} \tilde{p}^1) h_2 \\ & \quad \otimes (q^1 h_{(1,1)})_2 x^2 \cdot (\tilde{p}^2 \cdot m)_{(0)} \\ (1.22, 2.13) &= \sum h_1 m_{(-1)} \otimes h_2 \cdot m_{(0)}, \end{aligned}$$

for all $h \in H$ and $m \in M$, that is (2.3) holds for $F(M)$; since (2.2) is trivial, the proof is complete. \square

Remark 2.5. We would like to emphasize that (2.13), which can be rewritten as

$$\begin{aligned} \lambda_M(m) &= \sum Y^1 S(q^2 Y_2^2 (\tilde{p}^2 \cdot m)_{(1)} \tilde{p}^1) Y^3 \otimes q^1 Y_1^2 \cdot (\tilde{p}^2 \cdot m)_{(0)} \\ &= \sum Y^1 Z^1 \beta S(Y_1^2 Z^2) S((Y_2^2 Z^3 \cdot m)_{(1)}) S(X^2) \alpha X^3 Y^3 \otimes X^1 \cdot (Y_2^2 Z^3 \cdot m)_{(0)}, \end{aligned}$$

was chosen in such a way that (2.3) holds; it is remarkable that (2.1) then also holds.

We are now able to prove the main result of this Section; it generalizes [17, Prop. 6, (b)].

Theorem 2.6. *Let H be a quasi-Hopf algebra. Then the categories ${}_H \mathcal{YD}^H$ and ${}^H \mathcal{YD}$ are isomorphic.*

Proof. We will show that the functor F defined in Proposition 2.4 has an inverse. Applying Proposition 2.4 to the quasi-Hopf algebra H^{cop} , we find a functor $F^{\text{cop}} : {}_{H^{\text{cop}}} \mathcal{YD}^{H^{\text{cop}}} \rightarrow {}^H \mathcal{YD}$. After identifying ${}^H \mathcal{YD} \equiv {}_{H^{\text{cop}}} \mathcal{YD}^{H^{\text{cop}}}$ and ${}^H \mathcal{YD} \equiv {}_H \mathcal{YD}^H$, we obtain a functor $G : {}^H \mathcal{YD} \rightarrow {}_H \mathcal{YD}^H$. Explicitly, if $M \in {}^H \mathcal{YD}$ then $G(M) = M$ as a left H -module, with the right H -coaction given by

$$\begin{aligned} \rho_M(m) &= \sum m_{(0)} \otimes m_{(1)} \\ (2.14) \quad &= \sum \tilde{q}_1^2 X^2 \cdot (p^1 \cdot m)_{(0)} \otimes \tilde{q}_2^2 X^3 S^{-1}(\tilde{q}^1 X^1 (p^1 \cdot m)_{(-1)} p^2) \end{aligned}$$

for all $m \in M$, where $q_L = \sum \tilde{q}^1 \otimes \tilde{q}^2$ and $p_R = \sum p^1 \otimes p^2$ are the elements defined by (1.20) and (1.17), and $\lambda_M(m) = \sum m_{(-1)} \otimes m_{(0)}$ is the left coaction of H on M . We claim that the functors F and G are inverses. Indeed, let first $M \in {}_H \mathcal{YD}^H$, with right H -coaction denoted by $\rho_M(m) = \sum m_{(0)} \otimes m_{(1)}$. If we denote the right H -coaction on $G(F(M))$ by ρ'_M , then:

$$\begin{aligned} \rho'_M(m) &\stackrel{(2.14)}{=} \sum \tilde{q}_1^2 X^2 \cdot (p^1 \cdot m)_{(0)} \otimes \tilde{q}_2^2 X^3 S^{-1}(\tilde{q}^1 X^1 (p^1 \cdot m)_{(-1)} p^2) \\ (2.13) \quad &= \sum \tilde{q}_1^2 X^2 q_2^1 x^2 \cdot (\tilde{p}^2 p^1 \cdot m)_{(0)} \\ & \quad \otimes \tilde{q}_2^2 X^3 S^{-1}(p^2) q^2 x^3 (\tilde{p}^2 p^1 \cdot m)_{(1)} \tilde{p}^1 S^{-1}(\tilde{q}^1 X^1 q_1^1 x^1) \end{aligned}$$

$$\begin{aligned}
 (2.12, 1.1) &= \sum \tilde{q}_1^2 X^2(q^1 p_1^1)_2 x^2 \cdot (\tilde{p}^2 \cdot m)_{(0)} \\
 &\quad \otimes \tilde{q}_2^2 X^3 S^{-1}(p^2) q^2 p_2^1 x^3 (\tilde{p}^2 \cdot m)_{(1)} \tilde{p}^1 S^{-1}(\tilde{q}^1 X^1(q^1 p_1^1)_1 x^1) \\
 (1.26) &= \sum \tilde{q}_1^2 \cdot (\tilde{p}^2 \cdot m)_{(0)} \otimes \tilde{q}_2^2 (\tilde{p}^1 \cdot m)_{(1)} \tilde{p}^1 S^{-1}(\tilde{q}^1) \\
 (2.9) &= \sum (\tilde{q}_2^2 \tilde{p}^2 \cdot m)_{(0)} \otimes (\tilde{q}_2^2 \tilde{p}^2 \cdot m)_{(1)} \tilde{q}_1^2 \tilde{p}^1 S^{-1}(\tilde{q}^1) \\
 (1.28) &= \sum m_{(0)} \otimes m_{(1)} = \rho_M(m)
 \end{aligned}$$

for all $m \in M$. Hence, we have proved that $G(F(M)) = M$. Replacing H by H^{cop} , the same proof shows that $F(G(M)) = M$, and the proof is complete. \square

We will now show that categories of right-left and right Yetter-Drinfeld modules are also isomorphic to the category of left Yetter-Drinfeld modules. For a quasi-bialgebra H , we define

$${}^H \mathcal{YD}_H = {}_{H^{\text{op}}, \text{cop}} \mathcal{YD}^{H^{\text{op}}, \text{cop}} \quad \text{and} \quad \mathcal{YD}_H^H = {}_{H^{\text{op}}, \text{cop}}^{H^{\text{op}}, \text{cop}} \mathcal{YD}.$$

More explicitly:

Definition 2.7. Let H be a quasi-bialgebra.

a) A k -linear space M is called a right-left Yetter-Drinfeld module if M is a right H -module (with action denoted by $m \otimes h \mapsto m \cdot h$) and if H coacts on M from the left (denote the left coaction by $M \rightarrow H \otimes M$, $m \mapsto \sum m_{(-1)} \otimes m_{(0)}$) such that the following relations hold, for all $m \in M$ and $h \in H$:

$$\begin{aligned}
 (2.15) \quad &\sum m_{(-1)} x^1 \otimes x^3 (m_{(0)} \cdot x^2)_{(-1)} \otimes (m_{(0)} \cdot x^2)_{(0)} \\
 &= \sum y^2 (m \cdot y^1)_{(-1)_1} x^1 \otimes y^3 (m \cdot y^1)_{(-1)_2} x^2 \otimes (m \cdot y^1)_{(0)} \cdot x^3,
 \end{aligned}$$

$$(2.16) \quad \sum \varepsilon(m_{(-1)}) m_{(0)} = m,$$

$$(2.17) \quad \sum m_{(-1)} h_1 \otimes m_{(0)} \cdot h_2 = \sum h_2 (m \cdot h_1)_{(-1)} \otimes (m \cdot h_1)_{(0)}.$$

The category of right-left Yetter-Drinfeld modules is denoted by ${}^H \mathcal{YD}_H$.

b) A k -linear space M is called a right Yetter-Drinfeld module if M is a right H -module (with action denoted by $m \otimes h \mapsto m \cdot h$) and if H coacts on M from the right (denote the right coaction by $M \rightarrow M \otimes H$, $m \mapsto \sum m_{(0)} \otimes m_{(1)}$) such that the following relations hold, for all $m \in M$ and $h \in H$:

$$\begin{aligned}
 (2.18) \quad &\sum (m_{(0)} \cdot X^2)_{(0)} \otimes X^1 (m_{(0)} \cdot X^2)_{(1)} \otimes m_{(1)} X^3 \\
 &= \sum (m \cdot Y^3)_{(0)} \cdot X^1 \otimes Y^1 (m \cdot Y^3)_{(1)_1} X^2 \otimes Y^2 (m \cdot Y^3)_{(1)_2} X^3,
 \end{aligned}$$

$$(2.19) \quad \sum \varepsilon(m_{(1)}) m_{(0)} = m,$$

$$(2.20) \quad \sum m_{(0)} \cdot h_1 \otimes m_{(1)} h_2 = \sum (m \cdot h_2)_{(0)} \otimes h_1 (m \cdot h_2)_{(1)}.$$

The category \mathcal{YD}_H^H consists of such objects and morphisms which intertwine the H -action and the corresponding H -coaction.

Before we prove that all these categories are isomorphic, we first recall some generalities.

(i) Let H be a quasi-bialgebra and $\mathfrak{F} = \sum \mathfrak{F}^1 \otimes \mathfrak{F}^2 \in H \otimes H$ a twist with inverse $\mathfrak{F}^{-1} = \sum \mathfrak{G}^1 \otimes \mathfrak{G}^2$. Then there exists a braided isomorphism between the braided categories ${}^H \mathcal{YD}$ and ${}_{H^{\mathfrak{F}}} \mathcal{YD}$. Following [6], this functor is the identity on objects

(regarded as left H -modules) and morphisms, with the monoidal structure Ψ given by multiplication by \mathfrak{F}^{-1} , that is, for all $M, N \in {}^H_H\mathcal{YD}$ and $m \in M, n \in N$:

$$\Psi_{M,N} : M \otimes N \rightarrow M \otimes N, \quad \Psi_{M,N}(m \otimes n) = \sum \mathfrak{G}^1 \cdot m \otimes \mathfrak{G}^2 \cdot n.$$

If $M \in {}^H_H\mathcal{YD}$, then $M \in {}^{H_{\mathfrak{F}}}H_{\mathfrak{F}}\mathcal{YD}$ with the same H -action, and with coaction given by

$$(2.21) \quad \lambda_M^{\mathfrak{F}}(m) = \sum m_{(-1)}^{\mathfrak{F}} \otimes m_{(0)}^{\mathfrak{F}} := \sum \mathfrak{F}^1(\mathfrak{G}^1 \cdot m)_{(-1)} \mathfrak{G}^2 \otimes \mathfrak{F}^2 \cdot (\mathfrak{G}^1 \cdot m)_{(0)}.$$

The inverse of this functor is the following. If $M \in {}^{H_{\mathfrak{F}}}H_{\mathfrak{F}}\mathcal{YD}$, then $M \in {}^H_H\mathcal{YD}$ with the same H -action, and with coaction given by

$$(2.22) \quad \lambda_M^{\mathfrak{F}^{-1}}(m) = \sum m_{(-1)}^{\mathfrak{F}^{-1}} \otimes m_{(0)}^{\mathfrak{F}^{-1}} := \sum \mathfrak{G}^1(\mathfrak{F}^1 \cdot m)_{(-1)} \mathfrak{F}^2 \otimes \mathfrak{G}^2 \cdot (\mathfrak{F}^1 \cdot m)_{(0)}.$$

Now replace H by H^{cop} , and apply the above construction again. Then there exists a braided functor

$$F' : {}_H\mathcal{YD}^H \cong {}_{H^{\text{cop}}}^{H^{\text{cop}}}\mathcal{YD} \rightarrow ({}_{(H^{\text{cop}})_{\mathfrak{F}_{21}}}^{(H^{\text{cop}})_{\mathfrak{F}_{21}}}\mathcal{YD} = ({}_{(H_{\mathfrak{F}})^{\text{cop}}}^{(H_{\mathfrak{F}})^{\text{cop}}}\mathcal{YD} = {}_{H_{\mathfrak{F}}}\mathcal{YD}^{H_{\mathfrak{F}}},$$

where $\mathfrak{F}_{21} = \sum \mathfrak{F}^2 \otimes \mathfrak{F}^1$. Explicitly, if $M \in {}_H\mathcal{YD}^H$, then $M \in {}_{H_{\mathfrak{F}}}\mathcal{YD}^{H_{\mathfrak{F}}}$ with the same H -action, and with coaction given by

$$(2.23) \quad \rho_M^{\mathfrak{F}}(m) = \sum m_{(0)}^{\mathfrak{F}} \otimes m_{(1)}^{\mathfrak{F}} := \sum \mathfrak{F}^1 \cdot (\mathfrak{G}^2 \cdot m)_{(0)} \otimes \mathfrak{F}^2(\mathfrak{G}^2 \cdot m)_{(1)} \mathfrak{F}^1$$

for all $m \in M$. Moreover, this functor provides a braided isomorphism; its inverse is the following. If $M \in {}_{H_{\mathfrak{F}}}\mathcal{YD}^{H_{\mathfrak{F}}}$, then $M \in {}_H\mathcal{YD}^H$ with the same H -action, and with coaction given by

$$(2.24) \quad \rho_M^{\mathfrak{F}^{-1}}(m) = \sum m_{(0)}^{\mathfrak{F}^{-1}} \otimes m_{(1)}^{\mathfrak{F}^{-1}} := \sum \mathfrak{G}^1 \cdot (\mathfrak{F}^2 \cdot m)_{(0)} \otimes \mathfrak{G}^2(\mathfrak{F}^2 \cdot m)_{(1)} \mathfrak{F}^1.$$

(ii) The formulae (1.11) and (1.16) express the fact that the antipode S is an isomorphism between the quasi-bialgebras $H^{\text{op,cop}}$ and H_f , where H_f is the quasi-Hopf algebra H twisted by Drinfeld's element f defined in (1.13). Moreover, since

$$(2.25) \quad \sum S(g^1)\alpha g^2 = S(\beta), \quad \sum f^1\beta S(f^2) = S(\alpha),$$

$$(2.26) \quad \sum g^1 S(g^2\alpha) = \beta, \quad \sum S(\beta f^1)f^2 = \alpha$$

(see [5]), it follows that $S : H^{\text{op,cop}} \rightarrow H_f$ is a quasi-Hopf algebra isomorphism. Hence S induces braided isomorphisms between the braided categories ${}_{H^{\text{op,cop}}}^{H^{\text{op,cop}}}\mathcal{YD}$ and ${}_{H_f}^{H_f}\mathcal{YD}$, and ${}_{H^{\text{op,cop}}}\mathcal{YD}^{H^{\text{op,cop}}}$ and ${}_{H_f}\mathcal{YD}^{H_f}$ respectively.

Proposition 2.8. *Let H be a quasi-Hopf algebra. Then ${}^H\mathcal{YD}_H \cong {}_H\mathcal{YD}^H$ and $\mathcal{YD}_H^H \cong {}_H\mathcal{YD}$ as braided monoidal categories.*

Proof. We have seen above that

$${}^H\mathcal{YD}_H = {}_{H^{\text{op,cop}}}\mathcal{YD}^{H^{\text{op,cop}}} \cong {}_{H_f}\mathcal{YD}^{H_f} \cong {}_H\mathcal{YD}^H.$$

If we compute the connecting isomorphisms explicitly, then we obtain the following:

- If $M \in {}^H\mathcal{YD}_H$, then $M \in {}_H\mathcal{YD}^H$ with structure maps:

$$h \diamond m = m \cdot S^{-1}(h),$$

$$\rho_M(m) = \sum g^1 \cdot (f^2 \cdot m)_{(0)} \otimes g^2 S((f^2 \cdot m)_{(-1)}) f^1.$$

- Conversely, if $M \in {}_H\mathcal{YD}^H$, then $M \in {}^H\mathcal{YD}_H$ as follows:

$$m \diamond h = S(h) \cdot m,$$

$$\lambda_M(m) = \sum S^{-1}(f^2(g^2 \cdot m)_{(1)}g^1) \otimes f^1 \cdot (g^2 \cdot m)_{(0)}.$$

In a similar way, we have:

$$\mathcal{YD}_H^H = \frac{H^{\text{op, cop}}}{H^{\text{op, cop}}} \mathcal{YD} \cong \frac{H_f}{H_f} \mathcal{YD} \cong \frac{H}{H} \mathcal{YD}.$$

This isomorphism can be made explicit as follows.

- If $M \in \mathcal{YD}_H^H$, then $M \in \frac{H}{H} \mathcal{YD}$ with structure:

$$h \mapsto m = m \cdot S^{-1}(h),$$

$$\lambda_M(m) = \sum g^1 S((m \cdot S^{-1}(f^1))_{(1)}) f^2 \otimes (m \cdot S^{-1}(f^1))_{(0)} \cdot S^{-1}(g^2).$$

- Conversely, if $M \in \frac{H}{H} \mathcal{YD}$, then $M \in \mathcal{YD}_H^H$ with

$$m \leftarrow h = S(h) \cdot m,$$

$$\rho_M(m) = \sum f^2 \cdot (g^1 \cdot m)_{(0)} \otimes S^{-1}(f^1(g^1 \cdot m)_{(-1)}g^2),$$

completing the proof. \square

Corollary 2.9. *If H is a quasi-Hopf algebra then ${}^H\mathcal{YD}_H \cong \mathcal{YD}_H^H \cong \frac{H}{H} \mathcal{YD} \cong {}_H\mathcal{YD}^H$.*

3. THE RIGID BRAIDED CATEGORY $\frac{H}{H} \mathcal{YD}^{\text{fd}}$

Let H be a Hopf algebra with bijective antipode. It is well-known that the category of finite dimensional left Yetter-Drinfeld H -modules is a braided monoidal rigid category. In this Section we will show that the same result holds for the category of finite dimensional left Yetter-Drinfeld modules $\frac{H}{H} \mathcal{YD}^{\text{fd}}$ over a quasi-Hopf algebra H . We first need a Lemma.

Lemma 3.1. *Let H be a quasi-Hopf algebra and M a left H -module such that H coacts on M to the left (as usual, denote the left coaction by $\lambda_M : M \rightarrow H \otimes M$, $\lambda_M(m) = \sum m_{(-1)} \otimes m_{(0)}$). Then the compatibility relation (2.3) holds if and only if*

$$(3.1) \quad \sum (h \cdot m)_{(-1)} \otimes (h \cdot m)_{(0)} = \sum [q^1 h_1]_1 (p^1 \cdot m)_{(-1)} p^2 S(q^2 h_2) \otimes [q^1 h_1]_2 \cdot (p^1 \cdot m)_{(0)},$$

for all $h \in H$, $m \in M$, where $p_R = \sum p^1 \otimes p^2$ and $q_R = \sum q^1 \otimes q^2$ are the elements defined in (1.17-1.18).

Proof. Suppose that (2.3) holds. For any $h \in H$ and $m \in M$ we compute that

$$\sum (h \cdot m)_{(-1)} \otimes (h \cdot m)_{(0)}$$

$$(1.25) \quad = \sum (q_1^1 p^1 h \cdot m)_{(-1)} q_2^1 p^2 S(q^2) \otimes (q_1^1 p^1 h \cdot m)_{(0)}$$

$$(1.21) \quad = \sum ([q^1 h_1]_1 p^1 \cdot m)_{(-1)} [q^1 h_1]_2 p^2 S(q^2 h_2) \otimes ([q^1 h_1]_1 p^1 \cdot m)_{(0)}$$

$$(2.3) \quad = \sum [q^1 h_1]_1 (p^1 \cdot m)_{(-1)} p^2 S(q^2 h_2) \otimes [q^1 h_1]_2 \cdot (p^1 \cdot m)_{(0)}.$$

Conversely, assume that (3.1) holds; in particular we have that

$$(3.2) \quad \lambda_M(m) = \sum q_1^1 (p^1 \cdot m)_{(-1)} p^2 S(q^2) \otimes q_2^1 \cdot (p^1 \cdot m)_{(0)}, \quad \forall m \in M.$$

We then compute, for all $h \in H$ and $m \in M$:

$$\begin{aligned}
& \sum (h_1 \cdot m)_{(-1)} h_2 \otimes (h_1 \cdot m)_{(0)} \\
(3.1) \quad &= \sum [q^1 h_{(1,1)}]_1 (p^1 \cdot m)_{(-1)} p^2 S(q^2 h_{(1,2)}) h_2 \otimes [q^1 h_{(1,1)}]_2 \cdot (p^1 \cdot m)_{(0)} \\
(1.22) \quad &= \sum h_1 q_1^1 (p^1 \cdot m)_{(-1)} p^2 S(q^2) \otimes h_2 q_2^1 \cdot (p^1 \cdot m)_{(0)} \\
(3.2) \quad &= \sum h_1 m_{(-1)} \otimes h_2 \cdot m_{(0)}.
\end{aligned}$$

□

Proposition 3.2. *Let H be a quasi-Hopf algebra, $M \in {}^H_H\mathcal{YD}^{\text{fd}}$, $({}^i m)_{i=\overline{1,n}}$ a basis of M and $({}^i m)_{i=\overline{1,n}}$ its dual basis. Let M^* and *M be equal to the dual space $\text{Hom}(M, k)$ as a vector space, with the following H -action and H -coaction, defined for all $h \in H$, $m^* \in M^*$, ${}^*m \in {}^*M$ and $m \in M$.*

For M^* :

$$(3.3) \quad (h \cdot m^*)(m) = m^*(S(h) \cdot m);$$

$$\begin{aligned}
\lambda_{M^*}(m^*) &= \sum m_{(-1)}^* \otimes m_{(0)}^* \\
(3.4) \quad &= \sum_{i=1}^n \langle m^*, f^2 \cdot (g^1 \cdot {}^i m)_{(0)} \rangle S^{-1}(f^1 (g^1 \cdot {}^i m)_{(-1)} g^2) \otimes {}^i m.
\end{aligned}$$

For *M :

$$(3.5) \quad (h \cdot {}^*m)(m) := {}^*m(S^{-1}(h) \cdot m)$$

$$\begin{aligned}
\lambda_{{}^*M}({}^*m) &= \sum {}^*m_{(-1)} \otimes {}^*m_{(0)} \\
(3.6) \quad &= \sum_{i=1}^n \langle {}^*m, S^{-1}(f^1) \cdot (S^{-1}(g^2) \cdot {}^i m)_{(0)} \rangle g^1 S((S^{-1}(g^2) \cdot {}^i m)_{(-1)}) f^2 \otimes {}^i m.
\end{aligned}$$

Here $f = \sum f^1 \otimes f^2$ is the twist defined by (1.13), with inverse $f^{-1} = \sum g^1 \otimes g^2$. Then $M^*, {}^*M \in {}^H_H\mathcal{YD}$.

Proof. With the notation (2.21), (3.4) and (3.6) can be rewritten as

$$(3.7) \quad \lambda_{M^*}(m^*) = \sum_{i=1}^n \langle m^*, {}^i m_{(0)}^f \rangle S^{-1}({}^i m_{(-1)}^f) \otimes {}^i m;$$

$$(3.8) \quad \lambda_{{}^*M}({}^*m) = \sum_{i=1}^n \langle {}^*m, {}^i m_{(0)}^{\mathfrak{h}} \rangle S({}^i m_{(-1)}^{\mathfrak{h}}) \otimes {}^i m.$$

Here f is the Drinfeld's twist defined in (1.13) and $\mathfrak{h} = \sum S^{-1}(f^2) \otimes S^{-1}(f^1)$.

It follows that M^* with the H -action defined by (3.3) is a left H -module. We use the following notation for the comultiplication Δ_f and the reassociator Φ_f on H_f .

$$\Delta_f(h) = \sum h_{\underline{1}} \otimes h_{\underline{2}} = f \Delta(h) f^{-1};$$

$$\Phi_f = \sum X_f^1 \otimes X_f^2 \otimes X_f^3 = \sum Y_f^1 \otimes Y_f^2 \otimes Y_f^3.$$

For all $m^* \in M^*$, we have that

$$\sum X^1(Y^1 \cdot m^*)_{(-1)_1} Y^2 \otimes X^2(Y^1 \cdot m^*)_{(-1)_2} Y^3 \otimes X^3 \cdot (Y^1 \cdot m^*)_{(0)}$$

$$\begin{aligned}
(3.7) &= \sum_{i=1}^n \langle Y^1 \cdot m^*, {}_i m_{(0)}^f \rangle X^1 S^{-1}({}_i m_{(-1)}^f)_1 Y^2 \\
&\quad \otimes X^2 S^{-1}({}_i m_{(-1)}^f)_2 Y^3 \otimes X^3 \cdot {}_i m \\
(3.3, 1.11, 1.16) &= \sum_{i=1}^n \langle m^*, Y_f^3 \cdot (X_f^1 \cdot {}_i m)_{(0)}^f \rangle S^{-1}(Y_f^2 (X_f^1 \cdot {}_i m)_{(-1)_2}^f X_f^3) \\
&\quad \otimes S^{-1}(Y_f^1 (X_f^1 \cdot {}_i m)_{(-1)_1}^f X_f^2) \otimes {}_i m \\
(M \in {}_{H_f}^{\text{Hf}} \mathcal{YD}) &= \sum_{i=1}^n \langle m^*, (X_f^2 \cdot {}_i m_{(0)}^f)_{(0)}^f \rangle S^{-1}((X_f^2 \cdot {}_i m_{(0)}^f)_{(-1)}^f X_f^3) \\
&\quad \otimes S^{-1}(X_f^1 {}_i m_{(-1)}^f) \otimes {}_i m \\
(1.16, 3.7) &= \sum_{i=1}^n \langle m^*, (S(X^2) \cdot {}_i m)_{(0)}^f \rangle X^1 S^{-1}((S(X^2) \cdot {}_i m)_{(-1)}^f) \\
&\quad \otimes {}_i m_{(-1)} X^3 \otimes {}_i m_{(0)} \\
(3.3, 3.7) &= \sum X^1 m_{(-1)}^* \otimes (X^2 \cdot m_{(0)}^*)_{(-1)} X^3 \otimes (X^2 \cdot m_{(0)}^*)_{(0)},
\end{aligned}$$

which is the relation (2.1) for our structures on M^* . It is not hard to see that (1.11) implies that $\Delta_f(S(h)) = \sum S(h_2) \otimes S(h_1)$, for all $h \in H$, and therefore

$$\begin{aligned}
&\sum h_1 m_{(-1)}^* \otimes h_2 \cdot m_{(0)}^* \\
(3.7, 3.3) &= \sum_{i=1}^n \langle m^*, (S(h_2) \cdot {}_i m)_{(0)}^f \rangle h_1 S^{-1}((S(h_2) \cdot {}_i m)_{(-1)}^f) \otimes {}_i m \\
&= \sum_{i=1}^n \langle m^*, (S(h)_1 \cdot {}_i m)_{(0)}^f \rangle S^{-1}((S(h)_1 \cdot {}_i m)_{(-1)}^f S(h)_2) \otimes {}_i m \\
(M \in {}_{H_f}^{\text{Hf}} \mathcal{YD}) &= \sum_{i=1}^n \langle m^*, S(h)_2 \cdot {}_i m_{(0)}^f \rangle S^{-1}(S(h)_1 {}_i m_{(-1)}^f) \otimes {}_i m \\
&= \sum_{i=1}^n \langle m^*, S(h_1) \cdot {}_i m_{(0)}^f \rangle S^{-1}({}_i m_{(-1)}^f) h_2 \otimes {}_i m \\
(3.7, 3.3) &= \sum (h_1 \cdot m^*)_{(-1)} h_2 \otimes (h_1 \cdot m^*)_{(0)},
\end{aligned}$$

for all $h \in H$ and $m^* \in M^*$. Thus, $M^* \in {}_H^{\text{Hf}} \mathcal{YD}$. Similar arguments show that ${}^*M \in {}_H^{\text{Hf}} \mathcal{YD}$, the details are left to the reader. \square

Theorem 3.3. *Let H be a quasi-Hopf algebra. Then ${}_{H_f}^{\text{Hf}} \mathcal{YD}^{\text{fd}}$ is a braided monoidal rigid category.*

Proof. We have seen that ${}_{H_f}^{\text{Hf}} \mathcal{YD}$ is braided, and it follows that ${}_{H_f}^{\text{Hf}} \mathcal{YD}^{\text{fd}}$ is also braided. Let $M \in {}_{H_f}^{\text{Hf}} \mathcal{YD}^{\text{fd}}$, $({}_i m)_{i=\overline{1,n}}$ a basis of M , and $({}^i m)_{i=\overline{1,n}}$ its dual basis. Consider $M^*, {}^*M \in {}_{H_f}^{\text{Hf}} \mathcal{YD}$ as above and define maps $\text{ev}_M, \text{coev}_M, \text{ev}'_M$ and coev'_M , similar to the maps defined in (1.48) and (1.49). We claim that this makes ${}_{H_f}^{\text{Hf}} \mathcal{YD}^{\text{fd}}$ into a monoidal category with left and right duality, hence a rigid monoidal category. By the previous results we only have to prove that $\text{ev}_M, \text{coev}_M, \text{ev}'_M$ and coev'_M

intertwine the corresponding H -coactions.

First observe that the axioms of a quasi-Hopf algebra imply:

$$(3.9) \quad \sum q^1 U_1^1 \otimes S(q^2 U_2^1) U^2 = (id \otimes S)(q_L),$$

where $q_R = \sum q^1 \otimes q^2$, $q_L = \sum f^1 \otimes f^2$ and $f^{-1} = \sum g^1 \otimes g^2$ are the elements defined by (1.18), (1.20), (1.13) and (1.14), and

$$(3.10) \quad U = \sum U^1 \otimes U^2 = \sum g^1 S(q^2) \otimes g^2 S(q^1).$$

We write $q_L = \sum \tilde{q}^1 \otimes \tilde{q}^2 = \sum \tilde{Q}^1 \otimes \tilde{Q}^2$, and then compute for all $m^* \in M^*$ and $m \in M$:

$$\begin{aligned} & (id_H \otimes ev_M) \lambda_{M^* \otimes M}(m^* \otimes m) \\ (2.4, 3.7, 1.48) &= \sum_{i=1}^n \langle x^1 Y^1 \cdot m^*, {}_i m_{(0)}^f \rangle \langle X^2 \cdot {}_i m, \alpha X^3 x^3 \cdot (Y^2 \cdot m)_{(0)} \rangle \\ & \quad X^1 S^{-1}({}_i m_{(-1)}^f) x^2 (Y^2 \cdot m)_{(-1)} Y^3 \otimes 1 \\ (3.3, 3.4, 1.18) &= \sum \langle m^*, S(x^1 Y^1) f^2 \cdot (g^1 S(q^2) x^3 \cdot (Y^2 \cdot m)_{(0)})_{(0)} \rangle \\ & \quad S^{-1}(S(x^2) f^1 (g^1 S(q^2) x^3 \cdot (Y^2 \cdot m)_{(0)})_{(-1)} g^2 S(q^1)) \\ & \quad (Y^2 \cdot m)_{(-1)} Y^3 \otimes 1 \\ (3.10, 3.1) &= \sum \langle m^*, S(x^1 Y^1) f^2 [q^1 U_1^1 x_1^3]_2 \cdot (p^1 \cdot (Y^2 \cdot m)_{(0)})_{(0)} \rangle \\ & \quad S^{-1}(S(x^2) f^1 [q^1 U_1^1 x_1^3]_1 (p^1 \cdot (Y^2 \cdot m)_{(0)})_{(-1)}) \\ & \quad p^2 S(q^2 U_2^1 x_2^3) U^2 (Y^2 \cdot m)_{(-1)} Y^3 \otimes 1 \\ (3.9, 1.31) &= \sum \langle m^*, S(Y^1) \tilde{q}^1 \tilde{Q}_1^2 X^2 \cdot (p^1 \cdot (Y^2 \cdot m)_{(0)})_{(0)} \rangle \\ & \quad S^{-1}(\tilde{Q}^1 X^1 (p^1 \cdot (Y^2 \cdot m)_{(0)})_{(-1)} p^2 S(\tilde{q}^2 \tilde{Q}_2^2 X^3)) \\ & \quad (Y^2 \cdot m)_{(-1)} Y^3 \otimes 1 \\ (1.20, 1.3, 1.5) &= \sum \langle m^*, S(Y^1) \tilde{q}^1 x^3 y_2^2 \cdot (p^1 \cdot (Y^2 \cdot m)_{(0)})_{(0)} \rangle S^{-1}(S(x^1 y^1) \alpha x^2 y_1^2 \\ & \quad (p^1 \cdot (Y^2 \cdot m)_{(0)})_{(-1)} p^2 S(\tilde{q}^2 y^3)) (Y^2 \cdot m)_{(-1)} Y^3 \otimes 1 \\ (2.3, 1.17, 1.3) &= \sum \langle m^*, S(Y^1) \tilde{q}^1 x^3 \cdot (X^2 y_2^1 \cdot (Y^2 \cdot m)_{(0)})_{(0)} \rangle \\ & \quad S^{-1}(S(x^1 X^1 y_1^1 z^1) \alpha x^2 (X^2 y_2^1 z^2 \cdot (Y^2 \cdot m)_{(0)})_{(-1)}) \\ & \quad X^3 y^2 z_1^3 \beta S(\tilde{q}^2 y^3 z_2^3)) (Y^2 \cdot m)_{(-1)} Y^3 \otimes 1 \\ (1.5, 2.3, 2.1) &= \sum \langle m^*, S(Y^1) \tilde{q}^1 \cdot (X^1 y_1^1 Y^2 \cdot m)_{(0)} \rangle S^{-1}(\alpha (X^1 y_1^1 Y^2 \cdot m)_{(-1)_2} \\ & \quad X^3 y^2 \beta S(\tilde{q}^2 y^3)) (X^1 y_1^1 Y^2 \cdot m)_{(-1)_1} X^2 y_2^1 Y^3 \otimes 1 \\ (1.5, 2.2, 1.20) &= \sum \langle m^*, S(x^1 Y^1) \alpha x^2 X^1 y_1^1 Y^2 \cdot m \rangle \\ & \quad S^{-1}(S(X^2 y_2^1 Y^3) \alpha X^3 y^2 \beta S(x^3 y^3)) \otimes 1 \\ (1.3, 1.5) &= \sum \langle m^*, S(x^1 Y^1) \alpha x^2 Y^2 \cdot m \rangle S^{-1}(S(y^1 Y^3) \alpha y^2 \beta S(x^3 y^3)) \otimes 1 \\ (1.6, 1.48) &= \langle m^*, \alpha \cdot m \rangle 1 \otimes 1 = \lambda_k \circ ev_M(m^* \otimes m). \end{aligned}$$

In H_f we have the following properties (see (2.25) and (1.10)):

$$(3.11) \quad \alpha_f = \sum S(g^1) \alpha g^2 = S(\beta), \quad \beta_f = \sum f^1 \beta S(f^2) = S(\alpha).$$

Using (1.16-1.18) we obtain that

$$(3.12) \quad (p_R)_f = \sum S(\tilde{q}^2) \otimes S(\tilde{q}^1), \quad (q_R)_f = \sum S(\tilde{p}^2) \otimes S(\tilde{p}^1),$$

where $p_L = \sum \tilde{p}^1 \otimes \tilde{p}^2$ and $q_L = \sum \tilde{q}^1 \otimes \tilde{q}^2$ are the elements defined in (1.19-1.20). Specializing the relation (3.1) to the category $\frac{H_f}{H_f} \mathcal{YD}$, by (1.11) we conclude that

$$(3.13) \quad \sum (S(h) \cdot m)_{(-1)}^f \otimes (S(h) \cdot m)_{(0)}^f = \sum S(h_{(2,2)} \tilde{p}_2^2) (S(\tilde{q}^2) \cdot m)_{(-1)}^f \\ S(S(h_{1\tilde{p}^1}) \tilde{q}^1) \otimes S(h_{(2,1)} \tilde{p}_1^2) \cdot (S(\tilde{q}^2) \cdot m)_{(0)}^f,$$

for all $h \in H$ and $m \in M$. Now, if we denote by $\sum \tilde{P}^1 \otimes \tilde{P}^2$ another copy of p_L , then we compute:

$$\begin{aligned} & \lambda_{M \otimes M^*} \circ \text{coev}_M(1) \\ (1.48, 2.4, 3.7) &= \sum_{i,j=1}^n \langle Y^2 \cdot {}^i m, {}^j m_{(0)}^f \rangle X^1 (x^1 Y^1 \beta \cdot {}^i m)_{(-1)} x^2 \\ & \quad S^{-1}({}^j m_{(-1)}^f) Y^3 \otimes X^2 \cdot (x^1 Y^1 \beta \cdot {}^i m)_{(0)} \otimes X^3 x^3 \cdot {}^j m \\ (3.3, 1.19) &= \sum_{i=1}^n X^1 (x^1 S(\tilde{p}^1) \cdot (S(X^3 x^3) \cdot {}^i m)_{(0)}^f)_{(-1)} x^2 \\ & \quad S^{-1}((S(X^3 x^3) \cdot {}^i m)_{(-1)}^f) \tilde{p}^2 \\ & \quad \otimes X^2 \cdot (x^1 S(\tilde{p}^1) \cdot (S(X^3 x^3) \cdot {}^i m)_{(0)}^f)_{(0)} \otimes {}^i m \\ (3.13, 1.30, 1.1) &= \sum_{i=1}^n X^1 (g^1 S(Y^2 X_{(1,2)}^3 \tilde{p}_2^1) \cdot (S(\tilde{q}^2) \cdot {}^i m)_{(0)}^f)_{(-1)} g^2 \\ & \quad S(Y^1 X_{(1,1)}^3 \tilde{p}_1^1) \tilde{q}^1 S^{-1}((S(\tilde{q}^2) \cdot {}^i m)_{(-1)}^f) Y^3 X_2^3 \tilde{p}^2 \\ & \quad \otimes X^2 \cdot (g^1 S(Y^2 X_{(1,2)}^3 \tilde{p}_2^1) \cdot (S(\tilde{q}^2) \cdot {}^i m)_{(0)}^f)_{(0)} \otimes {}^i m \\ (1.11, 2.3, 1.16, 3.7) &= \sum_{i=1}^n X^1 S(X_1^3 \tilde{p}^1)_1 g^1 \cdot (Y_f^2 \cdot (S(\tilde{q}^2) \cdot {}^i m)_{(0)}^f)_{(-1)}^f Y_f^3 \tilde{q}^1 \\ & \quad S^{-1}(Y_f^1 (S(\tilde{q}^2) \cdot {}^i m)_{(-1)}^f) X_2^3 \tilde{p}^2 \\ & \quad \otimes X^2 S(X_1^3 \tilde{p}^1)_2 g^2 \cdot (Y_f^2 \cdot (S(\tilde{q}^2) \cdot {}^i m)_{(0)}^f)_{(0)}^f \otimes {}^i m \\ (M \in \frac{H_f}{H_f} \mathcal{YD}) &= \sum_{i=1}^n X^1 S(X_1^3 \tilde{p}^1)_1 g^1 Y_f^2 (X_f^1 S(\tilde{q}^2) \cdot {}^i m)_{(-1)_2}^f X_f^3 \tilde{q}^1 \\ & \quad S^{-1}(Y_f^1 (X_f^1 S(\tilde{q}^2) \cdot {}^i m)_{(-1)_1}^f X_f^2) X_2^3 \tilde{p}^2 \\ & \quad \otimes X^2 S(X_1^3 \tilde{p}^1)_2 g^2 Y_f^3 \cdot (X_f^1 S(\tilde{q}^2) \cdot {}^i m)_{(0)}^f \otimes {}^i m \\ (1.20, 1.16, 3.11) &= \sum_{i=1}^n X^1 S(X_1^3 \tilde{p}^1)_1 g^1 Y_f^2 S^{-1}(Y_f^1 {}^i m_{(-1)_1}^f \beta_f S({}^i m_{(-1)_2}^f)) X_2^3 \tilde{p}^2 \\ & \quad \otimes X^2 S(X_1^3 \tilde{p}^1)_2 g^2 Y_f^3 \cdot {}^i m_{(0)}^f \otimes {}^i m \end{aligned}$$

$$\begin{aligned}
(M \in {}_{H_f}^{\text{Hr}} \mathcal{YD}) &= \sum_{i=1}^n X^1 S(X_1^3 \tilde{p}^1)_1 g^1 Y_f^2 S^{-1}(Y_f^1 \beta_f) X_2^3 \tilde{p}^2 \\
&\quad \otimes X^2 S(X_1^3 \tilde{p}^1)_2 g^2 Y_f^3 \cdot {}_i m \otimes {}^i m \\
(1.16, 3.11, 1.11) &= \sum_{i=1}^n X^1 g^1 S(Y^2 X_{(1,2)}^3 \tilde{p}_2^1) \alpha Y^3 X_2^3 \tilde{p}^2 \\
&\quad \otimes X^2 g^2 S(Y^1 X_{(1,1)}^3 \tilde{p}_1^1) \cdot {}_i m \otimes {}^i m \\
(1.30, 1.5, 1.6, 1.48) &= \sum_{i=1}^n S(\tilde{p}_1^2 \tilde{P}^1) \alpha \tilde{p}_2^2 \tilde{P}^2 \otimes S(\tilde{p}^1) \cdot {}_i m \otimes {}^i m \\
&= (id \otimes coev_M) \circ \lambda_k(1),
\end{aligned}$$

as needed. Hence, the evaluation and coevaluation maps ev_M and $coev_M$ are morphisms in ${}_{H_f}^{\text{Hr}} \mathcal{YD}$. Using similar computations, we can check that the maps ev'_M and $coev'_M$ are morphisms in ${}_{H_f}^{\text{Hr}} \mathcal{YD}$. \square

4. THE CANONICAL ISOMORPHISMS IN ${}_{H_f}^{\text{Hr}} \mathcal{YD}^{\text{fd}}$

If \mathcal{C} is a braided rigid category then, for any two objects $M, N \in \mathcal{C}$, there exist two canonical isomorphisms in \mathcal{C} : $M \cong M^{**}$ and $(M \otimes N)^* \cong M^* \otimes N^*$. In this section our goal is to compute explicitly the above isomorphisms in the particular case $\mathcal{C} = {}_{H_f}^{\text{Hr}} \mathcal{YD}^{\text{fd}}$. Then we will specialize them for the category of finite dimensional modules over a quasitriangular quasi-Hopf algebra.

Let \mathcal{C} be a rigid monoidal category and M, N objects of \mathcal{C} . If $\nu : M \rightarrow N$ is a morphism in \mathcal{C} , following [14] we can define the transposes of ν as being the compositions

$$(4.1) \quad \begin{array}{ccccc}
\nu^* : N^* & \xrightarrow{l_{N^*}} & N^* \otimes \underline{1} & \xrightarrow{id_{N^*} \otimes coev_M} & N^* \otimes (M \otimes M^*) \\
& \xrightarrow{id_{N^*} \otimes (\nu \otimes id_{M^*})} & N^* \otimes (N \otimes M^*) & \xrightarrow{a_{N^*, N, M^*}^{-1}} & (N^* \otimes N) \otimes M^* \\
& \xrightarrow{ev_N \otimes id_{M^*}} & \underline{1} \otimes M^* & \xrightarrow{r_{M^*}^{-1}} & M^*,
\end{array}$$

and

$$(4.2) \quad \begin{array}{ccccc}
{}^* \nu : {}^* N & \xrightarrow{r_{{}^* N}} & \underline{1} \otimes {}^* N & \xrightarrow{coev'_M \otimes id_{{}^* N}} & ({}^* M \otimes M) \otimes {}^* N \\
& \xrightarrow{(id_{{}^* M} \otimes \nu) \otimes id_{{}^* N}} & ({}^* M \otimes N) \otimes {}^* N & \xrightarrow{a_{{}^* M, N, {}^* N}} & {}^* M \otimes (N \otimes {}^* N) \\
& \xrightarrow{id_{{}^* M} \otimes ev'_N} & {}^* M \otimes \underline{1} & \xrightarrow{l_{{}^* M}^{-1}} & {}^* M.
\end{array}$$

On the other hand, if \mathcal{C} is a rigid braided category then there exist functors $M \rightarrow M^*$ and $M \rightarrow {}^* M$ which are inverses to each other. Indeed, it is easy to see that $(M, coev_M : \underline{1} \rightarrow M \otimes M^*, ev_M : M^* \otimes M \rightarrow \underline{1})$ satisfies the axioms of left dual of M^* and $(M, coev'_M : \underline{1} \rightarrow {}^* M \otimes M, ev'_M : M \otimes {}^* M \rightarrow \underline{1})$ satisfies the axioms of right dual for ${}^* M$. Therefore we have an isomorphism $\theta_M : M \cong ({}^* M^*) \cong ({}^* M)^*$

defined as follows (see [16]):

$$(4.3) \quad \theta_M : M \begin{array}{ccc} \xrightarrow{l_M} & M \otimes \underline{1} & \xrightarrow{id_M \otimes \text{coev}_{*M}} \\ \xrightarrow{a_{M, *M, (*M)^*}^{-1}} & (M \otimes *M) \otimes (*M)^* & \xrightarrow{ev'_M \otimes id_{(*M)^*}} \\ \xrightarrow{r_{(*M)^*}^{-1}} & (*M)^* & \underline{1} \otimes (*M)^* \end{array} M \otimes (*M \otimes (*M)^*)$$

The inverse θ_M^{-1} is given by the composition

$$(4.4) \quad (*M)^* \begin{array}{ccc} \xrightarrow{l_{(*M)^*}} & (*M)^* \otimes \underline{1} & \xrightarrow{id_{(*M)^*} \otimes \text{coev}'_M} \\ \xrightarrow{a_{(*M)^*, *M, M}^{-1}} & ((*M)^* \otimes *M) \otimes M & \xrightarrow{ev_{*M} \otimes id_M} \\ & & \underline{1} \otimes M \xrightarrow{r_M^{-1}} M. \end{array} (*M)^* \otimes (*M \otimes M)$$

In a similar way, we have an isomorphism $\theta'_M : M \cong *(M^*)$. We also have a natural isomorphism $\Theta_M : M^* \rightarrow *M$ given by (see [2]):

$$(4.5) \quad \Theta_M : M^* \begin{array}{ccc} \xrightarrow{l_{M^*}} & M^* \otimes \underline{1} & \xrightarrow{id_{M^*} \otimes \text{coev}'_M} \\ \xrightarrow{a_{M^*, *M, M}^{-1}} & (M^* \otimes *M) \otimes M & \xrightarrow{c_{M^*, *M} \otimes id_M} \\ \xrightarrow{a_{*M, M^*, M}} & *M \otimes (M^* \otimes M) & \xrightarrow{id_{*M} \otimes ev_M} \\ \xrightarrow{l_{*M}^{-1}} & *M, & *M \otimes \underline{1} \end{array} M^* \otimes (*M \otimes M)$$

Its inverse is the following composition

$$(4.6) \quad \Theta_M^{-1} : *M \begin{array}{ccc} \xrightarrow{r_{*M}} & \underline{1} \otimes *M & \xrightarrow{\text{coev}_M \otimes id_{*M}} \\ \xrightarrow{a_{M, M^*, *M}} & M \otimes (M^* \otimes *M) & \xrightarrow{id_M \otimes c_{M^*, *M}^{-1}} \\ \xrightarrow{a_{M^*, *M, M}^{-1}} & (M \otimes *M) \otimes M^* & \xrightarrow{ev'_M \otimes id_{M^*}} \\ \xrightarrow{r_{M^*}^{-1}} & M^*. & \underline{1} \otimes M^* \end{array} (M \otimes M^*) \otimes *M$$

Thus the functors $M \rightarrow M^*$ and $M \rightarrow *M$ are naturally isomorphic. We conclude that

$$M^{**} = (M^*)^* \cong *(M^*) \cong M \cong (*M)^* \cong *(*M) = **M.$$

We will apply these results in the particular case where $\mathcal{C} = {}^H_H\mathcal{YD}^{\text{fd}}$.

1) The maps ν^* and $*\nu$ coincide with the usual transposed map of ν . Indeed, by (4.1) and (1.6), we have that

$$\begin{aligned} \nu^*(n^*) &= \sum_{i=1}^t \langle x^1 \cdot n^*, \alpha x^2 \beta \cdot \nu(i n) \rangle x^3 \cdot i n \\ &= \sum_{i=1}^t \langle n^*, S(x^1) \alpha x^2 \beta S(x^3) \cdot \nu(i n) \rangle i n = n^* \circ \nu, \end{aligned}$$

where $(i n)_{i=\overline{1, t}}$ is a basis of N and $(i n)_{i=\overline{1, t}}$ its dual basis. A similar computation shows that $*\nu(*n) = *n \circ \nu$ for any $*n \in *N$.

2) It is not hard to see that the map θ_M defined by (4.3) is given by

$$(4.7) \quad \theta_M(m) = \sum_{i=1}^n \langle i m, m \rangle^{*i} m$$

for all $m \in M$, where, if $({}^i m)_{i=\overline{1,n}}$ is a basis of M with dual basis $({}^i m)_{i=\overline{1,n}}$ in M^* , then ${}^*i m$ is the image of ${}^i m$ under the canonical map $M \rightarrow M^{**}$. Moreover, the morphism θ'_M is defined by the same formula (4.7) as θ_M . A straightforward but tedious computation proves that the morphism $\Theta_M : M^* \rightarrow {}^*M$ defined by (4.5) is given by

$$(4.8) \quad \Theta_M(m^*) = \sum_{i,j=1}^n \langle m^*, S(p^1)f^2 \cdot (g^1 \cdot {}_j m)_{(0)} \rangle \langle {}^j m, S(q^2)S^{-1}(q^1 S^{-1}(f^1(g^1 \cdot {}_j m)_{(-1)}g^2)p^2) \cdot {}^i m \rangle {}^i m,$$

for all $m^* \in M^*$. The inverse map $\Theta_M^{-1} : {}^*M \rightarrow M^*$ is given by (see (4.6))

$$(4.9) \quad \Theta_M^{-1}({}^*m) = \sum_{i,j=1}^n \langle {}^*m, V^2 \cdot {}_j m_{(0)} \rangle \langle {}^j m, \beta S(V^1 {}_j m_{(-1)}) \cdot {}^i m \rangle {}^i m,$$

for any ${}^*m \in {}^*M$, where

$$(4.10) \quad V = \sum V^1 \otimes V^2 = \sum S^{-1}(f^2 p^2) \otimes S^{-1}(f^1 p^1).$$

Therefore, we obtain the following result.

Proposition 4.1. *Let H be a quasi-Hopf algebra and $M, N \in {}^H_H \mathcal{YD}^{f.d.}$. Then ${}^r \Gamma_M = \theta'^{-1}_M \circ \Theta_{M^*} : M^{**} \rightarrow M$ is an isomorphism of Yetter-Drinfeld modules. Explicitly, ${}^r \Gamma_M$ is given by*

$$(4.11) \quad {}^r \Gamma_M(m^{**}) = \sum_{i=1}^n \langle m^{**}, {}^i m \rangle q^1 S^{-2}((S^2(p^1) \cdot {}^i m)_{(-1)}^{\mathcal{F}}) p^2 S^2(q^2) \cdot (S^2(p^1) \cdot {}^i m)_{(0)}^{\mathcal{F}}$$

for all $m^{**} \in M^{**}$, where

$$(4.12) \quad \mathcal{F} = \sum S(g^2)f^1 \otimes S(g^1)f^2.$$

The inverse of ${}^r \Gamma_M$ is given by ${}^r \Gamma_M^{-1} = \Theta_{M^*}^{-1} \circ \theta'_M$, that is,

$$(4.13) \quad {}^r \Gamma_M^{-1}(m) = \sum_{i=1}^n \langle {}^i m, S((p^1 \cdot m)_{(-1)} p^2) \alpha \cdot (p^1 \cdot m)_{(0)} \rangle {}^*i m.$$

Similarly, ${}^l \Gamma_M := \theta_M^{-1} \circ \Theta_{*M}^{-1} : {}^{**}M \rightarrow M$ provides an isomorphism of Yetter-Drinfeld modules. Explicitly, we have that

$$(4.14) \quad {}^l \Gamma_M({}^{**}m) = \sum_{i=1}^n \langle {}^{**}m, {}^i m \rangle V^1 g^1 S((S^{-1}(V^2 g^2) \cdot {}^i m)_{(-1)}) \alpha \cdot (S^{-1}(V^2 g^2) \cdot {}^i m)_{(0)},$$

for all ${}^{**}m \in {}^{**}M$. The inverse of ${}^l \Gamma_M$ is given by

$$(4.15) \quad {}^l \Gamma_M^{-1}(m) = \sum_{i=1}^n \langle {}^i m, S^{-2}(q^1(p^1 \cdot m)_{(-1)} p^2) q^2 \cdot (p^1 \cdot m)_{(0)} \rangle {}^*i m.$$

Proof. Straightforward, we leave the details to the reader. \square

Now, we will focus on the second isomorphism. Let \mathcal{C} be a braided rigid category and $M, N \in \mathcal{C}$. By [2] there exist natural isomorphisms

$$\begin{aligned} \sigma_{M,N}^* &: M^* \otimes N^* \rightarrow (M \otimes N)^* \\ {}^* \sigma_{M,N} &: {}^*M \otimes {}^*N \rightarrow {}^*(M \otimes N). \end{aligned}$$

In fact, $\sigma_{M,N}^* = \phi_{N,M}^* \circ c_{N^*,M^*}^{-1}$, where $\phi_{N,M}^* : N^* \otimes M^* \rightarrow (M \otimes N)^*$ is the composition:

$$\begin{aligned}
(4.16) \quad & r_{(M \otimes N)^*}^{-1} \circ \text{ev}_N \otimes (M \otimes N)^* \circ (N^* \otimes r_N^{-1}) \otimes (M \otimes N)^* \\
& \circ (N^* \otimes (\text{ev}_M \otimes N)) \otimes (M \otimes N)^* \circ N^* \otimes a_{M^*,M,N}^{-1} \otimes (M \otimes N)^* \\
& \circ a_{N^*,M^*,M \otimes N} \otimes (M \otimes N)^* \circ a_{N^* \otimes M^*,M \otimes N,(M \otimes N)^*}^{-1} \\
& \circ N^* \otimes M^* \otimes \text{coev}_{M \otimes N} \circ N^* \otimes M^* \otimes l_{N^* \otimes M^*}
\end{aligned}$$

The inverse of $\phi_{N,M}^*$ is given by the following composition.

$$\begin{aligned}
(4.17) \quad & r_{N^*}^{-1} \otimes M^* \circ (\text{ev}_{M \otimes N} \otimes N^*) \otimes M^* \circ a_{(M \otimes N)^*,M \otimes N,N^*}^{-1} \otimes \text{id}_{M^*} \\
& \circ ((M \otimes N)^* \otimes a_{M,N,N^*}^{-1}) \otimes M^* \circ ((M \otimes N)^* \otimes (M \otimes \text{coev}_N)) \otimes M^* \\
& \circ ((M \otimes N)^* \otimes l_M) \otimes M^* \circ a_{(M \otimes N)^*,M,M^*}^{-1} \\
& \circ (M \otimes N)^* \otimes \text{coev}_M \circ l_{(M \otimes N)^*}.
\end{aligned}$$

In a similar way, we can define ${}^* \phi_{N,M} : {}^* N \otimes {}^* M \rightarrow {}^*(M \otimes N)$ and its inverse ${}^* \phi_{N,M}^{-1}$; then we put ${}^* \sigma_{M,N} = {}^* \phi_{N,M} \circ c_{N^*,M^*}^{-1}$.

Proposition 4.2. *Let H be a quasi-Hopf algebra and $M, N \in {}_H^H \mathcal{YD}^{\text{fd}}$. Let $\{i m\}_{i=\overline{1,s}}$ and $\{i m\}_{i=\overline{1,s}}$ be dual bases in M and M^* , and $\{j n\}_{j=\overline{1,t}}$ and $\{j n\}_{j=\overline{1,t}}$ dual bases in N and N^* . Then the map $\sigma_{M,N}^* : M^* \otimes N^* \rightarrow (M \otimes N)^*$ defined by*

$$\begin{aligned}
(4.18) \quad \sigma_{M,N}^*(m^* \otimes n^*)(m \otimes n) &= \sum \langle m^*, f^2 \tilde{q}_2^2 X^3 S^{-1}(\tilde{q}^1 X^1(p^1 \cdot n)_{(-1)} p^2) \cdot m \rangle \\
& \langle n^*, f^1 \tilde{q}_1^2 X^2 \cdot (p^1 \cdot n)_{(0)} \rangle
\end{aligned}$$

is an isomorphism of Yetter-Drinfeld modules. Here $p_R = \sum p^1 \otimes p^2$ and $q_L = \sum \tilde{q}^1 \otimes \tilde{q}^2$ are the elements defined in (1.17) and (1.20). The inverse of $\sigma_{M,N}^*$ is given by

$$(4.19) \quad \sigma_{M,N}^{*-1}(\mu) = \sum_{i,j=1}^{s,t} \langle \mu, (g^1 \cdot j n)_{(-1)} g^2 \cdot i m \otimes (g^1 \cdot j n)_{(0)} \rangle i m \otimes j n.$$

In a similar way, the map ${}^* \sigma_{M,N} : {}^* M \otimes {}^* N \rightarrow {}^*(M \otimes N)$ defined by

$$\begin{aligned}
(4.20) \quad {}^* \sigma_{M,N}({}^* m \otimes {}^* n)(m \otimes n) &= \sum \langle {}^* m, S^{-1}(f^2 q^1(x^1 \tilde{p}^1 S^{-1}(f^1) \cdot n)_{(-1)} x^2 \tilde{p}_1^2) \cdot m \rangle \\
& \langle {}^* n, S^{-1}(x^3 \tilde{p}_2^2) q^2 \cdot (x^1 \tilde{p}^1 S^{-1}(f^1) \cdot n)_{(0)} \rangle
\end{aligned}$$

is an isomorphism of Yetter-Drinfeld modules; $q_R = \sum q^1 \otimes q^2$ and $p_L = \sum \tilde{p}^1 \otimes \tilde{p}^2$ are the elements defined in (1.18) and (1.19). The inverse of ${}^* \sigma_{M,N}$ is given by

$$(4.21) \quad {}^* \sigma_{M,N}^{-1}(\nu) = \sum_{i,j=1}^{s,t} \langle \nu, (S^{-1}(g^2) \cdot j n)_{(-1)} S^{-1}(g^1) \cdot i m \otimes (S^{-1}(g^2) \cdot j n)_{(0)} \rangle i m \otimes j n.$$

Proof. We will show that $\sigma_{M,N}^* = \phi_{N,M}^* \circ c_{N^*,M^*}^{-1}$, where $\phi_{N,M}^*$ is given by (4.16). We first calculate $\phi_{N,M}^*$. As before, we write $q_R = \sum q^1 \otimes q^2$ and $p_R = \sum p^1 \otimes p^2 =$

$\sum P^1 \otimes P^2$, and then compute for all $n^* \in N^*$, $m^* \in M^*$, $m \in M$ and $n \in N$ that:

$$\begin{aligned}
& \phi_{N,M}^*(n^* \otimes m^*)(m \otimes n) \\
(4.16) \quad &= \sum \langle m^*, S(x^1 X^2 y_2^1) \alpha x^2 (X^3 y^2 \beta S(y^3))_1 \cdot m \rangle \\
& \quad \langle n^*, S(X^1 y_1^1) \alpha x^3 (X^3 y^2 \beta S(y^3))_2 \cdot n \rangle \\
(1.12) \quad &= \sum \langle m^*, S(y_2^1) \gamma^1 (y^2 \beta S(y^3))_1 \cdot m \rangle \langle n^*, S(y_1^1) \gamma^2 (y^2 \beta S(y^3))_2 \cdot n \rangle \\
(1.11, 1.17) \quad &= \sum \langle m^*, f^1 S(p^1)_1 g^1 \gamma^1 p_1^2 \cdot m \rangle \langle n^*, f^2 S(p^1)_2 g^2 \gamma^2 p_2^2 \cdot n \rangle \\
(1.15) \quad &= \sum \langle m^*, f^1 (S(p^1) \alpha p^2)_1 \cdot m \rangle \langle n^*, f^2 (S(p^1) \alpha p^2)_2 \cdot n \rangle \\
(1.17, 1.6) \quad &= \sum \langle m^*, f^1 \cdot m \rangle \langle n^*, f^2 \cdot n \rangle.
\end{aligned}$$

Using (4.17) or by a direct computation it is easy to see that

$$(4.22) \quad \phi_{N,M}^{*-1}(\mu) = \sum_{i,j=1}^{s,t} \langle \mu, g^1 \cdot_i m \otimes g^2 \cdot_j n \rangle^j n \otimes^i m$$

for all $\mu \in (M \otimes N)^*$. Also, it is not hard to see that (1.9) and (1.16) imply that

$$(4.23) \quad \sum q^1 g_1^1 \otimes S(q^2 g_2^1) g^2 = \sum S(X^3) f^1 \otimes S(X^1 \beta S(X^2) f^2).$$

The same relations and (2.26) imply that

$$(4.24) \quad \sum S(p^2 f^1) F^1 f_1^2 \otimes S(p^1) F^2 f_2^2 = q_L.$$

Using (4.23), the axioms of a quasi-Hopf algebra, again (1.9) and (1.16), and finally (1.27) we obtain the following relation:

$$\begin{aligned}
(4.25) \quad & \sum q^1 [g^1 S(\tilde{q}_1^2 X^2) f^2]_1 \otimes S(q^2 [g^1 S(\tilde{q}_1^2 X^2) f^2]_2) g^2 S(\tilde{q}^1 X^1) \otimes S(\tilde{q}^2 X^3) f^1 \\
&= \sum f^2 F_2^2 x^2 \otimes S(F^2 x^3) \otimes f^1 F_1^1 x^1.
\end{aligned}$$

Let $f = \sum F^1 \otimes F^2 = \sum \mathbf{F}^1 \otimes \mathbf{F}^2$ and $p_R = \sum P^1 \otimes P^2$ be other copies of f and p_R . For $m^* \in M^*$, $n^* \in N^*$, $m \in M$ and $n \in N$ we calculate:

$$\begin{aligned}
& (\phi_{N,M}^* \circ c_{N^*,M^*}^{-1})(m^* \otimes n^*)(m \otimes n) \\
(2.6, 3.4) \quad &= \sum \langle m^*, \tilde{q}^1 X^1 S^{-1}(F^1 (g^1 S(\tilde{q}_1^2 X^2) f^2 \cdot n)_{(-1)} g^2) p^2 S(\tilde{q}_2^2 X^3) f^1 \cdot m \rangle \\
& \quad \langle n^*, S(p^1) F^2 \cdot (g^1 S(\tilde{q}_1^2 X^2) f^2 \cdot n)_{(0)} \rangle \\
(3.1, 4.25) \quad &= \sum \langle m^*, S^{-1}(\mathbf{F}^1 f_1^2 F_{(2,1)}^1 x_1^2 (P^1 \cdot n)_{(-1)} P^2 S(F^2 x^3)) p^2 f^1 F_1^1 x^1 \cdot m \rangle \\
& \quad \langle n^*, S(p^1) \mathbf{F}^2 f_2^2 F_{(2,2)}^1 x_2^2 \cdot (P^1 \cdot n)_{(0)} \rangle \\
(4.25, 4.24) \quad &= \sum \langle m^*, S^{-1}(\tilde{q}^1 F_{(2,1)}^1 x_1^2 (P^1 \cdot n)_{(-1)} P^2 S(F^2 x^3)) F_1^1 x^1 \rangle \\
& \quad \langle n^*, F_2^1 \tilde{q}^2 x_2^2 \cdot (P^1 \cdot n)_{(0)} \rangle \\
(1.24, 1.3) \quad &= \sum \langle m^*, F^2 \tilde{q}_2^2 X^3 S^{-1}(\tilde{q}^1 X^1 (P^1 \cdot n)_{(-1)} P^2) \cdot m \rangle \\
& \quad \langle n^*, F^1 \tilde{q}_1^2 X^2 \cdot (P^1 \cdot n)_{(0)} \rangle \\
(4.18) \quad &= \sigma_{M,N}^*(m^* \otimes n^*)(m \otimes n).
\end{aligned}$$

Obviously, the inverse of $\sigma_{M,N}^*$ is $\sigma_{M,N}^{*-1} = c_{N^*,M^*} \circ \phi_{N,M}^{*-1}$. By (4.22) it follows that $\sigma_{M,N}^{*-1}$ is defined by the formula given in (4.19). The assertion concerning the

morphism ${}^*\sigma_{M,N}$ can be proved in a similar way. We only notice that

$${}^*\phi_{N,M}({}^*n \otimes {}^*m)(m \otimes n) = \sum \langle {}^*m, S^{-1}(f^2) \cdot m \rangle \langle {}^*n, S^{-1}(f^1) \cdot n \rangle$$

for all ${}^*m \in {}^*M$, ${}^*n \in {}^*N$, $m \in M$ and $n \in N$. \square

Let (H, R) be a quasitriangular quasi-Hopf algebra. Then we have a monoidal functor $\mathbb{F} : {}_H\mathcal{M} \rightarrow {}_H^H\mathcal{YD}$ which acts as the identity on objects and morphisms. For $M \in {}_H\mathcal{M}$, $\mathbb{F}(M) = M$ as a left H -module, and with left H -coaction given by

$$(4.26) \quad \lambda_M(m) = \sum R^2 \otimes R^1 \cdot m,$$

for all $m \in M$. Moreover, this functor sends algebras, coalgebras, bialgebras etc. in ${}_H\mathcal{M}$ to the corresponding objects in ${}_H^H\mathcal{YD}$ (see [6, Proposition 2.4]).

Corollary 4.3. ([8, Lemma 3.2]). *Let H be a quasitriangular quasi-Hopf algebra and M a finite dimensional left H -module. Then $M \cong M^{**}$ and $M \cong {}^{**}M$ as H -modules.*

Proof. We have seen above that M can be viewed as an object in ${}_H^H\mathcal{YD}$, so $M \cong M^{**}$ and $M \cong {}^{**}M$ as Yetter-Drinfeld modules, cf. Proposition 4.1. Thus, $M \cong M^{**}$ and $M \cong {}^{**}M$ as H -modules. We will compute explicitly these isomorphisms. Applying (1.40) twice, we obtain that

$$(4.27) \quad \mathcal{F}_{21}R\mathcal{F}^{-1} = (S^2 \otimes S^2)(R),$$

where \mathcal{F} is the twist defined by (4.12) and $\mathcal{F}_{21} = \sum \mathcal{F}^2 \otimes \mathcal{F}^1$. We set $\mathcal{F}^{-1} = \sum \mathcal{G}^1 \otimes \mathcal{G}^2$ and then we compute the isomorphism ${}^r\Gamma_M : M^{**} \rightarrow M$:

$$\begin{aligned} {}^r\Gamma_M(m^{**}) &\stackrel{(4.11, 4.26)}{=} \sum_{i=1}^n \langle m^{**}, {}^i m \rangle q^1 S^{-2}(\mathcal{F}^1 R^2 \mathcal{G}^2) p^2 S^2(q^2) \mathcal{F}^2 R^1 \mathcal{G}^1 S^2(p^1) \cdot {}_i m \\ (4.27) \quad &= \sum_{i=1}^n \langle m^{**}, {}^i m \rangle q^1 R^2 p^2 S^2(q^2) S^2(R^1) S^2(p^1) \cdot {}_i m \\ (1.18, 1.38) \quad &= \sum_{i=1}^n \langle m^{**}, {}^i m \rangle u^{-1} \cdot {}_i m, \end{aligned}$$

for all $m^{**} \in M^{**}$. A similar computation tells us that

$${}^r\Gamma_M^{-1}(m) = \sum_{i=1}^n \langle {}^i m, u \cdot m \rangle {}^{*i} m,$$

for all $m \in M$, which is equivalent to ${}^r\Gamma_M^{-1}(m)(\varphi) = \varphi(u \cdot m)$ for all $\varphi \in M^*$ and $m \in M$. In this way we have recovered the isomorphism $\hat{\psi} : M \rightarrow M^{**}$ given in [8]. Moreover, in this particular case, ${}^l\Gamma_M : {}^{**}M \rightarrow M$ is given by ${}^l\Gamma({}^{**}m) = \sum_{i=1}^n \langle {}^{**}m, {}^i m \rangle u \cdot {}_i m$ for all ${}^{**}m \in {}^{**}M$. Indeed, by (4.14) and (4.26) we

have that

$$\begin{aligned}
{}^l\Gamma_M(**m) &\stackrel{(1.40)}{=} \sum_{i=1}^n \langle **m, {}^i m \rangle V^1 R^2 g^2 \alpha S^{-1}(V^2 R^1 g^1) \cdot {}_i m \\
(2.26, 4.10) &= \sum_{i=1}^n \langle **m, {}^i m \rangle S^{-1}(S(R^2) f^2 p^2) S^{-1}(\beta) S^{-2}(S(R^1) f^1 p^1) \cdot {}_i m \\
(1.40, 2.26) &= \sum_{i=1}^n \langle **m, {}^i m \rangle S^{-2}(S(R^2) p^2 \alpha R^1 p^1) \cdot {}_i m \\
(1.37) &= \sum_{i=1}^n \langle **m, {}^i m \rangle S^{-2}(u) \cdot {}_i m = \sum_{i=1}^n \langle **m, {}^i m \rangle u \cdot {}_i m.
\end{aligned}$$

Now, it follows that ${}^l\Gamma_M^{-1}(m) = \sum_{i=1}^n \langle {}^i m, u^{-1} \cdot m \rangle {}^* i m$ for all $m \in M$, and this finishes our computation. \square

Remark 4.4. The element u plays a central role in the theory of quasitriangular Hopf algebras. It is therefore natural to try to generalize its properties to the quasi-Hopf algebra setting. The major problem in [5] was to find the inverse of the element u defined in [1]. Now, if we forget the definitions of u and u^{-1} in the quasi-Hopf case, combining Corollary 4.3 with the similar result in the Hopf case we will obtain in a natural way the definitions for u and its inverse u^{-1} .

Corollary 4.5. ([8, Lemma 3.3]). *Let (H, R) be a quasitriangular quasi-Hopf algebra and M, N two finite dimensional left H -modules. Then $M^* \otimes N^* \cong (M \otimes N)^*$ and ${}^* M \otimes {}^* N \cong {}^*(M \otimes N)$ as H -modules.*

Proof. We regard M, N as objects in ${}^H_H \mathcal{YD}^{\text{fd}}$ via the functor \mathbb{F} defined in (4.26). Thus, by Proposition 4.2, we obtain that $M^* \otimes N^* \cong (M \otimes N)^*$ and ${}^* M \otimes {}^* N \cong {}^*(M \otimes N)$ as Yetter-Drinfeld modules, so they are isomorphic also as H -modules. Moreover, (1.42) implies that the isomorphism $\sigma_{M,N}^* : M^* \otimes N^* \rightarrow (M \otimes N)^*$ defined in (4.18) is given by

$$\sigma_{M,N}^*(m^* \otimes n^*)(m \otimes n) = \sum \langle m^*, f^2 \bar{R}^2 \cdot m \rangle \langle n^*, f^1 \bar{R}^1 \cdot n \rangle$$

for all $m^* \in M^*, n^* \in N^*, m \in M, n \in N$, where $R^{-1} := \sum \bar{R}^1 \otimes \bar{R}^2$ is the inverse of the R -matrix R . Note that it is just the isomorphism $\mu_{M,N}$ defined in [8]. Also, (1.36) implies that the isomorphism ${}^* \sigma_{M,N} : {}^* M \otimes {}^* N \rightarrow {}^*(M \otimes N)$ defined by (4.20) is given by

$${}^* \sigma_{M,N}({}^* m \otimes {}^* n)(m \otimes n) = \sum \langle {}^* m, S^{-1}(f^2 \bar{R}^2) \cdot m \rangle \langle {}^* n, S^{-1}(f^1 \bar{R}^1) \cdot n \rangle,$$

for all ${}^* m \in {}^* M, {}^* n \in {}^* N, m \in M$ and $n \in N$. Finally, it is not hard to see that

$$\sigma_{M,N}^{*-1}(\mu) = \sum_{i,j=1}^{s,t} \langle \mu, R^2 g^2 \cdot {}_i m \otimes R^1 g^1 \cdot {}_j n \rangle {}^i m \otimes {}^j n$$

for all $\mu \in (M \otimes N)^*$, and

$$\begin{aligned}
{}^* \sigma_{M,N}^{-1}(\nu) &\stackrel{(4.20,4.26)}{=} \sum_{i,j=1}^{s,t} \langle \nu, R^2 S^{-1}(g^1) \cdot {}_i m \otimes R^1 S^{-1}(g^2) \cdot {}_j n \rangle {}^i m \otimes {}^j n \\
(1.40) \quad &= \sum_{i,j=1}^{s,t} \langle \nu, S^{-1}(R^2 g^2) \cdot {}_i m \otimes S^{-1}(R^1 g^1) \cdot {}_j n \rangle {}^i m \otimes {}^j n
\end{aligned}$$

for any $\nu \in {}^*(M \otimes N)$, completing the proof. \square

Remark 4.6. Continuing the ideas of Remark 4.4, we notice that the above Corollary suggests the two formulae (1.36) and (1.42) for the inverse of the R -matrix R , and, also, the formula (1.40). All these formulae were first proved by Hauser and Nill [13] in the case that (H, R) is a finite dimensional quasitriangular quasi-Hopf algebra. They also used the bijectivity of the antipode.

5. APPLICATIONS

Let B be a Hopf algebra in a braided rigid category \mathcal{C} . Then B^* and *B are also Hopf algebras in \mathcal{C} , see [2, 19]. The structure maps on B^* are the following ones:

$$(5.1) \quad m_{B^*} : B^* \otimes B^* \xrightarrow{\sigma_{B,B}^*} (B \otimes B)^* \xrightarrow{\Delta_B^*} B^*,$$

$$(5.2) \quad \Delta_{B^*} : B^* \xrightarrow{m_B^*} (B \otimes B)^* \xrightarrow{\sigma_{B,B}^{*-1}} B^* \otimes B^*,$$

$$(5.3) \quad S_{B^*} = S_B^*, \quad u_{B^*} = \varepsilon_B^*, \quad \varepsilon_{B^*} = u_B^*.$$

The structure maps on the left dual can be obtained in a similar way.

We can consider algebras, coalgebras, bialgebras and Hopf algebras in the braided category ${}^H_H\mathcal{YD}$. More precisely, an algebra in ${}^H_H\mathcal{YD}$ is an object $B \in {}^H_H\mathcal{YD}$ with the additional structure:

- B is a left H -module algebra, i.e. B has a multiplication and a unit 1_B satisfying the conditions:

$$\begin{aligned}
(ab)c &= \sum (X^1 \cdot a)[(X^2 \cdot b)(X^3 \cdot c)], \\
h \cdot (ab) &= \sum (h_1 \cdot a)(h_2 \cdot b), \\
(5.4) \quad h \cdot 1_B &= \varepsilon(h)1_B,
\end{aligned}$$

for all $a, b, c \in B$ and $h \in H$.

- B is a quasi-comodule algebra, that is, the multiplication m and the unit η of B intertwine the H -coaction λ_B . By (2.4), this means:

$$\begin{aligned}
\lambda_B(bb') &= \sum X^1(x^1 Y^1 \cdot b)_{(-1)} x^2 (Y^2 \cdot b')_{(-1)} Y^3 \\
(5.5) \quad &\otimes [X^2 \cdot (x^1 Y^1 \cdot b)_{(0)}][X^3 x^3 \cdot (Y^2 \cdot b')_{(0)}],
\end{aligned}$$

for all $b, b' \in B$, and

$$(5.6) \quad \lambda_B(1_B) = 1_H \otimes 1_B.$$

A coalgebra in ${}^H_H\mathcal{YD}$ is an object B with

- B is a left H -module coalgebra, i.e. B has a comultiplication $\underline{\Delta}_B : B \rightarrow B \otimes B$ (we will denote $\underline{\Delta}(b) = \sum b_{\underline{1}} \otimes b_{\underline{2}}$) and a usual counit $\underline{\varepsilon}_B$ such that:

$$(5.7) \quad \sum X^1 \cdot b_{(\underline{1}, \underline{1})} \otimes X^2 \cdot b_{(\underline{1}, \underline{2})} \otimes X^3 \cdot b_{\underline{2}} = \sum b_{\underline{1}} \otimes b_{(\underline{2}, \underline{1})} \otimes b_{(\underline{2}, \underline{2})},$$

$$(5.8) \quad \underline{\Delta}_B(h \cdot b) = \sum h_1 \cdot b_{\underline{1}} \otimes h_2 \cdot b_{\underline{2}}, \quad \underline{\varepsilon}_B(h \cdot b) = \varepsilon(h) \underline{\varepsilon}_B(b),$$

for all $h \in H$, $b \in B$, where we adopt for the quasi-coassociativity of $\underline{\Delta}_B$ the same notations as in the Section 1.

- B is a quasi-comodule coalgebra, i.e. the comultiplication $\underline{\Delta}_B$ and the counit $\underline{\varepsilon}_B$ intertwine the H -coaction λ_B . Explicitly, for all $b \in B$ we must have that:

$$(5.9) \quad \begin{aligned} \sum b_{(-1)} \otimes b_{(0)\underline{1}} \otimes b_{(0)\underline{2}} &= \sum X^1(x^1 Y^1 \cdot b_{\underline{1}})_{(-1)} x^2(Y^2 \cdot b_{\underline{2}})_{(-1)} Y^3 \otimes \\ &X^2 \cdot (x^1 Y^1 \cdot b_{\underline{1}})_{(0)} \otimes X^3 x^3 \cdot (Y^2 \cdot b_{\underline{2}})_{(0)}, \end{aligned}$$

and

$$(5.10) \quad \sum \underline{\varepsilon}_B(b_{(0)}) b_{(-1)} = \underline{\varepsilon}_B(b) 1.$$

An object $B \in {}^H_H\mathcal{YD}$ is a bialgebra if it is an algebra and a coalgebra in ${}^H_H\mathcal{YD}$ such that $\underline{\Delta}_B$ is an algebra morphism, i.e. $\underline{\Delta}_B(1_B) = 1_B \otimes 1_B$ and, by (1.51) and (2.5), for all $b, b' \in B$ we have that:

$$(5.11) \quad \begin{aligned} \underline{\Delta}_B(bb') &= \sum [y^1 X^1 \cdot b_{\underline{1}}] [y^2 Y^1(x^1 X^2 \cdot b_{\underline{2}})_{(-1)} x^2 X_1^3 \cdot b'_{\underline{1}}] \\ &\otimes [y_1^3 Y^2 \cdot (x^1 X^2 \cdot b_{\underline{2}})_{(0)}] [y_2^3 Y^3 x^3 X_2^3 \cdot b'_{\underline{2}}]. \end{aligned}$$

Finally, a bialgebra B in ${}^H_H\mathcal{YD}$ is a Hopf algebra if there exists a morphism $\underline{S} : B \rightarrow B$ in ${}^H_H\mathcal{YD}$ such that $\sum \underline{S}(b_{\underline{1}}) b_{\underline{2}} = \sum b_{\underline{1}} \underline{S}(b_{\underline{2}}) = \underline{\varepsilon}_B(b) 1$ for all $b \in B$; we then say that B is a braided Hopf algebra.

If $B \in {}^H_H\mathcal{YD}^{\text{fd}}$ is a Hopf algebra then B^* and *B become Hopf algebras in ${}^H_H\mathcal{YD}$. We will compute the structure maps.

Proposition 5.1. *Let H be a quasi-Hopf algebra and B a finite dimensional Hopf algebra in ${}^H_H\mathcal{YD}$. Then B^* is a Hopf algebra in ${}^H_H\mathcal{YD}$ with the following structure maps:*

$$(5.12) \quad \begin{aligned} (a^* \circ b^*)(b) &= \sum \langle a^*, f^2 \tilde{q}_2^2 X^3 S^{-1}(\tilde{q}^1 X^1(p^1 \cdot b_{\underline{2}})_{(-1)} p^2) \cdot b_{\underline{1}} \rangle \\ &\langle b^*, f^1 \tilde{q}_1^2 X^2 \cdot (p^1 \cdot b_{\underline{2}})_{(0)} \rangle \end{aligned}$$

$$(5.13) \quad \Delta_{B^*}(b^*) = \sum_{i,j=1}^n \langle b^*, [(g^1 \cdot j b)_{(-1)} g^2 \cdot i b] (g^1 \cdot j b)_{(0)} \rangle^i b \otimes^j b$$

$$(5.14) \quad S_{B^*}(b^*) = b^* \circ \underline{S}, \quad u_{B^*}(1) = \underline{\varepsilon}_B, \quad \varepsilon_{B^*}(b^*) = b^*(1_B),$$

for all $a^*, b^* \in B^*$ and $b \in B$, where $\{ib\}_{i=\overline{1,n}}$ is a basis in B with dual basis $\{{}^i b\}_{i=\overline{1,n}}$ in B^* . Similarly, *B is a Hopf algebra in ${}^H_H\mathcal{YD}$ with the following structure

maps:

$$(5.15) \quad \begin{aligned} (*a \bar{\circ} *b)(b) &= \sum \langle *a, S^{-1}(f^2 q^1 (x^1 \tilde{p}^1 S^{-1}(f^1) \cdot b_2)_{(-1)} x^2 \tilde{p}_1^2) \cdot b_1 \rangle \\ &\quad \langle *b, S^{-1}(x^3 \tilde{p}_2^2) q^2 \cdot (x^1 \tilde{p}^1 S^{-1}(f^1) \cdot b_2)_{(0)} \rangle \end{aligned}$$

$$(5.16) \quad \begin{aligned} \Delta_{*B}(b^*) &= \sum_{i,j=1}^n \langle *b, [(S^{-1}(g^2) \cdot {}_j b)_{(-1)} \\ &\quad S^{-1}(g^1) \cdot {}_i b] (S^{-1}(g^2) \cdot {}_j b)_{(0)} \rangle {}^i b \otimes {}^j b \end{aligned}$$

$$(5.17) \quad S_{*B}(*b) = *b \circ \underline{S}, \quad u_{*B}(1) = \underline{\varepsilon}_B, \quad \varepsilon_{*B}(*b) = *b(1_B),$$

for all $*a, *b \in *B$ and $b \in B$.

Proof. Follows easily by computing the morphisms in (5.1-5.3), using (4.18-4.21); the details are left to the reader. \square

Let H be a Hopf algebra. It is well-known that H becomes an algebra in the monoidal category ${}^H_H\mathcal{YD}$, with Yetter-Drinfeld structure given by

$$h \triangleright h' = \sum h_1 h' S(h_2), \quad \lambda(h) = \sum h_1 \otimes h_2$$

for all $h, h' \in H$. Moreover, H is quantum commutative as an algebra in ${}^H_H\mathcal{YD}$ (see for example [9]). Now, let H be a quasi-Hopf algebra. In [7] a new multiplication \circ on H is introduced, given by the formula

$$(5.18) \quad h \circ h' = \sum X^1 h S(x^1 X^2) \alpha x^2 X_1^3 h' S(x^3 X_2^3)$$

Let H_0 be equal to H as a vector space, with multiplication \circ . Then H_0 is a left H -module algebra with unit β and left H -action given by

$$(5.19) \quad h \triangleright h' = \sum h_1 h' S(h_2),$$

for all $h, h' \in H$. It was also shown in [6] that H_0 is an algebra in the category ${}^H_H\mathcal{YD}$, with H -coaction given by

$$(5.20) \quad \lambda_H(h) = \sum h_{(-1)} \otimes h_{(0)} = \sum X^1 Y_1^1 h_1 g^1 S(q^2 Y_2^2) Y^3 \otimes X^2 Y_2^1 h_2 g^2 S(X^3 q^1 Y_1^2),$$

where $f^{-1} = \sum g^1 \otimes g^2$ and $q = q_R = \sum q^1 \otimes q^2$ are the elements defined by (1.14) and (1.18). In addition, in [4] it was shown that H_0 is actually quantum commutative as an algebra in ${}^H_H\mathcal{YD}$.

If (H, R) is quasitriangular, then H_0 is a Hopf algebra with bijective antipode in ${}_H\mathcal{M}$, with the additional structure (see [6]):

$$(5.21) \quad \begin{aligned} \underline{\Delta}(h) &= \sum h_{\underline{1}} \otimes h_{\underline{2}} \\ &= \sum x^1 X^1 h_1 g^1 S(x^2 R^2 y^3 X_2^3) \otimes x^3 R^1 \triangleright y^1 X^2 h_2 g^2 S(y^2 X_1^3), \end{aligned}$$

$$(5.22) \quad \underline{\varepsilon}(h) = \varepsilon(h),$$

$$(5.23) \quad \underline{S}(h) = \sum X^1 R^2 p^2 S(q^1 (X^2 R^1 p^1 \triangleright h) S(q^2) X^3),$$

for all $h \in H$, where $R = \sum R^1 \otimes R^2$ and $f^{-1} = \sum g^1 \otimes g^2$, $p_R = \sum p^1 \otimes p^2$ and $q_R = \sum q^1 \otimes q^2$ are the elements defined by (1.14), (1.17) and (1.18). If we consider the left H -coaction

$$(5.24) \quad \lambda_{H_0}(h) = \sum R^2 \otimes R^1 \triangleright h,$$

induced by (4.26), then H_0 becomes a Hopf algebra in ${}^H_H\mathcal{YD}$, with bijective antipode. From now on, we will refer to H_0 as a Hopf algebra in ${}^H_H\mathcal{M}$, and, via the monoidal functor $\mathbb{F} : {}^H_H\mathcal{M} \rightarrow {}^H_H\mathcal{YD}$, in ${}^H_H\mathcal{YD}$, with structure maps (5.18), (5.19) and (5.21-5.24).

If H is finite dimensional, then H_0^* is also a Hopf algebra in ${}^H_H\mathcal{YD}$. By Proposition 3.2 and (1.40), H_0^* is a Yetter-Drinfeld module via

$$(5.25) \quad (h \mapsto \varphi)(h') = \varphi(S(h) \triangleright h')$$

$$(5.26) \quad \lambda_{H_0^*}(\varphi) = \sum R^2 \otimes R^1 \mapsto \varphi$$

for all $\varphi \in H^*$ and $h, h' \in H$. The structure of H_0^* as a Hopf algebra in ${}^H_H\mathcal{YD}$ is given by:

$$(5.27) \quad (\varphi \circ \Psi)(h) = \sum \langle \varphi, f^2 \bar{R}^2 \triangleright h_1 \rangle \langle \Psi, f^1 \bar{R}^1 \triangleright h_2 \rangle,$$

$$(5.28) \quad 1_{H_0^*} = \underline{\varepsilon},$$

$$(5.29) \quad \Delta_{H_0^*}(\varphi) = \sum_{i,j=1}^n \langle \varphi, (R^2 g^2 \triangleright i e) \circ (R^1 g^1 \triangleright j e) \rangle^i e \otimes^j e,$$

$$(5.30) \quad \varepsilon_{H_0^*}(\varphi) = \varphi(\beta),$$

$$(5.31) \quad S_{H_0^*}(\varphi) = \varphi \circ \underline{S},$$

for all $h \in H$ and $\varphi, \Psi \in H^*$, where $R^{-1} = \sum \bar{R}^1 \otimes \bar{R}^2$, $\{i e\}_{i=1, \dots, n}$ is a basis of H and $\{^i e\}_{i=1, \dots, n}$ the corresponding dual basis of H^* . The left dual *H_0 of H_0 is also a Hopf algebra in ${}^H_H\mathcal{YD}$. First, by Proposition 3.2 and (1.40), *H_0 is a Yetter-Drinfeld module via

$$(5.32) \quad (h \succ \varphi)(h') = \varphi(S^{-1}(h) \triangleright h')$$

$$(5.33) \quad \lambda_{{}^*H_0}(\varphi) = \sum R^2 \otimes R^1 \succ \varphi$$

for all $\varphi \in H^*$ and $h, h' \in H$. Then the structure of *H_0 as a Hopf algebra in ${}^H_H\mathcal{YD}$ is given by the formulae

$$(5.34) \quad (\varphi \circ \Psi)(h) = \sum \langle \varphi, S^{-1}(f^2 \bar{R}^2) \triangleright h_1 \rangle \langle \Psi, S^{-1}(f^1 \bar{R}^1) \triangleright h_2 \rangle,$$

$$(5.35) \quad 1_{{}^*H_0} = \underline{\varepsilon},$$

$$(5.36) \quad \Delta_{{}^*H_0}(\varphi) = \sum_{i,j=1}^n \langle \varphi, [S^{-1}(R^2 g^2) \triangleright i e] \circ [S^{-1}(R^1 g^1) \triangleright j e] \rangle^i e \otimes^j e,$$

$$(5.37) \quad \varepsilon_{{}^*H_0}(\varphi) = \varphi(\beta),$$

$$(5.38) \quad S_{{}^*H_0}(\varphi) = \varphi \circ \underline{S},$$

for all $\varphi \in H^*$ and $h, h' \in H$. From (4.5) we know that $\Theta_{H_0} : H_0^* \rightarrow {}^*H_0$ is an isomorphism of Yetter-Drinfeld modules. In this particular case, Θ_{H_0} is given by

$$\begin{aligned} \Theta_{H_0}(\varphi) &\stackrel{(4.8, 4.26)}{=} \sum_{i,j=1}^n \langle \varphi, S(p^1) f^2 R^1 g^1 \triangleright j e \rangle \\ &\quad \langle j e, S(q^2) S^{-1}(q^1 S^{-1}(f^1 R^2 g^2) p^2) \triangleright i e \rangle^i e \end{aligned}$$

$$(1.40) = \sum_{i,j=1}^n \langle \varphi, S(R^1 p^1) \triangleright_j e \rangle \langle {}^j e, S^{-1}(q^1 R^2 p^2 S^2(q^2)) \triangleright_i e \rangle^i e$$

$$(1.38) = \sum_{i=1}^n \langle \varphi, S^{-1}(u^{-1}) \triangleright_i e \rangle^i e = u^{-1} \succ \varphi$$

for all $\varphi \in H_0^*$. Since $S^2(u) = u$ it is easy to see that $\Theta_{H_0}^{-1}(\varphi) = u \mapsto \varphi$ for all $\varphi \in {}^*H_0$.

Proposition 5.2. *Let (H, R) be a triangular quasi-Hopf algebra. With notation as above, $\Theta_{H_0} : H_0^* \rightarrow {}^*H_0$ is a braided Hopf algebra isomorphism.*

Proof. We only have to prove that Θ_{H_0} is an algebra and coalgebra morphism in ${}^H_H\mathcal{YD}$. We need the formula:

$$(5.39) \quad \Delta(S(u)) = \sum (R_{21}R)^{-1}(S(u) \otimes S(u))(S \otimes S)(f_{21}^{-1})f,$$

which can be found in [5]. Since (H, R) is triangular and $S(u^{-1}) = S^{-1}(u^{-1})$ it follows that

$$(5.40) \quad \Delta(S^{-1}(u^{-1})) = \sum f^{-1}(S \otimes S)(f_{21})(S^{-1}(u^{-1}) \otimes S^{-1}(u^{-1})).$$

Now, for all $\varphi, \Psi \in H_0^*$ and $h \in H$ we compute:

$$\begin{aligned} \Theta_{H_0}(\varphi \circ \Psi)(h) &= \sum \langle \varphi \circ \Psi, S^{-1}(u^{-1}) \triangleright h \rangle \\ (5.27, 5.8) &= \sum \langle \varphi, f^2 \bar{R}^2 S^{-1}(u^{-1})_1 \triangleright h_{\underline{1}} \rangle \langle \Psi, f^1 \bar{R}^1 S^{-1}(u^{-1})_2 \triangleright h_{\underline{2}} \rangle \\ (5.40, 1.40) &= \sum \langle \varphi, S(f^2 \bar{R}^2) S^{-1}(u^{-1}) \triangleright h_{\underline{1}} \rangle \langle \Psi, S(f^1 \bar{R}^1) S^{-1}(u^{-1}) \triangleright h_{\underline{2}} \rangle \\ (1.39) &= \sum \langle \varphi, S^{-1}(f^2 \bar{R}^2 u^{-1}) \triangleright h_{\underline{1}} \rangle \langle \Psi, S^{-1}(f^1 \bar{R}^1 u^{-1}) \triangleright h_{\underline{2}} \rangle \\ (5.34) &= ((u^{-1} \succ \varphi) \bar{\circ} (u^{-1} \succ \Psi))(h) = (\Theta_{H_0}(\varphi) \bar{\circ} \Theta_{H_0}(\Psi))(h), \end{aligned}$$

as needed. It is not hard to see that $\Theta_{H_0}(1_{H_0^*}) = 1_{{}^*H_0}$, so Θ_{H_0} is an algebra morphism. It is also a coalgebra morphism since

$$\begin{aligned} \Delta_{{}^*H_0} \Theta_{H_0}(\varphi) &= \Delta_{{}^*H_0}(u^{-1} \succ \varphi) \\ (5.36, 5.4) &= \sum_{i,j=1}^n \langle \varphi, (S^{-1}(u^{-1})_1 S^{-1}(R^2 g^2) \triangleright_i e) \\ &\quad \circ (S^{-1}(u^{-1})_2 S^{-1}(R^1 g^1) \triangleright_j e) \rangle^i e \otimes {}^j e \\ (5.40, 1.39) &= \sum_{i,j=1}^n \langle \varphi, (g^1 S^{-1}(R^2 u^{-1}) \triangleright_i e) \circ (g^2 S^{-1}(R^1 u^{-1}) \triangleright_j e) \rangle^i e \otimes {}^j e \\ (1.39, 1.40) &= \sum_{i,j=1}^n \langle \varphi, (R^2 g^2 \triangleright_i e) \circ (R^1 g^1 \triangleright_j e) \rangle u^{-1} \succ {}^i e \otimes u^{-1} \succ {}^j e \\ &= (\Theta_{H_0} \otimes \Theta_{H_0}) \Delta_{H_0^*}(\varphi) \end{aligned}$$

for any $\varphi \in H_0^*$, and since (1.5) and $\varepsilon(u) = 1$ imply $\varepsilon_{{}^*H_0} \circ \Theta_{H_0} = \varepsilon_{H_0^*}$. \square

Remark 5.3. If (H, R) is not triangular then Θ_{H_0} is, in general, not an algebra or a coalgebra morphism. Indeed, if (H, R) is an arbitrary quasitriangular quasi-Hopf

algebra then computations similar to the ones presented above show that

$$\begin{aligned} \sum \Theta_{H_0}((\bar{r}^1 \bar{R}^2 \mapsto \varphi) \circ (\bar{r}^2 \bar{R}^1 \mapsto \Psi)) &= \Theta_{H_0}(\varphi) \bar{\sigma} \Theta_{H_0}(\Psi), \\ (\Delta_{*H_0} \Theta_{H_0})(\varphi) &= R_{21} R \succ (\Theta_{H_0} \otimes \Theta_{H_0})(\Delta_{H_0^*}(\varphi)) \end{aligned}$$

for any $\varphi, \Psi \in H_0^*$, where $\sum \bar{r}^1 \otimes \bar{r}^2$ is another copy of R^{-1} and we extend the action of H on $*H_0$ to an action of $H \otimes H$ on $*H_0 \otimes *H_0$.

Let H be a quasi-Hopf algebra. Then H^* is an (H, H) -bimodule, by

$$\langle h \rightarrow \varphi, h' \rangle = \varphi(h'h), \quad \langle \varphi \leftarrow h, h' \rangle = \varphi(hh').$$

The convolution $\langle \varphi \Psi, h \rangle = \sum \varphi(h_1) \psi(h_2)$, $h \in H$, is a multiplication on H^* ; it is not associative, but only quasi-associative:

$$(5.41) \quad (\varphi \psi) \xi = \sum (X^1 \rightarrow \varphi \leftarrow x^1) [(X^2 \rightarrow \psi \leftarrow x^2) (X^3 \rightarrow \xi \leftarrow x^3)], \quad \forall \varphi, \psi, \xi \in H^*.$$

In addition, for all $h \in H$ and $\varphi, \psi \in H^*$ we have that

$$(5.42) \quad h \rightarrow (\varphi \psi) = \sum (h_1 \rightarrow \varphi) (h_2 \rightarrow \psi) \quad \text{and} \quad (\varphi \psi) \leftarrow h = \sum (\varphi \leftarrow h_1) (\psi \leftarrow h_2).$$

By [3], if H is a finite dimensional quasitriangular quasi-Hopf algebra then on the dual of H there exists another structure of Hopf algebra in ${}^H_H \mathcal{YD}$, denoted by \underline{H}^* . The structure of \underline{H}^* as a Yetter-Drinfeld module is given by the formulae

$$(5.43) \quad h \cdot \varphi = \sum h_1 \rightarrow \varphi \leftarrow S^{-1}(h_2),$$

$$(5.44) \quad \lambda_{\underline{H}^*}(\varphi) = \sum R^2 \otimes R^1 \cdot \varphi,$$

for all $h \in H$, $\varphi \in H^*$. The structure of \underline{H}^* as a Hopf algebra in ${}^H_H \mathcal{YD}$ is given by:

$$(5.45) \quad \begin{aligned} \varphi \bullet \Psi &= \sum (x^1 X^1 \rightarrow \varphi \leftarrow S^{-1}(f^2 x_2^3 Y^3 R^1 X^2)) \\ &\quad (x^2 Y^1 R_1^2 X_1^3 \rightarrow \Psi \leftarrow S^{-1}(f^1 x_1^3 Y^2 R_2^2 X_2^3)), \end{aligned}$$

$$(5.46) \quad \Delta_{\underline{H}^*}(\varphi) = \sum X_1^1 p^1 \rightarrow \varphi_2 \leftarrow S^{-1}(X_2^1 p^2) \otimes X^2 \rightarrow \varphi_1 \leftarrow S^{-1}(X^3),$$

$$(5.47) \quad \varepsilon_{\underline{H}^*}(\varphi) = \varphi(S^{-1}(\alpha)),$$

$$(5.48) \quad \begin{aligned} S_{\underline{H}^*}(\varphi) &= \sum Q^1 q^1 R^2 x^2 \\ &\quad \cdot [p^1 P^2 S(Q^2) \rightarrow \bar{S}^{-1}(\varphi) \leftarrow S(q^2 R^1 x^1 P^1) x^3 S^{-1}(p^2)], \end{aligned}$$

for all $\varphi, \Psi \in H^*$, where $\sum Q^1 \otimes Q^2$ is another copy of q_R and $\bar{S}^{-1}(\varphi) = \varphi \circ S^{-1}$ for any $\varphi \in H^*$. The unit element is ε .

Proposition 5.4. *Let (H, R) be a finite dimensional triangular quasi-Hopf algebra. Then $*H_0$ and \underline{H}^{*cop} are isomorphic as braided Hopf algebras.*

Proof. Since (H, R) is triangular it follows that $c_{N,M}^{-1} = c_{M,N}$, so \underline{H}^{*cop} is a Hopf algebra in ${}^H_H \mathcal{YD}$. On the other hand, both $*H_0$ and \underline{H}^{*cop} are Hopf algebras in ${}^H_H \mathcal{YD}$ as images, through the functor \mathbb{F} defined by (4.26), of corresponding objects in ${}_H \mathcal{M}$. Therefore, it suffices to prove that $*H_0$ and \underline{H}^{*cop} are isomorphic as Hopf algebras in ${}_H \mathcal{M}$. To this end we claim that $\mu : *H_0 \rightarrow \underline{H}^{*cop}$ given by

$$\mu(\varphi) = \sum g^1 \rightarrow \varphi \leftarrow S^{-1}(g^2)$$

is a Hopf algebra isomorphism in ${}_H\mathcal{M}$. In fact, μ is H -linear since

$$\begin{aligned} \mu(h \succ \varphi)(h') &= \sum \langle \varphi, S^{-1}(h) \triangleright (S^{-1}(g^2)h'g^1) \rangle \\ (1.11) &= \sum \langle \varphi, S^{-1}(h_2g^2)h'h_1g^1 \rangle \\ (5.43) &= \langle \mu(\varphi), S^{-1}(h_2)h'h_1 \rangle = (h \cdot \mu(\varphi))(h') \end{aligned}$$

for all $\varphi \in H^*$ and $h, h' \in H$. It is also an algebra morphism in ${}_H\mathcal{M}$ since

$$\begin{aligned} &\mu(\varphi \bar{\circ} \Psi)(h) \\ (5.34, 5.21, 1.11) &= \sum \langle \varphi, S^{-1}(f^2 \bar{R}^2) \triangleright x^1 X^1 S^{-1}(F^2 g_2^2 \mathbf{G}^2) h_1 g_1^1 G^1 \\ &\quad S(x^2 R^2 y^3 X_2^3) \rangle \langle \Psi, S^{-1}(f^1 \bar{R}^1) x^3 R^1 \triangleright y^1 X^2 \\ &\quad S^{-1}(F^1 g_1^1 \mathbf{G}^1) h_2 g_2^1 G^2 S(y^2 X_1^3) \rangle \\ (1.11, 1.9, 1.16) &= \sum \langle \varphi, S^{-1}(f^2 \bar{R}^2) \triangleright x^1 S^{-1}(F^2 X^3 g^2) h_1 X_1^1 g_{(1,1)}^1 \mathbf{G}_1^1 G^1 \\ &\quad S(x^2 R^2 y^3) \rangle \langle \Psi, S^{-1}(f^1 \bar{R}^1) x^3 R^1 \triangleright y^1 \\ &\quad S^{-1}(F^1 X^2 g_2^1 \mathbf{G}^2) h_2 X_2^1 g_{(1,2)}^1 \mathbf{G}_2^1 G^2 S(y^2) \rangle \\ (1.9, 1.16, 1.1) &= \sum \langle \varphi, S^{-1}(f^2 \bar{R}^2) \triangleright x^1 S^{-1}(F^2 X^3 g^2) h_1 X_1^1 y^1 g_1^1 \\ &\quad \mathbf{G}^1 S(x^2 R^2) \rangle \langle \Psi, S^{-1}(f^1 \bar{R}^1) \triangleright S^{-1}(F^1 X^2 y^3 g_{(2,2)}^1) \\ &\quad \mathbf{G}_2^2 G^2 S(x_1^3 R_1^1) \rangle h_2 X_2^1 y^2 g_{(2,1)}^1 \mathbf{G}_1^1 G^1 S(x_2^3 R_2^1) \rangle \\ (1.11, 1.40, 1.34) &= \sum \langle \varphi, S^{-1}(f^2 \bar{R}^2) \triangleright x^1 S^{-1}(F^2 X^3 g^2) h_1 X_1^1 y^1 R^2 g_2^1 \mathbf{G}^2 S(x^2) \rangle \\ &\quad \langle \Psi, S^{-1}(f^1 \bar{R}^1) \triangleright S^{-1}(F^1 X^2 y^3 R_2^1 (g_1^1 \mathbf{G}^1 S(x^3))_2 G^2) \\ &\quad h_2 X_2^1 y^2 R_1^1 (g_1^1 \mathbf{G}^1 S(x^3))_1 G^1 \rangle \\ (1.9, 1.16) &= \sum \langle \varphi, S^{-1}(f^2 \bar{R}^2) \triangleright S^{-1}(F^2 X^3 x^3 g_2^2 \mathbf{G}^2) h_1 X_1^1 y^1 R^2 \\ &\quad x^2 g_1^2 \mathbf{G}^1 \rangle \langle \Psi, S^{-1}(f^1 \bar{R}^1) \triangleright S^{-1}(F^1 X^2 y^3 R_2^1 x_2^1 g_2^1 G^2) h_2 X_2^1 y^2 R_1^1 x_1^1 g_1^1 G^1 \rangle \\ (1.11 \text{ twice}) &= \sum \langle \varphi, S^{-1}(F^2 X^3 x^3 \bar{R}_2^2 g^2) h_1 X_1^1 y^1 R^2 x^2 \bar{R}_1^2 g^1 \rangle \\ &\quad \langle \Psi, S^{-1}(R^1 x^1 \bar{R}^1) \triangleright S^{-1}(F^1 X^2 y^3 G^2) h_2 X_2^1 y^2 G^1 \rangle \\ (1.33, 1.3) &= \sum \langle \varphi, S^{-1}(F^2 x_2^3 X^3 \bar{R}^2 Y^2 g^2) h_1 x^1 Y^1 g^1 \rangle \\ &\quad \langle \Psi, S^{-1}(\bar{R}^1 Y^3) \triangleright S^{-1}(F^1 x_1^3 X^2 G^2) h_2 x^2 X^1 G^1 \rangle \\ (1.11, R^{-1} = R_{21}) &= \sum \langle \mu(\varphi), S^{-1}(F^2 x_2^3 X^3 R^1 Y^2) h_1 x^1 Y^1 \rangle \\ &\quad \langle \mu(\Psi), S^{-1}(F^1 x_1^3 X^2 R_2^2 Y_2^3) h_2 x^2 X^1 R_1^2 Y_1^3 \rangle \\ (5.45) &= (\mu(\varphi) \bullet \mu(\Psi))(h) \end{aligned}$$

for all $\varphi, \Psi \in H^*$, $h \in H$, and $\mu(1_{*H_0}) = \mu(\varepsilon) = \varepsilon = 1_{\underline{H}^{*cop}}$. It remains to show that μ is a coalgebra morphism and a bijection. To this end, we calculate for any $\varphi \in H^*$:

$$\begin{aligned} (\mu \otimes \mu) \Delta_{*H_0} &= \sum_{i,j=1}^n \langle \varphi, (S^{-1}(R^2 g^2) \triangleright i e) \circ (S^{-1}(R^1 g^1) \triangleright j e) \rangle \\ &\quad G^1 \rightharpoonup i e \leftarrow S^{-1}(G^2) \otimes G^1 \rightharpoonup j e \leftarrow S^{-1}(G^2) \end{aligned}$$

$$\begin{aligned}
(5.18, 1.40) &= \sum_{i,j=1}^n \langle \varphi, X^1 [R^2 S^{-1}(g^1) \triangleright S^{-1}(G^2)]_i e G^1] S(x^1 X^2) \alpha x^2 X_1^3 \\
&\quad [R^1 S^{-1}(g^2) \triangleright S^{-1}(G^2)]_j e G^1] S(x^3 X_2^3) \rangle^i e \otimes^j e \\
(1.11) &= \sum_{i,j=1}^n \langle \varphi, X^1 R_1^2 S^{-1}(g_2^1 G^2) \rangle_i e g_1^1 G^1 S(x^1 X^2 R_2^2) \alpha x^2 \\
&\quad [X^3 R^1 S^{-1}(g^2) \triangleright S^{-1}(G^2)]_j e G^1] S(x^3) \rangle^i e \otimes^j e \\
(1.33, 1.40, 1.34) &= \sum_{i,j=1}^n \langle \varphi, X^1 S^{-1}(y^2 r^2 g_2^2 G^2) \rangle_i e y^1 g^1 S(x^1 R^2 X^3) \alpha x^2 \\
&\quad [R^1 X^2 S^{-1}(y^3 r^1 g_1^2 G^1) \triangleright S^{-1}(G^2)]_j e G^1] S(x^3) \rangle^i e \otimes^j e \\
(1.9, 1.16) &= \sum_{i,j=1}^n \langle \varphi, S^{-1}(y^2 r^2 X^3 g^2) \rangle_i e y^1 X^1 g_1^1 G^1 S(x^1 R^2) \alpha x^2 \\
&\quad [R^1 S^{-1}(y^3 r^1 X^2 g_2^1 G^2) \triangleright S^{-1}(G^2)]_j e G^1] S(x^3) \rangle^i e \otimes^j e \\
(1.40, 1.34, 1.11) &= \sum_{i,j=1}^n \langle \varphi, S^{-1}(R_2^2 X^3 g^2) \rangle_i e R_1^2 X^2 g_2^1 G^2 S(x^1) \alpha x^2 \\
&\quad S^{-1}(R_2^1 X_2^1 g_{(1,2)}^1 G_2^1 G^2) \rangle_j e R_1^1 X_1^1 g_{(1,1)}^1 G_1^1 G^1 S(x^3) \rangle^i e \otimes^j e \\
(1.9, 1.16, 1.1) &= \sum_{i,j=1}^n \langle \varphi, S^{-1}(R_2^2 X^3 g^2) \rangle_i e R_1^2 X^2 x^3 g_{(2,2)}^1 G_2^2 G^2 \alpha \\
&\quad S^{-1}(R_2^1 X_2^1 x^2 g_{(2,1)}^1 G_1^2 G^1) \rangle_j e R_1^1 X_1^1 x^1 g_1^1 G^1 S(x^3) \rangle^i e \otimes^j e \\
(2.26, 1.5) &= \sum_{i,j=1}^n \langle \varphi, S^{-1}(R_2^2 X^3 g^2) \rangle_i e R_1^2 X^2 x^3 S^{-1}(R_2^1 X_2^1 x^2 \beta) \\
&\quad \rangle_j e R_1^1 X_1^1 x^1 g^1 \rangle^i e \otimes^j e \\
(1.17, 5.43) &= \sum_{i,j=1}^n \langle \mu(\varphi), S^{-1}(X^3) \rangle_i e X^2 S^{-1}(X_2^1 p^2) \rangle_j e X_1^1 p^1 \\
&\quad R^2 \cdot \cdot^i e \otimes R^1 \cdot \cdot^j e
\end{aligned}$$

$$(5.46, R^{-1} = R_{21}) = c_{\underline{H}^*, \underline{H}^*}^{-1}(\Delta_{\underline{H}^*}(\mu(\varphi))) = \Delta_{\underline{H}^{*cop}}(\mu(\varphi))$$

as needed. Obviously $\varepsilon\mu = \varepsilon$. It is easy to see that μ is bijective with inverse

$$\mu^{-1}(\varphi) = \sum f^1 \rightharpoonup \varphi \leftarrow S^{-1}(f^2)$$

for any $\varphi \in H^*$. Thus, the proof is complete. \square

Corollary 5.5. *If (H, R) is a finite dimensional triangular quasi-Hopf algebra then $H_0^* \cong {}^*H_0 \cong \underline{H}^{*cop}$ as braided Hopf algebras.*

REFERENCES

- [1] D. Altschuler and A. Coste, Quasi-quantum groups, knots, three-manifolds, and topological field theory, *Comm. Math. Phys.* **150** (1992), 83–107.
- [2] N. Andruskiewitsh and M. Graña, Braided Hopf algebras over abelian finite groups, *Bol. Acad. Ciencias (Córdoba)* **63** (1999), 45–78.
- [3] D. Bulacu and S. Caenepeel, The quantum double for quasitriangular quasi-Hopf algebras, *Comm. Algebra* **31** (2003), 1403–1425.

- [4] D. Bulacu, S. Caenepeel and F. Panaite, More properties of Yetter-Drinfeld modules over quasi-Hopf algebras, in “Hopf algebras in non-commutative geometry and physics”, Caenepeel S. and Van Oystaeyen, F. (eds.), Lecture Notes Pure Appl. Math., Dekker, New York, to appear.
- [5] D. Bulacu and E. Nauwelaerts, Quasitriangular and ribbon quasi-Hopf algebras, *Comm. Algebra* **31** (2003), 1–16.
- [6] D. Bulacu and E. Nauwelaerts, Radford’s biproduct for quasi-Hopf algebras and bosonization, *J. Pure Appl. Algebra* **174** (2002), 1–42.
- [7] D. Bulacu, F. Panaite and F. Van Oystaeyen, Quasi-Hopf algebra actions and smash products, *Comm. Algebra* **28** (2000), 631–651.
- [8] D. Bulacu, F. Panaite and F. Van Oystaeyen, Quantum traces and quantum dimensions for quasi-Hopf algebras, *Comm. Algebra* **27** (1999), 6103–6122.
- [9] S. Caenepeel, F. Van Oystaeyen and Y. H. Zhang, Quantum Yang-Baxter module algebras, *K-Theory* **8** (1994), 231–255.
- [10] R. Dijkgraaf, V. Pasquier and P. Roche, Quasi-Hopf algebras, group cohomology and orbifold models, *Nuclear Phys. B Proc. Suppl.* **18 B** (1990), 60–72.
- [11] V. G. Drinfeld, Quasi-Hopf algebras, *Leningrad Math. J.* **1** (1990), 1419–1457.
- [12] F. Hausser and F. Nill, Diagonal crossed products by duals of quasi-quantum groups, *Rev. Math. Phys.* **11** (1999), 553–629.
- [13] F. Hausser and F. Nill, Doubles of quasi-quantum groups, *Comm. Math. Phys.* **199** (1999), 547–589.
- [14] C. Kassel, “Quantum Groups”, *Graduate Texts in Mathematics* **155**, Springer Verlag, Berlin, 1995.
- [15] S. Majid, Quantum double for quasi-Hopf algebras, *Lett. Math. Phys.* **45** (1998), 1–9.
- [16] S. Majid, “Foundations of quantum group theory”, Cambridge Univ. Press, 1995.
- [17] D. E. Radford and J. Towber, Yetter-Drinfeld categories associated to an arbitrary bialgebra, *J. Pure Appl. Algebra* **87** (1993), 259–279.
- [18] M. E. Sweedler, “Hopf algebras”, Benjamin, New York, 1969.
- [19] M. Takeuchi, Finite Hopf algebras in braided tensor categories, *J. Pure Appl. Algebra* **138** (1999), 59–82.

FACULTY OF MATHEMATICS, UNIVERSITY OF BUCHAREST, RO-70109 BUCHAREST 1, ROMANIA
E-mail address: `dbulacu@al.math.unibuc.ro`

FACULTY OF APPLIED SCIENCES, VRIJE UNIVERSITEIT BRUSSEL, VUB, B-1050 BRUSSELS, BELGIUM
E-mail address: `scaenepe@vub.ac.be`
URL: `http://homepages.vub.ac.be/~scaenepe/`

INSTITUTE OF MATHEMATICS OF THE ROMANIAN ACADEMY, PO-BOX 1-764, RO-70700 BUCHAREST, ROMANIA
E-mail address: `florin.panaite@imar.ro`