### A NOTE ON PARA-QUATERNION MANIFOLDS

Adrian Mihai Ionescu, Gabriel Eduard Vîlcu Politehnica University Bucharest, Romania UPG University, Ploiești, Romania

### <u>Abstract:</u>

We show that a para-quaternion nearly Kahler manifold is necessarily a para-quaternion Kahler manifold

## Introduction

A para-quaternion Hermitian structure on a pseudo-Riemannian manifold (M,g) is a 3-dimensional subbundle V of End(TM), which locally admits bases of skew-symmetric I,J,K w.r.t. g that further satisfies  $I^2 = -J^2 = -K^2 = -Id$ , IJ = -JI = K, JK = -KJ = -I, KI = -IK = J.

Let (M,g,V) be an almost para-quaternion-Hermitian manifold (as above), then it is para-quaternion Kahler if V is invariant under covariant differentiation w.r.t. Levi-Civita conexion of (M,g).

Several authors studied para-quaternion geometry and its similitude to the quaternion geometry, see e.g. [1], [2].

It follows in particular that  $\dim M \stackrel{:}{:} 4$ , the metric is neutral, i.e. it has signature (2m, 2m), where  $\dim M = 4m$  and there is a globally well-defined 4-form  $\Omega$  on M, given by  $\Omega = \Omega_I \wedge \Omega_I - \Omega_J \wedge \Omega_J - \Omega_K \wedge \Omega_K$ , where  $\Omega_I(X,Y) = g(X,IY)$ ,  $\Omega_I(X,Y) = g(X,JY)$ ,  $\Omega_K(X,Y) = g(X,KY)$ 

We begin with the following

## Lemma 1:

For a (pseudo-)Riemannian manifold (M,g) and a (local) skewsymmetric endomorphism I of TM denote  $\Omega_I(X,Y) = g(X,IY)$  and  $M_I(X,Y) := (\nabla_X I)Y = \nabla I(Y;X)$ . Then:

a) 
$$(\Omega_I \wedge \Omega_I)(X, Y, Z, W) = 2\sum_{Y \in W} \Omega_I(X, Y)\Omega_I(Z, W)$$

 $\nabla_{U}(\Omega_{I} \wedge \Omega_{I})(X, Y, Z, W) = 2\sum_{Y} \sup_{Z} \{g(X, M_{I}(U, Y))\Omega_{I}(Z, W) + \Omega_{I}(X, Y))g(Z, M_{I}(U, W)\} \}, \text{ where the sums}$ 

are taken over cyclic permutations of  $\{Y, Z, W\}$ .

**b)**  $g(M_I(X,Y),Z) + g(M_I(X,Z),Y) = 0$ 

If  $I^2 = \pm Id$ , then

- c)  $M_{I}(X,IY) = -IM_{I}(X,Y)$ ,  $g(M_{I}(X,Y),IY) = g(M_{I}(X,Y),Y) = 0$
- d)  $M_I(X,Z) = 0$  for any pair of Hermitian-totally real vectors (i.e.  $Z \perp \{X, IX\}$ ) if and only if  $M_I = 0$ , i.e. M is Kahler.

If furthermore I, J, K satisfy the para-quaternionic identities, then

e)  $M_I(X,JY) = M_K(X,Y) - IM_J(X,Y)$ ,  $M_J(X,IY) = -M_K(X,Y) - JM_I(X,Y)$  (and 4 other similar relations hold)

We can state now the result, see [3]

## Theorem 1:

Let (M,g, V) be an almost para-quaternion-Hermitian manifold. Suppose that M is nearly-para-quaternion Kahler, that is  $(\nabla_X \Omega)(X,Y,Z,W) = 0$ , for any  $X,Y,Z,W \in TM$ . Then M is para-quaternion Kahler.

# Proof:

One must show that for any vectors X,Y it holds that  $M_I(X,Y),M_J(X,Y),M_K(X,Y) \in Span\{IY,JY,KY\}$ . This will be done by systematic use of formulae in Lemma1.

# Claim1:

For para-quaternion-Hermitian totally real vectors X,Y (i.e.  $X \perp \{Y, IY, JY, KY\}$ ) one has  $M_I(X,Y), M_I(X,Y), M_K(X,Y) \in Q(X) + Q(Y)$ .

This is true if dim M = 8; then for any triple  $\{X, Y, Z\}$  of pairwise para-quaternion-Hermitian totally real vectors, Lemma1 gives  $(\nabla_X \Omega)(X, Y, Z, IX) = -2 \|X\|^2 g(Y, M_I(X, Z))$  and similarly for J, K. Since M is nearly-para-quaternion Kahler, the Claim follows.

# Claim2:

For para-quaternion-Hermitian totally real vectors X,Y one has  $M_I(X,Y), M_J(X,Y), M_K(X,Y) \perp Q(X)$ .

Applying Lemmal again gives  $(\nabla_X \Omega)(X, Y, JX, IX) = -2 ||X||^2 g(Y, M_I(X, JX)) - 2 ||X||^2 g(IX, M_J(X, Y))$   $+ 2 ||X||^2 g(X, M_K(X, Y))$ , or after rearranging  $(M_K + IM_J - JM_I)(X, Y) \perp X$ . Replacing Y by KY gives again by Lemmal  $(-IM_I + JM_J - 3KM_K)(X, Y) \perp X$ .

Similarly one obtains

 $(M_I + KM_J - JM_K)(X, Y) \perp X$ ,  $(3IM_I + JM_J + KM_K)(X, Y) \perp X$ 

 $(M_J + KM_I - IM_K)(X, Y) \perp X$ ,  $(-IM_I - 3JM_J + KM_K)(X, Y) \perp X$ .

It follows that  $IM_I(X,Y), JM_J(X,Y), KM_K(X,Y) \perp X$ , equivalent to the Claim, again by use of Lemma1 and substitutions Y to IY, JY, KY.

Now one has  $M_I(X,Y), M_J(X,Y), M_K(X,Y) \in Span\{IY, JY, KY\}$ , for paraquaternion-Hermitian totally real vectors X, Y, since  $g(M_I(X,Y),Y) = g(M_I(X,Y),Y) = g(M_K(X,Y),Y) = 0$ .

The remaining cases to consider now follows easily since  $M_i$ -s are tensors while e.g.

 $g(M_I(X, IX), IX) = 0$ 

$$\begin{split} g(M_{I}(X,JX),Y) &= -g(M_{I}(X,Y),JX) = 0\\ g(M_{I}(X,JX),JX) &= 0\\ g(M_{I}(X,JX),KX) &= g(M_{K}(X,X) - IM_{J}(X,X),KX) = g(M_{K}(X,X),KX) - g(M_{J}(X,X),JX) = 0 \end{split}$$

This ends the proof of the Theorem.

## References:

[1] E. Garcia-Rio, Y. Mathushita, R.Vazquez-Lorenzo Paraquaternionic Kahler manifolds, Rocky Jour. Spring 2001
[2] S. Vukmirovic Para-quaternionic reduction, preprint
[3] A. Swann Some remarks on quaternion-hermitian manifolds, Archivum Mathematicum (Brno), Tomus 33(1997), 319-351