Bernoulli-Taylor formula of ψ -umbral difference calculus

A.K.Kwaśniewski

Institute of Computer Science, Białystok University PL-15-887 Białystok, ul.Sosnowa 64, POLAND e-mail: kwandr@uwb.edu.pl

November 7, 2018

Abstract

We shall present here the $*_{\psi}$ -Bernoulli-Taylor^{*} formula of a new sort with the rest term of the Cauchy type recently derived by the author in the case of ψ -difference calculus. The central importance of such a type formulas is beyond any doubt.

 * see: historical remark at the beginning of this note KEY WORDS: $\psi\text{-}calculus,$ Bernoulli-Taylor formula, Graves-Heisenberg-Weyl algebra AMS S.C. (1991) 17B01, 17B35, 33C45, 34A , 81S0505

1 One Historical Remark

Here are the famous examples of expansion

$$\partial_0 = \sum_{n=1}^{\infty} \frac{x^{n-1}}{n!} \frac{d^n}{dx^n}$$

or

$$\epsilon_0 = \sum_{n=0}^{\infty} (-1)^n \frac{x^n}{n!} \frac{d^n}{dx^n}$$

where ∂_0 is the divided difference operator while ϵ_0 is at the zero point evaluation functional. If one compares these with "series universalissima"

of J.Bernoulli from $Acta \ Erudicorum$ (1694) (see commentaries in [12]) and with

$$exp\{yD\} = \sum_{k=0}^{\infty} \frac{y^k D^k}{k!}, \quad D = \frac{d}{dx},$$

then confrontation with B.Taylor's "Methodus incrementorum directa et inversa" (1715), London; entitles one to call the expansion formulas considered in this note "Bernoulli - Taylor formulas" or (for $n \to \infty$) "Bernoulli - Taylor series" [1].

2 Introduction

While deriving the Bernoulli-Taylor ψ -formula one is tempted to adapt the ingenious Viskov's method [2] of arriving to formulas of such type for various pairs of operations. In our case these would be ψ -differentiation and ψ -integration (see: Appendix). However straightforward application of Viscov methods in ψ -extensions of umbral calculus leads to sequences which are not normal (Ward) hence a new invention is needed. This expected and verified here invention is the new specific $*_{\psi}$ product of analytic functions or formal series. This note is based on [3] where the derivation of this new form of Bernoulli-Taylor $*_{\psi}$ - formula was delivered due to the use of a specific $*_{\psi}$ product of formal series.

3 Classical Bernoulli-Taylor formulas with the rest term of the Cauchy type by Viskov method

Let us consider the obvious identity

$$\sum_{k=0}^{n} (\alpha_k - \alpha_{k+1}) = \alpha_0 - \alpha_{n+1}$$
(1)

in which (1) we now put $\alpha_k = a^k b^k; a, b \in \mathcal{A}$. \mathcal{A} is an associative algebra with unity over the field F=R,C. Then we get

$$\sum_{k=0}^{n} a^{k} (1-ab) b^{k} = 1 - a^{n+1} b^{n+1}; a, b \in \mathcal{A}$$
(2)

Numerous choices of $a, b \in \mathcal{A}$ result in many important specifications of (2)

Example 1. Let \mathcal{F} denotes the linear space of sufficiently smooth functions $f: F \longrightarrow F$. Let

$$a: \mathcal{F} \longrightarrow \mathcal{F}; \qquad (af)(x) = \int_{a}^{b} f(t)dt,$$

$$b: \mathcal{F} \longrightarrow \mathcal{F}; \qquad (bf)(x) = (\frac{d}{dx}f)(x); \qquad (3)$$

$$l: \mathcal{F} \longrightarrow \mathcal{F}; \qquad (lf)(x) = f(x).$$

Then $[b,a]=1-ab=\varepsilon_{\alpha}$ where ε_{α} is evaluation functional on \mathcal{F} i.e.

$$\varepsilon_{\alpha}(f) = f(\alpha) \tag{4}$$

Using now the text-book integral Cauchy formula (k > 0)

$$(a^k f)(x) = \int_a^x \frac{(x-t)^{k-1}}{(k-1)!} f(t) dt,$$
(5)

and under the choice (3) one gets from (2) the well-known Bernoulli-Taylor formula

$$f(x) = \sum_{k=0}^{n} \frac{(x-\alpha)^k}{k!} f^{(k)}(\alpha) + R_{n+1}(x)$$
(6)

with the rest term $R_{n+1}(x)$ in the Cauchy form

$$R_{n+1}(x) = \int_{a}^{x} \frac{(x-t)^{n}}{n!} f^{(n+1)}(t) dt$$
(7)

Example 2. [1] Let \mathcal{F} denotes the linear space of functions $f: \mathbb{Z}_+ \longrightarrow F; \mathbb{Z}_+ = \mathbb{N} \cup \{0\}$. Let

$$a: Z_{+} \longrightarrow \mathcal{F}; \qquad (af)(x) = \sum_{k=0}^{x-1} f(k),$$

$$b: Z_{+} \longrightarrow \mathcal{F}; \qquad (bf)(x) = f(x+1) - f(x), \qquad (8)$$

$$l: Z_{+} \longrightarrow \mathcal{F}; \qquad (lf)(x) = f(x).$$

It is easy to see that $[b,a]=1-ab=\varepsilon_0$ where ε_0 is evaluation functional i.e. $\varepsilon_0(f) = f(0)$. $b = \Delta$ is the standard difference operator with its left inverse

definite summation operator a. The corresponding Δ - calculus Cauchy formula is also known (see formula (31 p.310 in [5]);

$$(a^k f)(x) = \sum_{r=0}^{x-1} \frac{(x-r-1)^{k-1}}{(k-1)!} f(r); k > 0$$
(9)

where $x^{\underline{n}} = x(x-1)(x-2)...(x-n+1)$.

Under the choice (8) one gets from (2) the Δ - calculus Bernoulli - Taylor fomula [1]

$$f(x) = \sum_{k=0}^{n} \frac{x^{\underline{k}}}{k!} (\Delta^{k} f)(0) + R_{n+1}(x)$$
(10)

with the rest term $R_{n+1}(x)$ in the Cauchy Δ form

$$R_{n+1}(x) = \sum_{r=0}^{x-1} \frac{(x-r-1)^{\underline{n}}}{n!} (\Delta^{n+1} f)(r);$$
(11)

4 " $*_{\psi}$ realization" of Bernoulli identity.

Now a specifically new form of the Bernoulli-Taylor formula with the rest term of the Cauchy type as well as Bernoulli-Taylor series is to be supplied in the case of ψ -difference *umbral* calculus (see [5-8] and [9,10] and references therein). For that to do we use natural ψ -umbral representation [13,14] of Graves-Heisenberg-Weyl (GHW) algebra [11,12] generators \hat{p} and \hat{q} and then we use Bernoulli identity (12)

$$\hat{p}\sum_{k=0}^{n} \frac{(-\hat{q})^k \hat{p}^k}{k!} = \frac{(-\hat{q})^n \hat{p}^{n+1}}{n!}$$
(12)

derived by Viskov from (1) under the substitution (see (28) in [2])

$$\alpha_0 = 0, \ \alpha_k = (-1)^k (\hat{q})^{k-1} \hat{p}^k (k-1)!, \ k = 1, 2, \dots$$

due to $\hat{p}\hat{q}^n=\hat{q}^n\hat{p}+n\hat{q}^{n-1}$ (n=1,2,...) resulting by induction from

$$[\hat{p}, \hat{q}] = 1 \tag{13}$$

Example 1. The choice $\hat{p} = D \equiv \frac{d}{dx}$ and $\hat{q} = \hat{x} - y, y \in F$; $\hat{x}f(x) = xf(x)$ after substitution into Bernoulli identity (12) and integration $\int_{\alpha}^{x} dt$ gives the Bernoulli - Taylor formula (6).

Example 2. The choice [2] $\hat{p} = \Delta$ and $\hat{q} = \hat{x} \circ E^{-1}$ where $E^{\alpha}f(x) = f(x+\alpha)$ after substitution into Bernoulli identity (12) and " Δ - integration" $\sum_{r=0}^{\alpha-1}$ gives the Bernoulli - Mac laurin formula of the following form $(\alpha, x \in \mathbf{Z}, \nabla = 1 - E^{-1})$ with the rest term $R_{n+1}(x)$

$$f(0) = \sum_{k=0}^{n} \frac{\alpha^{k}}{k!} (-1)^{k+1} (\nabla^{k} f)(\alpha) + R_{n+1}(\alpha); \qquad (14)$$

$$R_{n+1}(\alpha) = (-1)^n \sum_{r=0}^{\alpha-1} \frac{r^n}{n!} (\nabla^{n+1} f)(r+1).$$
(15)

Example 3. Here $f^{(k)} \equiv \partial_{\psi}^k f$ and $f(x) *_{\psi} g(x) \equiv f(\hat{x}_{\psi})g(x)$ - see Appendix. The choice $\hat{p} = \partial_{\psi}$ and $\hat{q} = \hat{z}_{\psi}$ (z = x - y) where $\hat{x}_{\psi}x^n = \frac{n+1}{(n+1)_{\psi}}x^{n+1}$ after substitution into Bernoulli identity (12) and " ∂_{ψ} - integration" $\int_{\alpha}^{x} d_{\psi}t$ (see: Appendix) gives another Bernoulli - Taylor ψ -formula of the form:

$$f(x) = \sum_{k=0}^{n} \frac{1}{k!} (x - \alpha)^{k_{*\psi}} *_{\psi} f^{(k)}(\alpha) + R_{n+1}(x)$$
(16)

with the rest term $R_{n+1}(x)$ in the Cauchy-form

$$R_{n+1}(x) = \frac{1}{n!} \int_{\alpha}^{x} d_q t (x-t)^{n_{*q}} *_q f^{(n+1)}(t) dt$$
(17)

In the above notation $x^{0*_{\psi}} = 1$, $x^{n*_{\psi}} \equiv x *_{\psi} (x^{(n-1)*_{\psi}}) = x *_{\psi} \dots *_{\psi} x = \frac{n!}{n_{\psi}!}x^n$; $n \ge 0$.

Naturally $\partial_{\psi} x^{n*_{\psi}} = n x^{(n-1)*_{\psi}}$ and in general f, g - may be formal series for which

$$\partial_{\psi}(f *_{\psi} g) = (Df) *_{\psi} g + f *_{\psi} (\partial_{\psi} g)$$
(18)

i.e. Leibniz $*_{\psi}$ rule holds [13, 14, 15].

Summary: These another forms of both the Bernoulli -Taylor formula with the rest term of the Cauchy type [3] as well as Bernoulli - Taylor series are quite easily handy due to the technique developed in [13, 14] where one may find more on $*_{\psi}$ product devised perfectly suitable for the Ward's "calculus of sequences" [6] or more exactly $*_{\psi}$ is devised perfectly suitable for the socalled ψ - extension on Finite Operator Calculus of Rota (see [9, 10, 14, 15] and references therein)

5 Appendix

$*_{\psi}$ product

Let $n - \psi \equiv \psi_n$; $\psi_n \neq 0$: n > 0. Let ∂_{ψ} be a linear operator acting on formal series and defined accordingly by $\partial_{\psi} x^n = n_{\psi} x^{n-1}$.

We introduce now a intuition appealing ∂_{ψ} -difference-ization rules for a specific new $*_{\psi}$ product of functions or formal series. This $*_{\psi}$ product is what we call: the ψ -multiplication of functions or formal series as specified below. **Notation A.1.**

 $x *_{\psi} x^{n} = \hat{x}_{\psi}(x^{n}) = \frac{(n+1)}{(n+1)_{\psi}} x^{n+1}; \quad n \ge 0 \quad \text{hence } x *_{\psi} 1 = (1_{\psi})^{-1} x \not\equiv x$ therefore $x *_{\psi} \alpha 1 = \alpha 1 *_{\psi} x = x *_{\psi} \alpha = \alpha *_{\psi} x = \alpha (1_{\psi})^{-1} x$ and $\forall x, \alpha \in F;$ $f(x) *_{\psi} x^{n} = f(\hat{x}_{\psi}) x^{n}.$

For $k \neq n$ $\mathbf{x}^{n} *_{\psi} \mathbf{x}^{k} \neq \mathbf{x}^{k} *_{\psi} \mathbf{x}^{n}$ as well as $\mathbf{x}^{n} *_{\psi} \mathbf{x}^{k} \neq \mathbf{x}^{n+k}$ - in general.

In order to facilitate the formulation of observations accounted for on the basis of ψ -calculus representation of GHW algebra we shall use what follows. **Definition A.1.** With Notation A.1. adopted define the $*_{\psi}$ powers of x according to $x^{n*_{\psi}} \equiv x *_{\psi} x^{(n-1)*_{\psi}} = \hat{x}_{\psi}(x^{(n-1)*_{\psi}}) = x *_{\psi} x *_{\psi} \dots *_{\psi} x$ $= \frac{n!}{n_{\psi}!} x^n; \quad n \ge 0$. Note that $x^{n*_{\psi}} *_{\psi} x^{k*_{\psi}} = \frac{n!}{n_{\psi}!} x^{(n+k)*_{\psi}} \neq x^{k*_{\psi}} *_{\psi} x^{n*_{\psi}} = \frac{k!}{k_{\psi}!} x^{(n+k)*_{\psi}}$ for $k \ne n$ and $x^{0*_{\psi}} = 1$.

This noncommutative ψ -product $*_{\psi}$ is devised so as to ensure the following observations.

Observation A.1

- a) $\partial_{\psi} x^{n*_{\psi}} = n x^{(n-1)*_{\psi}}; \ n \ge 0$
- b) $\exp_{\psi}[\alpha x] \equiv \exp \{\alpha \hat{x}_{\psi}\} \mathbf{1}$
- c) $\exp [\alpha x] *_{\psi} (\exp_{\psi} \{\beta \hat{x}_{\psi}\}\mathbf{1}) = (\exp_{\psi} \{[\alpha + \beta] \hat{x}_{\psi}\})\mathbf{1}$
- $d) \ \partial_{\psi}(x^{k} *_{\psi} \quad x^{n*_{\psi}}) = (Dx^{k}) *_{\psi} x^{n*_{\psi}} + x^{k} *_{\psi} (\partial_{\psi} x^{n*_{\psi}})$
- e) $\partial_{\psi}(f *_{\psi} g) = (Df) *_{\psi} g + f *_{\psi} (\partial_{\psi} g) ; f, g$ formal series
- f) $f(\hat{x}_{\psi})g(\hat{x}_{\psi}) \mathbf{1} = f(x) *_{\psi} \tilde{g}(x) ; \tilde{g}(x) = g(\hat{x}_{\psi})\mathbf{1}.$

 ψ -Integration Let: $\partial_o x^n = x^{n-1}$. The linear operator ∂_o is identical with divided difference operator. Let $\hat{Q}f(x)f(qx)$. Recall also that to the " ∂_q difference-ization" there corresponds the q-integration which is a right inverse operation to "q-difference-ization". Namely

$$F(z) := \left(\int_{q} \varphi\right)(z) := (1-q) z \sum_{k=0}^{\infty} \varphi\left(q^{k} z\right) q^{k}$$
(19)

i.e.

$$F(z) \equiv \left(\int_{q} \varphi\right)(z) = (1-q) z \left(\sum_{k=0}^{\infty} q^{k} \hat{Q}^{k} \varphi\right)(z) = \left((1-q) z \frac{1}{1-q\hat{Q}} \varphi\right)(z). \quad (20)$$

Of course

$$\partial_q \circ \int_q = id$$
 (21)

as

$$\frac{1-q\hat{Q}}{(1-q)}\partial_0\left((1-q)\hat{z}\frac{1}{1-q\hat{Q}}\right) = id.$$
(22)

Naturally (22) might serve to define a right inverse operation to "q-differenceization" $(\partial_q \varphi)(x) = \frac{1-q\hat{Q}}{(1-q)}\partial_0 \varphi(x)$ and consequently the "q-integration" as represented by (19) and (20). As it is well known the definite q-integral is an numerical approximation of the definite integral obtained in the $q \to 1$ limit.

Finally we introduce the analogous representation for ∂_{ψ} difference-ization

$$\partial_{\psi} = \hat{n}_{\psi} \partial_o; \ \hat{n}_{\psi} x^{n-1} = n_{\psi} x^{n-1}; \ n \ge 1$$
(23)

Then

$$\int_{\psi} x^{n} = \left(\hat{x}\frac{1}{\hat{n}_{\psi}}\right) x^{n} = \frac{1}{(n+1)_{\psi}} x^{n+1}; \ n \ge 0$$
(24)

and of course $\left(\int_{\psi} \equiv \int d_{\psi}\right)$

$$\partial_{\psi} \circ \int_{\psi} = id \tag{25}$$

Naturally

$$\partial_{\psi} \circ \int_{a}^{x} f(t) d_{\psi} t = f(x)$$

The formula of "per partes" ψ -integration is easily obtainable from (Observation A.1 e) and it reads:

$$\int_{a}^{b} (f *_{\psi} \partial_{\psi} g)(t) d_{\psi} t = [(f *_{\psi} g)(t)]_{a}^{b} - \int_{a}^{b} (Df *_{\psi} g)(t) d_{\psi} t$$
(26)

Two Closing Remarks:

I. All these above may be quite easily extended [15] to the case of any $Q \in End(P)$ linear operator that reduces by one the degree of each polynomial [16]. Namely one introduces [15]:

Definition A.2.

$$\hat{x}_Q \in End(P), \hat{x}_Q : F[x] \to F[x]$$

such that $(x^n) = \frac{(n+1)}{(n+1)_{\psi}} q_{n+1}; n \ge 0$; where $Qq_n = nq_{n-1}$.

Then \star_Q product of formal series and Q-integration are defined analogously. (This has been accomplished by my student E. Krot).

II. In 1937 Jean Delsarte [17] had derived the general Bernoulli-Taylor formula for a class of linear operators δ including linear operators that reduce by one the degree of each polynomial. The rest term of the Cauchy-like type in his Taylor formula (I) is given in terms of the unique solution of a first order partial differential equation in two real variables. This first order partial differential equation is determined by the choice of the linear operator δ and the function f under expansion. In our Bernoulli-Taylor -formula (16)-(17) or in its straightforward \star_Q product of formal series and Q-integration generalization - there is no need to solve any partial differential equation.

References

- [1] N. Y. Sonin Izw. Akad. Nauk 7, 337 (1897). O.V.
- [2] Viskov: Trudy Matiematicz'eskovo Instituta AN SSSR 177, 21 (1986).
- [3] A. K. Kwaśniewski: q-difference calculus Bernoulli-Taylor formula Białystok Univ. Inst. Comp. Sci. UwB Preprint 32 (August 2001)
- [4] A. O. Gel'fond "Izczislenie koniecznych raznostiej" "Nauka" 1967 Moskva.
- [5] J. Thomae J.Reine Angew. Math. 70, 258-281 (1869).
- [6] M. Ward: Amer. J. Math. 58, 255-266 (1936).
- [7] O.V. Viskov: Soviet Math. Dokl 16, 1521-1524 (1975).
- [8] O.V. Viskov: Soviet Math. Dokl 19, 250-253 (1978) F.

- [9] A. K. Kwaśniewski: Integral Transforms and Special Functions 2(4), 333 (2001)
- [10] A. K. Kwaśniewski: Rep. Math. Phys. 47 (4), 305-342 (2001).
- [11] C. Graves: Proc. Royal Irish Academy 6, 144 (1853-1857).
- [12] O.V. Viskov: Acta. Sci. Math. 59, 585 (1994).
- [13] A. K. Kwaśniewski "Extended finite operator calculus an example of algebraization of analysis, Białystok Univ. Inst. Comp. Sci. UwB Preprint 28, (April 2001).
- [14] A. K. Kwaśniewski: On Simple Characterisations of Sheffer ψpolynomials and Related Propositions of the Calculus of Sequences Bulletin de la Soc. des Sciences et de Lettres de Lodz ; 52, Ser. Rech. Deform. 36 (2002): 45-65
- [15] A. K. Kwaśniewski: Main theorems of extended finite operator calculus Integral Transforms and Special Functions Vol 14, No 6, (2003): 499-516
- [16] G. Markowsky: Math. Anal. Appl 63, 145-155 (1978)
- [17] J. Delsarte C.R. Acad. Sc. Paris 204,(3) II,408-409 1937