A geometrical construction for the polynomial invariants of some reflection groups

A. Sarti

FBR 17 Mathematik, Universität Mainz Staudingerweg 9, 55099 Mainz Germany e-mail:sarti@mathematik.uni-mainz.de

November 13, 2018

Abstract

In these notes we investigate the ring of real polynomials in four variables, which are invariant under the action of the reflection groups [3, 4, 3] and [3, 3, 5]. It is well known that they are rationally generated in degree 2,6,8,12 and 2,12,20,30. We give a different proof of this fact by giving explicit equations for the generating polynomials.

0 Introduction

There are four groups generated by reflections which operate on the fourdimensional Euclidian space. These are the symmetry groups of some regular four dimensional polytopes and are described in [Co2] p. 145 and table I p. 292-295. With the notation there the groups and their orders are

Group	[3,3,3]	[3,3,4]	[3,4,3]	[3,3,5]
Order	120	384	1152	14400

They operate in a natural way on the ring of polynomials $R = \mathbb{R}[x_0, x_1, x_2, x_3]$ and it is well known that the ring of invariants R^G (*G* one of the groups above) is algebraically generated by a set of four independent polynomials (cf. [Bu] p. 357). Coxeter shows in [Co1] that the rings R^G , G = [3,3,3] or [3,3,4] are generated in degree 2, 3, 4, 5 resp. 2, 4, 6, 8 and since the product of the degrees is equal to the order of the group, any other invariant polynomial is a combination with real coefficients of products of these invariants (i.e., in the terminology of [Co1], the ring R^G is rationally generated by the polynomials). Coxeter also gives equations for the generators. In the case of the groups [3, 4, 3] and [3, 3, 5] he recalls a result of Racah, who shows with the help of the theory of Lie groups that the rings R^G are rationally generated in degree 2, 6, 8, 12 resp. 2, 12, 20, 30 (cf. [Ra]). Neither Coxeter nor Racah give equations for the polynomials. In these notes we construct the generators and give a different proof of the result of Racah. The invariant of degree two is well known (cf. [Co1]) and can be given as

$$q = x_0^2 + x_1^2 + x_2^2 + x_3^2.$$

We construct the other invariants in a completely geometrical way. For proving that our polynomials together with the quadric generate the ring R^G , we show some relations between them and the invariant forms of the binary tetrahedral group and of the binary icosahedral group.

It is a pleasure to thank W. Barth of the University of Erlangen for many helpful discussions.

1 Notations and preliminaries

Denote by R the ring of polynomials in four variables with real coefficients $\mathbb{R}[x_0, x_1, x_2, x_3]$, by G a finite group of homogeneous linear substitutions, and by R^G the ring of invariant polynomials.

1. A set of polynomials F_1, \ldots, F_n in R is called *algebraically dependent* if there is a non trivial relation

$$\sum \alpha_I(F_1^{i_1} \cdot \ldots \cdot F_n^{i_n}) = 0,$$

where $I = (i_1, \ldots, i_n) \in \mathbb{N}^n, \alpha_I \in \mathbb{R}$.

2. The polynomials are called *algebraically independent* if they are not dependent. For the ring R^G , there always exists a set of four algebraically independent polynomials (cf. [Bu], thm. I, p. 357).

3. We say that R^G is algebraically generated by a set of polynomials F_1, \ldots, F_4 , if for any other polynomial $P \in R^G$ we have an algebraic relation

$$\sum \alpha_I (P^{i_0} \cdot F_1^{i_1} \cdot \ldots \cdot F_4^{i_4}) = 0.$$

4. We say that the ring R^G is rationally generated by a set of polynomials F_1, \ldots, F_4 , if for any other polynomial $P \in R^G$ we have a relation

$$\sum \alpha_I(F_1^{i_1} \cdot \ldots \cdot F_4^{i_4}) = P, \ \alpha_I \in \mathbb{R}$$

5. The four polynomials of 3 are called a *basic set* if they have the smallest possible degree (cf. [Co1]).

6. There are two classical 2 : 1 coverings

$$\rho: SU(2) \to SO(3)$$
 and $\sigma: SU(2) \times SU(2) \to SO(4)$,

we denote by T, O, I the tetrahedral group, the octahedral group and the icosahedral group in SO(3) and by $\tilde{T}, \tilde{O}, \tilde{I}$ the corresponding binary groups in SU(2) via the map ρ . The σ -images of $\tilde{T} \times \tilde{T}, \tilde{O} \times \tilde{O}$ and $\tilde{I} \times \tilde{I}$ in SO(4)are denoted by G_6, G_8 and G_{12} . By abuse of notation we write (p,q) for the image in SO(4) of an element $(p,q) \in SU(2) \times SU(2)$. As showed in [Sa] (3.1) p. 436, the groups G_6 and G_{12} are subgroups of index four respectively two in the reflections groups [3, 4, 3] and [3, 3, 5].

2 Geometrical construction

Denote by \tilde{G} one of the groups \tilde{T} , \tilde{O} or \tilde{I} . Clearly, the subgroups $\tilde{G} \times 1$ and $1 \times \tilde{G}$ of SO(4) are isomorphic to \tilde{G} . Moreover, each of them operates on one of the two rulings of the quadric $\mathbb{P}_1 \times \mathbb{P}_1$ and leaves invariant the other ruling (as shown in [Sa]). We recall the lengths of the orbits of points under the action of the groups T, O and I

group	Т	0	Ι
lengths of the orbits	12, 6, 4	24, 12, 8, 6	$60, \ 30, \ 20, \ 12$

These lines are fixed by elements $(p, 1) \in \tilde{G} \times 1$ on one ruling, resp. $(1, p') \in 1 \times \tilde{G}$ on the other ruling of the quadric. Recall that these elements have two lines of fix points with eigenvalues α , $\bar{\alpha}$ which are in fact the eigenvalues of p and p'. We call two lines L, L' of $\mathbb{P}_1 \times \mathbb{P}_1$ a *couple* if L is fixed by (p, 1)with eigenvalue α and L' is fixed by (1, p) with the same eigenvalue.

2.1 The invariant polynomials of G_6 and of G_{12}

Consider the six couples of lines $L_1, L'_1, \ldots, L_6, L'_6$ in $\mathbb{P}_1 \times \mathbb{P}_1$ which form one orbit under the action of $\tilde{T} \times 1$, resp. $1 \times \tilde{T}$, and denote by $f_{11}^{(6)}, \ldots, f_{66}^{(6)}$ the six planes generated by such a couple of lines (and by abuse of notation their equation, too). Now set

$$F_6 = \sum_{g \in \tilde{T} \times 1} g(f_{11}^{(6)} \cdot f_{22}^{(6)} \cdot \ldots \cdot f_{66}^{(6)}) = \sum_{g \in \tilde{T} \times 1} g(f_{11}^{(6)}) \cdot g(f_{22}^{(6)}) \cdot \ldots \cdot g(f_{66}^{(6)}).$$

Observe that an element $g \in \tilde{T} \times 1$ leaves each line of one ruling invariant and operates on the six lines of the other ruling. A similar action is given by an element of $1 \times \tilde{T}$. Since we sum over all the elements of $\tilde{T} \times 1$, the action of $1 \times \tilde{T}$ does not give anything new, hence F_6 is G_6 -invariant. Furthermore observe that F_6 has real coefficients. In fact, in the above product, for each plane generated by the lines L_i, L'_i we also take the plane generated by the lines which consist of the conjugate points. The latter has equation $\bar{f}_{ii}^{(6)}$, i.e., we have an index $j \neq i$ with $f_{jj}^{(6)} = \bar{f}_{ii}^{(6)}$ and the products $f_{ii}^{(6)} \cdot \bar{f}_{ii}^{(6)}$ have real coefficients.

Consider now the orbits of lengths eight and twelve under the action of $O \times 1$ and $1 \times \tilde{O}$ and the planes $f_{ii}^{(8)}$, $f_{jj}^{(12)}$ generated by the eight, respectively by the twelve couples of lines. As before the polynomials

$$F_8 = \sum_{g \in \tilde{T} \times 1} g(f_{11}^{(8)} \cdot \ldots \cdot f_{88}^{(8)}),$$

$$F_{12} = \sum_{g \in \tilde{T} \times 1} g(f_{11}^{(12)} \cdot \ldots \cdot f_{1212}^{(12)})$$

are G_6 -invariant and have real coefficients.

Finally we consider the lines of $\mathbb{P}_1 \times \mathbb{P}_1$ which form orbits of length 12, 20 and 30 under the action of $\tilde{I} \times 1$ resp. $1 \times \tilde{I}$. The planes generated by the couples of lines produce the G_{12} -invariant real polynomials

$$\Gamma_{12} = \sum_{g \in \tilde{I} \times 1} g(h_{11}^{(12)} \cdot \ldots \cdot h_{1212}^{(12)}),$$

$$\Gamma_{20} = \sum_{g \in \tilde{I} \times 1} g(h_{11}^{(20)} \cdot \ldots \cdot h_{2020}^{(20)}),$$

$$\Gamma_{30} = \sum_{g \in \tilde{I} \times 1} g(h_{11}^{(30)} \cdot \ldots \cdot h_{3030}^{(30)}).$$

2.2 The invariant polynomials of the reflection groups

We consider the matrices

$$C = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & -1 \end{pmatrix}, \quad C' = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \end{pmatrix},$$

as in [Sa] (3.1) p. 436, the groups generated by G_6 , C, C' and G_{12} , C are the reflections groups [3, 4, 3] respectively [3, 3, 5].

Proposition 2.1 1. The polynomials F_6 , F_8 , F_{12} , Γ_{12} , Γ_{20} , Γ_{30} are C invariant.

2. The polynomials F_6 , F_8 , F_{12} are C' invariant.

Proof. 1. The matrix C interchanges the two rulings of the quadric, hence the polynomials F_i and Γ_j are invariant by construction. We prove 2 by a direct computation in the last section.

From this fact we obtain

Corollary 2.1 The polynomials q, F_6, F_8, F_{12} are [3, 4, 3]-invariant and the polynomials $q, \Gamma_{12}, \Gamma_{20}, \Gamma_{30}$ are [3, 3, 5]-invariant.

Here we denote by q the quadric $\mathbb{P}_1 \times \mathbb{P}_1$.

3 The rings of invariant forms

Identify \mathbb{P}_3 with $\mathbb{P}M(2 \times 2, \mathbb{C})$ by the map

$$(x_0:x_1:x_2:x_3) \quad \mapsto \quad \left(\begin{array}{cc} x_0+ix_1 & x_2+ix_3 \\ -x_2+ix_3 & x_0-ix_1 \end{array}\right).$$
 (1)

Furthermore consider the map

$$\begin{array}{cccc} \mathbb{C}^2 \times \mathbb{C}^2 & \longrightarrow & M(2 \times 2, \mathbb{C}) \\ ((z_0, z_1), (z_2, z_3)) & \longmapsto & \begin{pmatrix} z_0 z_2 & z_0 z_3 \\ z_1 z_2 & z_1 z_3 \end{pmatrix} = \mathcal{Z}.$$

$$(2)$$

Then \mathcal{Z} is a matrix of determinant $x_0^2 + x_1^2 + x_2^2 + x_3^2 = 0$ which is the equation of q. Now denote by $\mathcal{O}_{\mathbb{P}_3}(n)$ the sheaf of regular functions of degree n on \mathbb{P}_3 and by $\mathcal{O}_q(n,n)$ the sheaf of regular function of be-degree (n,n) on the quadric q. We obtain a surjective map between the global sections

$$\phi: H^0(\mathcal{O}_{\mathbb{P}_3}(n)) \longrightarrow H^0(\mathcal{O}_q(n,n))$$
(3)

by doing the substitution

$$x_0 = \frac{z_0 z_2 + z_1 z_3}{2}, \quad x_1 = \frac{z_0 z_2 - z_1 z_3}{2i}, x_2 = \frac{z_0 z_3 - z_1 z_2}{2}, \quad x_3 = \frac{z_0 z_3 + z_1 z_2}{2i}$$

in a polynomial $p(x_0, x_1, x_2, x_3) \in H^0(\mathcal{O}_{\mathbb{P}_3}(n))$. Observe that $\phi(q) = 0$. Now let

$$\begin{aligned} t &= z_0 z_1 (z_0^4 - z_1^4), \\ W &= z_0^8 + 14 z_0^4 z_1^4 + z_1^8, \\ \chi &= z_0^{12} - 33 (z_0^8 z_1^4 + z_0^4 z_1^8) + z_1^8 \end{aligned}$$

denote the \tilde{T} -invariant polynomials of degree 6,8 and 12 and let

$$\begin{array}{rcl} f &=& z_0 z_1 (z_0^{10} + 11 z_0^5 z_1^5 - z_1^{10}), \\ H &=& -(z_0^{20} + z_1^{20}) + 228 (z_0^{15} z_1^5 - z_0^5 z_1^{15}) - 494 z_0^{10} z_1^{10}, \\ \mathcal{T} &=& (z_0^{30} + z_1^{30}) + 522 (z_0^{25} z_1^5 - z_0^5 z_1^{25}) - 10005 (z_0^{20} z_1^{10} + z_0^{10} z_1^{20}) \end{array}$$

be the I-invariant polynomials of degree 12, 20, 30 given by Klein in [K] p. 51-58. Put $t_1 = t(z_0, z_1), t_2 = t(z_2, z_3), W_1 = W(z_0, z_1), W_2 = W(z_2, z_3)$ and analogously for the other invariants.

Proposition 3.1 If $p \in H^0(\mathcal{O}_{\mathbb{P}^3}(n))$ is G_6 -invariant, then:

$$\phi(p) = \sum_{I} \alpha_{I} t_{1}^{\alpha_{1}} t_{2}^{\alpha'_{1}} W_{1}^{\alpha_{2}} W_{2}^{\alpha'_{2}} \chi_{1}^{\alpha_{3}} \chi_{2}^{\alpha'_{3}}$$

If p is G_{12} -invariant, then:

$$\phi(p) = \sum_{J} \beta_{J} f_{1}^{\beta_{1}} f_{2}^{\beta_{1}'} H_{1}^{\beta_{2}} H_{2}^{\beta_{2}'} \mathcal{T}_{1}^{\beta_{3}} \mathcal{T}_{2}^{\beta_{3}'}$$

where

 $I = \{(\alpha_1, \alpha'_1, \alpha_2, \alpha'_2, \alpha_3, \alpha'_3) | \alpha_i, \alpha'_i \in \mathbb{N}, 6\alpha_1 + 8\alpha_2 + 12\alpha_3 = n, \ 6\alpha'_1 + 8\alpha'_2 + 12\alpha'_3 = n\}, \\ J = \{(\beta_1, \beta'_1, \beta_2, \beta'_2, \beta_3, \beta'_3) | \beta_i, \beta'_i \in \mathbb{N}, 12\beta_1 + 20\beta_2 + 30\beta_3 = n, \ 12\beta'_1 + 20\beta'_2 + 30\beta'_3 = n\}.$

Proof. Put

$$\phi(p) = p'(z_0, z_1, z_2, z_3).$$

An element $g = (g_1, g_2)$ in G_6 or G_{12} operates on $(x_0 : x_1 : x_2 : x_3) \in \mathbb{P}_3$ by the matrix multiplication

$$g_1 \left(\begin{array}{cc} x_0 + ix_1 & x_2 + ix_3 \\ -x_2 + ix_3 & x_0 - ix_1 \end{array} \right) g_2^{-1}$$

and on the matrix \mathcal{Z} of (2) by

$$g_1 \left(\begin{array}{ccc} z_0 z_2 & z_0 z_3 \\ z_1 z_2 & z_1 z_3 \end{array}\right) g_2^{-1} = g_1 \left(\begin{array}{ccc} z_0 \\ z_1 \end{array}\right) \cdot \left(\begin{array}{ccc} z_2 & z_3 \end{array}\right) g_2^{-1}.$$

Clearly if p is G_{6^-} or G_{12} -invariant then also the projection $\phi(p)$ with the previous operation is. In particular for $g = (g_1, 1)$ in $\tilde{T} \times 1$, resp. in $\tilde{I} \times 1$ the polynomial p' is $\tilde{T} \times 1$ -, respectively $\tilde{I} \times 1$ -invariant as polynomial in the coordinates $(z_0 : z_1) \in \mathbb{P}_1$ and for any $(z_2 : z_3) \in \mathbb{P}_1$. On the other hand for $g = (1, g_2)$ in $1 \times \tilde{T}$, resp. in $1 \times \tilde{I}$ the polynomial p' is $1 \times \tilde{T}$ -, respectively $1 \times \tilde{I}$ -invariant as polynomial in the coordinate $(z_2 : z_3) \in \mathbb{P}_1$ and for any $(z_0 : z_1) \in \mathbb{P}_1$. Hence p' must be in the form of the statement.

By a direct computation in section 4 we prove the following

Proposition 3.2 The quadric q does not divide the polynomials F_i , Γ_j . Moreover, F_6 does not divide F_{12} .

Corollary 3.1 We have $\phi(q) = 0$, $\phi(F_6) = t_1 \cdot t_2$, $\phi(F_8) = W_1 \cdot W_2$, $\phi(F_{12}) = \chi_1 \cdot \chi_2$, $\phi(\Gamma_{12}) = f_1 \cdot f_2$, $\phi(\Gamma_{20}) = H_1 \cdot H_2$, $\phi(\Gamma_{30}) = T_1 \cdot T_2$ (up to some scalar factor).

Proof. This follows from Proposition 3.1 and 3.2

Proposition 3.3 The polynomials q, F_6, F_8, F_{12} , resp. $q, \Gamma_{12}, \Gamma_{20}, \Gamma_{30}$ are algebraically independent.

Proof. Let $\sum_{I} \alpha_{I} q^{i_{1}} F_{6}^{i_{2}} F_{8}^{i_{3}} F_{12}^{i_{4}} = 0$ and $\sum_{J} \beta_{J} q^{j_{1}} \Gamma_{12}^{j_{2}} \Gamma_{20}^{j_{3}} \Gamma_{30}^{j_{4}} = 0$ be algebraic relations, $I = (i_{1}, i_{2}, i_{3}, i_{4}) \in \mathbb{N}^{4}$, $J = (j_{1}, j_{2}, j_{3}, j_{4}) \in \mathbb{N}^{4}$, $\alpha_{I}, \beta_{J} \in \mathbb{R}$, then

$$0 = \phi(\sum_{I} \alpha_{I} q^{i_{1}} F_{6}^{i_{2}} F_{8}^{i_{3}} F_{12}^{i_{4}}) = \sum_{I'} \alpha_{I'} \phi(F_{6})^{i'_{2}} \phi(F_{8})^{i'_{3}} \phi(F_{12})^{i'_{4}} = \sum_{I'} \alpha_{I'} t_{1}^{i'_{2}} t_{2}^{i'_{2}} W_{1}^{i'_{3}} W_{2}^{i'_{3}} \chi_{1}^{i'_{4}} \chi_{2}^{i'_{4}}$$

$$(4)$$

similarly

$$\begin{array}{rcl}
0 &=& \phi(\sum_{J} \beta_{J} q^{j_{1}} \Gamma_{12}^{j_{2}} \Gamma_{20}^{j_{3}} \Gamma_{30}^{j_{4}}) \\
&=& \sum_{J'} \beta_{J'} \phi(\Gamma_{12})^{j_{2}} \phi(\Gamma_{20})^{j_{3}'} \phi(\Gamma_{30})^{j_{4}'} \\
&=& \sum_{J'} \beta_{J'} f_{1}^{j_{2}'} f_{2}^{j_{2}'} H_{1}^{j_{3}'} H_{2}^{j_{3}'} \mathcal{T}_{1}^{j_{4}'} \mathcal{T}_{2}^{j_{4}'}.
\end{array} \tag{5}$$

If the polynomials t_1, W_1, χ_1 are fixed, we obtain a relation between t_2, W_2 and χ_2 , which is the same relation as for t_1, W_1 and χ_1 if we fix t_2, W_2 and χ_2 . The same holds for the polynomials f_1, H_1, \mathcal{T}_1 and f_2, H_2, \mathcal{T}_2 . From [K] p. 55 and p. 57 there are only the relations

$$108 t_1^4 - W_1^3 + \chi_1^2 = 0, \quad 108 t_2^4 - W_2^3 + \chi_2^2 = 0$$

and

$$\mathcal{T}_1^2 + H_1^3 - 1728f_1^5 = 0, \quad \mathcal{T}_2^2 + H_2^3 - 1728f_2^5 = 0$$

between these polynomials. By multiplying these relations, however, it is not possible to obtain expressions like (4) and (5). $\hfill \Box$

Corollary 3.2 The polynomials q, F_6, F_8, F_{12} , resp. $q, \Gamma_{12}, \Gamma_{20}, \Gamma_{30}$ generate rationally the ring of invariant polynomials of [3, 4, 3], resp. [3, 3, 5].

Proof. (cf. [Co1] p. 775) By Proposition 3.3 and Proposition 3.2 these are algebraically independent, moreover the products of their degrees are

$$2 \cdot 6 \cdot 8 \cdot 12 = 1152$$
 and $2 \cdot 12 \cdot 20 \cdot 30 = 14400$,

which are equal to the order of the groups [3, 4, 3] and [3, 3, 5]. By [Co1] this implies the assertion.

4 Explicit computations

We recall the following matrices of SO(4) (cf. [Sa]).

$$(q_2,1) = \begin{pmatrix} 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \end{pmatrix}, \quad (1,q_2) = \begin{pmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ -1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \end{pmatrix},$$

$$(p_3,1) = \frac{1}{2} \begin{pmatrix} 1 & -1 & 1 & -1 \\ 1 & 1 & -1 & -1 \\ -1 & 1 & 1 & -1 \\ 1 & 1 & 1 & 1 \end{pmatrix}, \quad (1,p_3) = \frac{1}{2} \begin{pmatrix} 1 & 1 & -1 & 1 \\ -1 & 1 & -1 & -1 \\ 1 & 1 & 1 & -1 \\ -1 & 1 & 1 & 1 \end{pmatrix},$$

$$(p_4,1) = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & -1 & 0 & 0 \\ 1 & 1 & 0 & 0 \\ 0 & 0 & 1 & -1 \\ 0 & 0 & 1 & 1 \end{pmatrix}, \quad (1,p_4) = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 1 & 0 & 0 \\ -1 & 1 & 0 & 0 \\ 0 & 0 & 1 & -1 \\ 0 & 0 & 1 & 1 \end{pmatrix},$$

$$(p_5,1) = \frac{1}{2} \begin{pmatrix} \tau & 0 & 1-\tau & -1 \\ 0 & \tau & -1 & \tau & -1 \\ \tau - 1 & 1 & \tau & 0 \\ 1 & 1-\tau & 0 & \tau \end{pmatrix},$$
$$(1,p_5) = \frac{1}{2} \begin{pmatrix} \tau & 0 & \tau -1 & 1 \\ 0 & \tau & -1 & \tau & 1 \\ 1-\tau & 1 & \tau & 0 \\ -1 & 1-\tau & 0 & \tau \end{pmatrix},$$

where $\tau = \frac{1}{2}(1 + \sqrt{5})$. Then we have

Group	Generators		
G_6	$(q_2, 1), (1, q_2), (p_3, 1), (1, p_3)$		
G_8	$(q_2, 1), (1, q_2), (p_3, 1), (1, p_3), (p_4, 1), (1, p_4)$		
G_{12}	$(q_2, 1), (1, q_2), (p_3, 1), (1, p_3), (p_4, 1), (1, p_4) (q_2, 1), (1, q_2), (p_3, 1), (1, p_3), (p_5, 1), (1, p_5)$		

Now we can write down the equations of the fix lines on $\mathbb{P}_1 \times \mathbb{P}_1$ and those of the planes which are generated by a couple of lines. The products of planes

of section 2.1 in the case of the group G_6 are

$$\begin{aligned} f_{11}^{(6)} \cdot f_{22}^{(6)} \cdot \ldots \cdot f_{66}^{(6)} &= (x_2 - ix_3)(x_1 + ix_3)(x_2 + ix_3)(x_1 - ix_2)(x_1 - ix_3)(x_1 + ix_2), \\ f_{11}^{(8)} \cdot f_{22}^{(8)} \cdot \ldots \cdot f_{88}^{(8)} &= (x_1 + ax_2 - bx_3)(x_1 + bx_2 - ax_3)(x_1 - ax_2 - bx_3)(x_1 - ax_3 - bx_2) \\ &\quad (x_2 + bx_1 - ax_3)(x_2 + ax_1 - bx_3)(x_2 - bx_1 + ax_3)(x_2 + bx_3 - ax_1), \\ f_{11}^{(12)} \cdot f_{22}^{(12)} \cdot \ldots \cdot f_{1212}^{(12)} &= (x_3 - x_1 + cx_2)(x_3 - x_1 - cx_2)(x_2 + x_3 - cx_1)(x_2 + x_3 + cx_1) \\ &\quad (x_3 - x_2 + cx_1)(x_3 - x_2 - cx_1)(x_1 + x_2 + cx_3)(x_1 + x_2 - cx_3) \\ &\quad (x_1 + x_3 - cx_2)(x_1 + x_3 + cx_2)(x_1 - x_2 + cx_3)(x_1 - x_2 - cx_3), \end{aligned}$$

with $a = (1/2)(1 + i\sqrt{3}), b = (1/2)(1 - i\sqrt{3}), c = i\sqrt{2}$. Then the G_6 -invariant polynomials F_6 , F_8 and F_{12} have the following expressions

$$\begin{split} F_6 &= x_0^6 + x_1^6 + x_2^6 + x_3^6 + 5x_0^2 x_1^2 (x_0^2 + x_1^2) + 5x_1^2 x_3^2 (x_1^2 + x_3^2) + 5x_1^2 x_2^2 (x_1^2 + x_2^2) \\ &\quad + 6x_0^2 x_2^2 (x_0^2 + x_2^2) + 6x_0^2 x_3^2 (x_0^2 + x_3^2) + 6x_3^2 x_2^2 (x_2^2 + x_3^2) + 2x_0^2 x_2^2 x_3^2, \end{split}$$

$$F_8 &= 3 \sum x_i^8 + 12 \sum x_i^6 x_j^2 + 30 \sum x_i^4 x_j^4 + 24 \sum x_i^4 x_j^2 x_k^2 + 144 x_0^2 x_1^2 x_2^2 x_3^2, \end{aligned}$$

$$F_{12} &= \frac{123}{8} \sum x_i^{12} + \frac{231}{4} \sum x_i^{10} x_j^2 + \frac{21}{8} \sum x_i^8 x_j^4 - \sum \frac{255}{2} \sum x_i^6 x_j^6 + \frac{949}{2} \sum x_i^8 x_j^2 x_k^2 \\ &\quad + \frac{1839}{2} \sum x_i^6 x_j^4 x_k^2 + \frac{6111}{4} \sum x_i^4 x_j^4 x_k^4 + 1809 \sum x_i^6 x_j^2 x_k^2 x_h^2 + \frac{7281}{2} \sum x_i^4 x_j^4 x_k^4 x_k^2 x_h^2. \end{split}$$

Here the sums run over all the indices i, j, k, h = 0, 1, 2, 3, always being different when appearing together. By applying the map ϕ , a computer computation with MAPLE shows that

$$\phi(F_6) = -\frac{13}{16}t_1 \cdot t_2,$$

$$\phi(F_8) = \frac{3}{64}W_1 \cdot W_2,$$

$$\phi(F_{12}) = \frac{3}{256}\chi_1 \cdot \chi_2$$

as claimed in Corollary 3.1.

Proof of Proposition 2.1, 2. The polynomials F_6 , F_8 , F_{12} remain invariant by interchanging x_2 with x_3 , which is what the matrix C' does.

Proof of Proposition 3.2. We write the computations just in the case of the [3, 4, 3]-invariant polynomials. Consider the points $p_1 = (i\sqrt{2}: 1: 1: 0)$ and $p_2 = (1: i: 0: 0)$, then $q(p_1) = q(p_2) = 0$ and by a computer computation with MAPLE we get $F_6(p_1) = 26$, $F_8(p_2) = 12$ and $F_{12}(p_2) = 32$. This shows that q does not divide the polynomials. Since $F_6(p_2) = 0$, F_6 does not divide F_{12} .

Remark 4.1 Observe that an equation for a [3, 4, 3]-invariant polynomial of degree six and for a [3, 3, 5]-invariant polynomial of degree twelve was given by the author in [Sa] by a direct computer computation with MAPLE.

References

- [Bu] Burnside, W.: Theory of groups of finite order, Dover Publications, Inc. (1955).
- [Co1] Coxeter, H. S. M.: The product of the generators of a finite group generated by reflections, Duke Math. J. Vol. 18 (1951) 765-782.
- [Co2] Coxeter, H. S. M.: Regular polytopes (second edition), The Macmillan company, New York (1963).
- [K] Klein, F.: Vorlesungen über das Ikosaeder und die Auflösung der Gleichungen vom fünften Grade, Nachdr. der Ausg. Leipzig, Teubner 1884, hrsg. mit einer Einführung und mit Kommentaren von Peter Slodowy, Birkhäuser-B. G. Teubner (1993).
- [Ra] Racah, G.: Sulla caratterizzazione delle rappresentazioni irriducibili dei gruppi semisemplici di Lie, Rend. Acad. Naz. dei Lincei, Classe di Scienze fisiche, matematiche e naturali (8), vol. 8 (1950) 108-112.
- [Sa] Sarti, A.: Pencils of symmetric surfaces in $\mathbb{P}^3(\mathbb{C})$, J. of Alg. 246, 429–452 (2001).