

B U L L E T I N

DE LA SOCIÉTÉ DES SCIENCES ET DES LETTRES DE LÓDŹ

2003

Vol. LIII

Recherches sur les déformations

Vol. XLII

pp. 39–41

Andrzej K. Kwaśniewski

INFORMATION ON COMBINATORIAL INTERPRETATION OF FIBONOMIAL COEFFICIENTS [LETTER TO THE EDITOR]

Summary

A combinatorial interpretation of the Fibonomial coefficients is given. (Cf.: binomial Newton and Gaußcoefficients, q -binomial coefficients – ref. [1] and refs. given therein.)

The Fibonacci sequence origin is attributed and referred to the first edition (lost) of “Liber abaci” (1202) by Leonardo Fibonacci [Pisano] (see second edition from 1228 reproduced as Il Liber Abaci di Leonardo Pisano pubblicato secondo la lezione Codice Maglibeciano by Baldassarre Boncompagni in Scritti di Leonardo Pisano vol. 1, (1857) Rome).

Very recently [1, 2] Fibonomial coefficients [–5] have been given a combinatorial interpretation as counting the number of specific finite “birth-selfsimilar” subposets of an infinite non-tree poset naturally related to the Fibonacci tree of rabbits growth process. (More: in [2] an explicit expression for ζ characteristic function in terms of Kronecker δ of the corresponding incidence algebra was provided- see below for further explanatory links).

In order to supply this combinatorial interpretation we use the following notation: *Fibonomial coefficients* [3–5] are defined as

$$\binom{n}{k}_F = \frac{F_n!}{F_k!F_{n-k}!} \equiv \frac{n_F^k}{k_F!}, \quad n_F \equiv F_n \neq 0,$$

where we make an analogy driven [6, 7] identifications ($n > 0$):

$$n_F! \equiv n_F(n-1)_F(n-2)_F(n-3)_F \dots 2_F 1_F;$$

$$0_F! = 1; \quad n_F^k = n_F(n-1)_F \dots (n-k+1)_F.$$

This is the specification of the notation from [6] for the purpose Fibonomial Calculus case (see Example 2.1 in this Bulletin paper [7]).

Define now a partially ordered infinite set P via its finite part – subposet P_m (*rooted at F_1 level subposet*) to be continued ad infinitum in an obvious way as seen from the figure of P_5 below. It looks like the Fibonacci tree with a specific “cobweb”.




Fig. 1. Combinatorial interpretation of Fibonomial coefficients.

If one defines this poset P with help of its incidence matrix ζ representing P uniquely then one arrives at ζ with easily recognizable staircase-like structure – of zeros in the upper part of this upper triangle matrix ζ ; see [1,2].

Recall: incidence matrix ζ is defined for any poset as follows ($x, y \in P$):

$$\zeta(x, y) = \begin{cases} 1 & \text{for } x \leq y, \\ 0 & \text{otherwise.} \end{cases}$$

Observation 1

The number of maximal chains starting from the root (level F_1) to reach any point at the n -th level labeled by F_n is equal to $n_F!$.

Observation 2 ($k > 0$)

The number of maximal chains starting from any fixed point at the level labeled by F_k to reach any point at the n -th level labeled by F_n is equal to n_F^m ($n = k + m$).

Observation 3 ($k > 0$)

Let $n = k + m$. The number of subposets P_m rooted at any fixed point at the level

labeled by F_k and ending at the n -th level labeled by F_n is equal to

$$\binom{n}{k}_F \equiv \frac{n_F^k}{k_F!}.$$

References

- [1] A. K. Kwaśniewski, *Combinatorial interpretation of Fibonomial coefficients*, Inst. Comp. Sci. UwB/Preprint no. 52, November 2003.
- [2] A. K. Kwaśniewski, *More on combinatorial interpretation of Fibonomial coefficients*, Inst. Comp. Sci. UwB/Preprints no. 56, November 2003.
- [3] E. Lucas, *Théorie des fonctions numériques simplement périodiques*, American Journal of Mathematics **1** (1878), 184–240; (Translated from the French by Sidney Kravitz), Ed. D. Lind, Fibonacci Association, 1969.
- [4] G. Fontené, *Généralisation d'une formule connue*, Nouvelles Annales de Mathématiques (4) **15** (1915), 112.
- [5] H. W. Gould, *The bracket function and Fontené-Ward generalized binomial coefficients with applications to Fibonomial coefficients*, The Fibonacci Quarterly **7** (1969), 23–40.
- [6] A. K. Kwaśniewski, *Towards ψ -extension of finite operator calculus of Rota*, Rep. Math. Phys. **47** no. 4 (2001), 305–342. ArXiv: math.CO/0402078 2004
- [7] A. K. Kwaśniewski, *On simple characterizations of Sheffer Ψ -polynomials and related propositions of the calculus of sequences*, Bull. Soc. Sci. Lettres Łódź **52**, Sér. Rech. Déform. **36** (2002), 45–65. ArXiv: math.CO/0312397 2003.
- [8] A. K. Kwaśniewski, *On duality triads*, ibid. **53** Sér. Rech. Déform. **42** (2003), 11–25. ArXiv: math.GM/0402260 v 1 Feb. 2004
- [9] A. K. Kwaśniewski, *On Fibonomial and other triangles versus duality triads*, ibid. **53** Sér. Rech. Déform. **42** (2003), 27–37. ArXiv: math.GM/0402288 v 1 Feb. 2004

High School of Mathematics and Applied Informatics
Kamienna 17, PL-15-021 Białystok
Poland

Presented by Julian Lawrynowicz at the Session of the Mathematical-Physical Commission of the Łódź Society of Sciences and Arts on December 17, 2003

O INTERPRETACJI KOMBINATORYCZNEJ WSPÓŁCZYNNIKÓW DWUMIĘNNYCH [LIST DO REDAKCJI]

S t r e s z c z e n i e

Przedstawiono poszukiwaną od dziesięcioleci kombinatoryczną interpretację współczynników fibonomialnych. (Porównaj: współczynniki binomialne Newtona czy Gaussa; współczynniki q -binomialne; patrz [1] i referencje tamże.)